

HIPK3 Promotes Growth and Metastasis of Esophageal Squamous Cell Carcinoma via Regulation of miR-599/c-MYC Axis

This article was published in the following Dove Press journal:
OncoTargets and Therapy

Yufeng Ba¹
Yining Liu²
Changsheng Li³
Yu Zhu⁴
Wenqun Xing¹

¹Department of Thoracic Surgery, Affiliated Cancer Hospital of Zhengzhou University, Zhengzhou City, Henan Province 450008, People's Republic of China; ²Department of Medical Records, Affiliated Cancer Hospital of Zhengzhou University, Zhengzhou City, Henan Province 450008, People's Republic of China; ³Department of Anesthesiology, Affiliated Cancer Hospital of Zhengzhou University, Zhengzhou City, Henan Province 450008, People's Republic of China; ⁴Department of Orthopedics, The Second Affiliated Hospital of Zhengzhou University, Zhengzhou City, Henan Province 450052, People's Republic of China

Background/Aims: this experimental design was based on HIPK3 to explore the pathogenesis of ESCC.

Methods: RT-qPCR was used to detect the expression of CircHIPK3 and miR-599 in ESCC tissues and cell lines. CCK-8, colony formation, flow cytometry and transwell assay were used to detect the effects of CircHIPK3 and miR-599 on tumor cell proliferation, apoptosis and migration and invasion. Target gene prediction and screening, luciferase reporter assays were used to validate downstream target genes of CircHIPK3 and miR-599. mRNA and protein expression of c-MYC were detected by RT-qPCR and Western blotting. The tumor changes in mice were detected by in vivo experiments in nude mice.

Results: HIPK3 was highly expressed in ESCC tissues and cell lines. In addition, HIPK3 expression levels were associated with advanced TNM stage, lymph node metastasis and tumor size. Moreover, HIPK3 was significantly promoted cell proliferation and migration of ESCC cells. In addition, HIPK3 was able to inhibit miRNA-599 expression and up-regulate the expression level of c-MYC. Finally, the results of in vivo animal models confirmed that HIPK3 promoted ESCC progression by modulating the miR-599/c-MYC axis.

Conclusion: HIPK3 can regulate the proliferation of esophageal squamous cell carcinoma cells by regulating miR-599/c-MYC axis, thereby inhibiting the occurrence and development of esophageal squamous cell carcinoma.

Keywords: esophageal squamous cell carcinoma, HIPK3, miR-599, c-MYC, proliferation

Introduction

Esophageal cancer (EC) is one of the common malignant tumors in humans.^{16,39} It has a high incidence and a poor prognosis, and its regional differences are obvious.⁴³ The primary histological type of primary EC is esophageal squamous cell carcinoma (ESCC).⁴² Esophageal squamous cell carcinoma is more common in China and other Eastern countries. Esophageal squamous cell carcinoma is more common in men over 50 years of age, and the incidence of men is much higher than that of women.^{4,31} Esophageal cancer mainly occurs in the middle and upper segments of the esophagus. Due to the anatomical location and the nature of the tumor, early esophageal cancer is not easy to be found, which affects the best time for treatment and belongs to refractory tumors. At present, patients with early and mid-stage esophageal cancer are treated with surgery or radiotherapy, supplemented by chemotherapy.^{21,35} However, after neoadjuvant chemotherapy for patients with locally advanced esophageal cancer of stage II B and stage III, the prognosis is significantly

Correspondence: Yufeng Ba
Department of Thoracic Surgery,
Affiliated Cancer Hospital of Zhengzhou
University, No. 127, Dongming Road,
Zhengzhou City, Henan Province 450008,
People's Republic of China
Tel + 86 371-65587242
Email klowpz882568@126.com

improved.^{11,28} Human esophageal cancer is a multi-step progression, and the process of esophageal epithelial malignancy develops from atypical hyperplasia, carcinoma in situ to advanced invasive carcinoma. The rapid advancement of molecular biology has proposed a new carcinogenic theory for the progression of esophageal cancer. The occurrence and development of tumors is a multi-gene, multi-stage, and multi-regulation process.^{17,19}

Circular RNA (circ RNA) is a type of non-coding RNA widely found in mammals and is mainly involved in the regulation of genes in organisms.^{13,15} Most of circRNA is derived from the exon region of the gene, and a small portion is formed by intron splicing. Circ RNA is widely involved in the process of physiological and pathological regulation of humans.^{3,40} In tumor research, the mechanism by which circular RNA acts as a miRNA “cavernous body” to regulate downstream target genes has been widely reported.⁴¹ It has been found that a plurality of circular RNAs themselves contain at least one miRNA binding site.⁸ Therefore, it can act as a “sponge” of RNA to adsorb miRNA, thereby regulating the expression of downstream target genes inhibited by miRNA through the mechanism of competitive endogenous RNAs (ce RNAs).³⁸ The circular RNA HIPK3 (circ HIPK3) is highly expressed in the liver, brain, and lung. It mainly originates from the second exon of the gene HIPK3. The circular RNA of HIPK3 origin has three kinds of splicing bodies, circ HIPK3 and circ HIPK3.1 and circ HIPK3.2.¹⁸ While only circ HIPK3 is highly abundant and has significant functions in cells.³⁷ However, the expression and regulation mechanism of circular RNA HIPK3 in ESCC has not been clarified.

miRNAs are endogenous non-coding single-stranded small RNA molecules 19 to 24 nucleotides in length in many organisms.³² miRNAs are involved in almost all tumor-related processes, including proliferation, differentiation, apoptosis, metastasis, and angiogenesis and so on.^{1,29} Many studies have shown that miRNA aberrant expression is closely related to the occurrence, development and prognosis of malignant tumors by regulating the expression of tumor suppressor genes and oncogenes.³⁰ miR-599 is a class of miRNA that promotes tumor cell proliferation. The oncogene c-MYC encodes a conserved protein molecule with a helix-loop-helix and leucine zipper structure, which is abnormally expressed in various cancer tissues and is closely related to cancer cell proliferation, tumor angiogenesis and cancer cell metastasis.⁶ As a transcriptional regulator, c-MYC can form heterodimers with other co-regulatory factors such as Max and regulate the expression of hundreds of genes.¹⁰ The

study found that c-MYC regulation of non-coding miRNAs was an important component of the c-MYC regulatory network, which can alter the expression profile of miRNAs and promote cancer.^{2,20} However, there are few studies in the ESCC. In view of the above problems, this study aimed to explore the effect of HIPK3 on the proliferation and migration of ESCC by detecting the expression of HIPK3 in LSCC tissues and cells, and further reveal the mechanism by which HIPK3 regulated cell proliferation and migration. Defining the pathogenesis of esophageal squamous cell carcinoma may provide a more accurate and effective new therapeutic target for ESCC.

Methods and Materials

Patient Samples and Cells

A total of 42 patients with postoperative ESCC and corresponding normal adjacent epithelial tissues (NATs, > 2 cm from tumor tissue) were selected from 2015 to 2017 in Zhengzhou, China. Informed consent was obtained from all patients. The study was approved by the Affiliated Cancer Hospital of Zhengzhou University Institutional Review Committee. The clinical pathological parameters and clinical features of all patients were shown in Table 1.

ESCC cell line (kyse-150, kyse-410, KYSE-510, ECA-109, EC-18 and TE-13) and normal immortalized cell line (NE1) were purchased from the Chinese Academy of Sciences Cell Bank (Shanghai, China). ESCC cell lines were subcultured in Eagle medium containing 10% fetal bovine serum.

RNA Extraction and Real-Time Quantitative Polymerase Chain Reaction

Total RNA was isolated from cells and tissues using TRIzol reagent in strict accordance with the instructions. Complementary DNA (cDNA) was then synthesized from total RNA using SuperScript III. The gene expression level of HIPK3 was determined by LightCycler™ 480 system (Roche Diagnostics), and the small-nuclear RNA U48 was selected as an internal standard control. The qPCR assay was performed in a LightCycler™ 480 system. The internal standard of miR-599 was U48. qRT-PCR analysis was performed using a SYBR Premix Ex Taq II kit (Takara, Otsu, Japan). The analysis was carried out using the $2^{-\Delta\Delta CT}$ method (Table 2).

Cell Transfection

miR-599 mime, miR-599 agomir and HIPK3 expressing vectors were purchased from Genecopia. In vitro cell experiments were transfected using Lipofectamine 2000

Table 1 Correlation Between HIPK3 Expression and Clinicopathological Features in ESCC Patients

Parameters	Group	N	HIPK3 Expression		P value
			High, n (%)	Low, n (%)	
Age (years)	≤60	12	5(41.7)	7(58.3)	0.522
	>60	30	16(53.3)	14(46.7)	
Gender	Female	11	6(54.5)	5(45.5)	0.656
	Male	31	15(48.4)	16(51.6)	
Smoking	Ever and current	23	13(56.5)	10(43.5)	0.331
	Never	19	8(42.1)	11(57.9)	
Drinking	Ever and current	26	14(53.8)	12(46.2)	0.218
	Never	16	7(43.8)	9(56.2)	
Tumor size (cm)	>5	15	11(73.3)	4(26.7)	0.011
	≤5	27	10(37.0)	17(63.0)	
Differentiation	Well or Moderate	28	13(46.4)	15(53.4)	0.087
	Poor	14	8(57.1)	6(42.9)	
TNM stage	A-IIB	26	17(65.4)	9(34.6)	0.021
	IIIA	16	4(25.0)	12(75.0)	
Lymph node metastasis	Positive	13	9	4	0.027
	Negative	29	12	17	

Table 2 The Oligonucleotides

Sense	Antisense
c-MYC 5'-CGAGTTTATTGTGCGAGCGAA -3'	5'-CTGTGGCATCGCCAGGCC-3'
U485'-GAGGAGAACAAGAGAGAGAGAAATG-3	5'-GCTGGGATGGAGACGCATGAGG-3
GAPDH 5'-CAAATGGAGCAGAGAGCCCT-3'	5'-CCCCATGATACCGTGGGTAG-3'

reagent (Invitrogen) and OptiMEM (Invitrogen).HIPK3 and the loader were first constructed to stably express TE-13 cells.Stable cell lines were then infected with miR-599 agomir or a negative control virus, followed by transfection using Lipofectamine 2000 reagent (Invitrogen) and OptiMEM (Invitrogen).

CCK-8 Method

Transfected cells were seeded into 96-well plates at a density of 1×10^3 cells per well. Three wells were replicated in each group and 10 μ L/well of CCK8 was added to each well of cells for the last 2 hrs of incubation. Finally, the OD value of the cell liquid was measured by an enzyme-linked immunosorbent assay.

Colony Formation

The transfected cells (1×10^3 cells/well) were placed in a 6-well plate to adhere to the cell wall within 24 hrs. After

24 hrs of exposure to ADR, the cells were washed and cultured in different media for 8 days.Finally, colonies were counted after fixation for 10 mins with 10% formaldehyde, and stained with 0.1% crystal violet for 10 min.Cell viability was plotted using GraphPad Prism 6 (GraphPadSoftware, Inc., San Diego, CA, USA).

Apoptosis Detection

Cell apoptosis was determined by FITC Annexin V Apoptosis Detection Kit (BD, Franklin Lakes, NJ, USA). After treated with transfection reagents for 48 h, cells were washed twice with cold PBS and then 1×10^5 cells were re-suspended in 100 μ L 1X binding buffer, then stained with 5 μ L FITC Annexin V and 5 μ L PI for 10 min at room temperature (25 °C) in the dark. 400 μ L 1X binding buffer was added to each tube. The apoptosis cells were then detected by flow cytometry using fluorescence-activated cell-sorting (FACS) flow cytometer (BD Biosciences, San Jose, CA, USA).

Cell Migration and Invasion Assay

Cell migration and invasion tests are shown in the literature.⁹

Luciferase Assay

The pmirGLO dual luciferase miRNA target expression vector was used in the 3'-untranslated region (UTR) luciferase assay (Promega, Madison, WI, USA). Cells were seeded at a density of 4×10^5 cells/mL in 24-well plates. After that, cells were co-transfected with hsa-miR-599 mimics and wild-type or mutant target sequences using Lipofectamine 2000 (Invitrogen). Luciferase assay are shown in the literature.²³

RNA Immunoprecipitation Assay (RIP)

RIP assay were performed using EZ-Magna RIP kit (Millipore, Billerica, MA, USA) and Argonaute 2 (Ago2) antibody (Abcam) to explore whether HIPK3 was existed in RNA induced silencing complex (RISC). Briefly, cells were lysed in RIP lysis buffer, followed by the incubation of protein A/G magnetic beads and antibody against IgG (Millipore) or Ago2 (Abcam). Then RNAs in magnetic beads-binding complexes were purified. At last, RT-qPCR assay was employed to measure the enrichment patterns of HIPK3 and miR-599 by IgG antibody or Ago2 antibody.

Western Blot

Western blot of c-MYC are shown in the literature.²⁵ In vivo antitumor activity.

The study was conducted in strict accordance with the reorganization of the National Institutes of Health Laboratory Animal Care and Use Guidelines. The agreement

was approved by the animal experiment ethics of the xxx hospital. BALB/c nude mice (4 weeks old, female) were maintained under pathogen free conditions.

Si-NC, si-HIPK3 or si-HIPK3/miR-599 cells were injected subcutaneously into the right flank of BALB/C nude mice ($n=10$ /group). Tumor volume of nude mice was measured every 7 days after injection. All nude mice were sacrificed 4 weeks later and the tumors were dissected for qPCR or Western blot analysis experiments.

Statistical Method

The monitoring data were analyzed by SPSS19.0 statistical software. The data analysis results were expressed as mean \pm standard deviation (mean \pm SD). The data analysis between the two groups was performed by *t* test. One-way variance analysis (ANOVA) was used for data analysis among multiple groups, and LSD test was used for subsequent analysis. $P < 0.05$ indicated that the difference was statistically significant.

Results

CircHIPK3 Was Up-Regulated in ESCC Tissues and Cell Lines

Firstly, the expression level of HIPK3 in ESCC was analyzed. It was found that HIPK3 was significantly up-regulated in cancer tissues compared with normal tissues (Figure 1A). Furthermore, similar to the tissue results, HIPK3 expression was significantly up-regulated in the ESCC cell line compared with NE1 cells (Figure 1B).

Patients were divided into high expression group ($n = 21$) and low expression group ($n = 21$) using the




Figure 1 HIPK3 was upregulated in ESCC tissues and cell lines. (A) The expression of HIPK3 in 42 pairs of ESCC tissues and in paired normal tissues. (B) The expression of HIPK3 in ESCC cell lines and NE1 cells. ** $p < 0.01$, *** $p < 0.001$.

median level (HIPK3) as the cutoff value. The results were shown in Table 1. The overexpression of HIPK3 was positively correlated with late TNM stage, lymph node metastasis and tumor size, but not with other parameters such as age, gender, alcohol

CircHIPK3 Promoted the Malignant Behaviors of ESCC Cells

As shown in Figure 2A and B, HIPK3 expression was significantly changed after siRNA or LV-HIPK3 transfection, indicating a successful transfection.




Figure 2 CircHIPK3 promoted proliferation and metastasis of ESCC cells. (A) HIPK3 levels in si-HIPK3 or LV-HIPK3 transfected cells. (B) Effect of HIPK3 on cell viability. (C) Effect of HIPK3 on cell proliferation. (D) Effect of HIPK3 on apoptosis. (E, F) Effect of HIPK3 on cell migration and invasion. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

As shown in **Figure 2C**, compared with the control group, the proliferation rate of the HIPK3 overexpression group was significantly enhanced in the ESCC cells, and the cell proliferation rate of the HIPK3 silenced group was significantly decreased. As shown by **Figure 2D**, the number of cell colonies in the HIPK3 overexpression group was significantly increased in ESCC cells, and the number of cell colonies in the HIPK3 silenced group was significantly reduced compared with the control group. As shown in **Figure 2E**, compared with the control group, the apoptosis rate of the HIPK3 overexpression group was significantly decreased in ESCC cells, and the apoptosis rate of the HIPK3 silenced group was significantly increased. As shown by **Figure 2E and F**, the number of cell migration and invasion in the HIPK3 overexpression group was significantly increased in ESCC cells compared with the control group, and the number of cell migration and invasion in the HIPK3 silenced group was significantly reduced. In summary, the above results indicated that circHIPK3 exerted a carcinogenic effect and promoted ESCC cell growth and metastasis.

miR-599 Was the Target Gene of CircHIPK3

In order to determine the underlying mechanism of action of CircHIPK3 in ESCC, it was predicted that miR-599 was

identified as a potential target for CircHIPK3 (**Figure 3A**) by bioinformatics. The WT-HIPK3 or mutant (Mut)-HIPK3 luciferase reporter plasmid for luciferase reporter assay in TE-13 cells was used to verify the predicted results. As shown in **Figure 3B**, ectopic expression of miR-599 significantly inhibited luciferase activity of WT-HIPK3, but had no significant effect on luciferase activity of Mut-HIPK3. To further test whether HIPK3 and miR-599 are associated through miRNA ribonucleoprotein complexes, Ago2 antibody was used to do the RNA immunoprecipitation experiment. The RT-PCR results showed that HIPK3 and miR-599 were preferentially enriched in the Ago2-containing miRNAs compared with the control IgG immune precipitates (**Figure 3C and D**). In addition, as shown in **Figure 3E**, HIPK3 knockdown can significantly increase the expression level of miR-599 in TE-13 cells. HIPK3 knockdown can significantly up-regulate the expression level of miR-599 in TE-13 cells, while HIPK3 overexpression can significantly down-regulate the expression level of miR-599 in TE-13 cells.

Next the expression pattern of miR-599 in ESCC was analyzed. As shown in **Figure 3F**, the expression level of miR-599 was significantly down-regulated in ESCC tissues compared with that in the normal tissues. In addition, it was found that there was a direct negative correlation between circHIPK3 expression and miR-599 levels in ESCC tissues




Figure 3 HIPK3 directly targeted miR-599 in ESCC. **(A)** Bioinformatics analysis revealed predicted binding sites between HIPK3 and miR-599. **(B)** Relative luciferase activity in TE-13 cells co-transfected with wild-type (HIPK3-WT) or mutant reporter plasmid (HIPK3-mut) and miR-599 mimic or negative control. **(C)** Cellular lysates from ESCC cells were used for RIP with an Ago2 antibody. The Ago2 protein level was detected by Western blot, **(D)** and the relative expression of HIPK3 and miR-599 in the immunoprecipitate was measured by RT-PCR. **(E)** The miR-599 expression in ESCC tissues and matched normal tissues. **(F)** Expression of miR-599 in the presence of si-HIPK3 or LV-HIPK3. **(G)** The relationship between HIPK3 and miR-599 in the ESCC organization. (** P < 0.01).

(Figure 3G). In summary, these results indicated that circHIPK3 may exert its biological function through miR-599.

CircHIPK3 Promoted the Malignant Behaviors of ESCC Cells by Inhibiting miR-599

Next, whether HIPK3 exerts its biological effects by inhibiting miR-599 was explored. As shown in Figure 4A, the cell proliferation rate of miR-599 mimetic + siHIPK3 group was significantly lower than that in HIPK3 silencing group. The cell proliferation rate of miR-599 mimetic + Lv HIPK3 group was significantly higher than that in HIPK3 overexpression group. As shown in Figure 4B, the number of cell colonies in the miR-599 mimetic + siHIPK3 group was significantly lower than that in the HIPK3 silencing group, while the number of cell colonies in the miR-599 mimetic + Lv HIPK3 group was significantly higher than that in the HIPK3 overexpression group. As shown in Figure 4C, the apoptosis rate of miR-599 mimetic + siHIPK3 group was

significantly higher than that of HIPK3 silenced group, and the apoptosis rate of miR-599 mimetic + Lv HIPK3 group was significantly lower compared with that in HIPK3 overexpression group. As shown by Figure 4D and E, the miR-599 mimetic + siHIPK3 group had significantly increased cell migration and invasion compared with that in HIPK3 silencing group. Cell migration and invasion of the miR-599 mimetic + Lv HIPK3 group were significantly reduced compared with that in the HIPK3 overexpressing group. In summary, these results indicated that HIPK3 exerted a biological role in ESCC cells via sponge miR-599.

HIPK3 Increased c-MYC Expression by Inhibiting miR-599

In order to determine the specific mechanism of action of HIPK3 and miR-599 in ESCC progression, it was predicted that c-MYC was identified as a potential target for miR-599 (Figure 5A) by bioinformatics. As shown in Figure 5B, ectopic expression of c-MYC significantly inhibited the




Figure 4 (A) HIPK3 promoted cell growth and metastasis by inhibiting miR-599. (B) Cell viability assay. (C) Cell proliferation assay. (D) Apoptosis assay. (E, F) Cell migration and invasion assays. * $p < 0.05$, ** $p < 0.01$.




Figure 5 HIPK3 increased c-MYC expression levels by inhibiting miR-599. **(A)** Bioinformatics analysis revealed predicted binding sites between c-MYC and miR-599. **(B)** Luciferase activity was detected in cells co-transfected with miR-599 and a luciferase reporter containing c-MYC or mutant forms. **(C)** c-MYC mRNA expression levels in TC-13 cells. **(D)** c-MYC protein expression levels in TC-13 cells. **(E, F)** Changes in expression of c-MYC protein in TE-13 cells. **(G)** A positive correlation between HIPK3 and c-MYC expression in ESCC tissues. **(H)** Negative correlation between c-MYC and miR-599 in ESCC organization. (#P <0.05, **P <0.01).

luciferase activity of WT-miR-559, but had no significant effect on the luciferase activity of mut-miR-559. Furthermore, in TE-13 cells, miR-559 mimics were able to significantly reduce c-MYC mRNA and protein expression compared with the control group (Figure 5C and D). These data indicated that c-MYC was a direct target of miR-559.

Next, whether HIPK3 promotes the development of ESCC by regulating the miR-559/c-MYC axis was further studied. As shown in Figure 5E and F, HIPK3 knockdown inhibited c-MYC expression levels, while miR-559 mimics further increased c-MYC expression levels. In contrast, overexpression of HIPK3 significantly enhanced c-MYC expression levels, while miR-559 mimics further inhibited HIPK3-induced c-MYC upregulation. As shown in Figure 5G and H, c-MYC expression was positively correlated with HIPK3 expression and negatively correlated with miR-559 expression. These results indicated that HIPK3 can increase c-MYC expression levels by inhibiting miR-559.

HIPK3 Promoted ESCC Progression in vivo by Modulating miR-599/c-MYC Axis

In order to further determine the effect of HIPK3 on ESCC progression in vivo, ESCC cells transfected with si-HIPK3

or si-HIPK3/miR-599 agomir were injected subcutaneously into nude mice. As shown in Figure 6A–C, the tumor size and tumor weight of the si-HIPK3 group were significantly lower compared with that in the control group. In the miR-599 agomir treatment group, the tumor growth of si-HIPK3 transfected cells was further inhibited. Furthermore, as shown in Figure 6D, HIPK3 silencing significantly inhibited c-MYC expression, while miR-599 agomir treatment further reduced c-MYC expression levels in tumors. These results indicated that HIPK3 promoted ESCC progression in vivo by inhibiting the miR-599/c-MYC axis.

Discussion

Esophageal cancer is a common malignant tumor of the digestive system that originates in the esophageal mucosal epithelial tissue.¹⁴ Esophageal adenocarcinoma (esophageal adenocarcinoma) occurring from submucosal or cardiac glands of the esophagus is the main pathological type of esophageal cancer in Western countries. Esophageal squamous cell carcinoma (ESCC), which originates from esophageal squamous cells, has always been the main pathological type of esophageal adenocarcinoma.³⁶ At the molecular biology level, the genetic susceptibility of




Figure 6 HIPK3 promoted ESCC progression in vivo by modulating the miR-599/c-MYC axis. (A–C) Tumor size and tumor weight. (D) Expression changes of c-MYC protein in each group. (#P <0.05, **P <0.01).

esophageal cancer is related to the activation and expression of oncogenes and the loss or inactivation of tumor suppressor genes.^{19,33} At present, the specific pathogenesis of esophageal cancer is still unclear. Molecular biology research on esophageal cancer, searching for early diagnostic markers of esophageal cancer and related factors and mechanisms of invasion and metastasis have become a hot topic in this field.

Circular RNAs (circRNAs) exhibiting different properties than linear RNA.²⁶ The current study found that circular RNA can play a role in miRNA molecular sponge and regulate gene transcription.²² CircRNA plays an important role in diseases.²⁷ Studies have shown that the circular RNA cir-ITCH is lowly expressed in esophageal squamous cell carcinoma. As sponges of mir-7, mR-17 and mir-214, cir-ITCH can increase the expression level of ITCH, promote ubiquitination, and thus inhibit esophageal squamous cell carcinoma.³⁴ Circ HIPK3 is another up-and-coming circular RNA that has received increasing attention. In normal cells, its abundance is comparable to or even higher than its linear RNA.¹⁸ In tumor research, it was found that circ HIPK3 can promote the proliferation of osteosarcoma.³⁷ The results of this study showed that the expression of HIPK3 in cancer tissues and cells was significantly lower than that in normal adjacent normal tissues and normal cells ($P < 0.05$). Moreover, overexpression of HIPK3 was positively correlated with advanced TNM stage and tumor size, but not with other parameters such as age, gender, alcohol consumption, smoking history or differentiation status. HIPK3 silencing inhibited cell viability, promoted apoptosis and reduced cell migration and invasiveness, while HIPK3 overexpression can increase cell viability, inhibit cell apoptosis and increase cell migration and invasion. In summary, HIPK3 can be used as a potential molecular target for the treatment of ESCC to control the development of tumors by inhibiting its expression.

As a target for circ RNA, mi RNA is the most widely studied class of non-coding RNAs.⁵ It can regulate cell proliferation, differentiation and apoptosis by degrading target mRNA or inhibiting its translation, and participate in processes such as individual development, body metabolism, tumorigenesis and development.³² Studies had found that overexpression of miR-599 inhibited the proliferation of breast cancer and lung cancer (!!! INVALID CITATION !!!). We screened miR-599 as a target gene for HIPK3 through a database. It regulated its expression by targeting the 3'UTR of the miR-199a-3p gene. HIPK3 knockdown can up-regulate the expression level of miR-599, and vice versa.

The expression level of miR-599 was down-regulated in ESCC tissues compared with that in normal tissues, and an inverse correlation between circHIPK3 expression and miR-599 levels was observed in ESCC tissues.

c-MYC is a proto-oncogene of MYC family cells, which expresses a nuclear protein and participates in the regulation of other genes by binding DNA.²⁴ The expression of c-MYC protein was higher in proliferating cells, which decreased with cell differentiation and lost the ability of promoting proliferation. When Down regulation is expressed, cells are induced to differentiate into terminal cells.⁷ c-MYC target-genes, which mediate these functions of c-MYC, represent a complex network of protein- and non-coding RNAs, including numerous miRNAs. For example, c-MYC directly regulates expression of the miR-17-92 cluster, miR-34a, miR-15a/16-1 and miR-9. Moreover, the expression and activity of c-MYC itself are under the control of miRNAs.¹² Here, for the first time, we identified MYC as a target of miR-599 and showed that the miR-599-MYC axis regulates cell proliferation and migration in ESCC. circHIPK3 regulates the proliferation and migration of ESCC cells by serving as a miR-599 sponge and thus modulates MYC expression. Further understanding of the circHIPK3- miR-599- MYC axis may provide a novel therapeutic strategy for ESCC in the future.

Conclusion

HIPK3 inhibited ESCC proliferation and invasion and metastasis by regulating miR-599/c-MYC axis, suggesting that HIPK3 may be a potential therapeutic target for ESCC. It provided experimental basis for the study of clinical prognosis of the tumor and further targeted intervention therapy.

Ethical Statement and Consent for Publication

Written informed consent was obtained from all patients. The study was approved by the Affiliated Cancer Hospital of Zhengzhou University Institutional Review Committee.

Disclosure

The authors declare that they have no competing interests in this work.

References

1. Anokyedanso F, Trivedi CM, Juhr D, et al. Highly efficient miRNA-mediated reprogramming of mouse and human somatic cells to pluripotency. *Cell Stem Cell*. 2011;8(4):376–388. doi:10.1016/j.stem.2011.03.001

2. Chang TC, Yu D, Lee YS, et al. Widespread microRNA repression by Myc contributes to tumorigenesis. *Nat Genet.* 2008;40(1):43–50. doi:10.1038/ng.2007.30
3. Chen G, Shi Y, Zhang Y, Sun J. CircRNA_100782 regulates pancreatic carcinoma proliferation through the IL6-STAT3 pathway. *Oncotargets Ther.* 2017;10:5783–5794. doi:10.2147/OTT.S150678
4. Chen Y, Yin D, Li L, Deng Y, Tian W. Screening aberrant methylation profile in esophageal squamous cell carcinoma for Kazakhs in Xinjiang area of China. *Mol Biol Rep.* 2015;42(2):457–464. doi:10.1007/s11033-014-3788-z
5. Cheng AM, Byrom M, Shelton J, Ford LP. Antisense inhibition of human miRNAs and indications for an involvement of miRNA in cell growth and apoptosis. *Nucleic Acids Res.* 2005;33(4):1290–1297. doi:10.1093/nar/gki200
6. Dang CV. MYC on the path to cancer. *Cell.* 2012;149(1):22–35. doi:10.1016/j.cell.2012.03.003
7. Delmore Jake E, Issa Ghayas C, Lemieux Madeleine E, et al. BET bromodomain inhibition as a therapeutic strategy to target c-Myc. *Cell.* 2011;146(6):904–917. doi:10.1016/j.cell.2011.08.017
8. Guo L, Zheng L, Zhao Y, Wang Q. Profiling and bioinformatic analyses indicate differential circRNA and miRNA/isomiR expression and interactions. *Biomed Res Int.* 2018;2018:1–9.
9. Guy JB, Espenel S, Vallard A, et al. Evaluation of the cell invasion and migration process: a comparison of the video microscope-based scratch wound assay and the boyden chamber assay. *J Visual Exp Jove.* 2017;129.
10. Hsieh AL, Walton ZE, Altman BJ, Stine ZE, Dang CV. MYC and metabolism on the path to cancer. *Semin Cell Dev Biol.* 2015;43:11–21. doi:10.1016/j.semedb.2015.08.003
11. Ikeguchi M, Kohno Y, Kihara K, et al. Neoadjuvant chemotherapy for clinical Stage II and III thoracic esophageal squamous cell carcinoma with curative esophagectomy. *J Cancer Ther.* 2015;06(15):1207–1213. doi:10.4236/jct.2015.615131
12. Jackstadt R, Hermeking H. MicroRNAs as regulators and mediators of c-MYC function. *Biochim Biophys Acta.* 2015;1849(5):544–553. doi:10.1016/j.bbagr.2014.04.003
13. Jeck WR, Sharpless NE. Detecting and characterizing circular RNAs. *Nat Biotechnol.* 2014;32(5):453–461. doi:10.1038/nbt.2890
14. Ji Y, Du X, Tian Y, et al. A Phase II study of S-1 with concurrent radiotherapy in elderly patients with esophageal cancer. *Oncotarget.* 2017;8(47):83022–83029. doi:10.18632/oncotarget.v8i47
15. Lasda E, Parker R. Circular RNAs: diversity of form and function. *RNA.* 2014;20(12):1829–1842. doi:10.1261/rna.047126.114
16. Li B, Xu WW, Han L, et al. MicroRNA-377 suppresses initiation and progression of esophageal cancer by inhibiting CD133 and VEGF. *Oncogene.* 2017a;36(28):3986–4000. doi:10.1038/onc.2017.29
17. Li Q, Zhang X, Li N, Liu Q, Chen D. miR-30b inhibits cancer cell growth, migration, and invasion by targeting homeobox A1 in esophageal cancer. *Biochem Biophys Res Commun.* 2017b;485(2):506–512. doi:10.1016/j.bbrc.2017.02.016
18. Li Y, Zheng F, Xiao X, et al. CircHIPK3 sponges miR-558 to suppress heparanase expression in bladder cancer cells. *EMBO Rep.* 2017c;18(9):1646–1659. doi:10.15252/embr.201643581
19. Long L, Pang X, Lei F, et al. SLC52A3 expression is activated by NF- κ B p65/Rel-B and serves as a prognostic biomarker in esophageal cancer. *Cell Mol Life Sci.* 2018;75(14):2643–2661. doi:10.1007/s00018-018-2757-4
20. Ma L, Young JJ, Prabhala H, et al. miR-9, a MYC/MYCN-activated microRNA, regulates E-cadherin and cancer metastasis. *Nat Cell Biol.* 2010;12(3):247–256. doi:10.1038/ncb2024
21. Ma X, Zhao K, Guo W, et al. Salvage lymphadenectomy versus salvage radiotherapy/chemoradiotherapy for recurrence in cervical lymph node after curative resection of esophageal squamous cell carcinoma. *Ann Surg Oncol.* 2015;22(2):624–629. doi:10.1245/s10434-014-4008-8
22. Militello G, Weirick T, John D, Doring C, Dimmeler S, Uchida S. Screening and validation of lncRNAs and circRNAs as miRNA sponges. *Brief Bioinform.* 2016;18(5):780–788.
23. Miyakawa H, Iguchi T. Comparative luciferase assay for establishing reliable in vitro screening system of juvenile hormone agonists. *J Appl Toxicol.* 2017;37(9):1082. doi:10.1002/jat.v37.9
24. O'Donnell KA, Al E. c-Myc-regulated microRNAs modulate E2F1 expression. *Nature.* 2005;435(7043):839–843. doi:10.1038/nature03677
25. O'donnell KA, Wentzel EA, Zeller KI, Dang CV, Mendell JT. c-Myc-regulated microRNAs modulate E2F1 expression. *Nature.* 2005;435(7043):839–843. doi:10.1038/nature03677
26. Pamudurti NR, Bartok O, Jens M, et al. Translation of CircRNAs. *Mol Cell.* 2017;66(1):9–21. doi:10.1016/j.molcel.2017.02.021
27. Patop IL, Kadener S. circRNAs in cancer. *Curr Opin Genet Dev.* 2018;48:121–127. doi:10.1016/j.gde.2017.11.007
28. Shen L, Wang H, Dong B, et al. Possible prediction of the response of esophageal squamous cell carcinoma to neoadjuvant chemotherapy based on gene expression profiling. *Oncotarget.* 2016;7(4):4531–4541. doi:10.18632/oncotarget.6554
29. Shimono Y, Zabala M, Cho RW, et al. Downregulation of miRNA-200c links breast cancer stem cells with normal stem cells. *Cell.* 2009;138(3):592–603. doi:10.1016/j.cell.2009.07.011
30. Smirnova L, Grafe A, Seiler A, Schumacher S, Nitsch R, Wulczyn FG. Regulation of miRNA expression during neural cell specification. *Eur J Neurosci.* 2005;21(6):1469–1477. doi:10.1111/ejn.2005.21.issue-6
31. Teng H, Li X, Liu X, Wu J, Zhang J. The absence of human papillomavirus in esophageal squamous cell carcinoma in East China. *Int J Clin Exp Pathol.* 2014;7(7):4184.
32. Ventura A, Young AG, Winslow MM, et al. Targeted deletion reveals essential and overlapping functions of the miR-17–92 family of miRNA clusters. *Cell.* 2008;132(5):875–886. doi:10.1016/j.cell.2008.02.019
33. Vrana D, Matzenauer M, Aujesky R, et al. Potential predictive role of MicroRNAs in the neoadjuvant treatment of esophageal cancer. *Anticancer Res.* 2017;37(2):403–412. doi:10.21873/anticancer
34. Wan L, Zhang L, Fan K, Cheng Z, Sun Q, Wang J. Circular RNA-ITCH suppresses lung cancer proliferation via inhibiting the Wnt/ β -catenin pathway. *Biomed Res Int.* 2016;2016:1579490. doi:10.1155/2016/1579490
35. Wang Q, Zhang W, Liu X, et al. Prognosis of esophageal squamous cell carcinoma patients with preoperative radiotherapy: comparison of different cancer staging systems. *Thorac Cancer.* 2014;5(3):204–210. doi:10.1111/tca.2014.5.issue-3
36. Wong CH, Ma BBY, Hui CWC, Tao Q, Chan ATC. Preclinical evaluation of afatinib (BIBW2992) in esophageal squamous cell carcinoma (ESCC). *Am J Cancer Res.* 2015;5(12):3588–3599.
37. Xiaolong M, Kumpeng Z, Chunlin Z. Circular RNA circ_HIPK3 is down-regulated and suppresses cell proliferation, migration and invasion in osteosarcoma. *J Cancer.* 2018;9(10):1856–1862. doi:10.7150/jca.24619
38. Xiong D, Dang Y, Lin P, et al. A circRNA–miRNA–mRNA network identification for exploring underlying pathogenesis and therapy strategy of hepatocellular carcinoma. *J Transl Med.* 2018;16(1):220. doi:10.1186/s12967-018-1593-5
39. Xu WW, Li B, Zhao JF, et al. IGF2 induces CD133 expression in esophageal cancer cells to promote cancer stemness. *Cancer Lett.* 2018;425:88–100. doi:10.1016/j.canlet.2018.03.039
40. Zhang H, Jiang L, Sun D, Hou J, Ji Z. CircRNA: a novel type of biomarker for cancer. *Breast Cancer.* 2018;25(1):1–7. doi:10.1007/s12282-017-0793-9
41. Zhang P, Meng X, Chen H, et al. PlantCircNet: a database for plant circRNA–miRNA–mRNA regulatory networks. *Database.* 2017;2017.
42. Zhao Y, Zhou W, Xue L, Zhang W, Zhan Q. Nicotine activates YAP1 through nAChRs mediated signaling in esophageal squamous cell cancer (ESCC). *PLoS One.* 2014;9:3.
43. Ziqiang T. Regional differences of methylation of metallothionein-3 gene in tissues of esophageal cancer. *Tumori.* 2009;1:1137–1139.

OncoTargets and Therapy

Dovepress

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic

agents and protocols on patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/oncotargets-and-therapy-journal>