

Research Article

The Relationship between Plants Used to Sustain Finches (Fringillidae) and Uses for Human Medicine in Southeast Spain

Antonio Belda,¹ Victoriano Peiró,^{2,3} and Eduardo Seva^{2,3}

¹Departamento Ciencias de la Tierra y del Medio Ambiente, Universidad de Alicante, Campus San Vicente, Apartado 99, 03080 Alicante, Spain

²Departamento de Ecología, Universidad de Alicante, Campus San Vicente, Apartado 99, 03080 Alicante, Spain

³IMEM, Universidad de Alicante, Campus San Vicente, Apartado 99, 03080 Alicante, Spain

Correspondence should be addressed to Antonio Belda, antonio.belda@ua.es

Received 29 December 2011; Revised 13 February 2012; Accepted 14 February 2012

Academic Editor: Andrea Pieroni

Copyright © 2012 Antonio Belda et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We analyzed plants that are traditionally used by wild bird hunters and breeders to capture and promote captive breeding of *Fringillidae* (finches or songbirds) in the province of Alicante, Spain. The majority of plants used in songbird breeding have medicinal properties in traditional human medicine (48 different uses); thus, another main goal was to show their relationships with human medical uses. We compiled a list of 97 plant species from 31 botanical families that are used to attract finches and identified 11 different use categories for these plants in finch keeping. The most common uses were for trapping birds and as a source of food for birds in captivity. *Cannabis sativa* has the greatest cultural importance index (CI = 1.158), and *Phalaris canariensis* (annual canary grass or al pist) was the most common species used to attract *Fringillidae* and was used by all informants ($n = 158$). Most of the 97 species are wild plants and mainly belong to the families Compositae, Gramineae, Cruciferae, and Rosaceae and also have medicinal properties for humans. In the study area, the intensification of agriculture and abandonment of traditional management practices have caused the population of many songbirds to decline, as well as the loss of popular ethnographic knowledge.

1. Introduction

Throughout the ages, the human race has used plants for various purposes [1], particularly those that are accessible. In the Iberian Peninsula, several studies have been developed on medicinal plants [2–8] and edible flora [9, 10], as well as some general ethnobotanical studies [11–15], and others about the importance of home gardens and cultivated areas in the evolution of useful flora [16]. However, few studies have described the use of plants in ethnoveterinary medicine [17, 18], or in attracting and maintaining birds of the *Fringillidae* family in captivity [19, 20]. Plants have been used in traditional medicine for several thousands of years to treat and cure diseases in domestic animals and human populations, especially native ones [21, 22]. Furthermore, in nature, wild birds use particular plant species, which possess insecticidal and bactericidal properties, to build their nests.

This practice creates optimal conditions for egg laying and incubation [23].

The ecological knowledge of local traditional uses that depend on the dynamics of natural resources has been reflected in numerous studies [24–26], considering the ecological knowledge of local communities of hunters, anglers, and gatherers [27].

The culture of capturing songbirds was introduced to the Iberian Peninsula by the Romans and had its beginnings, as did other forms of hunting, in the absolute necessity of human nutrition. Thus, these birds were traditionally caught as a source of food in Valencia, at least since the 17th century [28]. Today, following old customs and culinary habits, there are still hunters who hunt this group of birds in order to eat them. On the other hand, the term “*pajareros*” describes people who are dedicated to hunting, breeding, or selling birds [29]. Although these birds are not hunted excessively, it

is essential to monitor and control illegal methods of hunting Fringillidae and to conserve this family of birds [30].

The capture of birds using a hinged net assembly is a traditional hunting method that is widespread in the province of Alicante and elsewhere in the Iberian Peninsula. These game nets are made with cotton, hemp, or nylon mesh. They are placed on the ground and have a manual activation system; once a bird enters the net, a rope is pulled to trap the bird inside the net (see Photographic annex). The nets used since the middle ages to capture several species of Fringillidae intended for use as pets are well known among the inhabitants of this zone [31, 32]. These birds are relatively easy to maintain and rear in captivity, and it is easy to train them to participate in singing competitions. Thus, at present, the capture of five species of birds (*Serinus serinus*, “verdecillo”, *Carduelis carduelis*, “jilguero”, *Carduelis chloris*, “verderón”, *Carduelis cannabina*, “pardillo,” and *Fringilla coelebs*, “pinzón”) is authorized and regulated by law (Council Directive 79/409/EEC and national Laws 4/1989, 62/2006, and 13/2004). What is more, it is an important cultural movement around the Mediterranean Basin [33]. The current trend is to increase breeding in captivity and reduce the capture of wild birds. Therefore, it is important to acquire more knowledge about the traditional use of cultivated and wild plants.

The main aim of this paper is to document the cross-cultural comparison between plant uses for songbirds and humans in Mediterranean environments, relating an ethnoveterinary field study and its eventual link to folk therapies for humans, in order to preserve ethnological knowledge on European folk health. With this purpose in mind, the information on plant uses for songbirds (capturing, feeding, and breeding) gathered here was collected during fieldwork and complemented with ethnobotanical references. Finally, we would like to contribute to the dissemination of results within the scientific community in order to open a door to research in other disciplines.

2. Materials and Methods

2.1. Study Area. The province of Alicante is located in the southeast region of Spain, in the southern part of Valencia. It is geographically located between the coordinates 38°30'N and 0°50'E (Figure 1). The total area occupied by the province is 5,863 km², it has a population of 1,783,555 inhabitants, and there are 141 localities. The province has a very mountainous and rugged relief, except for some river valleys. Thus, approximately 60% of the study area is located between elevations of 200 and 1,500 m above sea level.

Due to its geographical location, the province of Alicante has a typical Mediterranean climate with mild temperatures. Thus, the average temperatures are between 6.2° and 16.8° in the coldest month (January) and 20.4° and 30.6° in the hottest (August), with an annual mean of 17.8°. The average annual rainfall is 336 mm, concentrated in spring and autumn, and there is a prominent dry period in summer. However, there are some climatic differences between the coast and the interior of the province, due to its topography

[34, 35]. The plant species in the province of Alicante include sclerophyllous shrubs and trees, which are adapted to Mediterranean stress conditions. Local flora, consisting of evergreen, coriaceous, glabrous, and aromatic plants, is adapted to conserve water for much of the year. Some qualities are common to many of these plants, including resistance to drought, adaptations to heat, and low tolerance to low temperatures. These bioclimatic and biogeographical conditions favour the development of rare, endemic, and endangered species [35, 36]. Considering its bioclimatic and biogeographical conditions, the province of Alicante may potentially give rise to vegetation that can be divided into three main types: evergreen oak forest (*Rubio longifolia-Quercetum rotundifoliae*), ash-maple forest (*Fraxino ornii-Aceretum granatensis*), and spiny maquis (*Chamaerops humilis—Rhamnus lycioides*) [37].

2.2. Ethnology. A total of 69 localities were prospected with oral interviews in all regions of the Alicante province (*El Comptat, L'Alcoia, Alt Vinalopo, Vinalopo Mitja, Marina Baixa, Marina Alta, L'Alacanti, Baix Vinalopo, and Baix Segura-Vega Baixa*) (Figure 1). Vernacular names of plant species traditionally used were obtained in the field by interviews with the local population. Ethnological information was based primarily on semistructured interviews, in which we gathered information. This ranged from the different plant species used to attract and maintain songbirds, the season of plant collection, traditional uses of the plant species collected, the composition of commercial mixtures used to feed captive birds, and folk remedies used to cure songbird illnesses, to the environmental problems faced by the community.

People with a specific profile were selected in order to obtain high-quality and reliable information. People interviewed were older (50–85 years old), living in a rural environment and from a variety of socioeconomic strata, who had captured and bred birds throughout their lives. We wanted to emphasize the ethnobotanical importance of local variations of plant names and the different applications of these species. We conducted 158 oral interviews; 95.57% ($n = 151$) of the informants were male and 4.43% ($n = 7$) female, and the mean age was 56.7 years. In 48 municipalities, inhabitants speak Valencian (variant of Catalan), and Castilian (standard Spanish) is spoken in the others.

Numerous folk botanical references were examined [38–41], including a variety of local books [35], magazines [7], and festivals, to obtain information on remedies for animal illnesses. Even though the information included in our analysis arose from an array of different spoken and written sources in the study area, the semistructured interviews revealed many important issues previously unidentified [20].

A digital sound recorder was used to record interviews and to create an audio record of the information. In addition, a photographic archive, with photographs of each of the species referred to by the informants, was constructed and deposited in the Ecology Department Archive of Alicante University.

TABLE 1: Plant species in the study area and their traditional uses in finches and humans. Finch uses: 1, facilitate breeding; 2, attract birds; 3, commercial seed mixes; 4, leafy vegetables; 5, birdliming; 6, material for cages and accessories; 7, catching tools; 8, vermifuge; 9, camouflage of capture nets; 10, provide pigments; 11, cure diseases. Medical human uses: 1, alteration of blood pressure; 2, haemorrhoids; 3, depurative; 4, anxiety; 5, diarrhoea; 6, heartburn; 7, indigestion; 8, liver disease; 9, loss of appetite; 10, constipation; 11, helminthiasis; 12, cough; 13, cold; 14, respiratory problems; 15, hyperglycemia; 16, anaemia; 17, hypercholesterolemia; 18, retention of liquids; 19, undefined symptom (tonic); 20, gout; 21, rheumatism; 22, inflammation of bones or joints; 23, undefined symptom (analgesic); 24, injury; 25, burns; 26, kidney stones; 27, menstruation; 28, lack of breast milk secretion; 29, ischocolia; 30, chilblains; 31, pimples; 32, skin diseases; 33, eczema; 34, skin fungus; 35, rubefaction; 36, calluses and skin hardness; 37, warts; 38, bacteria; 39, microbes; 40, headache; 41, inflammation; 42, fever; 43, alopecia; 44, flushing (refreshing); 45, alcoholism; 46, toothache; 47, mineral deficiency; 48, eye infection. Type: W, wild; C, cultivated. RFC: relative frequency of citation. CI: cultural importance index.

Scientific name	Herbarium voucher (ABH)	Family	Finch uses	RFC	CI	Medical human uses	Type	References
<i>Agave americana</i> L.	17879	Agavaceae	1	7.59	0.076	—	W	[39]
<i>Allium sativum</i> L.	Seen alive	Alliaceae	8	5.06	0.051	41, 13, 5, 21, 30	C	[35, 39, 44]
<i>Amaranthus blitum</i> L.	3989	Amaranthaceae	2,3	25.32	0.285	—	W	[41]
<i>Anagallis arvensis</i> L.	22647	Primulaceae	4	16.46	0.165	24, 3, 14	W	[40]
<i>Andryala ragulina</i> L.	4430	Compositae	5	72.15	0.722	31	W	[41]
<i>Arundo donax</i> L.	32085	Gramineae	6,7	58.23	0.810	1, 48, 12, 22, 18, 3	C	[39]
<i>Avena sativa</i> L.	10488	Gramineae	3	84.81	0.848	—	W	
<i>Avena sterilis</i> L.	1582	Gramineae	3	78.48	0.785	—	W	
<i>Beta vulgaris</i> L.	10652	Chenopodiaceae	4	30.38	0.304	37, 10, 18, 44, 1, 41, 6, 2	C	[41]
<i>Bituminaria bituminosa</i> (L.) C. H. Stirt.	50474	Leguminosae	2	32.91	0.329	—	W	
<i>Brachypodium retusum</i> (Pers.) Beauv.	31248	Gramineae	7	67.09	0.671	1	W	[39]
<i>Brassica napus</i> L.	39373	Cruciferae	3	89.87	0.899	—	C	
<i>Brassica oleracea</i> L. subsp. <i>oleracea</i>	34847	Cruciferae	4	21.52	0.215	24, 28, 45, 41, 21, 18, 10, 48, 4, 7	C	[41]
<i>Brassica oleracea</i> L. var. <i>italica</i> Plenck	Seen alive	Cruciferae	1,4	62.03	0.924	—	C	
<i>Brassica rapa</i> L.	7969	Cruciferae	3	88.61	0.886	—	C	
<i>Cannabis sativa</i> L.	32225	Cannabaceae	1,3	92.41	1.158	—	C	
<i>Capsella bursa-pastoris</i> (L.) Medicus	47380	Cruciferae	4	16.46	0.165	5, 18, 1, 27	W	[40]
<i>Carthamus tinctorius</i> L.	3894	Compositae	3	10.13	0.101	—	W	
<i>Centaurea aspera</i> L.	21338	Compositae	2	60.76	0.608	9, 7, 15, 24, 34, 38, 41, 1, 13, 39, 19, 29	W	[7, 35, 40]
<i>Centaurea calcitrapa</i> L.	36097	Compositae	2	46.84	0.468	15	W	[40]
<i>Centaurea mariolensis</i> Rouy	13242	Compositae	2	10.13	0.101	—	W	
<i>Centaurea meliensis</i> L.	36917	Compositae	2	11.39	0.114	—	W	
<i>Chamaerops humilis</i> L.	559	Palmae	1	13.92	0.139	9	W	[39]
<i>Chelidonium majus</i> L.	18328	Papaveraceae	11	3.8	0.038	—	C	
<i>Chondrilla juncea</i> L.	7142	Compositae	5	49.37	0.494	9	W	[39]
<i>Cicer arretinum</i> L.	17633	Leguminosae	11	3.8	0.038	31, 18, 13	W	[41]

TABLE 1: Continued.

Scientific name	Herbarium voucher (ABH)	Family	Finch uses	RFC	CI	Medical human uses	Type	References
<i>Cichorium intybus</i> L.	37547	Compositae	3,4	54.43	0.639	18, 19, 10, 9	W	[35, 39, 44]
<i>Cirsium arvense</i> (L.) Scop.	35007	Compositae	2	8.86	0.089	9, 2	C	[39]
<i>Cirsium monspessulanum</i> (L.) Hill	51477	Compositae	2	10.13	0.101	—	W	
<i>Citrus limon</i> (L.) Burm. Fil.	49856	Rutaceae	11	2.53	0.025	24, 5, 3, 18, 13	C	[39]
<i>Coryza bonariensis</i> (L.) Cronq.	17943	Compositae	2	12.66	0.127	—	C	
<i>Cynara cardunculus</i> L.	35991	Compositae	2	16.46	0.165	37, 27, 29, 32, 9, 3, 8, 18, 15	W	[7, 35, 41, 44]
<i>Cynara scolymus</i> L.	31715	Compositae	2	44.3	0.443	—	C	
<i>Daphne gnidium</i> L.	10830	Thymelaeaceae	6,8	41.77	0.481	37, 13, 8, 11, 35	W	[7, 35, 39]
<i>Daucus carota</i> L.	33104	Umbelliferae	4,10	18.99	0.222	18, 37, 9, 19	W	[35, 39]
<i>Diplotaxis erucoides</i> (L.) DC.	47963	Cruciferae	2,4	91.14	0.911	—	W	[40]
<i>Dittrichia viscosa</i> (L.) Greuter	39371	Compositae	2	68.35	0.684	24, 1, 26, 17, 46	W	[39]
<i>Echinochloa crus-galli</i> (L.) Beauv.	14692	Gramineae	2	6.33	0.063	—	W	
<i>Eruca vesicaria</i> (L.) Cav.	41713	Cruciferae	4	20.25	0.203	—	W	
<i>Erucastrum virgatum</i> C. Presl	4460	Cruciferae	2,4	15.19	0.190	—	W	
<i>Euphorbia characias</i> L.	7226	Euphorbiaceae	5	36.71	0.367	8	C	[39, 44]
<i>Foeniculum vulgare</i> Miller	23129	Umbelliferae	3	24.05	0.241	18, 33, 10, 14, 48, 13, 9, 8, 39, 28, 7	W	[7, 35, 39, 44]
<i>Fragaria vesca</i> L.	52157	Rosaceae	4,10	11.39	0.139	13, 30, 20	W	[44]
<i>Galactites tomentosa</i> Moench	42051	Compositae	2	11.39	0.114	—	C	
<i>Guizotia abyssinica</i> (L.f.) Cass.	9666	Compositae	3	87.34	0.873	—	W	
<i>Helianthus annuus</i> L.	5220	Compositae	3	87.34	0.873	—	W	
<i>Heliotropium europaeum</i> L.	14672	Boraginaceae	2	82.28	0.823	24, 29, 42, 27, 20, 37	W	[35, 41]
<i>Hyparrhenia hirta</i> (L.) Staff	41077	Gramineae	7	32.91	0.329	—	C	[39]
<i>Lactuca sativa</i> L.	Seen alive	Compositae	3,4	58.23	0.728	44, 47, 4, 46	C	[41]
<i>Lactuca serriola</i> L.	47376	Compositae	4	22.78	0.228	10, 4	W	[41]
<i>Laurus nobilis</i> L.	43242	Lauraceae	8	13.92	0.139	14, 13, 9, 7, 6, 31, 8	C	[35, 39, 44]
<i>Lavandula latifolia</i> Medicus	20246	Labiatae	4,11	37.97	0.456	19, 7, 39, 24, 21, 18, 33, 14, 41, 5	W	[7, 35, 39, 44]
<i>Linum usitatissimum</i> L.	32017	Linaceae	3	13.92	0.139	10, 33, 21, 31, 24, 14, 13, 41, 4	W	[35, 39]
<i>Lobularia maritima</i> (L.) Desv.	15843	Cruciferae	4	35.44	0.354	18, 26, 41, 23, 42	W	[41]
<i>Lygeum spartum</i> L.	8128	Gramineae	5	45.57	0.456	—	W	
<i>Malus domestica</i> (Borkh.) Borkh.	37495	Rosaceae	4,11	27.85	0.392	7, 37	W	[40]
<i>Mantisalca salmantica</i> (L.) Briq. and Cavill.	5273	Compositae	2	13.92	0.139	15	C	[41]

TABLE 1: Continued.

Scientific name	Herbarium voucher (ABH)	Family	Finch uses	RFC	CI	Medical human uses	Type	References
<i>Nerium oleander</i> L.	46139	Apocynaceae	6,8	30.38	0.411	32	C	[39]
<i>Nicotiana tabacum</i> L.	4391	Solanaceae	8,11	13.92	0.234	43, 40, 46	W	[39]
<i>Ocimum basilicum</i> L.	Seen alive	Labiatae	8	3.8	0.038	7, 4, 8, 13, 10, 5	C	[35, 39, 44]
<i>Olea europaea</i> L.	17212	Oleaceae	5	5.06	0.051	36, 10, 1, 41, 7, 29, 37, 24, 19, 25, 23, 8	W	[7, 35, 40, 44]
<i>Onopordum acanthium</i> L.	11328	Compositae	2	11.39	0.114	—	W	
<i>Panicum miliaceum</i> L.	36589	Gramineae	3	68.35	0.063	—	W	
<i>Papaver rhoeas</i> L.	37589	Papaveraceae	3,4	26.58	0.291	14, 12, 46, 23, 9, 18, 13, 4	C	[7, 35, 39, 44]
<i>Papaver somniferum</i> L.	10585	Papaveraceae	3,4	17.72	0.203	13, 4, 5, 12, 46	W	[39]
<i>Paronychia argentea</i> Lam.	13044	Caryophyllaceae	1	6.33	0.063	18, 24, 1	C	[35]
<i>Perilla frutescens</i> L.	Seen alive	Labiatae	3	60.76	0.608	—	C	
<i>Phagnalon saxatile</i> (L.) Cass.	49631	Compositae	2	10.13	0.101	—	W	
<i>Phalaris canariensis</i> L.	14955	Gramineae	3	100	1.000	—	W	
<i>Phoenix dactylifera</i> L.	14303	Palmae	1	10.13	0.101	9	W	[39]
<i>Phragmites australis</i> (Cav.) Steudel	40289	Gramineae	6,7	64.56	0.918	—	W	
<i>Picris echioides</i> L.	47438	Compositae	2	7.59	0.076	—	W	[40]
<i>Pinus halepensis</i> Miller	37506	Pinaceae	2,5,9	51.9	0.791	14, 13, 41, 19, 37, 24, 39	W	[35, 40, 44]
<i>Pinus pinea</i> L.	32768	Pinaceae	3,5,9	7.59	0.120	—	C	
<i>Piptatherum miliaceum</i> (L.) Coss.	6843	Gramineae	3	27.85	0.278	—	C	
<i>Pistacia lentiscus</i> L.	10319	Anacardiaceae	11	6.33	0.063	37	C	[39]
<i>Portulaca oleracea</i> L.	36619	Portulacaceae	2,3	72.15	0.759	41, 38, 15, 44, 33, 23, 18, 11, 4, 25	W	[40]
<i>Punica granatum</i> L.	46140	Punicaceae	9	8.86	0.089	11, 15	W	[40]
<i>Raphanus sativus</i> L.	51395	Cruciferae	3,4	49.37	0.741	29, 12, 3, 21, 13,	W	[41]
<i>Rosa aegrestis</i> Savi	51473	Rosaceae	11	6.33	0.063	18, 41, 24, 4	W	[35, 44]
<i>Rubus ulmifolius</i> Schott	40230	Rosaceae	4,10	11.39	0.133	24, 46, 15, 5, 39, 9, 3, 31	W	[7, 35, 39, 44]
<i>Scolymus hispanicus</i> L.	20754	Compositae	2	40.51	0.405	—	W	
<i>Scolymus maculatus</i> L.	20114	Compositae	2	10.13	0.101	—	W	
<i>Scorzonera hispanica</i> L.	4557	Compositae	4	16.46	0.165	—	W	
<i>Senecio vulgaris</i> L.	7527	Compositae	4	7.59	0.076	—	W	
<i>Setaria italica</i> (L.) P. Beauv.	16519	Gramineae	3	46.84	0.468	—	C	
<i>Silybum marianum</i> (L.) Gaertner	32020	Compositae	2	73.42	0.734	21, 8, 9, 3, 36, 37, 27, 29, 32, 15, 1, 42, 12, 13, 40	C	[7, 35, 41, 44]
<i>Sonchus oleraceus</i> L.	47365	Compositae	4	44.3	0.443	9	W	[39]

TABLE 1: Continued.

Scientific name	Herbarium voucher (ABH)	Family	Finch uses	RFC	CI	Medical human uses	Type	References
<i>Sonchus tenerrimus</i> L.	37483	Compositae	4	40.51	0.405	9	W	[39]
<i>Sorghum halepense</i> (L.) Pers.	3298	Gramineae	3	53.16	0.532	46	W	[41]
<i>Spinacia oleracea</i> L.	Seen alive	Chenopodiaceae	4	43.04	0.430	—	C	
<i>Stellaria media</i> (L.) Vill.	10674	Caryophyllaceae	4	11.39	0.114	—	W	
<i>Stipa tenacissima</i> L.	44375	Gramineae	5	92.41	0.924	37	W	[39]
<i>Taraxacum vulgare</i> (Lam.) Schrank	1808	Compositae	4	49.37	0.494	33, 8	W	[44]
<i>Thymelaea hirsuta</i> L.	41262	Thymelaeaceae	2	27.85	0.278	11, 10	C	[41]
<i>Urtica dioica</i> L.	40147	Urticaceae	1,4	26.58	0.316	1, 20, 5, 41, 16, 13	W	[7]
<i>Urtica urens</i> L.	33640	Urticaceae	1,4,11	34.18	0.424	21, 13, 41, 3, 30, 9, 18, 37, 1, 12, 16	W	[35, 39, 44]
<i>Viscum album</i> L.	49508	Viscaceae	7	60.76	0.608	—	W	

3.1. Uses in Finches. The most important plant species used by bird breeders are *Phalaris canariensis*, *Cannabis sativa*, *Stipa tenacissima*, *Diploptaxis eruroides*, and *Brassica napus*, representing more than 90% of relative citation frequency (RFC). Among the species with the greatest cultural importance, two species with values higher than 1 for the CI index are striking: *Cannabis sativa* (CI = 1.158) and *Phalaris canariensis* (CI = 1). In contrast, the lowest CI are in *Citrus limon* (CI = 0.025), *Chelidonium majus*, *Cicer arietinum*, *Ocimum basilicum* (CI = 0.038), *Allium sativum*, and *Olea europaea* (CI = 0.051).

Most of the plant species (24.75%) identified were placed inside nets to attract and capture wild birds in the field (Figure 2). Thus, once birds have entered the nets, the hunter pulls a rope, and the birds are trapped (Figures 3 and 4). The stems of some plants (e.g., *Lygeum spartum*, *Olea europaea* and *Stipa tenacissima*) are spread with an adhesive substance called birdlime (“envisque” or “liga” in local Spanish), obtained from a mixture of resins (e.g., resin from *Pinus halepensis* and *Pinus pinea*), olive oil (from *Olea europaea*), and some plants (e.g., *Andryala ragusina*, *Chondrilla juncea*, and *Euphorbia characias*). Birds that land on these stems while frequenting feeders or watering points are captured in this way. Catching tools include plants that are used to construct hunter refuges (e.g., *Arundo donax*, *Phragmites australis*, and *Viscum album*) or decoys that are used to attract other birds to the nets (e.g., *Brachypodium retusum* and *Hyparrhenia hirta*). Capture nets must blend in with the terrain conditions; therefore, they are dyed a matte colour that is as close as possible to the surrounding environment. Hunters use an infusion of certain plants (e.g., *Punica granatum*, *Pinus halepensis*, and *Pinus pinea*) to produce these dyes.

Furthermore, many of the species were used to produce the seeds and wild vegetables (18.81%) used to feed birds in captivity. Plants that facilitate breeding include the ones that are used by birds in captivity to build nests (e.g., *Agave americana*, *Cannabis sativa*, *Chamaerops humilis*, *Paronychia argentea*, and *Phoenix dactylifera*), feed their offspring (e.g., *Brassica oleracea* var. *italica*), and stimulate mating (e.g., *Urtica dioica* and *Urtica urens*). Breeders used the fruits and roots of some plants (e.g., *Daucus carota*, *Fragaria vesca*, and *Rubus ulmifolius*) to enhance the natural red factor in some species of birds, providing natural pigments, particularly in *Carduelis cannabina* and *Carduelis carduelis*. Currently, the cages are made principally from metal or synthetic materials; however, informants can identify the specific natural materials that are used to be used to build cages and cage accessories (e.g., *Arundo donax*, *Daphne gnidium*, *Phragmites australis*, and *Nerium oleander*).

Birds in captivity may suffer from certain diseases, and breeders often try to cure these birds by using natural, plant-based remedies. Thus, there are some vulnerary plants (e.g., *Chelidonium majus* and *Rosa agrestis*) and others that stop haemorrhages (e.g., ash of *Nicotiana tabacum*). Some species have antibacterial properties (e.g., *Cicer arietinum*, vinegar of *Malus domestica*, and *Citrus limon*), or they promote moulting (e.g., *Lavandula latifolia*), have disinfectant functions to eliminate microbes (e.g., *Pistacia lentiscus*), or can

host beneficial probiotic bacteria or tonic (e.g., vinegar of *Malus domestica*). Some plants have been used as vermifuge, placed in the breeding carrier, in order to expel parasites (e.g., worms) from the intestines, such as mites (especially *Syringophilus* sp., *Dermoglyphus* sp., and *Dermanyssus* sp.) and lice (*Menacanthus* sp. and *Goniocotes* sp.) that affect this group of birds. Leafy vegetables are used as a laxative treatment, the juice of *Urtica urens* to prevent anaemia, and *Cicer arietinum* is used to stop diarrhoea. To sum up, we show the number of species that are used with specific bird veterinarian uses in Table 2.

3.2. Human Medicine Uses. According to the ethnobotanical references consulted, we found 57 plants used in finches that have medical properties in humans. These species are used to cure some ailments related to each pathological group (Table 2). Thus, 48 human uses have been detected in the 97 plant species collected in the study area. *Silybum marianum* (15), *Olea europaea* (12), and *Centaurea aspera* (12) are the species with greater therapeutic uses. We found that 48 uses were related to medical properties: alteration of blood pressure ($n = 9$), haemorrhoids ($n = 3$), depurative ($n = 9$), anxiety ($n = 9$), diarrhoea ($n = 9$), heartburn ($n = 2$), indigestion ($n = 8$), liver disease ($n = 9$), loss of appetite ($n = 19$), constipation ($n = 11$), helminthiasis ($n = 8$), cough ($n = 7$), cold ($n = 16$), respiratory problems ($n = 8$), hyperglycemia ($n = 9$), anaemia ($n = 2$), hypercholesterolemia ($n = 1$), retention of liquids ($n = 16$), undefined symptom (tonic) ($n = 7$), gout ($n = 3$), rheumatism ($n = 5$), inflammation of bones or joints ($n = 11$), undefined symptom (analgesic) ($n = 4$), injury ($n = 13$), burns ($n = 4$), kidney stones ($n = 2$), menstruation ($n = 5$), lack of breast milk secretion ($n = 2$), ischocholia ($n = 6$), chilblains ($n = 3$), pimples ($n = 5$), skin diseases ($n = 3$), eczema ($n = 5$), skin fungus ($n = 1$), rubefaction ($n = 1$), calluses and skin hardness ($n = 3$), warts ($n = 11$), bacteria ($n = 2$), microbes ($n = 5$), headache ($n = 2$), inflammation ($n = 11$), fever ($n = 3$), alopecia ($n = 1$), flushing (refreshing) ($n = 3$), alcoholism ($n = 1$), toothache ($n = 8$), mineral deficiency ($n = 1$), and eye infection ($n = 2$).

We only found three vulnerary plants for finches; however, there are 13 species of the total used for this use in humans. There are three antibacterial plants in birds, while in humans we found two different species (*Portulaca oleracea* and *Centaurea aspera*). One plant is disinfectant for finches, while in humans there are 5 antiseptics to eliminate microbes (*Foeniculum vulgare*, *Centaurea aspera*, *Pinus halepensis*, *Lavandula latifolia*, and *Rubus ulmifolius*) and fungal species (*Centaurea aspera*). Twenty eight species are used as a laxative treatment in birds, whereas only eleven have the same medical use for humans. Conversely, we found no plants that are probiotic or that stop bleeding in humans.

4. Discussion

Traditionally, nutritive uses [45] and curative applications [46] of ethnobotanical knowledge have been linked to women. They have demonstrated a high knowledge of both



FIGURE 2: Local bird breeder.


FIGURE 4: Goldfinch claim in *Centaurea aspera*.

FIGURE 3: Traditional hunting method using nets.

wild and cultivated species [47, 48], especially in rural areas [7]. In contrast, wild bird hunting is traditionally a male-dominated pastime. Therefore, we want to highlight that the stakeholders have high know-how, which reflects their identification of different species and their applications. The names and traditional uses can vary depending on geographical location, as vernacular names serve as intangible heritage. Thus, it is necessary to preserve this heritage and promote educational and awareness programmes [20].

The cultural importance index corresponds with an interest in detailing the specific uses of plants that better reflect the cultural aspects of plant utilization. In fact, ethnobotanical publications usually present plant uses in tables or catalogues, where the information is grouped by species, indicating their particular uses and, commonly, the number of informants who mentioned them. This way of grouping is much more reasonable for evaluating the importance of each plant species by its cultural consensus [8]. This additive index takes into account not only the spread of use (number of informants) for each species, but also its versatility, that is, the diversity of its uses [17]. Thus, *Cannabis sativa* and *Phalaris canariensis* have the greatest CI, being the principal commercial seed and, moreover, *Cannabis sativa* has other uses. In contrast, the lowest CI are in plants that are used to cure or have no typical uses and are not used by informants to breed songbirds.

Various mixes of dried seeds, composed of seeds from different species, both wild and cultivated, are used to feed birds in captivity [49]. Each bird breeder uses the mixture

of seeds that he/she deems most appropriate. However, some breeders use leafy vegetables to feed birds and supplement their diet of dried seeds. These plants are used mainly in summer, during the birds' moulting period and as a laxative. Other species not cited in this study, such as *Ilex aquifolium*, *Viscum cruciatum*, or *Onopordum nervosum* [11–13, 50–52], are used to capture birds in other areas. Moreover, some plants also have different veterinary uses in other Mediterranean regions. Thus, some authors show that several species, such as *Stellaria media*, *Avena sativa*, and *Urtica dioica*, are used to increase fertility and egg production in chickens. *Urtica urens* is mixed with feed for hens so that they lay eggs earlier in their lifespan and as a result, the eggshells will be harder. *Cirsium arvense*, *Daphne gnidium*, *Phragmites australis*, and *Linum usitatissimum* are anti-diarrhoeal and have been used to favour digestion. *Allium sativum*, *Daphne gnidium*, *Nerium oleander*, and *Nicotiana tabacum* are useful against parasites on farms, and *Cicer arietinum* is used to facilitate the expulsion of the placenta and for purgation in goats and sheep. *Olea europaea* is used to treat mastitis or to detoxicate, and latex from *Chelidonium majus* and *Pinus halepensis* is used to treat wounds [17, 19, 21, 53]. With these data, we can verify that there is a popular tradition for the use of ethnoveterinary plants in Mediterranean areas.

Furthermore, some species identified without human medicine use in the study area have them in other Spanish regions [5, 54–57], such as *Avena sativa* (toothache and quitting smoking), *Bituminaria bituminosa* (vulnerary), *Brassica oleracea* var. *italica* (vulnerary, remineralizing, headache, and anthelmintic), *Brassica rapa* (culinary), *Cannabis sativa* (refreshing and relaxing), *Chelidonium majus* (anti-cholagogue, hepatoprotective, anti-inflammatory, antiseptic, warts, laxative, and vulnerary), *Conyza bonariensis* (digestive), *Helianthus annuus* (febrifuge), *Phagnalon saxatile* (carminative, analgesic, and cholesterol levels), *Phalaris canariensis* (cholesterol), *Scorzonera hispanica* (diuretic, uric acid, and cholesterol), *Senecio vulgaris* (anti-inflammatory and antiseptic), and *Viscum album* (anticatarrhal, antiseptic, antivariolous, parasiticide, salutiferous, and sedative). Other species, such as *Carthamus tinctorius*, *Centaurea mariolensis*, *Centaurea melitensis*, *Guizotia abyssinica*, *Panicum miliaceum*, *Perilla frutescens*, *Setaria italica*, and *Spinacia oleracea*, do not present other applications in humans, according to

TABLE 2: Number and frequency of plants used for a specific human use.

Pathologic group	Human use	Medical code	No. of species	Frequency	Bird veterinarian
Circulatory system	Alteration of blood pressure	1	9	9.28	
	Haemorrhoids	2	3	3.09	
Mental illness	Undefined symptom (depurative)	3	9	9.28	
	Anxiety	4	9	9.28	
	Diarrhoea	5	9	9.28	1
	Heartburn	6	2	2.06	
	Indigestion	7	8	8.25	1 probiotic
Digestive system	Liver disease	8	4	4.12	
	Loss of appetite	9	19	19.59	
	Constipation	10	11	11.34	28
	Helminthiasis	11	8	8.25	6
	Cough	12	7	7.22	
Respiratory system	Cold	13	16	16.49	
	Respiratory problems	14	8	8.25	
	Hyperglycemia	15	9	9.28	
	Anaemia	16	2	2.06	1
Metabolism, nutrition, and so forth	Hypercholesterolemia	17	1	1.03	
	Retention of liquids	18	16	16.49	
	Undefined symptom (Tonic)	19	7	7.22	1
	Gout	20	3	3.09	
Bones, joints, and so forth	Rheumatism	21	5	5.15	
	Inflammation	22	1	1.03	
	Undefined symptom (analgesic)	23	4	4.12	
Traumatic injuries and poisoning	Injury	24	13	13.40	3
	Burns	25	4	4.12	
	Kidney stones	26	2	2.06	
Genital urinary	Menstruation	27	5	5.15	
	Lack of breast milk secretion	28	2	2.06	
	Ischocholia	29	6	6.19	
	Chilblain	30	3	3.09	
	Pimples	31	5	5.15	
Skin and subcutaneous tissues	Skin problems	32	3	3.09	
	Eczema	33	5	5.15	
	Skin fungus	34	1	1.03	
	Rubefaction	35	1	1.03	
	Calluses and skin hardness	36	2	2.06	
Infectious and parasitic diseases	Warts	37	11	11.34	
	Bacteria	38	2	2.06	3
	Microbes	39	5	5.15	1
	Headache	40	2	2.06	
	Inflammation	41	11	11.34	
Symptoms, signs, and poorly defined morbid states	Fever	42	3	3.09	
	Alopecia	43	1	1.03	
	Flushing	44	3	3.09	
	Alcoholism	45	1	1.03	
	Toothache	46	8	8.25	
	Mineral deficiency	47	1	1.03	1 molting
	Nervous system and sensory organs	Eye infection	48	2	2.06

these references. This may be due to the rarity of these species or that they are not traditionally cultivated species in the area.

However, some of the species studied in this project are in the group of the top vascular plants in traditional phytotherapy in other regions, such as *Allium sativum* (antiparasitic, anthelmintic, anti-inflammatory/analgesic, antiverucose, and anti-bronchitic), *Foeniculum vulgare* (carminative, cold, intestinal anti-inflammatory, laxative, gastralgia, diuretic, and antihalitosis), and *Olea europaea* (antihypertensive, hyperglycemia, hernia, food poisoning, heartburn, warts, cough, erysipelas, sores, psoriasis, burns, hoarseness, baldness, rheumatism, antipyretic, antiseptic, laxative, and antinostalgic) [18, 58].

On the other hand, bird populations have declined, mainly due to the abandonment of crops, the use of pesticides, predation of nests, poaching, increased predation due to changes in their natural habitat, uncontrolled development, and in general socioeconomic changes in recent decades [59]. In this aspect, the mechanization of agricultural practices has changed the structure of these agrarian ecosystems, accompanied by a steady degradation and loss of landscape elements with important ecological functions [60]. To preserve bird populations, it is essential to maintain fields active. There are many plants linked to these environments that birds use daily, such as for food or other purposes.

5. Conclusions

In conclusion, data obtained in this research are scarcely known and show many details of plants related to songbirds, facilitating access to interesting and novel information. This allows recovery of forgotten uses and traditions, highlighting the utilization of different species to attract and cure birds and their relation to human medicine, and resulting in a very interesting contribution to ethnobotanical bibliography.

We found that the majority of the plant species related to songbirds were wild, reflecting that the wild bird hunters are aware of this preference and exploit this knowledge of wild flora in their hunting. This demonstrates that informants have great knowledge of the plants used in traditional medicine and finch keeping. Also, the majority of species have medicinal properties that can be used for informants to cure different pathologies.

Acknowledgments

The authors thank the Consellería de Medio Ambiente, Agua, Urbanismo y Vivienda, and the rangers who worked with them on the project. They would also like to thank SEPRONA (Department of Nature Protection), Cultural Spanish Wildlife Ornithological Federation (FOCSE), and all collaborators who have contributed information to their study, particularly Miguel Segarra Teruel. In addition, they thank José Carlos Cristobal and Antonio De la Torre for the vouchers provided. This study was supported by the WADI Project (INCO2003-MPC2-015226), financed by the European Commission.

References

- [1] A. Agelet, M. A. Bonet, and J. Vallès, "Homegardens and their role as a main source of medicinal plants in mountain regions of Catalonia (Iberian Peninsula)," *Economic Botany*, vol. 54, no. 3, pp. 295–309, 2000.
- [2] D. Raja, C. Blanche, and J. Xirau, "Contribution to the knowledge of the pharmaceutical ethnobotany of La Segarra region (Catalonia, Iberian Peninsula)," *Journal of Ethnopharmacology*, vol. 57, no. 3, pp. 149–160, 1997.
- [3] M. À. Bonet, M. Parada, A. Selga, and J. Vallès, "Studies on pharmaceutical ethnobotany in the regions of L'Alt Empordà and Les Guilleries (Catalonia, Iberian Peninsula)," *Journal of Ethnopharmacology*, vol. 68, no. 1–3, pp. 145–168, 1999.
- [4] A. Agelet and J. Vallès, "Studies on pharmaceutical ethnobotany in the region of Pallars (Pyrenees, Catalonia, Iberian Peninsula). Part I. General results and new or very rare medicinal plants," *Journal of Ethnopharmacology*, vol. 77, no. 1, pp. 57–70, 2001.
- [5] A. Agelet and J. Vallès, "Studies on pharmaceutical ethnobotany in the region of Pallars (Pyrenees, Catalonia, Iberian Peninsula). Part II. New or very rare uses of previously known medicinal plants," *Journal of Ethnopharmacology*, vol. 84, no. 2–3, pp. 211–227, 2003.
- [6] J. Camejo-Rodrigues, L. Ascensão, M. A. Bonet, and J. Vallès, "An ethnobotanical study of medicinal and aromatic plants in the Natural Park of "Serra de São Mamede" (Portugal)," *Journal of Ethnopharmacology*, vol. 89, no. 2–3, pp. 199–209, 2003.
- [7] A. Belda, F. J. Bellod, and S. Ríos, "Avance sobre la flora medicinal en la sierra de Mariola (Valencia-Alicante)," *Flora Montiberica*, vol. 28, pp. 29–48, 2004.
- [8] J. Tardío and M. Pardo-de-Santayana, "Cultural importance indices: a comparative analysis based on the useful wild plants of Southern Cantabria (Northern Spain)," *Economic Botany*, vol. 62, no. 1, pp. 24–39, 2008.
- [9] M. À. Bonet and J. Vallès, "Use of non-crop food vascular plants in Montseny biosphere reserve (Catalonia, Iberian Peninsula)," *International Journal of Food Sciences and Nutrition*, vol. 53, no. 3, pp. 225–248, 2002.
- [10] J. Tardío, H. Pascual, and R. Morales, "Wild food plants traditionally used in the province of Madrid, Central Spain," *Economic Botany*, vol. 59, no. 2, pp. 122–136, 2005.
- [11] L. Mulet, *Estudio Etnobotánico de la Provincia de Castellón*, Diputación Provincial de Castellón, 1991.
- [12] M. Gil-Pinilla, *Estudio etnobotánico de la flora aromática y medicinal del término municipal de Cantalojas (Guadalajara)*, Doctoral thesis, Universidad Complutense de Madrid, 1995.
- [13] A. Verde, J. Fajardo, D. Rivera, and C. Obón, *Etnobotánica en el Entorno del Parque Nacional de Cabañeros*, Organismo Autónomo Parques Nacionales, Madrid, Spain, 2000.
- [14] R. Y. Cavero, S. Akerreta, and M. I. Calvo, "Pharmaceutical ethnobotany in Northern Navarra (Iberian Peninsula)," *Journal of Ethnopharmacology*, vol. 133, no. 1, pp. 138–146, 2011.
- [15] R. Morales, J. Tardío, L. Aceituno, M. Molina, and M. Pardo de Santayana, "Biodiversity and ethnobotany in Spain," *Memorias de la Real Sociedad Española de Historia Natural*, vol. 9, pp. 157–207, 2011.
- [16] V. Reyes-García, S. Vila, L. Aceituno-Mata et al., "Gendered homegardens: a study in three mountain areas of the Iberian peninsula," *Economic Botany*, vol. 64, no. 3, pp. 235–247, 2010.
- [17] J. A. González, M. García-Barriuso, and F. Amich, "Ethnoveterinary medicine in the Arribes del Duero, western Spain,"

- Veterinary Research Communications*, vol. 35, no. 5, pp. 283–310, 2011.
- [18] G. Benítez, M. R. González-Tejero, and J. Molero-Mesa, “Knowledge of ethnoveterinary medicine in the Province of Granada, Andalusia, Spain,” *Journal of Ethnopharmacology*, vol. 139, no. 2, pp. 429–439, 2012.
- [19] R. E. Uncini-Manganelli, F. Camangi, and P. E. Tomei, “Curing animals with plants: traditional usage in Tuscany (Italy),” *Journal of Ethnopharmacology*, vol. 78, no. 2–3, pp. 171–191, 2001.
- [20] A. Belda, J. E. Martínez-Pérez, C. Martín, V. Peiró, and E. Seva, “Plants used to capture and sustain wild finches (Fringillidae) in Southeast Spain,” *Economic Botany*, vol. 64, no. 4, pp. 367–373, 2010.
- [21] R. E. Uncini-Manganelli and P. E. Tomei, *Documenti per la Conoscenza delle Tradizioni Etno-Farmacobotaniche in Toscana*, S. Marco litotipo, Accademia Lucchese di Scienze, Lettere ed Arti, Lucca, Italy, 1999.
- [22] L. Viegi, A. Pieroni, P. M. Guarrera, and R. Vangelisti, “A review of plants used in folk veterinary medicine in Italy as basis for a databank,” *Journal of Ethnopharmacology*, vol. 89, no. 2–3, pp. 221–244, 2003.
- [23] L. Clark and J. R. Mason, “Use of nest material as insecticidal and anti-pathogenic agents by the European Starling,” *Oecologia*, vol. 67, no. 2, pp. 169–176, 1985.
- [24] E. Eythorson, “Sami Fjord fishermen and the state: traditional knowledge and resource management in northern Norway,” in *Traditional Ecological Knowledge. Concepts and Cases*, J. T Inglis, Ed., pp. 133–143, International Development Research Centre, Ottawa, Canada, 1993.
- [25] R. E. Johannes, “The need for a centre for the study of indigenous fishers’ knowledge,” Contribution to Wise Coastal Practices for Sustainable Human Development, 2001, <http://www.csiwisepractices.org/>.
- [26] I. G. Baird, “Local ecological knowledge and small-scale freshwater fisheries management in the Mekong river in Southern Laos,” in *Putting Fishers’ Knowledge to Work*, N. Haggan, C. Brignall, and L. Wood, Eds., pp. 87–99, University of British Columbia, 2007.
- [27] A. Davis and J. R. Wagner, “Who knows? On the importance of identifying “experts” when researching local ecological knowledge,” *Human Ecology*, vol. 31, no. 3, pp. 463–489, 2003.
- [28] F. Martínez, *Arte de Cocina, Pastelería, Bizcochería y Conservería*, Extramuros, Madrid, Spain, Facsimile edition published in 1778, Imprenta de Pantaleón Aznar, 2009.
- [29] Real Academia Española, *Diccionario de la Lengua Española*, 22rd edition, 2001.
- [30] A. Belda, J. E. Martínez, J. Arques, V. Peiró, E. Seva, and D. Jiménez, “Métodos de caza tradicionales empleados en el Carrascal de la Font Roja,” *Mediterranea*, vol. 19, pp. 9–39, 2008.
- [31] E. J. Belda, La captura con red de fringílidos en la Comunidad Valenciana en el 2003: estudio científico de dicha modalidad de caza. Informe para la Conselleria de Territorio y Vivienda de la Generalitat Valenciana, pp. 102, 2003.
- [32] M. D. Boza, *El trampeo y Demás Artes de Caza Tradicionales en la Península Ibérica*, Editorial Hispano Europea, 2004.
- [33] M. Natale and L. G. Pidalà, *El Jilguero*, Croma press S. A., 2004.
- [34] A.J. Pérez, *Atlas climático de la Comunidad Valenciana (1961–1990)*, Conselleria de de Obras Públicas, Urbanismo y Transporte, 1994.
- [35] A. Belda and F. J. Bellod, *Plantas Medicinales de la Sierra de Mariola*, Servicio de Publicaciones de la Universidad de Alicante, 2007.
- [36] L. Serra, *Orquídeas Silvestres de la Comunidad Valenciana*, Conselleria de Medio Ambiente de la Generalidad Valenciana, 2001.
- [37] L. Serra, *Estudio Crítico de la Flora Vasculare de la Provincia de Alicante: Aspectos Nomenclaturales, Biogeográficos y de Conservación*, vol. 19, Ruizia, 2007.
- [38] D. Climent, *Les Nostres Plantes: (una aproximació multidisciplinaria al món vegetal de les nostres terres)*, Alacant Aguaclara, 1992.
- [39] J. Pellicer, *Costumari Botànic: Recerques Etnobotàniques a Les Comarques Centrals Valencianes*, vol. 1, Edicions del Bullent, Valencia, Spain, 2000.
- [40] J. Pellicer, *Costumari Botànic: Recerques Etnobotàniques a Les Comarques Centrals Valencianes*, vol. 2, Edicions del Bullent, Valencia, Spain, 2000.
- [41] J. Pellicer, *Costumari Botànic: Recerques Etnobotàniques a Les Comarques Centrals Valencianes*, vol. 3, Edicions del Bullent, Valencia, Spain, 2004.
- [42] O. Guasch, *Observación Participante. Cuadernos Metodológicos*, vol. 20, Centro de Investigaciones Sociológicas, Madrid, Spain, 1997.
- [43] G. Mateo and M. B. Crespo, *Manual Para La Determinación De La Flora Valenciana*, Monografías de Flora Montiberica 4, Valencia, Spain, 3th edition, 2003.
- [44] J. C. Belda and A. Belda, *Naturaleza y Salud: Aceites, Cataplasmas y Tisanas*, Carena Editors S.L., 2008.
- [45] M. Sundriyal and R. C. Sundriyal, “Wild edible plants of the Sikkim Himalaya: nutritive values of selected species,” *Economic Botany*, vol. 55, no. 3, pp. 377–390, 2001.
- [46] E. Dopico, J. L. San Fabian, and E. Garcia-Vazquez, “Traditional medicine in twenty-first Spain,” *Human Ecology*, vol. 36, no. 1, pp. 125–129, 2008.
- [47] F. A. J. de Clerck and P. Negreros-Castillo, “Plant species of traditional Mayan homegardens of Mexico as analogs for multistrata agroforests,” *Agroforestry Systems*, vol. 48, no. 3, pp. 303–317, 2000.
- [48] A. C. Hamilton, “Medicinal plants, conservation and livelihoods,” *Biodiversity and Conservation*, vol. 13, no. 8, pp. 1477–1517, 2004.
- [49] H. G. Adams, *Cage and Singing Birds: How to Catch, Keep, Breed, and Rear Them. With Full Directions as to Their Nature, Habits, Food, Diseases, Etc*, George Routledge, London, UK, 1854.
- [50] D. Rivera and C. Obón, *Las Plantas Medicinales de Nuestra Región*, Regional de Murcia, 1991.
- [51] D. Rivera, A. Verde, J. Fajardo, C. Inocencio, C. Obón, and M. Heinrich, *Guía Etnobotánica de Los Alimentos Locales Recolectados en la Provincia de Albacete*, Instituto de Estudios Albacetenses “Don Juan Manuel”, Excm. Diputación de Albacete, 2006.
- [52] M. Pardo-de-Santayana, *Estudios Etnobotánicos en Campoo (Cantabria): Conocimiento y Uso Tradicional de Las Plantas*, Consejo Superior de Investigaciones Científicas, 2008.
- [53] A. Pieroni, M. E. Giusti, C. de Pasquale et al., “Circum-mediterranean cultural heritage and medicinal plant uses in traditional animal healthcare: a field survey in eight selected areas within the RUBIA project,” *Journal of Ethnobiology and Ethnomedicine*, vol. 2, article 16, 2006.

- [54] M. Pardo-de-Santayana, J. Tardío, and R. Morales, “The gathering and consumption of wild edible plants in the Campoo (Cantabria, Spain),” *International Journal of Food Sciences and Nutrition*, vol. 56, no. 7, pp. 529–542, 2005.
- [55] M. Rigat, T. Garnatge, and J. Vallès, *Plantes I Gent. Estudi Etnobotànic de l’Alta Vall del Ter*, Centre d’Estudis Comarcals del Ripollès, 2006.
- [56] A. Agelet, *Plantes Medicinals del Pallars*, Impremta Aubert, 2008.
- [57] P. Font, *Plantas Medicinales: El Dioscórides Renovado*, Ediciones Peninsula S.A, 1999.
- [58] M. Parada, E. Carrió, M. A. Bonet, and J. Vallès, “Ethnobotany of the Alt Empordà region (Catalonia, Iberian Peninsula). Plants used in human traditional medicine,” *Journal of Ethnopharmacology*, vol. 124, no. 3, pp. 609–618, 2009.
- [59] J. E. Martínez-Pérez, A. Belda-Antolí, C. Martín-Cantarino, and E. Seva-Román, “Long-term (1883 to 2008) historical changes in agricultural landscape patterns in a traditional irrigated system of Alicante province, Spain: the case of Lo Contador property (Carrizales of Elche),” in *Sustainable Management of Mediterranean Coastal Fresh and Transitional Water Bodies: a Socio-Economic and Environmental Analysis of Changes and Trends to Enhance and Sustain Stakeholder Benefits*, F. Scapini, J. M. Boffa, E. Conrad, L. F. Cassar, and M. Nardi, Eds., pp. 29–36, Firenze University Press, 2009.
- [60] R. B. Sage, D. M. B. Parish, M. I. A. Woodburn, and P. G. L. Thompson, “Songbirds using crops planted on farmland as cover for game birds,” *European Journal of Wildlife Research*, vol. 51, no. 4, pp. 248–253, 2005.