



ELSEVIER

Contents lists available at ScienceDirect

Data in brief

journal homepage: www.elsevier.com/locate/dib

Data Article

Dataset on marine ecosystem services supplied by coral reefs, sandy beaches and coastal lagoons in different eutrophication states



Charlène Kermagoret ^{a, b, *}, Joachim Claudet ^{c, d},
Valérie Derolez ^e, Maggy M. Nugues ^{f, d}, Vincent Ouisse ^e,
Nolwenn Quillien ^g, Denis Bailly ^a

^a Univ Brest, Ifremer, CNRS, UMR 6308, AMURE, IUEM, 29280, Plouzane, France

^b Département des Sciences Naturelles, Institut des Sciences de la Forêt Tempérée, Université du Québec en Outaouais, Gatineau, Canada

^c National Center for Scientific Research, PSL Université Paris, CRIOBE, USR 3278 CNRS-EPHE-UPVD, Maison des Océans, 195 rue Saint-Jacques 75005 Paris, France

^d Labex Corail, CRIOBE, 98729 Moorea, French Polynesia

^e MARBEC, Ifremer, IRD, Univ Montpellier, CNRS, Av. Jean Monnet, 30171 - 34203 Sète Cedex, CS, France

^f EPHE, PSL Research University, UPVD-CNRS, USR3278 CRIOBE, F-66860 Perpignan, France

^g France Energies Marines, 29200 Brest, France

ARTICLE INFO

Article history:

Received 9 April 2019

Received in revised form 10 May 2019

Accepted 22 May 2019

Available online 29 May 2019

Keywords:

Ecosystem services

Marine ecosystems

Eutrophication

Marine biodiversity

ABSTRACT

This data article provides indicators of Ecosystem Service (ES) supply for coral reefs, sandy beaches and coastal lagoons in different ecological states regarding eutrophication. 14 ES are considered: food through fisheries; material; molecules; coastal protection; nutrient regulation; pathogen regulation; climate regulation; support of recreational and leisure activities; contribution to a pleasant landscape; contribution to culture and territorial identity; emblematic biodiversity; habitat; trophic networks; recruitment. For each ecosystem 3 to 4 eutrophication states are described. Indicators of ES supply are filled on the basis of a literature review supplemented with expert-knowledge. A semi-quantification of the indicator value is finally provided. Tendencies and trade-offs between ES are analyzed in How does

DOI of original article: <https://doi.org/10.1016/j.ococoaman.2019.03.028>.

* Corresponding author. Univ Brest, Ifremer, CNRS, UMR 6308, AMURE, IUEM, 29280, Plouzane, France.

E-mail address: Charlene.Kermagoret@ifremer.fr (C. Kermagoret).

<https://doi.org/10.1016/j.dib.2019.104078>

2352-3409/© 2019 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

eutrophication impact bundles of ecosystem services in multiple coastal habitats using state-and-transition models [1].

© 2019 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Specifications Table [please fill in right-hand column of the table below]

Subject area	<i>Environmental management</i>
More specific subject area	<i>Ecosystem services and biodiversity</i>
Type of data	<i>Table</i>
How data was acquired	<i>Literature review and expert-knowledge coded on Microsoft Excel 2016</i>
Data format	<i>Raw</i>
Experimental factors	<i>For each ecosystem, 3 to 4 eutrophication states are considered and described by supply indicators of 14 ES</i>
Experimental features	<i>Eutrophication states are described by dominant species and associated biodiversity. Supply indicators of ES are qualitative or quantitative information, depending on available data. Information is then coded to assess the variation of ES supply between eutrophication states (five levels from very low to very high).</i>
Data source location	<i>The literature review encompassed knowledge obtained and disseminated on a global scale, while expert knowledge focused on data observed on a more local scale, based on their field studies. However, experts had a good understanding of these ecosystems that allowed them to pronounce in a qualitative way where data gaps were identified.</i>
Data accessibility	<i>Data are provided in this article</i>
Related research article	C. Kermagoret, J. Claudet, V. Derolez, M. Nugues, V. Ouisse, N. Quillien, Y. Baulaz, P. Le Mao, P. Scemama, D. Vaschalde, D. Bailly, R. Mongruel, Comparison of how eutrophication affects bundles of ecosystem services in multiple coastal habitats using state-and-transition models, <i>Ocean Coast. Manage.</i> 174 (2019) 144–153. https://doi.org/10.1016/j.ocecoaman.2019.03.028

Value of the data

- The dataset is useful for understanding the relations between ecological functions, ecological states and ES.
- Scientific can benefit from this data to carry out ES assessment or to inform decision-making processes and management strategies regarding eutrophication.
- Practitioners and policy-makers can benefit from this data to adopt management strategies or measures for biodiversity conservation.
- The data provide a first overview of the effects of eutrophication on ES bundles and can serve as a basis for a larger database.
- The data can be used for further insight the trade-offs and synergies between ES and can be used for modelling.
- The dataset helps to build relationships between disparate data on the effect of eutrophication on ecological function and ES.

1. Data

The dataset contains qualitative description of ES provided by coral reefs, sandy beaches and coastal lagoons in different eutrophication states. Three to four eutrophication states are considered, from non-eutrophic to hyper-eutrophic state. 14 ES are considered: food through fisheries; material; molecules; coastal protection; nutrient regulation; pathogen regulation; climate regulation; support of recreational and leisure activities; contribution to a pleasant landscape; contribution to culture and territorial identity; emblematic biodiversity; habitat; trophic networks; recruitment. The capacity of the ecosystem to provide ES is described for each ES in each eutrophication state using qualitative information collected in the literature completed by expert-knowledge. Data are presented in double entry matrix (ES/eutrophication states) for coral reefs (Table 1), sandy beaches (Table 2), coastal lagoons (Table 3). These capacities - or supplies - are finally semi-quantified in order to compare them in the bundle.

Table 1

Indicators and semi-quantification (SQ) of ES supply for coral reefs in 3 eutrophication states ("0: inexistent", "1: very low", "2: low", "3: medium", "4: high", "5: very high").

Eutrophication states	Non-eutrophic Dominance of hard coral		Eutrophic Dominance of algal turf and macroalgae		Hyper-eutrophic Dominance of cyanobacterial mats	
Human food through fisheries	Coral-dominated reefs are the most productive [12]	5	Depends on structural complexity. Multi-species fisheries can shift towards herbivorous fish species dominated yields [13]	3	Bottoms lacking structural complexity become very poor in target species.	1
Material (organism for domestic uses, industry, agriculture ...)	Ornamental purposes	2	Ornamental purposes	1	Ornamental purposes	1
Molecules (organisms from which are extracted molecules potentially useful for medicine)	Living coral reef supports a high diversity of organisms and thus potential biomolecules	4	Lower diversity of organisms and thus lower potential for biomolecules	2	Cyanobacteria and sponges are chemically rich and can be used in medical chemistry, pharmacology or phytopharmacy. ^a	4
Coastal protection (vegetal or animal reef supplying a protection against erosion and submersion)	Coral reef can dissipate 97% of the wave energy that would otherwise impact shorelines [15]	5	The loss of corals, the increased water depth between the reef crest and the surface should result in a less effective ES.	1	The loss of corals, the increased water depth between the reef crest and the surface should result in a nonexistent ES.	0
Nutrient regulation (ecosystem capacity to supply a "good quality water", limiting the risk of eutrophication, encouraging shellfish farming...)	Intensively performed by zooxanthellae in living coral reef.	5	The risk of hyper-eutrophication is greater in altered or dead coral reefs.	3	Cyanobacterial mats and sponges, able to provide the ES, can be easily washed away by storms.	1
Pathogen regulation (ability of ecosystems to purify the environment through hyperfiltration processes)	High capacity of pathogen regulation thanks coral microbiome.	4	Macroalgae alter the coral microbiome and elevate putative pathogen loads [16,17]	2	Macroalgae alter the coral microbiome and elevate putative pathogen loads [16,17]	1
Climate regulation (GES storage or sequestration)	Living coral reefs widely contribute to climate regulation, stocking GES through the production of carbonates [18]	4	The loss of corals results in a less effective ES.	2	The loss of corals results in a less effective, or even nonexistent, ES.	1
Support of recreational and leisure activities	In many tropical societies, relations to nature are often very different from those related to the Western lifestyle and the distinction between culture and nature is sometimes blurred.	5	In these tropical contexts, the difficult resilience and adaptive capacity to abrupt changes in coral reefs (eutrophication and other pressures) can alter cultural ES [19]	3	In these tropical contexts, the difficult resilience and adaptive capacity to abrupt changes in coral reefs (eutrophication and other pressures) can alter cultural ES [19]	1
Contribution to a pleasant landscape		5	Emblematic biodiversity also decreases.	3	Emblematic biodiversity also decreases.	1
Contribution to culture and territorial identity		5	Algal-dominated state can benefit some herbivorous fishes, but large fleshy macroalgae	2	changes in coral reefs (eutrophication and other pressures) can alter cultural ES [19]	1
Emblematic biodiversity (i.e. protected or rared species)		5		2	Emblematic biodiversity also decreases.	0
Habitat (nursery, reproduction area...)	Corals provide shelter and food for a large diversity of benthic organisms and allow	5		3	Cyanobacterial mats are often unpalatable to fishes. Mesopredators can switch prey,	1
Trophic networks		5		3		1
Recruitment		5		3		1

(continued on next page)

Table 1 (continued)

Eutrophication states	Non-eutrophic Dominance of hard coral	Eutrophic Dominance of algal turf and macroalgae	Hyper-eutrophic Dominance of cyanobacterial mats
	the creation of complex trophic networks. The three dimensional structure of corals are important to fish recruitment, which can, in turn, increase herbivory and favor coral dominance via positive feedback mechanisms [20]	are often unpalatable to fishes. Mesopredators can switch prey, shortening food chains, in response to coral reef degradation [21]	shortening food chains, in response to coral reef degradation [21]

^a However, chemical defense could lessen in absence of consumers. For example, sponge communities have become dominated by fast-growing species that lack chemical defenses on reefs where sponge-eating angelfishes and parrotfishes have been removed by overfishing [14].

2. Experimental design, materials and methods

For each ecosystem, eutrophication states are described by dominant species and associated biodiversity. First eutrophication state of coral reefs is characterized by the dominance of hard corals [1]. Other benthic communities present are algal turfs, crustose coralline algae, sponges and benthic cyanobacterial mats [2]. The increasing pressure leads to the development of algal turfs and fleshy macroalgae which are fast-growing organisms and a gradual decline of coral cover [3], in particular from competitive losses against algae under conditions of reduced herbivory [4]. With a continuous

Table 2

Indicators and semi-quantification (SQ) of ES supply for sandy beaches in 3 eutrophication states ("0: inexistent", "1: very low", "2: low", "3: medium", "4: high", "5: very high").

Eutrophication states	Non-eutrophic Reference species (Tellinidae, Spionidae, Amphiuroidae, Nephtyidae)	Eutrophic Green algae proliferation (Donacidae, Oweniidae, Magelonidae, Nephtyidae)	Hyper-eutrophic Green tide (Donacidae, Oweniidae)	
Human food through fisheries	Sandy beaches support professional fisheries of the bivalve <i>Donax trunculus</i> , which is of commercial importance [22] They are also a high spot for surf casting targeting Moronidae (<i>Dicentrarchus labrax</i>).	5 A decrease in <i>Donax trunculus</i> density has been shown [7]. However, the presence of floating mats (within 0.5-1.5m of water) seemed to form the perfect hunting ground for the seabass (pers. obs.).	3 At strong eutrophication state, a decrease in <i>Donax trunculus</i> density has been shown [7]	1
Material (organisms for domestic uses, industry, agriculture ...)	Driftwood and seashell for ornamental purposes	1 Driftwood and seashell for ornamental purposes	1 Driftwood and seashell for ornamental purposes	1
Molecules (organisms from which are extracted molecules potentially useful for medicine)	The polychaete <i>Arenicola marina</i> is collected and bred to get hemoglobin for medical uses. As the diversity that is harboured by sandy beaches is high, specialized and unique, there are potentially lots of useful molecules.	3 <i>Arenicola marina</i> is collected and bred to get hemoglobin for medical uses.	3 <i>Arenicola marina</i> can be impacted by green tides.	2

Table 2 (continued)

Eutrophication states	Non-eutrophic Reference species (Tellinidae, Spionidae, Amphiuroidae, Nephtyidae)		Eutrophic Green algae proliferation (Donacidae, Oweniidae, Magelonidae, Nephtyidae)	Hyper-eutrophic Green tide (Donacidae, Oweniidae)	
Coastal protection (vegetal or animal reef supplying a protection against erosion and submersion)	Specific fauna and flora (mainly bioturbating organisms) able to reduce the hydrodynamics and stabilize the substrate.	3	Ulva mats impact the hydrodynamics [23] thus affecting the sediment transport between subtidal sands, sandy shores and dunes and ultimately the function of protection	1	Ulva mats impacts the hydrodynamics [23] thus affecting the sediment transport between subtidal sands, sandy shores and dunes and ultimately the function of protection
Nutrient regulation (ecosystem capacity to supply a "good quality water", limiting the risk of eutrophication, encouraging shellfish farming...)	Beach ecosystems are important in processing large quantities of organic material and recycling nutrients back to coastal waters [24]	4	The release in excess of nutrients from lands and the presence of heterogeneous cover of green macroalgae probably affect the filtering function of the system.	3	The release in excess of nutrients from lands and the presence of heterogeneous cover of green macroalgae probably affect the filtering function of the system.
Pathogen regulation (ability of ecosystems to purify the environment through hyperfiltration processes)	Specific literature does not exist but as sandy beaches are made of porous sands that form an excellent "digestive and incubating system" [24], they most probably filter and purify the environment	4		3	
Climate regulation (GES storage or sequestration)	Autotroph systems [25] Sequestration through the phytoplanktonic and microphytobenthic activity (short term).	3	Probably short term C sequestration through the photosynthetic activity of green algae, phytoplankton and microphytobenthos	3	Probably short term C sequestration mainly through the photosynthetic activity of green algae
Support of recreational and leisure activities	Landscape, leisure activities (e.g. go to the beach), territorial activities (e.g. recreational fisheries) and emblematic biodiversity (avifauna) are important [22,24]	5	Landscape, leisure activities, territorial activities and emblematic biodiversity are impacted by green algae [22,24,26]	4	Landscape, leisure activities, territorial activities and emblematic biodiversity are impacted by green tides [22,24,26]
Contribution to a pleasant landscape		5		3	
Contribution to culture and territorial identity		5		3	
Emblematic biodiversity (i.e. protected or rared species)		5		4	
Habitat (nursery, reproduction area...)	Nursery function [22]	5	Eutrophication impacts nursery function [27]	3	Eutrophication impacts nursery function [27,28]
Trophic networks	The food web is complex, showing several potential carbon pathways and diverse trophic niches [27]	5	The trophic network is in process of homogenization (27)	3	The trophic network is homogenized/ simplified and shows less niche differentiation [27]
Recruitment	High recruitment for flatfish and other species [29]	4	Ulva mats influence local hydrodynamics, which in turn influence the recruitment of some species. In Brittany, the heterogenous cover of Ulva enhances the recruitment of <i>Donax vitattus</i> [7]	5	When the Ulva biomass is too high, macroalgae affect the recruitment, community structure and production of benthic fauna, including meiofauna, macrofauna and flatfish [7]

Table 3

Indicators and semi-quantification (SQ) of ES supply for coastal lagoons in 4 eutrophication states ("0: inexistent", "1: very low", "2: low", "3: medium", "4: high", "5: very high").

Eutrophication states	Non-eutrophic	4	Eutrophic	5	Eutrophic	4	Hyper-eutrophic	4
Human food through fisheries	Carrying capacity for shellfish farming of oligotrophic lagoons is questioned.	4	Carrying capacity for shellfish farming of slightly eutrophic lagoons is better.	5	Impact of anoxic crises on shellfish stocks (death).	4	Impact of anoxic crises on shellfish stocks (death).	4
Material (organisms for domestic uses, industry, agriculture ...)	NA	0	NA	0	NA	0	NA	0
Molecules (organisms from which are extracted molecules potentially useful for medicine)	Potential	1	Potential	1	Potential	1	Potential	1
Coastal protection (vegetal or animal reef supplying a protection against erosion and submersion)	Seagrass meadows have the capacity to attenuate waves and to slow down currents [30]	3	Decrease with the alteration and decline of seagrass meadows [30]	2	Decrease with the alteration and decline of seagrass meadows [30]	1	Decrease with the alteration and decline of seagrass meadows [30]	1
Nutrient regulation (ecosystem capacity to supply a "good quality water", limiting the risk of eutrophication, encouraging shellfish farming...)	Seagrass beds play an important role in regulating benthic nutrient fluxes in lagoons as they increase the ability to store nutrients sustainably.	5	The flow of nutrients from the sediment to the water column and, at the same time, eutrophication levels are thus greater in lagoons without seagrass [31,32]	3	Eutrophication levels are greater in lagoons without seagrass [31,32]	2	Eutrophication levels are greater in lagoons without seagrass [31,32]	1
Pathogen regulation (ability of ecosystems to purify the environment through hyperfiltration processes)	Algicidal effects of <i>Zostera marina</i> L. and <i>Zostera noltii</i> Hornem. on <i>Alexandrium catenella</i> [33] Seagrass ecosystems reduce exposure to bacterial pathogens [34] but oligotrophication can lead to emergence of toxic dinoflagellate [35]	3	Algicidal effects of <i>Zostera marina</i> L. and <i>Zostera noltii</i> Hornem. on <i>Alexandrium catenella</i> [33]	4	A decrease of seagrass leads to a stronger exposure to bacterial pathogens of humans, fishes, and invertebrates [34]	2	A decrease of seagrass leads to a stronger exposure to bacterial pathogens of humans, fishes, and invertebrates [34]	1
Climate regulation (GES storage or sequestration)	Potential long-term sequestration in the sediment through perennial macrophytes	5	A decrease of perennial macrophytes leads to reduce the sequestration in the sediment	4	A decrease of perennial macrophytes leads to reduce the sequestration in the sediment	2	A decrease of perennial macrophytes leads to reduce the sequestration in the sediment	1

Table 3 (continued)

Eutrophication states	Non-eutrophic		Eutrophic		Eutrophic		Hyper-eutrophic	
Support of recreational and leisure activities	Scuba-diving, snorkeling, sailing, wind-surf	4	Scuba-diving, snorkeling, sailing, wind-surf	4	Avifauna observation	3	Avifauna observation	3
Contribution to a pleasant landscape	Avifauna, and elements of underwater seascape [36]	5	Avifauna, and elements of underwater seascape [36]	4	Avifauna	2	Avifauna	2
Contribution to culture and territorial identity	Biodiversity of coastal lagoons are a socialization area, sometimes assimilated to an urban park [36]	4	Socialization area [36]	4	Socialization area [36]	4	Socialization area [36]	4
Emblematic biodiversity (i.e. protected or rare species)	Protected and rare species (<i>Zostera sp.</i> , <i>Hippocampus sp.</i> , avifauna)	4	Protected and rare species (<i>Zostera sp.</i> , <i>Hippocampus sp.</i> , avifauna)	4	Avifauna (flamingos) [37]	3	Avifauna (flamingos) [37]	3
Habitat (nursery, reproduction area...)	Coastal lagoons provide higher temperature during growth and food to <i>Sparus aurata</i> , which allow good lipid reserves, and large sizes of juveniles, which may be important to their survival over winter [38] <i>Zostera meadows</i> are a habitat for many species.	5	A decrease of <i>Zostera meadows</i> impacts the ES	4	A decrease of <i>Zostera meadows</i> impacts the ES	4	Coastal lagoons provide higher temperature during growth and food to <i>Sparus aurata</i> , which allow good lipid reserves, and large sizes of juveniles, which may be very important to their survival over winter [38]	5
Trophic networks	High complexity [39]	5	A decrease of <i>Zostera meadows</i> leads to less complexity	4	A decrease of <i>Zostera meadows</i> leads to less complexity	2	A decrease of <i>Zostera meadows</i> leads to less complexity	1
Recruitment	Carrying capacity for juvenile oysters of oligotrophic lagoons is questioned [40]	3	Capacity for juvenile oysters	4	Capacity for juvenile oysters	4	Capacity for juvenile oysters	4

and increasing pressure, benthic cyanobacterial mats increase and become dominant at the expense of algal turfs and macroalgae while sponges showed a more limited but significant increase. Benthic cyanobacteria mats benefit from increased levels of nutrient [5] but also from high grazing pressure and elevated water temperature [6].

First eutrophication state of sandy beaches is described by groups of species defined as reference species living in a non-eutrophic ecosystem, where no green tides occur [1]. In some French Atlantic sandy beaches, these reference species, for marine benthic macrofauna, are part of Tellinidae, Spionidae, Amphiuroidae and Nephtyidae families [7]. The continuous supply of nutrients (exogenous inputs or release from sediments) causes a slight excess and leads to the gradual development of green algae. As a response, dominant species change in this eutrophic ecosystem with an appearance of new dominant species (Donacidae, Oweniidae, Magelonidae) and the decrease, even the disappearance, of some reference species (Tellinidae, Spionidae, Amphiuroidae) [7]. Where hydrodynamic conditions are

favorable, the massive supply of nutrients leads to the massive and rapid development of green algae forming green tides. Species of reference have disappeared in favor of species (Donacidae, Oweniidae) better adapted to eutrophic conditions.

First eutrophication state of primary production of coastal lagoon is characterized by a dominance of reference species that are typical of a lagoon environment in oligotrophic conditions [1]. For French Mediterranean coastal lagoons, the reference genus are the marine phanerogams *Zostera* and *Ruppia* which form seagrass beds, and perennial benthic macroalgae (eg. *Cystoseira* sp., *Acetabularia* sp.) [8,9]. The continuous supply of nutrients causes a slight excess and leads to the gradual disappearance of the reference species and the slow and sustainable development of algae [10]. The second state is dominated by a dominance of opportunistic and epiphytic macroalgae. The massive supply of nutrients leads to the massive and rapid dominance of free-floating blooming opportunistic algae. In the most eutrophicated systems, phytoplankton community dominates the water column.

We used the classification of the Common International Classification of Ecosystem Services (CICES) and the list of marine ES defined by Liquete et al. [11] to defined the ES constituting bundles. The main distinction between these classifications concerns supporting services or ecological functions. These latter are the underpinning structures and processes that ultimately give rise to ecosystem services - sometimes defined as 'intermediate services'. They are not covered in CICES which seeks to only identify the final services that link to the goods and benefits that are valued by people (i.e. demand). Since we focus here on the ES supply, main ecological functions are considered as recommended by Liquete et al. [11].

Each step involved a literature review regarding ecosystem responses to eutrophication that was supplemented with expert-knowledge. The literature review encompassed knowledge obtained and disseminated on a global scale, while expert knowledge focused on data observed on a more local scale, based on their field studies. However, experts had a good understanding of these ecosystems which allowed them to pronounce in a qualitative way where data gaps were identified. All information were compiled within Table 1. Information was then coded to summarize the variation of ES supply between states. Five levels of ES supplied were considered: "0: inexistent", "1: very low", "2: low", "3: medium", "4: high", "5: very high".

Conflict of Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- [1] C. Kermagoret, J. Claudet, V. Derolez, M. Nugues, V. Ouisse, N. Quillien, Y. Baulaz, P. Le Mao, P. Scemama, D. Vaschalde, D. Bailly, R. Mongruel, Comparison of how eutrophication affects bundles of ecosystem services in multiple coastal habitats using state-and-transition models, *Ocean Coast, Manage* 174 (2019) 144–153. <https://doi.org/10.1016/j.ocecoaman.2019.03.028>.
- [2] D.M. de Bakker, F.C. van Duyl, R.P.M. Bak, M.M. Nugues, G. Nieuwland, E.H. Meesters, 40 Years of benthic community change on the Caribbean reefs of Curaçao and Bonaire: the rise of slimy cyanobacterial mats, *Coral Reefs* 36 (2017) 355–367. <https://doi.org/10.1007/s00338-016-1534-9>.
- [3] T.P. Hughes, K.D. Anderson, S.R. Connolly, S.F. Heron, J.T. Kerry, J.M. Lough, A.H. Baird, J.K. Baum, M.L. Berumen, T.C. Bridge, D.C. Claar, C.M. Eakin, J.P. Gilmour, N.A.J. Graham, H. Harrison, J.A. Hobbs, A.S. Hoey, M. Hoogenboom, R.J. Lowe, M.T. McCulloch, J.M. Pandolfi, M. Pratchett, V. Schoepf, G. Torda, S.K. Wilson, Spatial and temporal patterns of mass bleaching of corals in the Anthropocene, *Science* 359 (2018) 80–83. <https://doi.org/10.1126/science.aan8048>.
- [4] M.J.A. Vermeij, I. Van Moorselaar, S. Engelhard, C. Hörnlein, S.M. Vonk, P.M. Visser, The effects of nutrient enrichment and herbivore abundance on the ability of turf algae to overgrow coral in the Caribbean, *PLoS One* 5 (2010) e14312, <https://doi.org/10.1371/journal.pone.0014312>.
- [5] H.J. Brocke, L. Polerecky, D. de Beer, M. Weber, J. Claudet, M.M. Nugues, Organic matter degradation drives benthic cyanobacterial mat abundance on caribbean coral reefs, *PLoS One* 10 (2015) e0125445, <https://doi.org/10.1371/journal.pone.0125445>.
- [6] D. Bender, G. Diaz-Pulido, S. Dove, Warming and acidification promote cyanobacterial dominance in turf algal assemblages, *Mar. Ecol. Prog. Ser.* 517 (2014) 271–284. <https://doi.org/10.3354/meps11037>.
- [7] N. Quillien, M.C. Nordström, O. Gauthier, E. Bonsdorff, Y.-M. Paulet, J. Grall, Effects of macroalgal accumulations on the variability in zoobenthos of high-energy macrotidal sandy beaches, *Mar. Ecol. Prog. Ser.* 522 (2015) 97–114. <https://doi.org/10.3354/meps11151>.

- [8] C. Christia, G. Giordani, E. Papastergiadou, Environmental Variability and Macrophyte Assemblages in Coastal Lagoon Types of Western Greece (Mediterranean Sea), *Water* 10, 2018. <https://doi.org/10.3390/w10020151>.
- [9] I. Le Fur, R. De Wit, M. Plus, J. Oheix, M. Simier, V. Ouisse, Submerged benthic macrophytes in Mediterranean lagoons: distribution patterns in relation to water chemistry and depth, *Hydrobiologia* 808 (2018) 175–200. <https://doi.org/10.1007/s10750-017-3421-y>.
- [10] W. Schramm, Factors influencing seaweed responses to eutrophication : some results from EU-project EUMAC, *J. Appl. Phycol.* 11 (1999) 69–78. <https://doi.org/10.1023/A:1008076026792>.
- [11] C. Liqueste, C. Piroddi, E.G. Drakou, L. Gurney, S. Katsanevakis, A. Charef, B. Egoth, Current Status, Future, Prospects for the assessment of marine and coastal ecosystem services: a systematic review., *PLoS One* 8 (2013) e67737, <https://doi.org/10.1371/journal.pone.0067737>.
- [12] T.P. Hughes, M.L. Barnes, D.R. Bellwood, J.E. Cinner, G.S. Cumming, J.B.C. Jackson, J. Kleypas, I.A. Van De Leemput, J.M. Lough, T.H. Morrison, S.R. Palumbi, E.H. Van Nes, M. Scheffer, Coral reefs in the anthropocene, *Nature* 546 (2017) 82–90. <https://doi.org/10.1038/nature22901>.
- [13] C.C. Hicks, L.B. Crowder, N.A.J. Graham, J.N. Kittinger, E. Le Cornu, Social drivers forewarn of marine regime shifts, *Front. Ecol. Environ.* 14 (2016) 252–260. <https://doi.org/10.1002/fee.1284>.
- [14] T.-L. Loh, J.R. Pawlik, Chemical defenses and resource trade-offs structure sponge communities on Caribbean coral reefs, *Proc. Natl. Acad. Sci. Unit. States Am.* 111 (2014) 4151–4156. <https://doi.org/10.1073/pnas.1321626111>.
- [15] F. Ferrario, M.W. Beck, C.D. Storlazzi, F. Micheli, C.C. Shepard, L. Airoidi, The effectiveness of coral reefs for coastal hazard risk reduction and adaptation, *Nat. Commun. Now.* 5 (2014) 1–9. <https://doi.org/10.1038/ncomms4794>.
- [16] R.L. Vega Thurber, D.E. Burkpile, C. Fuchs, A.A. Shantz, R. McMinds, J.R. Zaneveld, Chronic nutrient enrichment increases prevalence and severity of coral disease and bleaching, *Glob. Chang. Biol.* 20 (2013) 544–554. <https://doi.org/10.1111/gcb.12450>.
- [17] J.R.R. Zaneveld, D. Burkpile, A.A. Shantz, C.E. Pritchard, R. Mc Minds, J.P. Payet, R. Welsh, A.M.S. Correa, N.P. Lemoine, S.M. Rosales, C. Fuchs, J.A. Maynard, R. Vega Thurber, Overfishing and nutrient pollution interact with temperature to disrupt coral reefs down to microbial scales, *Nat. Commun.* 7 (2016) 11833. <https://doi.org/10.1038/ncomms11833>.
- [18] C.M. Roberts, B.C. O'Leary, D.J. McCauley, P.M. Cury, C.M. Duarte, J. Lubchenco, D. Pauly, A. Sáenz-Arroyo, U.R. Sumaila, R.W. Wilson, B. Worm, J.C. Castilla, Marine reserves and climate change, *PNA* 114 (2017) 6167–6175. <https://doi.org/10.1073/pnas.1701262114>.
- [19] E. Sterling, T. Ticktin, T. Kipa Kipa Morgan, G. Cullman, D. Alvira, P. Andrade, N. Bergamini, E. Betley, K. Burrows, S. Caillon, J. Claudet, R. Dacks, P. Eyzaguirre, C. Filardi, N. Gazit, C. Giardina, S. Jupiter, K. Kinney, J. McCarter, M. Mejia, K. Morishige, J. Newell, L. Noori, J. Parks, P. Pascua, A. Ravikumar, J. Tanguay, A. Sigouin, T. Stege, M. Stege, A. Wali, Culturally grounded indicators of resilience in social-ecological systems, environment and society, *Advances in Research* 8 (2017) 63–95. <https://doi.org/10.3167/ares.2017.080104>.
- [20] P.J. Mumby, R.S. Steneck, Coral reef management and conservation in light of rapidly evolving ecological paradigms, *Trends Ecol. Evol.* 10 (2008) 555–563. <https://doi.org/10.1016/j.tree.2008.06.011>.
- [21] T.N. Hempson, N.A.J. Graham, M.A. MacNeil, D.H. Williamson, G.P. Jones, G.R. Almany, Coral reef mesopredators switch prey, shortening food chains, in response to habitat degradation, *Ecol. Evol.* 7 (2017) 2626–2635. <https://doi.org/10.1002/ece3.2805>.
- [22] A. McLachlan, A. Brown, *The Ecology of Sandy Shores*, 2nd Ed, Elsevier, Ed., 2006.
- [23] S.C. Hull, Macroalgal mats and species abundance: a field experiment, *Estuar. Coast Shelf Sci.* 25 (1987) 519–532.
- [24] T.A. Schlacher, L.M.C. Thompson, Physical impacts caused by off-road vehicles to sandy beaches: spatial quantification of car tracks on an Australian barrier island, *J. Coast. Res.* 24 (2008) 234–242. <https://doi.org/10.2112/06-0691.1>.
- [25] A. Migné, D. Davout, N. Spilmont, V. Ouisse, G. Boucher, Spatial and temporal variability of CO2 fluxes at the sediment-air interface in a tidal flat of a temperate lagoon (Arcachon Bay, France), *J. Sea Res.* 109 (2016) 13–19. <https://doi.org/10.1016/j.seares.2016.01.003>.
- [26] A. Levain, Faire face aux « marées vertes », penser les crises du vivant, *ethnographiques.org*, Numéro 27 (2013).
- [27] N. Quillien, Dynamic ecosystems under anthropogenic stress : how does macrotidal sandy beach fauna respond to green tides ? PhD thesis. *Earth Sciences, Université de Bretagne occidentale – Brest* (2016).
- [28] E. Le Luherne, O. Le Pape, L. Murillo, M. Randon, C. Lebot, E. Reveillac, Influence of green tides in coastal nursery grounds on the habitat selection and individual performance of juvenile fish, *PLoS One* 12 (2017) e0170110, <https://doi.org/10.1371/journal.pone.0170110>.
- [29] B. Beyst, A. Cattrijsse, M. Mees, Feeding ecology of juvenile flatfishes of the surf zone of a sandy beach, *J. Fish Biol.* 55 (1999) 1171–1186.
- [30] A.E. Paquier, S. Meulé, E.J. Anthony, G. Bernard, Sedimentation and erosion patterns in a low shoot-density *Zostera noltii* meadow in the fetch-limited Berre lagoon, Mediterranean France, *J. Coast. Res.* 70 (2014) 563–567. <https://doi.org/10.2112/SI70-095.1>.
- [31] P. Viaroli, M. Bartoli, G. Giordani, M. Naldi, Community shifts, alternative stable states, biogeochemical controls and feedbacks in eutrophic coastal lagoons : a brief overview, *Aquat. Conserv.* 18 (2008) 105–117. <https://doi.org/10.1002/aqc.956>.
- [32] V. Ouisse, A. Fiandrino, R. De Wit, N. Malet, Restauration des écosystèmes lagunaires : évaluation du rôle du sédiment et des herbiers à phanérogames, *RST/LERLR*, 2013, 13–09.
- [33] M. Laabir, M. Grignon-Dubois, E. Masseret, B. Rezzonico, G. Soteras, M. Rouquette, F. Rieuvilleneuve, P. Cecchi, Algicidal effects of *Zostera marina* L. and *Zostera noltii* Hornem. extracts on the neuro-toxic bloom-forming dinoflagellate *Alexandrium catenella*, *Aquat. Bot.* 111 (2013) 16–25. <https://doi.org/10.1016/j.aquabot.2013.07.010>.
- [34] J.B. Lamb, J.A.J.M. van de Water, D.G. Bourne, C. Altier, M.Y. Hein, E.A. Fiorenza, N. Abu, J. Jompa, C.D. Harvell, Seagrass ecosystems reduce exposure to bacterial pathogens of humans, fishes and invertebrates, *Science* 355 (2017) 731–733. <https://doi.org/10.1126/science.aal1956>.
- [35] Y. Collos, B. Bec, C. Jauzein, E. Abadie, T. Laugier, Oligotrophication and emergence of picocyanobacteria and a toxic dinoflagellate in Thau lagoon, southern France Regional index term : Mediterranean, *J. Sea Res.* 61 (2009) 68–75. <https://doi.org/10.1016/j.seares.2008.05.008>.

- [36] L. Chazée, M. Réquier Desjardins, W. Khechimi, F.Z. Najjar, D. Moisan, M. Suc, N. Bendjedda, W. Benbelgacem, F. Allouche Khebour, A. Belarbi, F. Khaloul, O. Brahimi, S. Maza, L. Moussouni, M. Amara, Y. Madjbar, N. Ait Iftene, L. Khellouf, S. Bakour, Les services culturels récréatifs et éducatifs des zones humides en Méditerranée – des services sous-estimés malgré les avantages qu'ils procurent: résultats d'études en Méditerranée, Observatoire des zones humides méditerranéennes, 2017, 50pp.
- [37] R. De Wit, H. Rey-Valette, J. Balavoine, V. Ouisse, R. Lifran, Restoration ecology of coastal lagoons: new methods for the prediction of ecological trajectories and economic valuation, *Aquat. Conserv.* 27 (2015) 137–157. <https://doi.org/10.1002/aqc.2601>.
- [38] J. Tournois, F. Ferraton, L. Velez, D.J. McKenzie, C. Aliaume, L. Mercier, A.M. Darnaude, Temporal stability of otolith elemental fingerprints dis-criminates among lagoon nursery habitats, *Estuar. Coast Shelf Sci.* 131 (2013) 182–193. <https://doi.org/10.1016/j.ecss.2013.07.006>.
- [39] T.H. Pearson, R. Rosenberg, Macrobenthic succession in relation to organic enrichment and pollution of the marine environment, *Oceanogr. Mar. Biol.* 16 (1978) 229–311.
- [40] F. Lagarde, E. Roque D'orbcastel, M. Ubertini, S. Mortreux, I. Bernard, A. Fiandrino, C. Chiantella, B. Bec, C. Roques, D. Bonnet, G. Miron, M. Richard, S. Pouvreau, C. Lett, Recruitment of the Pacific oyster *Crassostrea gigas* in a shellfish-exploited Mediterranean lagoon: discovery, driving factors and a favorable environmental window, *Mar. Ecol.: Prog. Ser.* 578 (2017) 1–17. <https://doi.org/10.3354/meps12265>.