



Photosynthetic efficiency of young *Rhizophora mangle* L. in a mangrove in southeastern Brazil

S.S. PASCOALINI^{*,+} , M.M.P. TOGNELLA^{**,*} , A.R. FALQUETO^{**,*} , and M.L.G. SOARES^{***} 

*Department of Oceanography and Ecology, Federal University of Espírito Santo, Vitória, ES, Brazil**

*Department of Agricultural and Biological Sciences, Federal University of Espírito Santo, São Mateus, ES, Brazil***

*Department of Biological Oceanography, Rio de Janeiro State University, Rio de Janeiro, RJ, Brazil****

Abstract

The study proposes to evaluate the photosynthetic plasticity of *Rhizophora mangle* L. in four mangrove sites distributed along with the Great Vitória Estuarine System. The variation in organic matter content, which implies the higher essential nutrient availability, contributed to better energy flux performance related to electron transport. Furthermore, salinity damaged the reaction centers (RC), since the site with the highest salinity showed changes in the number and size of active photosynthetic RC and in the specific energy flows per active RC (absorption flux, trapped energy flux, and dissipated energy flux), but the plasticity of the species in response to salt stress was confirmed by the increase of performance index for energy conservation (PI_{Total}), net photosynthetic rate (P_N), and the water-use efficiency (WUE). Also, the results showed that the luminous intensity available compromises the functionality of PSII, in turn, it increases WUE. The results indicate the effect of the chlorophyll *a* content, which provides more substrate for light absorption, on the electron flow and PI_{Total} is related to P_N and WUE. The study indicates the ecological plasticity of *R. mangle* to the conditions of the evaluated area.

Keywords: gas exchange; JIP test; photosystems; tropical region.

Introduction

Mangrove is a halophytic forest ecosystem growing in intertidal zones along tropical and subtropical coasts

(Alongi 2009). Mangrove ecosystems are threatened all over the world by human pressures, leading to a demographic increase in the coastal zone, which, together with urbanization and industrialization, change the water

Highlights

- Energy flux corresponds to electron transport associated with organic matter
- Adjustment of photosynthetic parameters occurs at high salinity
- Photochemical performance is related to net photosynthesis and conservative water use

Received 1 November 2021

Accepted 26 April 2022

Published online 17 May 2022

*Corresponding author

e-mail: savia.pascoalini@gmail.com

Abbreviations: ABS/RC – absorption flux per reaction center; Chl – chlorophyll; C_i – intercellular CO_2 concentration; DI_0/RC – dissipated energy flux per RC; E – transpiration rate; ET_0/RC – electron transport flux per RC; F_0 – initial fluorescence; F_m – maximum fluorescence; F_v/F_0 – number and size of active photosynthetic RC; g_s – stomatal conductance; GVES – Great Vitória Estuarine System; OM – organic matter; PI_{abs} – performance index for energy conservation from exciton to the reduction of intersystem electron acceptors; PI_{Total} – performance index for energy conservation from exciton to the reduction of PSI end acceptors; P_N – net photosynthetic rate; Q_{leaf} – luminous intensity incident on the leaf surface; TR_0/RC – trapped energy flux per RC; WUE – water-use efficiency; δR_0 – efficiency with which an electron from the intersystem reduces the final acceptor of PSI; ϕD_0 – maximum efficiency of nonphotochemical de-excitation; ϕE_0 – quantum yield for electron transport; ϕP_0 – maximal quantum yield of PSII photochemistry; Ψ_0 – the probability of electron transport further than Q_A .

Acknowledgements: This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brazil (CAPES) – Finance Code 001. The second author was supported by FAPES research grants no. 60127627/2012 and no. 263/2016.

Conflict of interest: The authors declare that they have no conflict of interest.

regime and sediment/nutrient flow into the environment. Moreover, in some locations, these stressors may act synergistically with changes in the relative mean sea level, and, ultimately, climate change (Alongi 2015) may compromise ecosystem services. Understanding the *in situ* ecophysiology of mangrove plant species is important in determining how the ecosystem will behave in the face of environmental change and its long-term prospect of maintenance (Schaeffer-Novelli *et al.* 1990, Pascoalini *et al.* 2014), and contributing to coastal management.

Pioneering studies on the ecophysiology of mangroves focused on understanding plant behavior at different salinity levels (Ball *et al.* 1984, Ball 1986, 1988; Naidoo *et al.* 2002, Parida *et al.* 2004, Barr *et al.* 2009), because the salinity is considered the main abiotic factor determining the development of these forests (Parida and Jha 2010). Field and laboratory studies were carried out on various plant species, including *Avicennia marina* (Forssk.) Vierh., *Avicennia germinans* (L.) L., *Bruguiera gymnorhiza* (L.) Lam., *Kandelia candel* (L.) Druce, *Laguncularia racemosa* (L.) C. F. Gaertn., *Rhizophora mangle* L., and *Rhizophora mucronata* Lam. (Ball *et al.* 1984, Sobrado 1999, 2000; Naidoo *et al.* 2002, Krauss *et al.* 2006, Barr *et al.* 2009, Ru *et al.* 2009, Hoppe-Speer *et al.* 2011, Bompny *et al.* 2014). The relationship with salt differentiates mangrove species into salt-secreter, salt-accumulator, and salt-excluder (Parida and Jha 2010). However, more recent approaches have assessed species tolerance as a function of daily salinity variation (Bompny *et al.* 2014), contributing to an understanding of these species' ecological niches.

In addition to the influence of salinity on plant development, several studies address other variables that can influence biomass acquisition. Nonetheless, in general, these studies conducted in the laboratory or the field address extremes such as hypersaline and/or nutrient deficient mangrove areas (Lovelock *et al.* 2006a,b; Martin *et al.* 2010). Therefore, studies approaching the *in situ* behavior of species subjected to a variety of environmental conditions independently of stress occurrence are necessary.

R. mangle, a mangrove tree widely distributed on the American Atlantic coast, is among the species that have been the subject of ecophysiological studies. In general, *R. mangle* colonizes the low and middle intertidal zone, with a high frequency of flooding, and muddy substrates subjected to low salinity (Alongi 2009, Barr *et al.* 2009, Bompny *et al.* 2014, Lima *et al.* 2018, Lopes *et al.* 2019). *R. mangle* is considered a salt-excluder species, preventing a considerable amount of sodium and chlorine from being absorbed (Parida and Jha 2010). This is a strategy of less salt-tolerant plants. Moreover, the species exhibits the least conservative water- and nutrient-use efficiency compared to other neotropical mangroves (McKee 1995, Parida and Jha 2010, Soares *et al.* 2015, Tognella *et al.* 2016).

Chlorophyll (Chl) *a* fluorescence is a technique used to evaluate the plant photochemical performance providing us with information about the structure and functioning of PSII, the reaction center (RC), and the electron transport between PSII and PSI. By inducing fluorescence in dark-

adapted leaves, Strasser *et al.* (2000) derived several equations for the photochemical apparatus that are translated through the JIP test. Although the technique has been applied to several studies related to water and salt stress, its use in mangal areas is still restricted, but it can provide relevant information on the ecophysiological dynamics of mangrove species under field conditions (Naidoo *et al.* 2002, Gonzalez-Mendoza *et al.* 2007, Falqueto *et al.* 2008, 2012; Pascoalini 2014, Lopes *et al.* 2019, Tognella *et al.* 2022). Furthermore, gas-exchange measurements of mangroves have been widespread and very useful because they provide information on photosynthesis and water-use efficiency quickly and reliably in field conditions (Medina 1999, Soares *et al.* 2015, Larcher *et al.* 2016). Assessing the photochemical process through Chl *a* fluorescence measurements and photosynthesis through gas exchange allow the understanding of sink effects on photosynthesis, from leaf light capture to release of chemical energy. Falqueto *et al.* (2008, 2012) and Pascoalini (2014) conducted studies in the mangrove swamp of the Great Vitória Estuarine System (GVES) and found increased photochemical efficiency with salinity, however lower carbon assimilation for *R. mangle* (Pascoalini 2014).

Mangrove photosynthesis is mainly controlled by salinity. But, the availability of nutrients and light, substrate oxygenation, and sediment contamination also affect the photosynthetic process to a greater or a lesser extent, depending on the limitation imposed. This paper proposes to identify and better understand the photosynthetic plasticity of *R. mangle* in field conditions and some of the factors controlling it, including salinity, organic matter – which indicates nutrient availability, luminosity – related to canopy variation, chlorophyll content, and photochemical performance. The study is guided by the hypothesis that biotic and abiotic factors act directly or indirectly on parameters of the photosynthetic process, reflecting on the photosynthetic efficiency of *R. mangle* growing in the GVES.

Materials and methods

Study area: The study was conducted in the Great Vitória Estuarine System (Fig. 1), on the central Brazilian east coast (20°10'44.0"S and 20°16'31.1"S–40°15'11.0"W and 40°20'44.0"W). The region's climate is Am (Köppen's climate classification), *i.e.*, tropical with seasonal precipitation, the annual mean rainfall is approximately 1,350 mm (Alvares *et al.* 2013). The wettest and driest months extend from October to April, and from May to September, respectively (Alvares *et al.* 2013, INCAPER 2018). Rebello *et al.* (2011) indicate that the area experiences water scarcity almost every month of the year and a water surplus above 100 mm in November and December. The estuary presents dynamics dominated by the tides, mainly by the ebb tide, and they are classified as microtides with semidiurnal frequency, which is less than 2 m in amplitude (Rigo and Chacaltana 2006, Neves *et al.* 2012). The Santa Maria da Vitória River is the main freshwater source of the estuary (Teubner Júnior *et al.* 2018).




Fig. 1. Study sites in the Great Vitória Estuarine System (GVES), Espírito Santo, Brazil, which are located in the anticlockwise direction from the north entrance (S1) to the south entrance (S4). F – fringe; B – basin.

The GVES mangrove has an extension of 24.8 km² (Teubner Júnior *et al.* 2018) and is inserted in a wide protection area together with its buffer areas that make up the Vitória Bay Mangrove Protected Areas Mosaic established by state Decree no. 2625-R of 23 November 2010 (Espírito Santo 2010). The mangrove structural diversity is associated with the frequency of flooding, salinity, sediment characteristics, and anthropogenic pressures (Lima 2011, Zamprogno *et al.* 2016, Pascoalini *et al.* 2019).

Study sites and plant material: All parameters were measured at four sampling sites (S1, S2, S3, and S4) in the GVES, according to their location in the bay, *i.e.*, in the anticlockwise direction from the north entrance to the south entrance. Each site was divided by physiographic type as fringe (F) and basin (B) (Schaeffer-Novelli *et al.* 2000), totaling eight sampling points (S1F, S1B, S2F, S2B, S3F, S3B, S4F, and S4B). At each site, five completely expanded leaves (the second pair from the apex) were sampled from twelve individuals. Sample collections were carried out at low tide in spring between July and October 2016 and March and June 2017, starting at 08:00 h. The samples characterize the dry period, as the region went through a period of precipitation below expectations for two years (INCAPER 2018).

Abiotic variables: Organic matter (OM) analysis and the interstitial water salinity measurements were performed concurrently with the measurements of photosynthetic parameters and pigment indices. The salinity of interstitial

water was determined using a 5-cm diameter and 0.5 m in length PVC tube inserted at least –0.3-m sediment depth according to the method of Zamprogno *et al.* (2016) using a *Hach Multiparameter Meter (model HQ40d)* previously calibrated with a standard solution.

For OM analysis, samples of surface sediment (first 2 cm) were collected after the removal of the deposited macroscopic material (standing crop). The samples were kept at –20°C until the lyophilization procedure. The content of lyophilized organic matter was determined using dry mass after samples were ignited and ashed in a muffle furnace for 4 h at 550°C (Mook and Hoskin 1982). The samples were treated individually, consisting of three samples per site and year.

Chl index was measured using a portable Chl meter *ClorofiLOG (Falker, model CFL 1030)*; this equipment provides reliable and dimensionless information and its results are strongly related to those obtained by the classical method (Barbieri Júnior *et al.* 2012). The parameters obtained through the *ClorofiLOG* are Chl index *a* (Chl_a), *b* (Chl_b), and total (Chl_{total}). These variables were collected only in 2016.

Chl *a* fluorescence was measured using a *Plant Efficiency Analyzer (Handy-PEA, Hansatech, King's Lynn, Norfolk, UK)* with a high-resolution time (10 μs). Leaves were dark-adapted for 30 min using leaf clips (Falqueto *et al.* 2008, 2012) for oxidation of all reaction centers (RC) (Strasser *et al.* 2000). Light intensity reaching the leaf was 3,000 μmol(photon) m⁻² s⁻¹ (650 nm) which was sufficient

to generate maximal fluorescence for all the samples. The fast fluorescence kinetics (F_0 to F_m) was recorded from 10 μ s to 1 s. The initial fluorescence intensity at 50 μ s (F_0), 100 μ s, 300 μ s, 2 ms (F_1), 30 ms (F_1), and maximum fluorescence (F_m) were recorded and used to obtain the parameters of JIP test (Strasser *et al.* 2000, Yusuf *et al.* 2010). The selected parameters were: Area, F_0 , F_m , F_v/F_0 (number and size of active photosynthetic RC), specific energy flows per active RC, *i.e.*, ABS/RC (absorption flux), TR_0/RC (trapped energy flux), ET_0/RC (electron transport flux), and DI_0/RC (dissipated energy flux); ϕP_0 (maximal quantum yield of PSII photochemistry), Ψ_0 (probability of electron transport further than quinone A – Q_A^-), ϕE_0 (quantum yield for electron transport), δR_0 (efficiency with which an electron from the intersystem reduces the final acceptor of PSI), ϕD_0 (maximum efficiency of nonphotochemical de-excitation), PI_{abs} (performance index for energy conservation from exciton to the reduction of intersystem electron acceptors), PI_{Total} (performance index for energy conservation from exciton to the reduction of PSI end acceptors).

Gas exchange: Net photosynthetic rate (P_N [μ mol $m^{-2} s^{-1}$]), stomatal conductance (g_s [$mol(H_2O) m^{-2} s^{-1}$]), intercellular CO_2 concentration (C_i [μ mol(CO_2) mol^{-1}]), transpiration rate (E [$mmol(H_2O) m^{-2} s^{-1}$]), and luminous intensity incident on the leaf surface (Q_{leaf} [μ mol(photon) $m^{-2} s^{-1}$]) were measured using infrared gas analyzer *LCi* (*ADC, BioScientific Ltd.*, Hoddesdon, UK). The gas chamber was maintained in ambient conditions, the average photon flux density in the chamber was 257 μ mol $m^{-2} s^{-1}$, with an average CO_2 concentration of 372 μ mol(CO_2) mol^{-1} , and leaf temperature reaching 32°C on average. P_N and g_s were used to calculate and determine water-use efficiency ($WUE = P_N/g_s$ [μ mol(CO_2) $mol(H_2O)^{-1}$]) (Sobrado 2005).

Data analysis: Because the abiotic and physiological data were not normally distributed, the nonparametric *Kruskal–Wallis* test was used for comparisons between study sites and followed by multiple posterior comparisons. The *Mann–Whitney's* test was used in the comparison between the physiographic types (fringe and basin) (Zar 1996). The relationship between the variables was assessed by the correlation analysis, for this, the mean of the variables in each site was obtained, followed by a normality test (*Shapiro–Wilk*), taking into account this assumption, a *Pearson's* correlation analysis was conducted. The PI_{Total} variable did not meet normality, being normalized through the natural logarithm and, consequently, its pair in the correlation was also transformed. Correlations were calculated between the abiotic variables (Q_{leaf} , interstitial water salinity, OM) and biotic variables, and also between biotic variables. The data were presented as the mean \pm standard error (SE). The significance level was $p < 0.05$.

Results

Abiotic variables: The salinity was higher in the fringe forest in S4 (Table 1). The highest content of organic matter

Table 1. Mean (\pm standard error) of sediment variables including the parameters interstitial salinity and organic matter referring to sampling sites (S1 to S4) and physiographic types (F – fringe; B – basin) in the Great Vitória Estuarine System. *Capital letters* indicate difference between sites and *small letters* indicate difference between physiographic types ($p < 0.05$).

Sites	Interstitial salinity	Organic matter [%]
S1F	27.85 \pm 0.94 ^B	13.14 \pm 0.99 ^B
S1B	32.28 \pm 0.84 ^B	11.90 \pm 2.06 ^B
S2F	32.00 \pm 0.99 ^B	34.02 \pm 0.86 ^A
S2B	27.92 \pm 1.59 ^B	49.55 \pm 0.82 ^A
S3F	32.42 \pm 1.32 ^B	32.57 \pm 0.67 ^A
S3B	30.93 \pm 0.60 ^B	31.58 \pm 1.97 ^A
S4F	37.88 \pm 0.22 ^A	34.29 \pm 1.06 ^A
S4B	35.38 \pm 1.40 ^A	34.91 \pm 2.42 ^A

in the sediment occurred in S2 (basin forest). Statistically, differences in salinity occurred between the sites (Table 1), confirming the highest mean observed for S4, which is located near the south entrance of the system. OM percentage varied among the sites, with the lowest values observed in S1 (north entrance). No statistical difference was obtained when the physiographic types (fringe and basin) were compared (Table 1).

Chl index: The Chl_a and Chl_b index differed between the sites (Fig. 2A,B). The Chl_a index was the highest in S4 and the Chl_b index was in S1 and S2. No statistical difference was recorded for the Chl_{total} index between the sites. Plants of the basin forest had higher values of Chl_a and Chl_{total} (Fig. 2C).

Chl a fluorescence – JIP test: The results of the JIP test are shown in Fig. 3. The area above the curve decreased in S1, S3, and S4 (Fig. 3A). Sites S1, S3, and S4 exhibited the highest F_0 values. The F_m and F_v/F_0 were highest at S1. For the physiographic types, the highest F_0 value was recorded in the fringe forest and F_v/F_0 was the highest in the basin forest (Fig. 3B,D). The values of the specific energy fluxes expressed per RC, ABS/RC, TR_0/RC , ET_0/RC , and DI_0/RC were the highest in S3 and S4 (Fig. 4). S1 also had a higher mean for TR_0/RC , along with S3 and S4 (Fig. 4B). The highest values of ABS/RC, TR_0/RC , and DI_0/RC were recorded for fringe forests compared to basin forests (Fig. 4).

The maximum quantum yield of PSII photochemistry (ϕP_0) was the highest in S1 and S2 and higher Ψ_0 and ϕE_0 values were obtained in S2, S3, and S4 (Fig. 5A–C). The efficiency with which an electron from the intersystem reduces the final acceptor of PSI (δR_0) was the greatest in both S2 and S4 followed by S3 (Fig. 5D). The maximum efficiency of nonphotochemical de-excitation (ϕD_0) values was the highest in S3 and S4 and the lowest in S1 (Fig. 5E). The highest PI_{abs} and PI_{Total} values occurred in S2 and S4 (Fig. 5F,G). The comparison between the basin and fringe




Fig. 2. Mean (\pm standard error) of chlorophyll (Chl) index collected in the sampling sites (S1 to S4) and physiographic types (F – fringe; B – basin) in the Great Vitória Estuarine System. (A) Chl_a index, (B) Chl_b index, (C) Chl_{total} index. Capital letters indicate difference between sites and small letters indicate difference between physiographic types ($p < 0.05$).

forests indicates that the plants of the basin forest had higher values of ϕP_0 , Ψ_0 , ϕE_0 , PI_{abs} , and PI_{Total} . In contrast, the fringe forest showed a higher value of ϕD_0 (Fig. 5).

Gas exchange: The net photosynthetic rate (P_N) was significantly the highest in S2 and S4, followed by S3, with the basin forest exhibiting a greater value than that of the fringe (Fig. 6A). The highest values of C_i were reported in S1 and it was statistically similar to those values obtained in S2 and S3 (Fig. 6B). The highest g_s was recorded in S2. No statistical difference was recorded for E . Concerning the water economy, the WUE was statistically similar among S2, S3, and S4 (Fig. 6E). Statistical analysis for Q_{leaf} is displayed in Fig. 1S (supplement). Difference between sites was recorded, with S1 having the lowest value.

Linear correlation analysis: Table 2 shows a significant linear correlation between abiotic (interstitial salinity, OM, and Q_{leaf}) and biotic variables, and also between biotic variables. The correlation established between OM and Chl_a and the parameters related to the electron flow stands out; Q_{leaf} is intrinsically correlated with several steps in the photosynthetic process. PI_{Total} seems to act on P_N and, consequently, on WUE.

Discussion

Understanding the ecophysiology of plant species is important in the context of climate change as the rise in temperature and relative mean sea level will induce expansion/shrinkage of biomes and ecosystems (Wong *et al.* 2014). In addition, they can also lead to loss of habitat quality through the introduction/extinction of populations and communities. The species distribution responses in mangrove systems are closely related to environmental variables, such as flood frequency, salinity, and nutrient distribution (Cunha *et al.* 2006, Krauss *et al.* 2006, Lovelock *et al.* 2006b, Medina *et al.* 2010, Tognella *et al.* 2016). Yet, very few studies have examined the interactions between biochemical processes at energy flux level, photochemical efficiency, carbon, and nutrient assimilation that influence the incorporation of biomass by the individual.

Chl *a* fluorescence assessment determines the light-use efficiency in plants under ambient light conditions. The variable Area is proportional to the size of the electron acceptor pool on the reducing side of the PSII. If the electron transfer from the reaction center to the plastoquinone pool is blocked, the Area will be drastically reduced (Mehta *et al.* 2010). In this study, we found the lowest values for the Area of S3, which was followed by S1 and S4. Reductions in Area have already been reported in plants growing under high irradiance and salinity (Gonçalves and Santos Júnior 2005, 2007; Mehta *et al.* 2010). In this study, note that S4 had the highest salinity level, S1 had the lowest OM content, and, consequently, lower nutrient availability (Bai *et al.* 2005, Khadka 2016), and S3 had the highest mean of Q_{leaf} , indicating that the parameters interfere with light-use efficiency. Furthermore, S1 and S3, followed by S4, showed the highest F_0 values. Increased F_0 values may result from the impaired energy transfer between the light-harvesting complex and the reaction center, induced by low energy connectivity and damage to PSII reaction centers (Yamane *et al.* 1997, Tóth *et al.* 2007). Thus, increased F_0 may indicate some environmental stress. Falqueto *et al.* (2008, 2012) observed that salinity-tolerant species, *e.g.*, *R. mangle*, may show decreased F_0 values with an increase in salinity, but this pattern was not observed here. Reduction in maximal fluorescence in S2, S3, and S4 may indicate inhibition of PSII donor side electron transport resulting in P_{680}^* accumulation and/or decreased Q_A pool (Mehta *et al.* 2010).

The ABS/RC parameter is calculated as the total number of photons absorbed by chlorophyll molecules from all RCs divided by the total number of active RCs. Thus, ABS/RC values are affected by the ratio of active to




Fig. 3. Mean (\pm standard error) of Area, initial fluorescence (F_0), maximum fluorescence (F_m), number and size of active photosynthetic reaction centers (F_v/F_0) collected in the sampling sites (S1 to S4) and physiographic types (F – fringe; B – basin) in the Great Vitória Estuarine System. *Capital letters* indicate difference between sites and *small letters* indicate difference between physiographic types ($p < 0.05$).




Fig. 4. Mean (\pm standard error) of specific fluxes (ABS/RC – absorption; TR_0/RC – trapping, ET_0/RC – transport; and DI_0/RC – dissipation) per reaction center (RC) collected in the sampling sites (S1 to S4) and physiographic types (F – fringe; B – basin) in the Great Vitória Estuarine System. *Capital letters* indicate difference between sites and *small letters* indicate difference between physiographic types ($p < 0.05$).




Fig. 5. Mean (\pm standard error) of JIP test parameters collected in the sampling sites (S1 to S4) and physiographic types (F – fringe; B – basin) in the Great Vitória Estuarine System. (A) Maximum quantum yield of PSII photochemistry (ϕP_0), (B) probability of electron transport further than quinone A – Q_A^- (Ψ_0), (C) quantum yield for electron transport (ϕE_0), (D) efficiency with which an electron from the intersystem reduces the final acceptor of PSI (δR_0), (E) maximum efficiency of nonphotochemical de-excitation (ϕD_0), (F) performance index for energy conservation from exciton to the reduction of intersystem electron acceptors (PI_{abs}), (G) performance index for energy conservation from exciton to the reduction of PSI end acceptors (PI_{Total}). Capital letters indicate difference between sites and small letters indicate difference between physiographic types ($p < 0.05$).

inactive RCs (Mehta *et al.* 2010). High values of ABS/RC are interpreted in two ways: (1) increase in antenna size, which is responsible for photon absorption and transfer of the excitation energy to the reaction center; (2) reduction in stress-active reaction centers (Strasser and Stirbet

1998, Yusuf *et al.* 2010). In the present study, increases in ABS/RC values observed in S3 and S4 indicate a reduction in active reaction centers, considering the highest DI_0/RC values and the lowest F_v/F_0 values recorded in both sites. DI_0/RC indicates energy loss as heat or fluorescence or




Fig. 6. Mean (\pm standard error) parameters of gas exchange collected in the sampling sites (S1 to S4) and physiographic types (F – fringe; B – basin) in the Great Vitória Estuarine System. (A) Net photosynthetic rate (P_N), (B) intercellular CO_2 concentration (C_i), (C) stomatal conductance (g_s), (D) transpiration rate (E), and (E) water-use efficiency (WUE). Capital letters indicate difference between sites and *small letters* indicate difference between physiographic types ($p < 0.05$).

energy transfer to other systems such as nitrate reduction or generation of reactive oxygen species (Strasser *et al.* 2000). In contrast, reduction in F_v/F_0 expresses the inactivation of RC. So, these three parameters (DI_0/RC , F_v/F_0 , and ABS/RC) may indicate susceptibility to high irradiance in S3 and S4. Similar results have also been associated with water stress, as recorded by Mehta *et al.* (2010) and Falqueto *et al.* (2017). However, we observed an apparent recovery of the energy flux due to the increases in energy flux corresponding to electron transport (ET_0/RC , Ψ_0 , and ϕE_0).

The maximal quantum yield of PSII ($\phi P_0 = F_v/F_m$) is used to evaluate plant stress. In this study, ϕP_0 was the highest at S1. Moreover, S1 exhibited the lowest Ψ_0 , indicating less efficient production of reducing power (NADPH), as well as reduction of the oxidized plastoquinone pool, and inhibition of reoxidation of the Q_A (ϕE_0). The lowest ET_0/RC , Ψ_0 , ϕE_0 , and δR_0 values obtained in S1 suggest damage to energy flux corresponding to electrons transport (Gonçalves and Santos Júnior 2005). These parameters can be considered indicators of the nutritional status of plants (Swoczyna *et al.* 2019, Meng *et al.* 2021), since the reduction of nutrients directly influences the photosynthetic apparatus, mainly through the biosynthesis

and functioning of the main photosynthetic components, with a direct effect on the synthesis of proteins involved in photosynthetic reactions and synthesis of chlorophyll (Kalaji *et al.* 2014). The performance indices (PI_{abs} and PI_{Total}) have been used to identify strategies for energy use in plants (Gonçalves *et al.* 2010). PI_{abs} evaluates the variables ϕP_0 and ABS/RC , as well as $\Psi_0 = \text{ET}_0/\text{TR}_0$ (Mehta *et al.* 2010, Yusuf *et al.* 2010). PI_{Total} also considers the reduction of PSI acceptors by incorporating the δR_0 parameter (Yusuf *et al.* 2010). Thus, analysis of ϕP_0 alone can disguise the actual condition of the plant. For example, the highest ϕP_0 values were found in S1, but this site had the lowest PI_{abs} and PI_{Total} values indicating that electron flow after Q_A^- and from the intersystem for PSI final acceptor reduction is not in good condition, which was evidenced by the analysis of Ψ_0 and δR_0 .

Plants subjected to water and salt stresses lead to a reduction in Ψ_0 , ϕE_0 , PI_{abs} , and PI_{Total} and an increase in ABS/RC and DI_0/RC (Mehta *et al.* 2010, Redillas *et al.* 2011, Silvestre *et al.* 2014, Falqueto *et al.* 2017), as well as high irradiance stress. In the present study, we found that the site with the highest salinity level (*i.e.*, S4) exhibited the highest specific energy fluxes per reaction center (ABS/RC , TR_0/RC , ET_0/RC , and DI_0/RC). But also,

Table 2. Linear correlation between abiotic and biotic variables and between biotic variables collected in the mangrove forest of the Great Vitória Estuarine System. ABS/RC – absorption flux per reaction center; C_i – intercellular CO_2 concentration; DI_0/RC – dissipated energy flux per RC; ET_0/RC – electron transport flux per RC; F_m – maximum fluorescence; F_v/F_0 – number and size of active photosynthetic RC; P_N – net photosynthetic rate; PI_{Total} – performance index for energy conservation from exciton to the reduction of PSI end acceptors; Q_{leaf} – luminous intensity incident on the leaf surface; WUE – water-use efficiency; δR_0 – efficiency with which an electron from the intersystem reduces the final acceptor of PSI; ϕD_0 – maximum efficiency of nonphotochemical de-excitation; ϕE_0 – quantum yield for electron transport; ϕP_0 – maximal quantum yield of PSII photochemistry; Ψ_0 – probability of electron transport further than Q_A^- .

Variables		<i>r</i>	<i>p</i>
Organic matter	F_m	-0.76	0.0274
	Ψ_0	0.73	0.0393
	ϕE_0	0.74	0.0354
Q_{leaf}	F_v/F_0	-0.86	0.0054
	ϕP_0	-0.86	0.0056
	ϕD_0	0.86	0.0056
	ABS/RC	0.76	0.0267
	ET_0/RC	0.85	0.0075
	DI_0/RC	0.85	0.0062
	δR_0	0.93	0.0007
	C_i	-0.91	0.0015
	WUE	0.89	0.0028
Chl _a index	Ψ_0	0.79	0.0188
	ϕE_0	0.74	0.0339
	ET_0/RC	0.80	0.0166
	C_i	-0.73	0.0394
	WUE	0.75	0.0320
PI_{Total}	P_N	0.82	0.0126
	WUE	0.78	0.0222

the highest energy flux corresponds to electron transport (ET_0/RC , Ψ_0 , and ϕE_0). This finding shows the sensitivity of the species to salinity, at least concerning the reduction in the active reaction centers (ABS/RC and F_v/F_0) and greater entrapment in energy flux (TR_0/RC) (Strasser *et al.* 2004), since S4 exhibited the highest PI_{Total} value.

In situ studies are difficult to identify a single cause of stress observed, although there is an indication of irradiance stress at S3, this evaluation can be done taking into account the parameters ABS/RC, DI_0/RC , F_v/F_0 , ϕP_0 , PI_{abs} , and PI_{Total} , as well as the negative linear correlation between Q_{leaf} and F_v/F_0 . None of the evaluated sites presented high Q_{leaf} values. In any case, S1 and S3 are assumed to have some energy sink. S1 is more closely linked to the energy flux corresponding to electron transport beyond Q_A^- and, and S3 to PSII and PSI, in addition to the lowest Chl_a and Chl_b content. Reduction in photosynthetic pigments may represent a protective strategy in which photon trapping is reduced (Christian 2005). The forest at S3 showed the highest Q_{leaf} , which

compromises photosynthetic efficiency. In addition, Teubner Júnior *et al.* (2018) point out that the Bubu River, the location of S3, has one of the lowest nutrient inputs (nitrogen and phosphorus) to the GVES. Farnworth and Ellison (1996) considered *R. mangle* as a sun-and-shade flexible species, with ontogenetic variations.

High sand content and lower MO content in sediment were observed by Zamprogno *et al.* (2016) in the mangrove area around S1. In general, sites with coarser sediment favor *L. racemosa* because the species can more efficiently use nutrients, which are scarcer in sandy areas (Cintrón and Schaeffer-Novelli 1983).

The Chl index is largely used in agronomy because it provides a reliable relationship between chlorophyll content and leaf nitrogen concentration (Pariz *et al.* 2011, Haim *et al.* 2012, Conforto *et al.* 2014, Schlichting *et al.* 2015). Nitrogen is considered one of the most important nutrients for plants and around the world, there are mangrove areas with limited growth/structure due to this nutrient deficiency (Lovell *et al.* 2006a,b; Alongi 2011). In the present study, a relationship was established between the Chl_a and the electron flux variables (Ψ_0 , ϕE_0 , ET_0/RC) and WUE by linear correlation, showing the highest values for the variables in S4 compared with the other sites evaluated. Thus, we can consider that a greater nutrient contribution to a given region may favor energy flux and water-use efficiency. In contrast, for S1, which is considered a nutrient-poor region within the study due to its sediment characteristics, this would explain the reductions in Chl_a and Chl_b, electron transport chain energy fluxes, and performance indices (PI_{abs} and PI_{Total}). In addition, the linear correlation also confirmed the relationship between OM content and variables related to the energy flux in the electron transport chain.

Mangrove species are recognized for their conservative behavior in water use, *i.e.*, an increase in salinity is usually accompanied by a reduction in water loss, reflecting greater efficiency in water use. The intensity of this relationship depends on the species (Sobrado 2000, Soares *et al.* 2015) and the level of salinity and its variation (Ball 1988, Bompy *et al.* 2014). *R. mangle* is described in the literature as the most sensitive species to salinity among neotropical species. In general, it shows less assimilation and low water-use efficiency under high salinity conditions compared to other species (Ball 1988, Sobrado 2000). However, new studies have shown that in some sites, it is the dominant species in high-salinity conditions (Estrada *et al.* 2013, Bompy *et al.* 2014). Therefore, regarding *R. mangle*, more research on photosynthetic behavior in field conditions is needed, as well as contemplating a greater population diversity.

Pascoalini (2014) evaluated net photosynthesis by *R. mangle* in the Vitória Bay and found a reduction in carbon assimilation with increasing salinity (average salinity of 21 and 28). The present study found no similar pattern, but found responses to new stressors in the system, for example, the lesser P_N at the site with the lowest organic matter content (S1). P_N was statistically similar at both sites S2 and S4, followed by S3. However, two locations stand out: S2 and S4. S4 is the site with the highest salinity

level. It has a mixed forest including *A. schaueriana*, *L. racemosa*, and *R. mangle* and is subject to anthropic interference (Pascoalini *et al.* 2019). This mangrove area is adjacent to a port complex and, consequently, it is near the dredging along the estuary's central channel, as well as all contamination from the port activities. It also receives urban and industrial effluents from a metropolitan municipality (Vila Velha) via the Aribiri River and through direct sewage disposal in the mangrove. The literature highlights the presence of heavy metals above the expected for the region in both the mangrove and estuary sediment samples, with the presence of organic contaminants in the mangrove's sediment (Jesus *et al.* 2004, Zamprogno 2015). Nonetheless, individuals at S4 were photosynthetically efficient for both chlorophyll fluorescence parameters and gas exchange, a reflection of the species' photosynthetic plasticity. One of the characteristics of *R. mangle* is the possibility of excluding salts and contaminants at the root level (Souza *et al.* 2014), preserving the photosynthetic apparatus of energy sinks.

S2 also stands out. It is located in a region of the estuary in which the organic matter content reaches 50%, although being statistically similar to S3 and S4. In addition, the site has been undergoing estuarine salinization from climate variability. In the years 2015, 2016, and 2017, the municipality of Vitória had rainfall below 1,000 mm per year, with 2016 being the most critical (INCAPER 2018). PI_{abs} , PI_{Total} , P_N , and WUE were similar to site S4 and the vegetation would be expected to respond to increased salinity, since the low rainfall affected the salinity of the estuary, as pointed out by Leite (2018) and Teubner Júnior (2016). According to the literature, in the years 2000/2001, the salinity of the estuarine region near S2 was lower than 10 (Jesus *et al.* 2004) and in 2015 reached about 30 (Leite 2018). Zamprogno *et al.* (2016) measured the salinity of the interstitial water in the mangrove sediment and recorded values ranging from 11 to 21 for the years 2012 and 2013.

Among the variables analyzed, photosynthetic efficiency seems to be controlled by Q_{leaf} , since there is a relationship between it and the variables of chlorophyll *a* fluorescence and WUE, which was confirmed by linear correlation. The Chl concentration, observed in this study through the index Chl_a , controls the performance of plants at the level of energy flux in the electron transport chain and WUE. Variables related to the energy flux in the electron transport chain are also controlled by OM. Furthermore, PI_{Total} controls net photosynthesis and WUE. The results indicate that salinity has less control over photosynthetic efficiency and is visible only in variables related to reaction centers, showing negative interaction. On the other hand, the site (S4), with the highest salinity level, has the highest photosynthetic performance, same as S2, reaffirming the role of mangrove plants' efficiency in water economy, under conditions of significant nutrient input.

The input of nitrogen ($5.6 \text{ t km}^{-2} \text{ year}^{-1}$) and phosphorus ($2.8 \text{ t km}^{-2} \text{ year}^{-1}$) to GVES is high when compared to other intermediate-size watersheds in Brazil. The contribution of nutrients from the Santa Maria River to Vitória Bay is

more significant when compared to the other basins that contribute to the system. Cattle livestock is the largest source of nutrients for GVES and then is followed by domestic effluents (Teubner Júnior *et al.* 2018). However, considering the areas of the basins evaluated, Teubner Júnior *et al.* (2018) pointed out that domestic effluents account for the highest pressure in the system. The authors recognize the Aribiri River (S4) among other urban basins of the site as the region with the highest population density and highest nutrient yield, especially nitrogen.

Nitrogen-enriched sediments favor the increase of electron transport efficiency in *A. germinans* and *R. mangle*, as well as accumulation of the element in senescent leaves (Feller *et al.* 2003). S4 had the highest Chl_a values and carbon assimilation, and the literature points to the eutrophication of the estuary. Still, if there is a relationship between chlorophyll index and leaf nitrogen concentration in mangrove plants, one can assume that vegetation is responding to nutritional enrichment.

At first, the mangrove is favored by the nutritional enrichment as already reported in the literature (Lovelock *et al.* 2006a). However, Lovelock *et al.* (2009) found an increase in mortality with long-term nutritional enrichment, and this increment of mortality was observed in the Aribiri River mangrove, through the density of dead individuals in S4 (Zamprogno *et al.* 2016, Pascoalini *et al.* 2019). As previously mentioned, this mangrove area has high levels of organic and metal contaminants and the short-term response of vegetation, especially in the young compartment, may be indifferent to these high concentrations. This unresponsive behavior is related to the fact that *R. mangle* has morphophysiological mechanisms such as precipitated iron plates in the roots. These occur since plants exposed to anoxic sediments, such as mangrove plants, release oxygen from the roots to tolerate flooding, which causes precipitation of metals in the sediment and on the root surface. This iron-rich root precipitate forms a physical barrier to heavy metal absorption (Souza *et al.* 2014).

In the present work, light saturation was not achieved, but, possibly, individuals can increase their assimilation under more favorable conditions. The low assimilation recorded in this work might be related to light attenuation by the vegetation occurring in the study sites, in addition to the other environmental factors mentioned. Attenuation of visible light confers protection to Chl_a from photooxidation, a strategy the species uses to remain long periods under the canopy until gaps are formed increasing intraspecific competition and reducing growth in low light conditions (Lima *et al.* 2018). The species will remain in the system as a source of contribution to processes of regeneration/replacement of individuals. The leaf area index of mangroves is one of the lowest among tropical forests and their leaves are mostly clustered in the upper part of the crown, with a phyllotactic pattern that minimizes leaf overlapping (Tomlinson 1986). Moreover, the ecosystem lacks the understory characteristic of humid tropical forests (Cintrón and Schaeffer-Novelli 1983). Despite these singularities, some forests are more closed than others,

resulting in light control for young individuals. Therefore, this forest compartment may exhibit less photosynthesis when compared with the canopy (Alongi 2009).

In the comparison of the physiographic types, the plants in the basin showed higher values for Chl_a and $\text{Chl}_{\text{total}}$, F_v/F_0 , ϕP_0 , Ψ_0 , ϕE_0 , PI_{abs} , and PI_{Total} . The plants of the basin had higher photochemical performance than those in the fringe. This behavior is reflected in the P_N , with the basin plants recording the highest value. In contrast, the parameters with higher values recorded in the fringe are related to stress in plants and include F_0 , ϕD_0 , ABS/RC , TR_0/RC , and DI_0/RC . The mangrove area around GVES has the basin forests more structurally developed and the fringe areas are more vulnerable, having characteristics of forest-changing variations in flood patterns, erosive and depositional processes, and irregular freshwater flow (Zamprognó 2015, Zamprognó *et al.* 2016, Teubner Júnior *et al.* 2018).

The combination of gas exchange, chlorophyll *a* fluorescence, and chlorophyll index measurements in association with abiotic data that control mangrove growth and species competition reinforce the results obtained among the different sites and comparatively analyzed to assess the species plasticity to occupy different habitats in the mangrove. These findings corroborate the importance of the management of these species in metropolitan regions, contributing to the maintenance of ecosystem services and, consequently, biological, habitat, and genetic diversity (Millennium Ecosystem Assessment 2005).

Conclusion: The study indicates the photosynthetic plasticity of *R. mangle* in a complex mangrove area. The photosynthetic efficiency of the species is controlled by the available incident radiation and organic matter content. Salinity acts by reducing or inactivating the reaction centers. On the other hand, concerning gas exchange, salinity acts on water loss, which leads to reduced stomatal conductance and, thus, increased water-use efficiency in high salinity conditions. The higher availability of OM seems to be directly related to nutrients and, consequently, to a greater energy flux to the electron transport chain in the photochemical phase of photosynthesis. Q_{leaf} is an important factor controlling photosynthetic efficiency by mainly acting on chlorophyll fluorescence variables and water-use efficiency. The biotic variables identified as controlling the photosynthetic efficiency are Chl_a concentration (observed here through the Chl index) and PI_{Total} . The former for contributing to the performance of plants at energy flux level to the electron transport chain and conservative use of water and the latter for acting on P_N and also conservative use of water. The results obtained show that the fluorescence of Chl *a*, gas exchange, and Chl index, together with sediment analysis, are suitable parameters to evaluate *R. mangle* photosynthetic plasticity in metropolitan mangroves.

References

Alongi D.M.: The Energetics of Mangrove Forests. Pp. 216. Springer, Dordrecht 2009.

- Alongi D.M.: Carbon payments for mangrove conservation: ecosystem constraints and uncertainties of sequestration potential. – *Environ. Sci. Policy* **14**: 462-470, 2011.
- Alongi D.M.: The impact of climate change on mangrove forests. – *Curr. Clim. Change Rep.* **1**: 30-39, 2015.
- Alvares C.A., Stape J.L., Sentelhas P.C. *et al.*: Köppen's climate classification map for Brazil. – *Meteorol. Z.* **22**: 711-728, 2013.
- Bai J., Ouyang H., Deng W. *et al.*: Spatial distribution characteristics of organic matter and total nitrogen of marsh soils in river marginal wetlands. – *Geoderma* **124**: 181-192, 2005.
- Ball M.C.: Photosynthesis in mangroves. – *Wetlands* **6**: 12-22, 1986.
- Ball M.C.: Ecophysiology of mangroves. – *Trees-Struct. Funct.* **2**: 129-142, 1988.
- Ball M.C., Taylor S.E., Terry N.: Properties of thylakoid membranes of the mangroves, *Avicennia germinans* and *Avicennia marina*, and the sugar beet, *Beta vulgaris*, grown under different salinity conditions. – *Plant Physiol.* **76**: 531-535, 1984.
- Barbieri Júnior E., Rossiello R.O.P., Silva R.V.M.M. *et al.*: [A new chlorophyll meter to estimate chlorophyll contents in leaves of Tifton 85 bermudagrass.] – *Cienc. Rural* **42**: 2242-2245, 2012. [In Portuguese]
- Barr J.G., Fuentes J.D., Engel V., Zieman J.C.: Physiological responses of red mangroves to the climate in the Florida Everglades. – *J. Geophys. Res.* **114**: G02008, 2009.
- Bompy F., Lequeue G., Imbert D., Dulormne M.: Increasing fluctuations of soil salinity affect seedling growth performances and physiology in three Neotropical mangrove species. – *Plant Soil* **380**: 399-413, 2014.
- Christian R.: Interactive effects of salinity and irradiance on photoprotection in acclimated seedlings of two sympatric mangroves. – *Trees-Struct. Funct.* **19**: 596-606, 2005.
- Cintrón G., Schaeffer-Novelli Y.: Introducción a la ecología del manglar. [Introduction to mangrove ecology.] Pp. 109. ROSTLAC, Montevideo 1983. [In Spanish]
- Conforto E.C., Cornélio M.L., Andreoli R.P. *et al.*: [Validation of arbitrary units of chlorophyll content obtained in intact rubber tree leaves.] – *Revista Agro@mbiente On-line* **8**: 288-292, 2014. [In Portuguese]
- Cunha S.R., Tognella-De-Rosa M.M.P., Costa C.S.B.: Salinity and flooding frequency as determinant of mangrove forest structure in Babitonga Bay, Santa Catarina state, southern Brazil. – *J. Coast. Res.* **39**: 1175-1180, 2006.
- Espírito Santo, Decree 2625-R, November 23, 2010. Official State Gazette, Executive Branch, Vitória, ES. Pp. 3, 2010. [In Portuguese]
- Estrada G.C.D., Soares M.L.G., Chaves F.O., Cavalcanti V.F.: Analysis of the structural variability of mangrove forests through the physiographic types approach. – *Aquat. Bot.* **111**: 135-143, 2013.
- Falqueto A.R., Santos P.N., Fontes R.V., Silva D.M.: Analysis of chlorophyll *a* fluorescence of two mangrove species of Vitória Bay (ES, Brazil) to natural variation of tide. – *Rev. Biociênc.* **18**: 14-23, 2012.
- Falqueto A.R., Silva D.M., Fontes R.V.: Photosynthetic performance of mangroves *Rhizophora mangle* and *Laguncularia racemosa* under fields conditions. – *Rev. Árvore* **32**: 577-582, 2008.
- Falqueto A.R., Silva Júnior R.A., Gomes M.T.G. *et al.*: Effects of drought stress on chlorophyll *a* fluorescence in two rubber tree clones. – *Sci. Hortic.-Amsterdam* **224**: 238-243, 2017.
- Farnworth E.J., Ellison A.M.: Sun-shade adaptability of the red mangrove, *Rhizophora mangle* (Rhizophoraceae): changes

- through ontogeny at several levels of biological organization. – *Am. J. Bot.* **83**: 1131-1143, 1996.
- Feller I.C., McKee K.L., Whigham D.F., O'Neill J.P.: Nitrogen vs. phosphorus limitations across an ecotonal gradient in a mangrove. – *Biogeochemistry* **62**: 145-175, 2003.
- Gonçalves J.F.C., Santos Júnior U.M.: Utilization of the chlorophyll *a* fluorescence technique as a tool for selecting tolerant species to environments of high irradiance. – *Braz. J. Plant Physiol.* **17**: 307-313, 2005.
- Gonçalves J.F.C., Santos Júnior U.M., Nina Júnior A.R., Chevreuil L.R.: Energetic flux and performance index in copaiba (*Copaifera multijuga* Hayne) and mahogany (*Swietenia macrophylla* King) seedlings grown under two irradiance environments. – *Braz. J. Plant Physiol.* **19**: 171-184, 2007.
- Gonçalves J.F.C., Silva C.E., Guimarães D.G., Bernardes R.S.: [Analysis of chlorophyll *a* fluorescence transients of young plants of *Carapa guianensis* and *Dipteryx odorata* submitted to two light environments.] – *Acta Amaz.* **40**: 89-98, 2010. [In Portuguese]
- Gonzalez-Mendoza D., Gil F.E., Santamaría J.M., Zapata-Perez O.: Multiple effects of cadmium on the photosynthetic apparatus of *Avicennia germinans* L. as probed by OJIP chlorophyll fluorescence measurements. – *Z. Naturforsch.* **939**: 265-272, 2007.
- Haim P.G., Zoffoli B.C., Zonta E., Araújo A.P.: [Nutritional diagnosis of nitrogen in bean leaves by digital image analysis.] – *Pesqui. Agropecu. Bras.* **47**: 1546-1549, 2012. [In Portuguese]
- Hoppe-Speer S.C.L., Adams J.B., Rajkaran A., Bailey D.: The response of the red mangrove *Rhizophora mucronata* Lam. to salinity and inundation in South Africa. – *Aquat. Bot.* **95**: 71-76, 2011.
- INCAPER (Instituto Capixaba de Pesquisa, Assistência Técnica e Extensão Rural). Accessed: 2018. <http://www.incaper.es.gov.br/>
- Jesus H.C., Costa E.A., Mendonça A.S.F., Zandonade E.: [Distribution and abundance of heavy metals in sediments from Vitória Island estuarine system.] – *Quím. Nova* **27**: 378-386, 2004. [In Portuguese]
- Kalaji H.M., Oukarroum A., Alexandrov V.: Identification of nutrient deficiency in maize and tomato plants by *in vivo* chlorophyll *a* fluorescence measurements. – *Plant Physiol. Bioch.* **81**: 16-25, 2014.
- Khadka W.: Assessment of relationship between soil organic matter and macronutrients, western Nepal. – *J. Biol. Pharm. Chem. Res.* **3**: 4-12, 2016.
- Krauss K.W., Twilley R.R., Doyle T.W., Gardiner E.S.: Leaf gas exchange characteristics of three neotropical mangrove species in response to varying hydroperiod. – *Tree Physiol.* **26**: 959-968, 2006.
- Larcher L., Boeger M.R.T., Sternberg L.S.L.O.: Gas exchange and isotopic signature of mangrove species in Southern Brazil. – *Aquat. Bot.* **133**: 62-69, 2016.
- Leite T.A.: [Filtering bivalves and the regulation of estuarine eutrophication – biofiltration of the mangrove oyster, *Crassostrea rhizophorae*, in the northern section of the Estuarine System of Baía de Vitoria – ES, Brazil.] Pp. 212. PhD dissertation. Federal University of Espírito Santo 2018. [In Portuguese]
- Lima K.O.O., Tognella M.M.P., Cunha S.R., Andrade H.A.: Growth models of *Rhizophora mangle* L. seedlings in tropical southwestern Atlantic. – *Estuar. Coast. Shelf Sci.* **207**: 154-163, 2018.
- Lima T.M.J.: [Effect of flood frequency on mangrove structure in Vitória Bay, Espírito Santo, Brazil.] Pp. 84. Master thesis. Federal University of Espírito Santo 2011. [In Portuguese]
- Lopes D.M.S., Tognella M.M.P., Falqueto A.R., Soares M.L.G.: Salinity variation effects on photosynthetic responses of the mangrove species *Rhizophora mangle* L. growing in natural habitats. – *Photosynthetica* **57**: 1142-1155, 2019.
- Lovelock C.E., Ball M.C., Choat B. *et al.*: Linking physiological processes with mangrove forest structure: phosphorus deficiency limits canopy development, hydraulic conductivity and photosynthetic carbon gain in dwarf *Rhizophora mangle*. – *Plant Cell Environ.* **29**: 793-802, 2006b.
- Lovelock C.E., Ball M.C., Feller I.C. *et al.*: Variation in hydraulic conductivity of mangroves: influence of species, salinity, and nitrogen and phosphorus availability. – *Physiol. Plantarum* **127**: 457-464, 2006a.
- Lovelock C.E., Ball M.C., Martin K.C., Feller I.C.: Nutrient enrichment increases mortality of mangroves. – *PLoS ONE* **4**: e5600, 2009.
- Martin K.C., Bruhn D., Lovelock C.E. *et al.*: Nitrogen fertilization enhances water-use efficiency in a saline environment. – *Plant Cell Environ.* **33**: 344-357, 2010.
- McKee K.L.: Interspecific variation in growth, biomass partitioning, and defensive characteristics of neotropical mangrove seedlings: response to light and nutrient availability. – *Am. J. Bot.* **82**: 299-307, 1995.
- Medina E.: Mangrove physiology: the challenge of salt, heat, and light stress under recurrent flooding. – In: Yáñez-Arancibia A., Lara-Domínguez A.L. (ed.): *Ecosistemas de manglar en América Tropical*. [Mangrove ecosystems in Tropical America.] Pp. 109-126. Instituto de Ecología A.C., México, UICN/ORMA, Costa Rica, NOAA/NMFS Silver Spring 1999. [In Spanish]
- Medina E., Cuevas E., Lugo A.E.: Nutrient relations of dwarf *Rhizophora mangle* L. mangroves on peat in eastern Puerto Rico. – *Plant Ecol.* **207**: 13-24, 2010.
- Mehta P., Jajoo A., Mathur S., Bharti S.: Chlorophyll *a* fluorescence study revealing effects of high salt stress on Photosystem II in wheat leaves. – *Plant Physiol. Bioch.* **48**: 16-20, 2010.
- Meng X., Chen W.W., Wang Y.Y. *et al.*: Effects of phosphorus deficiency on the absorption of mineral nutrients, photosynthetic system performance and antioxidant metabolism in *Citrus grandis*. – *PLoS ONE* **16**: e0246944, 2021.
- Millennium Ecosystem Assessment: *Ecosystems and Human Well-being: Synthesis*. Pp. 155. Island Press, Washington 2005.
- Mook D.H., Hoskin C.M.: Organic determinations by ignition: Caution advised. – *Estuar. Coast. Shelf Sci.* **15**: 697-699, 1982.
- Naidoo G., Tuffers A.V., von Willert D.J.: Changes in gas exchange and chlorophyll fluorescence characteristics of two mangroves and a mangrove associate in response to salinity in the natural environment. – *Trees-Struct. Funct.* **16**: 140-146, 2002.
- Neves R.C., Quaresma V.S., Bastos A.C., Silva J.C.R.: Sedimentary transport in coastal bays: case study of Vitória and Espírito Santo bays – ES – Brazil. – *Braz. J. Geophys.* **30**: 181-189, 2012.
- Parida A.K., Das A.B., Mitra B.: Effects of salt on growth, ion accumulation, photosynthesis and leaf anatomy of the mangrove, *Bruguiera parviflora*. – *Trees-Struct. Funct.* **18**: 167-174, 2004.
- Parida A.K., Jha B.: Salt tolerance mechanisms in mangroves: a review. – *Trees-Struct. Funct.* **24**: 199-217, 2010.
- Pariz C.M., Andreotti M., Bergamaschine A.F. *et al.*: Yield, chemical composition and chlorophyll relative content of Tanzania and Mombasa grasses irrigated and fertilized with

- nitrogen after corn intercropping. – R. Bras. Zootec. **40**: 728-738, 2011.
- Pascoalini S.S.: [Photosynthetic efficiency in mangroves at Baía de Vitória, ES.] Pp. 69. Master thesis. Federal University of Espírito Santo 2014. [In Portuguese]
- Pascoalini S.S., Lopes D.M.S., Falqueto A.R., Tognella M.M.P.: [Ecophysiological approach to mangroves: a review.] – Biotemas **27**: 1-11, 2014. [In Portuguese]
- Pascoalini S.S., Tognella M.M.P., Lima K.O.O., Falqueto A.R.: Structural plasticity and species distribution in a periurban mangrove of Southeastern Brazil. – Sci. Res. Essays **14**: 129-144, 2019.
- Rebello E.R.G., Silva S.M., Silva J.F.: [Vitória-ES water balance for climate change scenarios.] Pp. 1-5. 2011. <http://sbagro.org/files/biblioteca/3467.pdf> Accessed: 12 October, 2018. [In Portuguese]
- Redillas M.C.F.R., Strasser R.J., Jeong J.S. *et al.*: The use of JIP test to evaluate drought-tolerance of transgenic rice overexpressing *OsNAC10*. – Plant Biotechnol. Rep. **5**: 169-175, 2011.
- Rigo D., Chacaltana J.T.A.: Computational modelling of mangrove effects on the hydrodynamics of Vitoria Bay, Espírito Santo – Brazil. – J. Coast. Res. **39**: 1543-1545, 2006.
- Ru Q.M., Xiao Q., Lin P. *et al.*: Short- and long-term effects of NaCl on physiological and biochemical characteristics in leaves of a true mangrove, *Kandelia candel*. – Russ. J. Plant Physiol. **56**: 363-369, 2009.
- Schaeffer-Novelli Y., Cintrón-Molero G., Adaime R.R., Camargo T.M.: Variability of mangrove ecosystems along the Brazilian coast. – Estuaries **13**: 204-218, 1990.
- Schaeffer-Novelli Y., Cintrón-Molero G., Soares M.L.G., De-Rosa T.: Brazilian mangroves. – Aquat. Ecosyst. Health Manag. **3**: 561-570, 2000.
- Schlichting A.F., Bonfim-Silva E.M., Silva M.C. *et al.*: Efficiency of portable chlorophyll meters in assessing the nutritional status of wheat plants. – Ver. Bras. Eng. Agric. Ambient. **19**: 1148-1151, 2015.
- Silvestre S., Araújo S.S., Patto M.C.V., Silva J.M.: Performance index: an expeditious tool to screen for improved drought resistance in the *Lathyrus* genus. – J. Integr. Plant Biol. **56**: 610-621, 2014.
- Soares M.L.G., Tognella M.M.T., Cuevas E., Medina E.: Photosynthetic capacity and intrinsic water-use efficiency of *Rhizophora mangle* at its southernmost western Atlantic range. – Photosynthetica **53**: 464-470, 2015.
- Sobrado M.A.: Leaf photosynthesis of the mangrove *Avicennia germinans* as affected by NaCl. – Photosynthetica **36**: 547-555, 1999.
- Sobrado M.A.: Relation of water transport to leaf gas exchange properties in three mangrove species. – Trees-Struct. Funct. **14**: 258-262, 2000.
- Sobrado M.A.: Leaf characteristics and gas exchange of the mangrove *Laguncularia racemosa* as affected by salinity. – Photosynthetica **43**: 217-221, 2005.
- Souza I.C., Morozesk M., Duarte I.D. *et al.*: Matching pollution with adaptive changes in mangrove plants by multivariate statistics. A case study, *Rhizophora mangle* from four neotropical mangroves in Brazil. – Chemosphere **108**: 115-124, 2014.
- Strasser R.J., Srivastava A., Tsimilli-Michael M.: The fluorescence transient as a tool to characterize and screen photosynthetic samples. – In: Yunus M., Pathre U., Mohanty P. (ed.): Probing Photosynthesis: Mechanisms, Regulation and Adaptation. Pp. 445-483. Taylor & Francis, London 2000.
- Strasser R.J., Stirbet A.D.: Heterogeneity of photosystem II probed by the numerically simulated chlorophyll *a* fluorescence rise (O–J–I–P). – Math. Comput. Simulat. **48**: 3-9, 1998.
- Strasser R.J., Tsimilli-Michael M., Srivastava A.: Analysis of the chlorophyll *a* fluorescence transient. – In: Papageorgiou G.C., Govindjee (ed.): Chlorophyll *a* Fluorescence: A Signature of Photosynthesis. Advances in Photosynthesis and Respiration. Pp. 321-362. Springer, Dordrecht 2004.
- Swoczyna T., Łata B., Stasiak A. *et al.*: JIP-test in assessing sensitivity to nitrogen deficiency in two cultivars of *Actinidia arguta* (Siebold et Zucc.) Planch. ex Miq. – Photosynthetica **57**: 646-658, 2019.
- Teubner Júnior F.J.: [Supply of water and nutrients to the estuarine system of Baía de Vitória (ES): Subsidies for integrated environmental management.] Pp. 304. PhD dissertation. Federal University of Espírito Santo 2016. [In Portuguese]
- Teubner Júnior F.J., Lima A.T.M., Barroso G.F.: Emission rates of nitrogen and phosphorus in a tropical coastal river basin: a strategic management approach. – Environ. Monit. Assess. **190**: 747, 2018.
- Tognella M.M.P., Falqueto A.R., Espinoza H.D.C.F. *et al.*: Mangroves as traps for environmental damage to metals: The case study of the Fundão Dam. – Sci. Total Environ. **806**: 150452, 2022.
- Tognella M.M.P., Soares M.L.G., Cuevas E. *et al.*: Heterogeneity of elemental composition and natural abundance of stable isotopes of C and N in soils and leaves of mangroves at their southernmost West Atlantic range. – Braz. J. Biol. **76**: 994-1003, 2016.
- Tomlinson P.B.: The Botany of Mangroves. Pp. 419. Cambridge University Press, Cambridge 1986.
- Tóth S.Z., Schansker G., Strasser R.J.: A non-invasive assay of the plastoquinone pool redox state based on the OJIP-transient. – Photosynth. Res. **93**: 193-203, 2007.
- Wong P.P., Losada I.J., Gattuso J.P. *et al.*: Coastal systems and low-lying areas. – In: Field C.B., Barros V.R., Dokken D.J. *et al.* (ed.): Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Pp. 361-409. Cambridge University Press, Cambridge-New York 2014.
- Yamane Y., Kashino Y., Koike H., Satoh K.: Increases in the fluorescence *F_o* level and reversible inhibition of photosystem II reaction center by high-temperature treatments in higher plants. – Photosynth. Res. **52**: 57-64, 1997.
- Yusuf M.A., Kumar D., Rajwanshi R. *et al.*: Overexpression of γ -tocopherol methyl transferase gene in transgenic *Brassica juncea* plants alleviates abiotic stress: physiological and chlorophyll *a* fluorescence measurements. – BBA-Bioenergetics **1797**: 1428-1238, 2010.
- Zamprogno G.C.: [Use of biological and environmental factors to assess levels of vulnerability to impacts arising from urban occupation in the mangroves of Baía de Vitória, ES.] Pp. 216. PhD dissertation. Federal University of Espírito Santo 2015. [In Portuguese]
- Zamprogno G.C., Tognella M.M.P., Quaresma V.S. *et al.*: The structural heterogeneity of an urbanised mangrove forest area in southeastern Brazil: influence of environmental factors and anthropogenic stressors. – Braz. J. Oceanogr. **64**: 157-172, 2016.
- Zar J.H.: Biostatistical Analysis. Pp. 662. Prentice Hall, New Jersey 1996.