

MEETING ABSTRACT

Open Access

# TRPC3 overexpression promotes angiotensin II-induced cardiac dysfunction

Bernhard Doleschal<sup>1</sup>, Stefan Wolf<sup>1</sup>, Marie-Sophie Huber<sup>1</sup>, Gerald Wölkart<sup>1</sup>, Gudrun Antoons<sup>2</sup>, Klaus Groschner<sup>3\*</sup>

From 18th Scientific Symposium of the Austrian Pharmacological Society (APHAR). Joint meeting with the Croatian, Serbian and Slovenian Pharmacological Societies.

Graz, Austria. 20-21 September 2012

## Background

TRPC3 was recently demonstrated as a player in pathogenesis of cardiac hypertrophy, while the potential pro-arrhythmogenic role of TRPC3 is incompletely understood. Using a TRPC3 transgenic overexpression mouse model, we examined the involvement of TRPC3 in cardiac actions of angiotensin II (AngII).

## Methods

AngII effects on cardiac functions were characterized in Langendorff perfused hearts. Single ventricular myocytes were isolated and field-stimulated to measure effects on sarcomere shortening and  $\text{Ca}^{2+}$  transients. Furthermore, L-type  $\text{Ca}^{2+}$  channel current, action potentials and non-selective ion currents were analyzed electrophysiologically.

## Results

AngII (100 nM) reduced left ventricular pressure (LVP) within 2 min to 64%,  $+\text{dP}/\text{dt}$  to 50% and  $-\text{dP}/\text{dt}$  to 55% of control in TRPC3(+/−) hearts, while even producing a positive inotropic effect in wild-type (WT) hearts. Simultaneously, ECG recordings demonstrated AngII-induced episodes of acute arrhythmogenicity in all TRPC3(+/−) hearts ( $n = 6$ ), whereas rhythm of WT hearts ( $n = 6$ ) remained unaffected. The AngII-induced impairment of cardiac functions in TRPC3(+/−) hearts was partially reversed by Pyr3 (30  $\mu\text{M}$ ). The amplitude of  $\text{Ca}^{2+}$  transient was significantly higher ( $p < 0.05$ ;  $n = 60$ ) in myocytes from TRPC3(+/−) mice ( $[\text{Ca}^{2+}] \text{F/F}_0 0.354 \pm 0.024$ ) as compared to WT ( $[\text{Ca}^{2+}] \text{F/F}_0 0.262 \pm 0.021$ ). Also, the time constant ( $\tau$ ) of  $\text{Ca}^{2+}$  decline was different between WT ( $0.196 \pm 0.009$  ms;  $n = 61$ ) and TRPC3(+/−) ( $0.170 \pm 0.008$ ;  $n = 67$ ;  $p < 0.05$ ). Sarcomere shortening showed no

significant difference between the two groups ( $3.80 \pm 0.69\%$  vs.  $3.52 \pm 0.65\%$ ;  $n = 10$ ) whereas the SR-loading estimated from rapid application of caffeine (20 mM) revealed an increased SR loading of up to 40% in TRPC3(+/−) myocytes as compared to WT ( $p < 0.05$ ;  $n = 43$ ). The time constant of  $\text{Ca}^{2+}$  decline during caffeine challenge was also significantly changed ( $p < 0.05$ ) in TRPC3(+/−) myocytes ( $3.04 \pm 0.44$  ms;  $n = 11$ ) as compared to WT cells ( $1.65 \pm 0.158$  ms;  $n = 16$ ). Importantly, AngII (100 nM) induced a rise in diastolic  $\text{Ca}^{2+}$  levels, which was accompanied by irregular contractions in TRPC3(+/−) but not in WT myocytes. The rise in the diastolic  $\text{Ca}^{2+}$  levels was significantly suppressed by Pyr3 (10  $\mu\text{M}$ ;  $n = 16$ ), SEA 0400 (1  $\mu\text{M}$ ;  $n = 14$ ) and KN-93 (1  $\mu\text{M}$ ;  $n = 12$ ). Electrophysiological characterization of L-type voltage-gated  $\text{Ca}^{2+}$  currents and action potentials revealed that baseline electrophysiological parameters were not affected by TRPC3 overexpression, while AngII induced a transient prolongation of action potential duration only in TRPC3(+/−) myocytes. This TRPC3-dependent response was associated with a higher incidence of delayed afterdepolarizations.

## Conclusions

Our results demonstrate that AngII modulation of cardiac functions is strictly dependent on TRPC3 expression and suggest a key role of TRPC channels in AngII-mediated arrhythmogenicity.

## Acknowledgements

Supported by the Austrian Science Fund FWF and the International PhD Program Metabolic and Cardiovascular Disease DK-MCD.

## Author details

<sup>1</sup>Department of Pharmacology and Toxicology, Institute of Pharmaceutical Sciences, University of Graz, 8010 Graz, Austria. <sup>2</sup>Division of Cardiology, Medical University of Graz, 8036 Graz, Austria. <sup>3</sup>Institute of Biophysics, Medical University of Graz, 8010 Graz, Austria.

\* Correspondence: klaus.groschner@medunigraz.at

<sup>3</sup>Institute of Biophysics, Medical University of Graz, 8010 Graz, Austria  
Full list of author information is available at the end of the article

Published: 17 September 2012

doi:10.1186/2050-6511-13-S1-A83

Cite this article as: Doleschal et al.: TRPC3 overexpression promotes angiotensin II-induced cardiac dysfunction. *BMC Pharmacology and Toxicology* 2012 13(Suppl 1):A83.

Submit your next manuscript to BioMed Central  
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

