



The application of artificial intelligence in lung cancer: a narrative review

Huixian Zhang^{1,2,3#}, Die Meng^{1,2#}, Siqi Cai^{1,2#}, Haoyue Guo^{1,2}, Peixin Chen^{1,2}, Zixuan Zheng^{1,2}, Jun Zhu^{1,2}, Wencheng Zhao¹, Hao Wang^{1,2}, Sha Zhao¹, Jia Yu¹, Yayi He¹

¹Department of Medical Oncology, Shanghai Pulmonary Hospital, Tongji University Medical School Cancer Institute, Tongji University School of Medicine, Shanghai, China; ²Tongji University, Shanghai, China; ³Department of Medical Oncology, The First Affiliated Hospital of Zhengzhou University, Zhengzhou, China

Contributions: (I) Conception and design: Y He, H Zhang, D Meng, S Cai; (II) Administrative support: None; (III) Provision of study materials or patients: Y He, H Zhang, W Zhao; (IV) Collection and assembly of data: All authors; (V) Data analysis and interpretation: All authors; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

[#]These authors contributed equally to this work.

Correspondence to: Yayi He. Department of Medical Oncology, Shanghai Pulmonary Hospital, Tongji University Medical School Cancer Institute, Tongji University School of Medicine, No. 507 Zhengmin Road, Shanghai 200433, China. Email: 2250601@qq.com.

Objective: This review was conducted to systematically summarize the progress made by artificial intelligence technology in early screening based medical imaging, pathological diagnosis, genomics inspection, prognostic evaluation, and individual treatment of lung cancer.

Background: Lung cancer has a high mortality rate in China, which is closely related to the fact that most lung cancer patients are not diagnosed until the malignancy is advanced. Challenges remain in the early detection, accurate diagnosis, monitoring and individual treatment of lung cancers. Artificial intelligence has developed to process large amounts of data, from clinical presentations to physiological images, which is essential for solving the complex issues with clinical medicine. Increasing evidence has suggested that artificial intelligence technology provides novel, promising strategies for the diagnosis and treatment of lung cancer.

Methods: A review of literature was conducted in PubMed, EMBASE and Cochrane to identify the latest research on artificial intelligence and lung cancer, and ultimately to generate a narrative review.

Conclusions: Artificial intelligence plays an important role in the imaging inspection, histopathology examination and genomics inspection of lung cancer. In addition, artificial intelligence has the ability to detect a small number of biomarkers, which is conducive to lung cancer monitoring. Moreover, the intelligent treatment of lung cancer has gradually become the trend of future development, whether in internal medicine or surgical treatment. It is believed that artificial intelligence could improve the early diagnosis of lung cancer and assist doctors in treating lung cancer patients individually.

Keywords: Lung cancer; artificial intelligence; early diagnosis; individual treatment

Submitted Dec 07, 2020. Accepted for publication Mar 24, 2021.

doi: 10.21037/tcr-20-3398

View this article at: <http://dx.doi.org/10.21037/tcr-20-3398>

Introduction

The leading cause of cancer-related deaths across the world is lung cancer (1-4). Most lung cancer patients are subject to late diagnosis and the lack of efficient

treatment (5). More attention needs to be paid to rapid, sensitive and accurate detection techniques. At present, doctors mainly rely on the clinical features of patients and imaging characters to diagnose lung cancer. However, these methods are ineffective in finding lesions timely




Figure 1 The application of AI in lung cancer: AI plays an important role in the detection, dynamical monitor, and treatment of lung cancer. AI, artificial intelligence; TAAs, tumor-associated antigens; ctDNA, circulating tumor DNA; TEP, tumor educated platelets.

and completely. Artificial intelligence (AI) is referred to as the ability of a digital computer or a robot controlled by a digital computer to carry out some assignments that meet people’s needs (6). A large number of studies have demonstrated that AI is not only valid in detection, dynamical monitor, but also in quantitative treatment during the process of health protection, thus revealing a major capacity for cancer care (7,8). In this review, we will discuss the application of AI in lung cancer and the challenges in clinical application (*Figure 1*). We present the following article in accordance with the Narrative Review reporting checklist (available at <http://dx.doi.org/10.21037/tcr-20-3398>).

AI and detection

Imaging inspection

Clinically, when patients feel uncomfortable or have suspected lesions, their whole body or a certain part will be scanned by means of digital radiography (DR), computed

tomography (CT), positron emission computed tomography (PETCT), or magnetic resonance imaging (MRI) (9,10). These imaging techniques as mentioned above play an important role in cancer diagnosis. The main practical problem facing doctors is that their access to information and the integration of digital images of visible anomalies cannot be consistent and accurate (9,11,12).

In an attempt to obtain information instantly with no omission, AI developed its logical algorithm to detect pictures fast. Computer-aided detection (CAD) systems are applied to determine the target site in clinical pictures (13). AI-based detection tools can mark the lesion sites, reduce observational oversights, and make it convenient for readers. For example, CAD systems are easy to distinguish between bronchioles, lung wall and parenchyma clearly while highlighting the lesions that are different than normal, which enables the physicians to ascertain changes without error (12). On the other hand, computer-aided diagnosis systems have paid attention to classifying benign and malignant nodules (14). Up to now,

new approaches have been designed to detect and diagnose the lung nodules sized less than 3 mm (15). AI aims to tailor specific solutions to a certain person based on the texture of tumor, character, staging and invasion area.

The imaging features of early lung cancer are pure ground-glass nodule (GGN), partially solid or solid nodule (16). The early detection and surgical resection of lung cancer, presented as GGN or partial solid nodules, can increase the 5-year survival rate to over 95% (17). Cancerous pulmonary nodules are difficult to distinguish from benign ones in the early stages. The most suitable solution to lung cancer screening is low-dose CT (LDCT) (18). However, for radiologists, the early detection of lung cancer nodules is very challenging.

The application of AI can overcome the shortage of radiologists and is conducive to the early diagnosis of lung cancer. In recent years, there has been remarkable progress made by the CAD system of pulmonary nodules using traditional machine learning (ML) approaches (19). Recent studies have shown that AI technology in pulmonary nodule imaging is superior to radiologists in detecting the GGN of more than 5 mm and screening the nodules of 0~3 mm (20). However, CAD systems cannot be used alone in clinical practice. Vasopressors (VS)-CAD system, also known as the ClearRead CT software, which was approved by the US Food and Drug Administration (FDA) in 2016, can improve the detectability of small pulmonary nodules by inhibiting the normal structures in the lung, mainly blood vessels (21).

With the development of AI, deep learning (DL) and convolutional neural networks (CNNs) have shown their massive potential in the detection of pulmonary nodules and are effective in segmenting and classifying pulmonary nodules (22). Hua *et al.* established a DL-based CT detection system for the pulmonary nodules with a sensitivity of 73% and specificity of 80%, which is superior to the traditional CAD system (23). Ardila and his colleagues developed a DL-based model to predict the risk of developing the lung cancer by analyzing the recent and previous CT volume of patients (24). Nevertheless, detecting pulmonary nodules by AI still needs to be verified in practice and it requires more cases for rectification (25).

Histopathology inspection

Traditionally, pathological examination has been identified as the most important criterion for the diagnosis of lung cancer, which is an effective approach to digging out changes compared with the surrounding tissue and

to identifying the nature of the biopsy species (26-28). However, most patients who undergo biopsies are at an advanced stage and cannot be cured. In addition, what has been taken is just a part of tumor that can't indicate lesions entirely (29). The vast majority of pathological methods are limited to discriminating equivocal histological features, especially in poorly differentiated tumors (9). Drawing upon multiple strands of research into AI, histological analysis can be conducted to develop exhaustive and intelligent diagnostic tactics for biopsy sample (29).

The studies on identifying sensitive and specific method to make diagnosis is widely established. A novel source of large data for personalized medicine was collected by using high-throughput whole-slide scanning technologies and through image computing. Compared with routine clinical images, AI gets integrated with data information prognostically and predicatively to make accurate histology slides analysis (30). Moreover, pathologists can speed up workflow by using the automatic machines that may offer large volume slides and conduct analysis excellently (9,31). Immunohistochemical (IHC) staining that works in mucin staining is an easily available technique that helps to standardize the assessment of the subtype of lung cancer (29,32). In the study of Koh *et al.*, a reliable IHC panel was applied to diagnose the histological subtype of lung cancer thoroughly. They employed two AI approaches including a support vector machine (SVM) and a decision tree to ascertain ambiguous morphology from 30 small biopsy cases of lung cancer. In the absence of 3-marker IHC panel (i.e., TTF-1, Napsin A and p40), they achieved the highest accuracy of approximately 72% based on the decision tree model that performed two markers (p63 and CK5/6) (33). Although AI expedites workflow and provides precious images to help diagnose lung cancer, the application of AI in clinical settings is rare. The most important reason is that the technology algorithm has not been improved sufficiently to meet the needs of extended patients. In addition, the high price is also a barrier to the widespread application of AI.

Genomics inspection

With the release of the whole genome sequence and functional genomics data, individualized medical genetics assists with therapy decisions and helps practitioners to prescribe an appropriate therapy for a certain patient. The overexpression of a proto-oncogene or disruption caused by a tumor suppressor gene may lead to cell transformation and tumor formation (34-36). In recent researches, proto-

oncogenes and tumor-suppressor genes have been identified as markers for the diagnosis of cancer (30). For example, the mutations in the epidermal growth factor receptor (EGFR) gene or rearrangement of the anaplastic lymphoma kinase (ALK) receptor gene are conspicuous in lung cancer, and they are the molecular targets for custom therapeutic selection (37).

The gene conveys semblance of lung cancer by establishing microarrays and AI understands genomic pathways by optimizing algorithm (38,39). Cancer subtype, tumor growth and metastatic potential can be predicted using AI classifiers (40,41). With the growth of the next generation and other methods to reveal the sequences of gene appearance, clinicians will be able to rely on the genomic data to make individual treatment decisions (42). Watson for Genomics (WfG) is being developed to analyze the results of genetic sequencing for patients (43). A study was conducted to evaluate the effectiveness of WfG, in which the gene sequencing results of patients, including lung cancer patients with lung cancer, were first analyzed by hospital multidisciplinary experts and then reanalyzed by WfG (44). According to the results reported, for the 249 oncogenic alterations identified using both approaches, 84.6% of the targeted therapies proposed by experts were covered by WfG which also provided an additional 225 treatment options.

AI and monitor

Liquid biopsy, which is distinct from the traditional approaches retrieving biopsy arduously and huge invasively, is developed as a tool for screening. Liquid biopsy makes great contribution to determining the possible genomics alterations which will guide treatment options (6). In recent years, some studies have been conducted to reveal a variety of biomarkers, such as tumor-associated antigens (TAAs), circulating tumor DNA (ctDNA), tumor-associated autoantibodies (TAAbs), tumor educated platelets (TEP), and others (45).

Currently, the studies on liquid biopsy in cancer diagnosis are focused on the AI developed ML to detect a small number of biomarkers. ML is a part of AI and it is a great algorithm to find millions of microcirculatory genomics fragments that are more sensitive and earlier to be found compared with traditional methods (46). However, due to the limitation of tumor interstitial heterogeneity, there is still a long way to go for the application of liquid biopsy in clinical practice (11,47).

AI and treatment

With the development of AI, the intelligent diagnosis and treatment of lung cancer has gradually become the trend of future development. Clinical decision support system (CDSS) is a tool to assist clinicians in making clinical decisions through the comprehensive analysis of clinical knowledge (including clinical guidelines, evidence-based medicine, drug instructions, etc.) as well as the basic and clinical information on patients through AI (48). Watson for Oncology (WFO), as the best representative of CDSS, has been used in the diagnosis and treatment of some cancers, including lung cancer. A retrospective study was conducted to evaluate the consistency of WFO and multidisciplinary teams (MDT) in the treatment of lung cancer cases (49). The results suggested that the overall coincidence rate between MDT and WFO was 92.4% in lung cancer patients. The treatment recommendations for stage I and IV lung cancers are highly consistent. However, it is relatively low in stage II-III NSCLC and localized small cell lung cancer. In particular, the compliance rate for stage III NSCLC was only 80.8%.

Nevertheless, CDSS cannot completely replace doctors. Instead, it can play a supplementary role, especially for the patients with complex conditions. Moreover, due to the complexity of clinical medicine, doctor-patient communication and shared decision making may play a more important role in treatment.

Surgery

Video-assisted thoracic surgery (VATS) has been widely applied to treat lung cancer (50,51). A large number of previous studies have demonstrated that compared with thoracotomy, thoracoscopic lobectomy is characterized by less trauma, faster postoperative recovery and fewer complications (52). However, due to the limitations of thoracoscopy and surgical instruments, there will be some blind spots in the operation. In addition, the instrument arm is not as flexible as required to carry out more complex thoracoscopic surgery (53).

AI promises to revolutionize the way surgery is taught and practiced (54). Surgeons play a key role in integrating AI into clinical practice. The application of AI is beneficial for surgeons to analyze intraoperative progress in real time, improve the ability of intraoperative decision-making ability and finally improve the surgical outcomes (55). Robot surgery, which belongs to the physical branch of AI,

is a new direction of development for minimally invasive surgery (56). As a representative of robot surgery, the Da Vinci-robot-assisted thoracic surgery (RATS) has a precise system and convenient operation function, thus breaking the limits of hands and eyes while reducing pain and trauma for patients. The safety and feasibility of the Da Vinci surgical system in lobectomy have been demonstrated in several studies (57-59). A meta-analysis reveals that RATS and VATS are equally effective and safe in treating non-small-cell lung cancer, but RATS lasts longer. In addition, there were no statistically significant differences observed between RATS and VATS in terms of intraoperative conversion to thoracotomy rate, the incidence of postoperative complications, postoperative in-hospital mortality, intraoperative blood loss, postoperative drainage days and postoperative hospital stay (58). On the other hand, despite the above-mentioned advantages of RATS, there are some controversies surrounding this technology, such as the higher hospital costs and longer operation time, which may limit the development of RATS.

Radiotherapy

Chemoradiotherapy is the standard treatment for non-resectable advanced non-small cell lung cancer (NSCLC) (60,61). The combination of the AI and radiation contributes much in the development of radiology, which is conducive to realizing precision treatment (62,63). In a study, a clinical model of stereotactic body radiotherapy (SBRT) was designed for lung cancer patients using knowledge-based algorithms to plan treatment (64). In this study, 105 SBRT plans to treat lung cancer patients were involved, including 97 intensity modulated radiotherapy (IMRT), 6 volumetric arc radiotherapy (VMAT), and 2 three-dimensional conformal radiationtherapy (CRT). Knowledge-based model (KBM) training was carried out by combining the forementioned technologies, and multiple verification results showed that SBRT plans for lung cancer generated by KBM was comparable to clinical plans.

Existing radiation techniques are designed to eliminate tumors, but they often cause damage to healthy tissue. The application of AI will solve the problem effectively and be able to adjust the volume of treatment. In addition, the optimal angle and radiation beam can be selected automatically without being influenced by the posture of patients (65). Furthermore, intelligent system has the ability to regulate irradiation time, dose rate and images. It is believed that physicians will make plans rapidly and

precisely using AI (66,67).

Targeted therapy

Targeted therapy is a new approach to treat cancer with minimal damage to normal tissues by targeting molecules on tumor cells and developing effective blockers to intervene in the carcinogenesis process (68-70). Determining which location changed and what happened in the mutation region is an important step when we want to proceed. AI provides a large amount of data and enables an in-depth investigation, which supports us in finding various ways to prepare for useful option. In addition, intelligent machine establishes related model to find targeted gene or relational molecular pathway and make it visible.

With the assistance of AI, newly discovered therapeutic targets can be virtually screened in a very short time to find targeted old drugs or new compounds (71). AI-assisted reverse docking could be highly effective in reapplying old drugs or natural products to new indications (72). The characteristics of absorption, distribution, metabolism, excretion and toxicity (ADMET) can also be predicted by the DL model of AI, so as to improve the success rate of *in vivo* experiments (73).

Immunotherapy

The evaluation of immunotherapy for immune-responsive malignancies, particularly lung cancer, shows significant promise (74). Immune checkpoint inhibitors (ICIs), such as inhibitors of programmed death-1 (PD-1), cytotoxic correlative antigen-4 (CTLA-4), and programmed death-ligand 1 (PD-L1), exert prospective effects in advanced lung cancer (75-77).

However, some challenging problems arising from this domain are patient selection and immune response (78,79). Most of prior researches focused on predictive biomarkers about histological and genetic markers from biopsies (80). Charoentong *et al.* made a major contribution to setting up a score scheme using AI that can be applied to predict the response of patients to ICIs (81). In addition, Coroller *et al.* used CT data to demonstrate the radiomic feature after treatment (82).

The Keynote-042 study revealed that the expression of PD-L1 was closely related to the curative effect, and that the higher the expression level of PD-L1, the more significant the benefit of immunotherapy in NSCLC patients (83). The detection method of PD-L1 is simple,

convenient, straightforward, and less demanding on dyeing technology. However, there are still many problems, for example, the imbalance of detection method, strong subjectivity in the doctor's interpretation and the difficulty in giving quantitative data. AI technology has been successfully applied to the segmentation and recognition of tumor cells in tumor pathological images. Besides, it can also accurately quantify the results of IHC staining. Therefore, AI can detect the expression of PD-L1 in lung cancer tissues objectively and repeatably, which is conducive to predicting the efficacy of immunotherapy (84).

More effective biomarkers than PD-L1 are needed to predict efficacy. The toughest challenge for immunotherapy is that the process of treatment response is difficult to monitor because of dynamic changes. Once someone accepts immunotherapy, the following variation can hardly be ascertained accurately and completely. Although immunotherapy is hard to meet the final goals currently, precision treatment is made promising by the rapid development of AI (85).

Future challenges

The application of AI in understanding and treating lung cancer is advancing, but there still remain many obstacles to overcome. In healthcare settings, AI should follow unified planning and total target, as well as normalized and standard principle established by the authority that provides refined details to practice in real situation easily (62).

ML is a way to implement AI, which learns from examples rather than being programmed. Generally speaking, more examples are required for ML than human to master the same skill (56). Therefore, data acquisition is also of great importance. It is an easy way to obtain the medical records information of patients from electronic health records, which demonstrate all detailed information of the patient in hospital. Because of different recording requirements in disparate departments or districts, however, a large number of fragmented records present a challenge to the clinical application of AI (86). Moreover, language and culture barriers, patient privacy, sample volumes are all obstacles to obtaining available data (87,88).

Conclusions

The aspiration to improve the efficacy and efficiency of health care continues to drive diverse creations into practice, including AI. It has developed widespread and

is essential for solving complex issues in multiple areas. Herein, we explained multiple applications of AI in the care of lung cancer patients. Currently, the researches on AI in oncology mainly focus on diagnosis, dynamic monitoring and personal treatment. By using intelligent machines to process vast amounts of data, from clinical presentations to physiological images, physicians can make decisions quickly and accurately. Moreover, computer-assisted diagnosis will help to detect early lesions, which is beneficial to the prognosis of patients. Due to the lack of standard databases, standardization principles formulated by authoritative institutions, and corresponding national policies and regulations, the application of AI in lung cancer has a broad prospect, despite a long way to go. To realize the goal of individual treatment, more sensitive and precise machines or algorithms are needed.

Acknowledgments

Funding: This study was supported in part by a grant of National Natural Science Foundation of China (81802255), Clinical research project of Shanghai Pulmonary Hospital (FKLY20010), Young Talents in Shanghai (2019 QNB), 'Dream Tutor' Outstanding Young Talents Program (fkyq1901), Clinical research project of Shanghai Pulmonary Hospital (FKLY20001), Respiratory medicine, a key clinical specialty construction project in Shanghai, Promotion and application of multidisciplinary collaboration system for pulmonary non infectious diseases, Clinical Research Project of Shanghai Pulmonary Hospital (fk18005), Key Discipline in 2019 (oncology), Project of Shanghai Municipal Science and Technology Commission (Project of Municipal Science and Technology Commission), Scientific research project of Shanghai Pulmonary Hospital (fkcx1903), Shanghai Municipal Commission of Health and Family Planning (2017YQ050), Innovation Training Project of SITP of Tongji University, and key projects of leading talent (19411950300). Youth project of hospital management research fund of Shanghai Hospital Association (Q1902037).

Footnote

Reporting Checklist: The authors have completed the Narrative Review reporting checklist. Available at <http://dx.doi.org/10.21037/tcr-20-3398>

Conflicts of Interest: All authors have completed the ICMJE

uniform disclosure form (available at <http://dx.doi.org/10.21037/tcr-20-3398>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that the questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Sonogo M, Pellizzari I, Dall'Acqua A, et al. Common biological phenotypes characterize the acquisition of platinum-resistance in epithelial ovarian cancer cells. *Sci Rep* 2017;7:7104.
2. Krasnov GS, Puzanov GA, Afanasyeva MA, et al. Tumor suppressor properties of the small C-terminal domain phosphatases in non-small cell lung cancer. *Biosci Rep* 2019;39:BSR20193094.
3. Ferlay J, Soerjomataram I, Dikshit R, et al. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 2015;136:E359-86.
4. Roth C, Kasimir-Bauer S, Pantel K, et al. Screening for circulating nucleic acids and caspase activity in the peripheral blood as potential diagnostic tools in lung cancer. *Mol Oncol* 2011;5:281-91.
5. Guo H, Chen X, Su C, et al. Challenges and countermeasures of thoracic oncology in the epidemic of COVID-19. *Transl Lung Cancer Res* 2020;9:337-47.
6. O'Keeffe M, Barratt A, Maher C, et al. Media Coverage of the Benefits and Harms of Testing the Healthy: a protocol for a descriptive study. *BMJ Open* 2019;9:e029532.
7. Viscaino M, Maass JC, Delano PH, et al. Computer-aided diagnosis of external and middle ear conditions: A machine learning approach. *PLoS One* 2020;15:e0229226.
8. Li SM, Chen CH, Chen YW, et al. Upregulation of CISD2 augments ROS homeostasis and contributes to tumorigenesis and poor prognosis of lung adenocarcinoma. *Sci Rep* 2017;7:11893.
9. Rabbani M, Kanevsky J, Kafi K, et al. Role of artificial intelligence in the care of patients with nonsmall cell lung cancer. *Eur J Clin Invest* 2018;48.
10. Shi Z, Song T, Wan Y, et al. A systematic review and meta-analysis of traditional insect Chinese medicines combined chemotherapy for non-surgical hepatocellular carcinoma therapy. *Sci Rep* 2017;7:4355.
11. Bi WL, Hosny A, Schabath MB, et al. Artificial intelligence in cancer imaging: Clinical challenges and applications. *CA Cancer J Clin* 2019;69:127-57.
12. Fraioli F, Serra G, Passariello R. CAD (computed-aided detection) and CADx (computer aided diagnosis) systems in identifying and characterising lung nodules on chest CT: overview of research, developments and new prospects. *Radiol Med* 2010;115:385-402.
13. Brinkløv S, Kalko EK, Surlykke A. Intense echolocation calls from two 'whispering' bats, *Artibeus jamaicensis* and *Macrophyllum macrophyllum* (Phyllostomidae). *J Exp Biol* 2009;212:11-20.
14. Keshani M, Azimifar Z, Tajeripour F, et al. Lung nodule segmentation and recognition using SVM classifier and active contour modeling: a complete intelligent system. *Comput Biol Med* 2013;43:287-300.
15. Gong J, Liu J, Hao W, et al. Computer-aided diagnosis of ground-glass opacity pulmonary nodules using radiomic features analysis. *Phys Med Biol* 2019;64:135015.
16. Matsuguma H, Mori K, Nakahara R, et al. Characteristics of subsolid pulmonary nodules showing growth during follow-up with CT scanning. *Chest* 2013;143:436-43.
17. Henschke CI, Yip R, Smith JP, et al. CT Screening for Lung Cancer: Part-Solid Nodules in Baseline and Annual Repeat Rounds. *AJR Am J Roentgenol* 2016;207:1176-84.
18. Aberle DR, Adams AM, Berg CD, et al. Reduced lung-cancer mortality with low-dose computed tomographic screening. *N Engl J Med* 2011;365:395-409.
19. Goo JM. Computer-aided detection of lung nodules on chest CT: issues to be solved before clinical use. *Korean J Radiol* 2005;6:62-3.
20. Lo SB, Freedman MT, Gillis LB, et al. JOURNAL CLUB: Computer-Aided Detection of Lung Nodules on CT With a Computerized Pulmonary Vessel Suppressed Function. *AJR Am J Roentgenol* 2018;210:480-8.
21. Milanese G, Eberhard M, Martini K, et al. Vessel suppressed chest Computed Tomography for semi-automated volumetric measurements of solid pulmonary nodules. *Eur J Radiol* 2018;101:97-102.

22. Tandon YK, Bartholmai BJ, Koo CW. Putting artificial intelligence (AI) on the spot: machine learning evaluation of pulmonary nodules. *J Thorac Dis* 2020;12:6954-65.
23. Hua KL, Hsu CH, Hidayati SC, et al. Computer-aided classification of lung nodules on computed tomography images via deep learning technique. *Onco Targets Ther* 2015;8:2015-22.
24. Ardila D, Kiraly AP, Bharadwaj S, et al. End-to-end lung cancer screening with three-dimensional deep learning on low-dose chest computed tomography. *Nat Med* 2019;25:954-61.
25. Wan YL, Wu PW, Huang PC, et al. The Use of Artificial Intelligence in the Differentiation of Malignant and Benign Lung Nodules on Computed Tomograms Proven by Surgical Pathology. *Cancers (Basel)* 2020;12:2211.
26. Yang D, Li R, Zhang XH, et al. Perfusion Characteristics of Hepatocellular Carcinoma at Contrast-enhanced Ultrasound: Influence of the Cellular differentiation, the Tumor Size and the Underlying Hepatic Condition. *Sci Rep* 2018;8:4713.
27. Yuan Z, Quan J, Yunxiao Z, et al. Diagnostic value of contrast-enhanced ultrasound parametric imaging in breast tumors. *J Breast Cancer* 2013;16:208-13.
28. Liang H, Fu M, Zhou J, et al. Evaluation of 3D-CPA, HR-HPV, and TCT joint detection on cervical disease screening. *Oncol Lett* 2016;12:887-92.
29. Shulimzon TR. Endomicroscopy, Not "Optical Biopsy" (Yet). *Am J Respir Crit Care Med* 2017;195:962.
30. Djuric U, Zadeh G, Aldape K, et al. Precision histology: how deep learning is poised to revitalize histomorphology for personalized cancer care. *NPJ Precis Oncol* 2017;1:22.
31. Shillan D, Sterne JAC, Champneys A, et al. Use of machine learning to analyse routinely collected intensive care unit data: a systematic review. *Crit Care* 2019;23:284.
32. Jia K, He Y, Dziadziszko R, et al. T cell immunoglobulin and mucin-domain containing-3 in non-small cell lung cancer. *Transl Lung Cancer Res* 2019;8:895-906.
33. Koh J, Go H, Kim MY, et al. A comprehensive immunohistochemistry algorithm for the histological subtyping of small biopsies obtained from non-small cell lung cancers. *Histopathology* 2014;65:868-78.
34. Sun Y, Zhu S, Ma K, et al. Identification of 12 cancer types through genome deep learning. *Sci Rep* 2019;9:17256.
35. Wang Z, Sun Y. Targeting p53 for Novel Anticancer Therapy. *Transl Oncol* 2010;3:1-12.
36. Tanaka M, Kato K, Gomi K, et al. NUT midline carcinoma: report of 2 cases suggestive of pulmonary origin. *Am J Surg Pathol* 2012;36:381-8.
37. Lawrence MS, Stojanov P, Polak P, et al. Mutational heterogeneity in cancer and the search for new cancer-associated genes. *Nature* 2013;499:214-8.
38. Ohashi K, Sequist LV, Arcila ME, et al. Lung cancers with acquired resistance to EGFR inhibitors occasionally harbor BRAF gene mutations but lack mutations in KRAS, NRAS, or MEK1. *Proc Natl Acad Sci U S A* 2012;109:E2127-33.
39. Koivunen JP, Mermel C, Zejnullahu K, et al. EML4-ALK fusion gene and efficacy of an ALK kinase inhibitor in lung cancer. *Clin Cancer Res* 2008;14:4275-83.
40. Kikuchi T, Daigo Y, Katagiri T, et al. Expression profiles of non-small cell lung cancers on cDNA microarrays: identification of genes for prediction of lymph-node metastasis and sensitivity to anti-cancer drugs. *Oncogene* 2003;22:2192-205.
41. Haack H, Johnson LA, Fry CJ, et al. Diagnosis of NUT midline carcinoma using a NUT-specific monoclonal antibody. *Am J Surg Pathol* 2009;33:984-91.
42. Podolsky MD, Barchuk AA, Kuznetsov VI, et al. Evaluation of Machine Learning Algorithm Utilization for Lung Cancer Classification Based on Gene Expression Levels. *Asian Pac J Cancer Prev* 2016;17:835-8.
43. Chen Y, Yan W, Xie Z, et al. Comparative analysis of target gene exon sequencing by cognitive technology using a next generation sequencing platform in patients with lung cancer. *Mol Clin Oncol* 2021;14:36.
44. Itahashi K, Kondo S, Kubo T, et al. Evaluating Clinical Genome Sequence Analysis by Watson for Genomics. *Front Med (Lausanne)* 2018;5:305.
45. Revelo AE, Martin A, Velasquez R, et al. Liquid biopsy for lung cancers: an update on recent developments. *Ann Transl Med* 2019;7:349.
46. Chew RF, Amer S, Jones K, et al. Residential scene classification for gridded population sampling in developing countries using deep convolutional neural networks on satellite imagery. *Int J Health Geogr* 2018;17:12.
47. Lim ZF, Ma PC. Emerging insights of tumor heterogeneity and drug resistance mechanisms in lung cancer targeted therapy. *J Hematol Oncol* 2019;12:134.
48. Klarenbeek SE, Weekenstroom HHA, Sedelaar JPM, et al. The Effect of Higher Level Computerized Clinical Decision Support Systems on Oncology Care: A Systematic Review. *Cancers* 2020;12:1032.
49. Kim MS, Park HY, Kho BG, et al. Artificial intelligence and lung cancer treatment decision: agreement with recommendation of multidisciplinary tumor board. *Transl*

- Lung Cancer Res 2020;9:507-14.
50. Raman V, Christopher JK, Jawitz OK, et al. Robot- vs Video-Assisted Thoracoscopic Lobectomy for Early Lung Cancer. *JNCI Cancer Spectr* 2020;4:pkaa031.
 51. Wu L, Wang H, Cai H, et al. Comparison of Double Sleeve Lobectomy by Uniportal Video-Assisted Thoracic Surgery (VATS) and Thoracotomy for NSCLC Treatment. *Cancer Manag Res* 2019;11:10167-74.
 52. Paul S, Altorki NK, Sheng S, et al. Thoracoscopic lobectomy is associated with lower morbidity than open lobectomy: a propensity-matched analysis from the STS database. *J Thorac Cardiovasc Surg* 2010;139:366-78.
 53. Arad T, Levi-Faber D, Nir RR, et al. The learning curve of video-assisted thoracoscopic surgery (VATS) for lung lobectomy--a single Israeli center experience. *Harefuah* 2012;151:261-5, 320.
 54. Hashimoto DA, Rosman G, Rus D, et al. Artificial Intelligence in Surgery: Promises and Perils. *Ann Surg* 2018;268:70-6.
 55. Mangano A, Valle V, Dreifuss NH, et al. Role of Artificial Intelligence (AI) in Surgery: Introduction, General Principles, and Potential Applications. *Surg Technol Int* 2020;38:sti38/1369.
 56. Rajkomar A, Dean J, Kohane I. Machine Learning in Medicine. *N Engl J Med* 2019;380:1347-58.
 57. Veronesi G, Galetta D, Maisonneuve P, et al. Four-arm robotic lobectomy for the treatment of early-stage lung cancer. *J Thorac Cardiovasc Surg* 2010;140:19-25.
 58. Guo F, Ma D, Li S. Compare the prognosis of Da Vinci robot-assisted thoracic surgery (RATS) with video-assisted thoracic surgery (VATS) for non-small cell lung cancer: A Meta-analysis. *Medicine (Baltimore)* 2019;98:e17089.
 59. Liu X, Xu S, Liu B, et al. Survival Analysis of Stage I Non-small Cell Lung Cancer Patients Treated with Da Vinci Robot-assisted Thoracic Surgery. *Zhongguo Fei Ai Za Zhi* 2018;21:849-56.
 60. Wang CH, Lin CY, Chen JS, et al. Karnofsky Performance Status as A Predictive Factor for Cancer-Related Fatigue Treatment with Astragalus Polysaccharides (PG2) Injection-A Double Blind, Multi-Center, Randomized Phase IV Study. *Cancers (Basel)* 2019;11:128.
 61. Nakashima T, Huang C, Liu D, et al. Neural-cadherin expression associated with angiogenesis in non-small-cell lung cancer patients. *Br J Cancer* 2003;88:1727-33.
 62. Lambin P, Leijenaar RTH, Deist TM, et al. Radiomics: the bridge between medical imaging and personalized medicine. *Nat Rev Clin Oncol* 2017;14:749-62.
 63. Aerts HJ. The Potential of Radiomic-Based Phenotyping in Precision Medicine: A Review. *JAMA Oncol* 2016;2:1636-42.
 64. Chin Snyder K, Kim J, Reding A, et al. Development and evaluation of a clinical model for lung cancer patients using stereotactic body radiotherapy (SBRT) within a knowledge-based algorithm for treatment planning. *J Appl Clin Med Phys* 2016;17:263-75.
 65. Amit G, Purdie TG, Levinshtein A, et al. Automatic learning-based beam angle selection for thoracic IMRT. *Med Phys* 2015;42:1992-2005.
 66. Liu Z, Wang S, Dong D, et al. The Applications of Radiomics in Precision Diagnosis and Treatment of Oncology: Opportunities and Challenges. *Theranostics* 2019;9:1303-22.
 67. Hosny A, Parmar C, Quackenbush J, et al. Artificial intelligence in radiology. *Nat Rev Cancer* 2018;18:500-10.
 68. Jackisch C, Ruschoff J, Ullrich A. Development of Targeted Therapies in ErbB2-Positive Breast Cancer. *Breast Care (Basel)* 2008;3:3-6.
 69. Sai C, Wang J, Li B, et al. Isolation and identification of alkaloids from *Macleaya microcarpa* by UHPLC-Q-TOF-MS and their cytotoxic activity in vitro, antiangiogenic activity in vivo. *BMC Chem* 2020;14:5.
 70. He Y, Zhou C. Tyrosine kinase inhibitors interstitial pneumonitis: diagnosis and management. *Transl Lung Cancer Res* 2019;8:S318-20.
 71. Zhavoronkov A, Ivanenkov YA, Aliper A, et al. Deep learning enables rapid identification of potent DDR1 kinase inhibitors. *Nat Biotechnol* 2019;37:1038-40.
 72. Talele TT, Khedkar SA, Rigby AC. Successful applications of computer aided drug discovery: moving drugs from concept to the clinic. *Curr Top Med Chem* 2010;10:127-41.
 73. Liu B, He H, Luo H, et al. Artificial intelligence and big data facilitated targeted drug discovery. *Stroke Vasc Neurol* 2019;4:206-13.
 74. Teeling EC, Springer MS, Madsen O, et al. A molecular phylogeny for bats illuminates biogeography and the fossil record. *Science* 2005;307:580-4.
 75. Evans AG, French CA, Cameron MJ, et al. Pathologic characteristics of NUT midline carcinoma arising in the mediastinum. *Am J Surg Pathol* 2012;36:1222-7.
 76. Finlay WJJ, Coleman JE, Edwards JS, et al. Anti-PD1 'SHR-1210' aberrantly targets pro-angiogenic receptors and this polyspecificity can be ablated by paratope refinement. *MAbs* 2019;11:26-44.
 77. Prendergast CT, Sanin DE, Cook PC, et al. CD4+ T cell hyporesponsiveness after repeated exposure to *Schistosoma*

- mansoni larvae is dependent upon interleukin-10. *Infect Immun* 2015;83:1418-30.
78. Wolchok JD, Hoos A, O'Day S, et al. Guidelines for the evaluation of immune therapy activity in solid tumors: immune-related response criteria. *Clin Cancer Res* 2009;15:7412-20.
 79. Hodi FS, Hwu WJ, Kefford R, et al. Evaluation of Immune-Related Response Criteria and RECIST v1.1 in Patients With Advanced Melanoma Treated With Pembrolizumab. *J Clin Oncol* 2016;34:1510-7.
 80. He Y, Zhang X, Jia K, et al. OX40 and OX40L protein expression of tumor infiltrating lymphocytes in non-small cell lung cancer and its role in clinical outcome and relationships with other immune biomarkers. *Transl Lung Cancer Res* 2019;8:352-66.
 81. Charoentong P, Finotello F, Angelova M, et al. Pan-cancer Immunogenomic Analyses Reveal Genotype-Immunophenotype Relationships and Predictors of Response to Checkpoint Blockade. *Cell Rep* 2017;18:248-62.
 82. Coroller TP, Agrawal V, Narayan V, et al. Radiomic phenotypic features predict pathological response in non-small cell lung cancer. *Radiother Oncol* 2016;119:480-6.
 83. Mok TSK, Wu YL, Kudaba I, et al. Pembrolizumab versus chemotherapy for previously untreated, PD-L1-expressing, locally advanced or metastatic non-small-cell lung cancer (KEYNOTE-042): a randomised, open-label, controlled, phase 3 trial. *Lancet* 2019;393:1819-30.
 84. Koelzer VH, Sirinukunwattana K, Rittscher J, et al. Precision immunoprofiling by image analysis and artificial intelligence. *Virchows Arch* 2019;474:511-22.
 85. Mehnert JM, Monjazeb AM, Beerthuijzen JMT, et al. The Challenge for Development of Valuable Immunology Biomarkers. *Clin Cancer Res* 2017;23:4970-9.
 86. Schmidt C. M. D. Anderson Breaks With IBM Watson, Raising Questions About Artificial Intelligence in Oncology. *J Natl Cancer Inst* 2017;109.
 87. Leyens L, Reumann M, Malats N, et al. Use of big data for drug development and for public and personal health and care. *Genet Epidemiol* 2017;41:51-60.
 88. Auffray C, Balling R, Barroso I, et al. Making sense of big data in health research: Towards an EU action plan. *Genome Med* 2016;8:71.

Cite this article as: Zhang H, Meng D, Cai S, Guo H, Chen P, Zheng Z, Zhu J, Zhao W, Wang H, Zhao S, Yu J, He Y. The application of artificial intelligence in lung cancer: a narrative review. *Transl Cancer Res* 2021;10(5):2478-2487. doi: 10.21037/tcr-20-3398