

RESEARCH ARTICLE

Open Access

Epidemic and virulence characteristic of *Shigella* spp. with extended-spectrum cephalosporin resistance in Xiaoshan District, Hangzhou, China

Chuan-Ling Zhang^{1†}, Qing-Zhong Liu^{2*†}, Juan Wang², Xu Chu¹, Li-Meng Shen¹ and Yuan-Yu Guo¹

Abstract

Background: *Shigellae* have become increasingly resistant to the extended-spectrum cephalosporin (ESC) worldwide and pose a great challenge to anti-infection treatment options. The purpose of this study was to determine the resistance, cephalosporin resistance mechanisms, virulence characteristic and genotype of ESC-resistant *Shigella*.

Methods: From 2008 to 2012, *Shigella* isolates collected from diarrhea patients were detected for antibiotics sensitivity by disk diffusion, cephalosporin resistance determinants and virulence genes using polymerase chain reaction (PCR) and genotyping through enterobacterial repetitive intergenic consensus sequence PCR (ERIC-PCR).

Results: A total of 356 *Shigella* isolates were gathered, and 198 (55.6%, 58 *S. flexneri* and 140 *S. sonnei*) were resistant to ESC. All ESC-resistant isolates were susceptible to imipenem, and only 0.5% isolate was resistant to piperacillin/tazobactam. ESC-resistant *S. flexneri* showed high degrees of resistance to ampicillin (100%), ampicillin/sulbactam (96.6%), piperacillin (100%), trimethoprim/sulfamethoxazole (74.1%), ciprofloxacin (74.1%), levofloxacin (53.4%), ceftazidime (58.6%) and cefepime (58.6%). ESC-resistant *S. sonnei* exhibited high resistance rates to ampicillin (100%), piperacillin (100%) and trimethoprim/sulfamethoxazole (96.4%). Cephalosporin resistance genes were confirmed in 184 ESC-resistant isolates. *bla*_{CTX-M} types (91.8%, mainly *bla*_{CTX-M-14}, *bla*_{CTX-M-15} and *bla*_{CTX-M-57}) were most prevalent, followed by *bla*_{OXA-30} (26.3%). Over 99.0% ESC-resistant isolates harbored virulence genes *ial*, *ipaH*, *virA* and *sen*. However, *set1* were more prevalent in ESC-resistant *S. flexneri* isolates than in *S. sonnei* isolates. ERIC-PCR results showed that 2 and 3 main genotypes were detected in ESC-resistant *S. flexneri* and *S. sonnei*, respectively.

Conclusion: Our findings indicated that a high prevalence of ESC-resistant *Shigella* mediated mainly by *bla*_{CTX-M} with stronger resistance and virulence, and the existence of specific clones responsible for these *Shigella* infection in the region studied.

Keywords: *Shigella*, Extended-spectrum cephalosporin, Resistance, Cephalosporin resistance mechanism, Virulence markers, Genotyping

* Correspondence: zzzz2000@tom.com

†Equal contributors

²Department of Clinical Laboratory, Shanghai First People's Hospital, School of Medicine, Shanghai Jiaotong University, 100 Haining Road, Shanghai 200080, People's Republic of China

Full list of author information is available at the end of the article

Background

Shigellosis continues to be a major public health problem in non-industrialized countries. It was reported that about 91 million cases of shigellosis occurred, with 414,000 people dying in Asia annually [1]. In China, the morbidity of *Shigella* infections each year is unusually high. Wang et al. [2] have reported that in 2000 0.8-1.7 million episodes of shigellosis occurred in seven different geographical regions, China.

For shigellosis, antibiotic therapy can reduce the duration and severity of the illness [3]. However, with the long-term overuse of antimicrobials, the resistant *Shigella* has prevailed all over the world, and production of extended-spectrum β -lactamases (ESBLs) or AmpC enzymes is one of the most important resistance mechanisms. In China extended-spectrum cephalosporin (ESC)-resistant *Shigella* spp. has also been described in recent years [3-7]. However, as China has a vast territory area, the spectrum of antimicrobial resistance and the mechanisms of resistance may vary in different regions. Therefore, it is necessary to explore the variations of ESC resistance in clinical isolates of *Shigella* spp. from different geographical areas.

The ability of *Shigella* spp. to cause shigellosis is attributed to the expression of arrays of virulence genes associated with colonization, invasion/penetration and toxin-mediated disease, such as the invasion-associated locus (*ial*), the invasion plasmid antigen H gene (*ipaH*), *Shigella* enterotoxin 1 (ShET-1) gene (*set1A* and *set1B*), *Shigella* enterotoxin 2 (ShET-2) gene (*sen*) and *virA* gene [8,9]. Estimating the existence of virulence determinants in *Shigella* would help us better understand its pathogenicity. However, investigations into clinical *Shigella* spp. virulence factors are still rare in the world.

The purpose of this study was to investigate the resistance, the cephalosporin resistance mechanisms and the prevalence of putative virulence genes in ESC-resistant *Shigella* from patients with dysentery in Xiaoshan District, suburban of Hangzhou city, Zhejiang province, China. Finally we further ascertained the genotypes of these strains by ERIC-PCR (enterobacterial repetitive intergenic consensus sequence PCR) typing.

Methods

Bacterial isolates

Between January 2008 and December 2012, a total of 356 nonduplicate *Shigella* isolates were obtained from the faeces of different diarrhea patients at Zhejiang Xiaoshan hospital, Hangzhou City, Zhejiang Province, China. All isolates were identified by VITEK 60 microbial identification system (BioMérieux Corp., Lyon, France) and serotyped using commercial antisera (Lanzhou Institute of Biological Products, Lanzhou, China). Isolates studied were taken as part of standard patient care. Because of being focused on bacteria, no ethical approval was required for this study.

Antimicrobial susceptibility testing

The *Shigella* isolates were detected for antibiotic susceptibility by Kirby-Bauer method for the following antibiotics (Oxoid, Hampshire, UK), according to Clinical and Laboratory Standards Institute (CLSI), 2012 [10]: ampicillin (AMP), piperacillin (PIP), ciprofloxacin (CIP), trimethoprim/sulfamethoxazole (SXT), levofloxacin (LEV), ampicillin/sulbactam (SAM), piperacillin/tazobactam (TZP), imipenem (IMP), cefotaxime (CTX), ceftazidime (CAZ), cefepime (FEP) and cefoxitin (FOX). ESC-resistant organisms were defined as isolates showing resistance or intermediate to cefotaxime (zone diameter ≤ 25 mm) or ceftazidime (zone diameter ≤ 20 mm) or cefepime (zone diameter ≤ 17 mm) [10]. *Escherichia coli* ATCC 25922 was used for antibiotic susceptibility testing quality control.

ESBL and AmpC detection

ESC-resistant isolates were screened for ESBL production by the phenotypic confirmatory test according to CLSI [10]. AmpC examination was performed by the three-dimensional extract test as described by Philip et al. [11] for cefoxitin resistant or intermediate isolates.

ESBL, AmpC and virulence genes amplification

Detection of the ESBL genes (*bla*_{TEM}, *bla*_{OXA}, *bla*_{CTX-M-1, 2, 8, 9} and *bla*_{SHV}) and virulence determinants (*ial*, *iapH*, *set1A*, *set1B*, *sen* and *virA*) were performed by PCR for ESC-resistant isolates, as described previously [3,12,13]. AmpC genes (*bla*_{MOX}, *bla*_{CIT}, *bla*_{DHA}, *bla*_{ACC}, *bla*_{EBC} and *bla*_{FOX}) were examined by multiplex-PCR for cefoxitin resistant or intermediate isolates according to Perez-Perez et al. [14]. Sequences of the primers were given in Table 1. The PCR products for ESBL and AmpC genes were sequenced on both DNA strands with ABI 3730XL sequencer (Perkin-Elmer) by Shanghai Sangon Biotech. The nucleotide sequences were analyzed by the BLAST program. For virulence genes, a representative amplicon for the each gene detected was also sequenced to validate that primers amplified the target genes.

ERIC-PCR

ERIC-PCR was carried out on ESC-resistant *Shigella* isolates using primer ERIC2: 5'-AAGTAAGTGACTGGGGT GAGCG-3', as described by Navia et al. [15].

Clinical data

Clinical information of the patients was collected from the case notes. The data included date of onset, symptoms (such as times of diarrhea, nausea, bellyache, vomiting, fever and bloody stools), results of microscopic examination of stool (white blood cell and red blood cell), treatment and duration of illness.

Table 1 Primers used for PCR to detect *Shigella* resistance and virulence genes

Primer	Sequence (5'-3')	Amplicon size (bp)	Reference
ESBL genes			
CTX-M-1 group	F-GGCCCATGGTTAAAAAATCACTGC	891	[3]
	R-CCGTTTCCGCTATTACAAACCGTTG		
CTX-M-2 group	F-CTCAGAGCATTGCGCGCTCA	843	
	R-CCGCCGAGCCAGAATATCC		
CTX-M-8 group	F-ATGATGAGACATCGCGTTAAC	864	
	R-CGGTGACGATTTTCGCGGCAG		
CTX-M-9 group	F-GTGACAAAGAGAGTGCAACGG	856	
	R-ATGATTCTCGCCGCTGAAGCC		
OXA group	F-ACACAATACATATCAACTTCGC	813	
	R-AGTGTGTTTAGAATGGTGATC		
SHV group	F-GCCTTTTCGGCCTTCACTCAAG	989	
	R-TTAGCGTTGCCAGTGCTCGATCA		
TEM group	F-ATAAAATCTTGAAGACGAAA	1076	
	R-GACAGTTACCAATGCTTAATC		
AmpC β-lactamase genes			
MOX group	F-GCTGCTCAAGGAGCACAGGAT	520	
	R-CACATTGACATAGTGTGGTGTC		
CIT group	F-TGGCCAGAAGTACAGGCAAAA	462	
	R-TTCTCCTGAACGTGGCTGGC		
DHA group	F-AACTTTCACAGGTGTGCTGGGT	405	
	R-CCGTACGCATACTGGCTTTGC		
ACC group	F-AACAGCCTCAGCAGCCGGTTA	346	
	R-TTCGCCGAATCATCCTTAGC		
EBC group	F-TCGGTAAAGCCGATGTTGCGG	302	
	R-CTTCCACTGCGGCTGCCAGTT		
FOX group	F-AACATGGGTATCAGGGAGATG	190	
	R-CAAAGCGCGTAACCGGATTGG		
Virulence gene			
ial	F-CTGGATGGTATGGTGAGG	320	
	R-GGAGGCCAACAAATTATTTCC		
ipaH	F-TGGAAAACTCAGTGCCTCT	423	
	R-CCAGTCCGTAATTCATTCT		
set1A	F-TCACGCTACCATCAAAGA	309	
	R-TATCCCCCTTTGGTGTA		
set1B	F-GTGAACCTGCTGCCGATATC	147	
	R-ATTTGTGGATAAAAATGACG		
sen	F-ATGTGCCTGCTATTATTAT	799	
	R-CATAATAATAAGCGGTGACG		

Table 1 Primers used for PCR to detect *Shigella* resistance and virulence genes (Continued)

virA	F-CTGCATTCTGGCAATCTCTTCA	215	[13]
	CATC		
	R-TGATGAGCTAACTTCGTAAGCC		
	CTCC		

Statistical analysis

Pearson's chi-square test or Fisher's exact test if necessary was used to analyze the differences of antibiotics resistance, and virulence genes, and the relationship of virulence factors, resistance and ERIC-PCR genotypes (SPSS 11.5). All statistical tests were two tailed, with $p < 0.05$ considered statistically significant.

Results

Distribution of *Shigella* spp.

Among 356 *Shigella* isolates, *S. sonnei* was the most prevalent species ($n = 244$, 68.5%, 244/356), followed by *S. flexneri* ($n = 112$, 31.5%, 112/356). No sample was positive for *S. dysenteriae* and *S. boydii*. In *S. flexneri*, seven different serotypes were found: 63 type 2a, 15 type 2c, 18 type 4c, 6 type 2b, 5 type 1a, 3 type x and 2 type 3a. 35.7% (40/112) of *S. flexneri* and 47.1% (115/244) of *S. sonnei* cases were found among patients under 5 years old; 20.5% (23/112) of *S. flexneri* and 23.4% (57/244) of *S. sonnei* isolates were from victims aged from 5 to 14 years; 43.8% (49/112) of *S. flexneri* and 29.5% (72/244) of *S. sonnei* isolates were from patients aged ≥ 15 years.

Antimicrobial resistance

Of the 356 *Shigella* isolates, 198 (55.6%, 58 *S. flexneri* and 140 *S. sonnei*) were resistant to cefotaxime, 59 (16.6%) to ceftazidime, 66 (18.5%) to cefepime (Table 2). The strains with resistance to ceftazidime or cefepime were all resistant to cefotaxime. Thus, these cefotaxime resistant strains were referred to as ESC-resistant isolates in this study. In addition, in these ESC-resistant isolates, 8 (2.2%) were resistant or intermediate to cefoxitin.

Compared to cefotaxime susceptible *S. flexneri* isolates, the ESC-resistant *S. flexneri* isolates were more resistant to PIP (100% versus 85.2%, $p = 0.002$), SAM (96.6% versus 85.2%, $p = 0.035$), SXT (74.1% versus 46.3%, $p = 0.003$), CAZ (58.6% versus 0%, $p < 0.001$) and FEP (58.6% versus 0%, $p < 0.001$). In *S. sonnei*, the ESC-resistant strains were more resistant to AMP (100% versus 95.2%, $p = 0.009$), PIP (100% versus 51.9%, $p < 0.001$), SAM (24.3% versus 0%, $p < 0.001$), CAZ (17.9% versus 0%, $p < 0.001$) and FEP (22.9% versus 0%, $p < 0.001$) than the cefotaxime susceptible isolates (Table 2). Compared to ESC-resistant *S. sonnei* isolates, the ESC-resistant *S. flexneri* isolates were significantly

Table 2 Antibiotic resistance of ESC-resistant and susceptible *Shigella* isolates

Antibiotic	Number of isolates resistant to antibiotics (%)							ESC-resistant <i>S. flexneri</i> vs ESC-resistant <i>S. sonnei</i> (<i>p</i> value)
	Total (n = 356)	<i>S. flexneri</i> (n = 112)			<i>S. sonnei</i> (n = 244)			
		ESC-resistant (n = 58)	Cefotaxime susceptible (n = 54)	ESC-resistant vs cefotaxime susceptible (<i>p</i> value)	ESC-resistant (n = 140)	Cefotaxime susceptible (n = 104)	ESC-resistant vs cefotaxime susceptible (<i>p</i> value)	
AMP	348 (97.8)	58 (100)	51(94.4)	0.069	140(100)	99(95.2)	0.009	NA
PIP	298 (83.7)	58(100)	46(85.2)	0.002	140(100)	54 (51.9)	<0.001	NA
SAM	136 (38.2)	56(96.6)	46(85.2)	0.035	34(24.3)	0 (0.0)	<0.001	<0.001
TZP	1 (0.3)	1 (1.7)	0 (0.0)	3.332	0 (0.0)	0 (0.0)	NA	0.119
CAZ	59 (16.6)	34(58.6)	0 (0.0)	<0.001	25(17.9)	0 (0.0)	<0.001	<0.001
FEP	66 (18.5)	34(58.6)	0 (0.0)	<0.001	32(22.9)	0 (0.0)	<0.001	<0.001
FOX	8 (2.2)	3 (5.2)	0 (0.0)	0.090	5 (3.6)	0 (0.0)	0.056	0.618
IPM	0(0.0)	0(0)	0 (0.0)	NA	0 (0.0)	0 (0.0)	NA	NA
SXT	304 (85.4)	43(74.1)	25(46.3)	0.003	135(96.4)	101(97.1)	0.766	<0.001
CIP	86 (24.2)	43 (74.1)	39 (72.2)	0.819	3 (2.1)	1 (1.0)	0.485	<0.001
LEV	57 (16.0)	31(53.4)	20(37.0)	0.081	3 (2.1)	3 (2.9)	0.692	<0.001
β-lactam antibiotic + SXT + CIP or LEV	69 (19.4)	36 (62.1)	23 (42.6)	0.039	6 (4.3)	4 (3.8)	0.864	<0.001

AMP, ampicillin; PIP, piperacillin; SAM, ampicillin/sulbactam; TZP, piperacillin/tazobactam; FEP, cefepime; FOX, ceftioxitin; IMP, imipenem; SXT, trimethoprim/sulfamethoxazole; CIP, ciprofloxacin; LEV, levofloxacin.
p < 0.05 were considered statistically significant.
 NA, not available.

more likely to be resistant to CAZ (58.6% versus 17.9%, *p* < 0.001), FEP (58.6% versus 22.9%, *p* < 0.001), CIP (74.1% versus 2.1%, *p* < 0.001) and LEV (53.4% versus 2.1%, *p* < 0.001), and less likely to be resistant to SXT (74.1% versus 96.4%, *p* < 0.001). No *Shigella* isolates were resistant to IMP, and Only one *S. flexneri* isolates was resistant to TZP (Table 2).

Multiresistance among ESC-resistant *Shigella* isolates

Among the 356 *Shigella* isolates, 69 (36 ESC-resistant *S. flexneri*, 23 cefotaxime susceptible *S. flexneri*, 6 ESC-resistant *S. sonnei* and 4 cefotaxime susceptible *S. sonnei*) were multidrug resistant (MDR, resistant to 3 or more classes of antimicrobial agents), and the multi-resistance was more frequent among *S. flexneri* isolates than among *S. sonnei* isolates (Table 2).

Production of ESBL and AmpC

The phenotypic confirmatory test displayed that 96.0% (190/198) ESC-resistant isolates were ESBLs positive (Table 3). Of 8 ceftioxitin resistant or intermediate isolates, 2 were the three-dimensional extract test positive.

Distribution of ESBL and AmpC genes

Of the 198 ESC-resistant isolates, 105 (53.0%) were positive for *bla*_{CTX-M-14}, 28 (14.1%) for *bla*_{CTX-M-15}, 22 (13.6%) for *bla*_{CTX-M-57}, 8 (4.0%) for *bla*_{CTX-M-28}, 7 (3.3%) for *bla*_{CTX-M-22}, 52 (26.3%) for *bla*_{OXA-30}, 1 (0.5%) for *bla*_{CMY-2}, and 1 (0.5%) for *bla*_{DHA-1}; 28 (14.1%) isolates carried non-ESBL gene *bla*_{TEM-1}; 8 (4.0%) strains were negative for all β-lactamase genes detected. No isolate hosted *bla*_{SHV}, *bla*_{ACC}, *bla*_{EBC} and *bla*_{FOX} genes. And 44 (23.2%) isolates were found to harbor ≥2 β-lactamase genes (Table 3).

Distribution of virulence genes

All ESC-resistant *Shigella* isolates harbored the *ial*, *ipaH* and *virA* genes. The gene *sen* existed in 99.0% *Shigella* isolates. The genes *set1A* and *set1B* were more prevalent in *S. flexneri* strains than in *S. sonnei* isolates (*p* < 0.05) (Table 4). Of 198 ESC-resistant *Shigella* strains, 123 (*S. sonnei* isolates accounting for 92.7%, 114/123) possessed the most prevalent virulence gene combination of *ial* + *ipaH* + *virA* + *sen*, followed by the combination of *ial* + *ipaH* + *virA* + *set1A* + *set1B* + *sen* (*S. flexneri* isolates accounting for 86.8%, 46/53) (Table 4).

Table 3 Distribution of genotypes of ESBL and AmpC in 198 ESC-resistant *Shigella* isolates

ESBL phenotypic confirmatory test (n)	Genotype		<i>S. flexneri</i> (n = 58)	<i>S. sonnei</i> (n = 140)	Total (n = 198)	
	ESBL or AmpC group	ESBL or AmpC gene				
Positive (n = 190)	CTX-M-1	CTX-M-15	5	11	16	
		CTX-M-28	2	6	8	
		CTX-M-57	7	2	9	
		CTX-M-22	3	2	5	
	CTX-M-9	CTX-M-14	16	69	85	
		TEM	TEM-1	2	4	6
	OXA	OXA-30	4	11	15	
		CTX-M-1+CTX-M-9+OXA	CTX-M-15+CTX-M-14+OXA-30	1	0	1
	CTX-M-1+ OXA+TEM	CTX-M-22+OXA-30+TEM-1	0	2	2	
		CTX-M-57+ OXA-30+TEM-1	9	0	9	
	CTX-M-1+ TEM	CTX-M-57+ TEM-1	2	0	2	
	CTX-M-1+ OXA	CTX-M-57+ OXA-30	1	1	2	
	CTX-M-9+ OXA+TEM	CTX-M-15+ OXA-30	2	8	10	
		CTX-M-14+ OXA-30+ TEM-1	1	3	4	
	CTX-M-9+ OXA	CTX-M-14+ OXA-30	2	6	8	
	CTX-M-9+ TEM	CTX-M-14+ TEM-1	0	6	6	
	Not detect the ESBL or AmpC genotype			0	2	2
	Negative (n = 8)	CTX-M-9+CIT	CTX-M-14+CMY-2	0	1	1
		CTX-M-1+DHA+OXA	CTX-M-15+OXA-30+DHA-1	1	0	1
		Not detect the ESBL or AmpC genotype			0	6

ERIC-PCR

The 58 ESC-resistant *S. flexneri* isolates were classified into 8 types by ERIC-PCR. The distribution of isolates based on this epidemiological analysis was as follows: 26 isolates in type A, 11 isolates in type B, 5 isolates in type C, 4 isolates each in type D and type E, 3 isolates each in type F and type G, and 2 isolates in type H. Among

the 140 ESC-resistant *S. sonnei* isolates, 13 different genotypes (type A to M) were obtained. The major genotype was type A (66 isolates), followed by type B (19 isolates), type C (12 isolates), type D (8 isolates), type E (7 isolates), type F (7 isolates), type G (6 isolates), type H (6 isolates), type I (3 isolates), type J (2 isolates), type K (2 isolates), type L (1 isolates) and type M (1 isolates).

Table 4 Prevalence and combination of virulence genes among 198 ESC-resistant *Shigella* isolates

Isolate	Serotype	No. (%) of isolates that produced the following:						No. (%) of isolates that contained the following combination of genes				
		<i>ial</i>	<i>ipaH</i>	<i>virA</i>	<i>set1A</i>	<i>set1B</i>	<i>sen</i>	I	II	III	IV	V
<i>S. flexneri</i> (n = 58)	1a (n = 2)	2 (3.4)	2 (3.4)	2 (3.4)	0 (0.0)	0 (0.0)	2 (3.4)	0 (0.0)	2 (3.4)	0 (0.0)	0 (0.0)	0 (0.0)
	2a (n = 37)	37(63.8)	37 (63.8)	37 (63.8)	36 (62.1)	36 (62.1)	37 (63.8)	0 (0.0)	1 (1.7)	0 (0.0)	0 (0.0)	36 (62.1)
	2b (n = 2)	2 (3.4)	2 (3.4)	2 (3.4)	0 (0.0)	0 (0.0)	2 (3.4)	0 (0.0)	2(3.4)	0 (0.0)	0 (0.0)	0 (0.0)
	2c (n = 5)	5 (8.6)	5 (8.6)	5 (8.6)	0 (0.0)	0 (0.0)	5 (8.6)	0 (0.0)	5 (8.6)	0 (0.0)	0 (0.0)	0 (0.0)
	4c (n = 12)	12(20.7)	12 (20.7)	12 (20.7)	10 (17.2)	10 (17.2)	12 (20.7)	0 (0.0)	2 (3.4)	0 (0.0)	0 (0.0)	10 (17.2)
<i>S. sonnei</i> (n = 140)		58 (100)	58 (100)	58 (100)	46 (79.3)	46 (79.3)	58 (100)	0 (0.0)	12 (20.7)	0 (0.0)	0 (0.0)	46 (79.3)
		140(100)	140(100)	140(100)	12 (8.6)	19 (13.6)	138(98.6)	2 (1.4)	114 (81.4)	5 (3.6)	12 (8.6)	7 (5.0)
<i>S. flexneri</i> vs <i>S. sonnei</i> (p value)		NA	NA	NA	<0.001	0.003	0.360	0.364	<0.001	0.603	0.021	<0.001
Total		198(100)	198(100)	198(100)	58 (29.3)	65 (32.8)	196 (99.0)	2 (1.0)	123(62.1)	8 (4.0)	12 (6.1)	53 (26.8)

p < 0.05 were considered statistically significant.

I, *ia1* + *ipaH* + *virA*; II, *ia1* + *ipaH* + *virA* + *sen*; III, *ia1* + *ipaH* + *virA* + *set1A* + *sen*; IV, *ia1* + *ipaH* + *virA* + *set1B* + *sen*; V, *ia1* + *ipaH* + *virA* + *set1A* + *set1B* + *sen*. NA, not available.

Discussion

This investigation reported the prevalence and resistance of ESC-resistant *Shigella*, and the molecular analysis of cephalosporin resistance genes and virulence determinants in clinical isolates from Xiaoshan District, Hangzhou, China collected over a period of 5 years. In the economic undeveloped regions, *S. flexneri* is the most frequently isolated *Shigella* species. A similar situation also exists in China, according to previous data [2,4,6,16-18]. However, in our study *S. sonnei* was the most common cause of bacterial dysentery, which was consistent with the findings in industrialized countries. In recent years, the data from Kaengkhoi District of Thailand, Ho Chi Minh City of Vietnam, South Korea, Taiwan and the East, North and Northeast regions of China, which are the newly industrialized regions, also displayed a striking species shift from *S. flexneri* to *S. sonnei* [18-22]. Therefore, the species transition of shigellosis in the present study may be related to the economic growth in Xiaoshan District, suburban of Hangzhou city with higher economic indicators. Of course, other factors may also play a role and need further researches.

Through the resistance data analysis, we found that more than half of the *S. flexneri* and *S. sonnei* isolates were resistant to ESC (cefotaxime). Data from 8 Asian countries showed a high prevalence of resistance to the frontline antibiotics AMP (53.0%) and SXT (81.0%) among *Shigella* isolates [21]. However, the resistance rates to the both drugs among all our *Shigella* isolates were more higher (AMP, 97.8%; SXT, 85.4%), and were consistent with the data from other investigations, mainland China [6,16,17]. The data from Table 2 showed that SAM was not appropriate for the treatment of diarrhea caused by *S. flexneri*, regardless of the strains with ESC-resistance (96.6%) or susceptibility (85.2%); On the contrary, it can be used to prescribe for *S. sonnei* infections, especially for the cefotaxime susceptible strains. Although the resistance rate of *Shigella* to PIP was high (83.7%) in this study, TZP was of very high anti-*Shigella* activity (Table 2).

If the inhibition zone diameter of CAZ is ≥ 21 mm or that of FEP ≥ 18 mm, the two antibiotics can be reported susceptibility for enterobacteriaceae, irrespective of the isolates producing ESBLs or not, according to the CLSI [10]. In all *Shigella* isolates studied, a similar results of resistance to CAZ and FEP (16.6% and 18.5%) were observed, which were higher than that reported by Yang et al. (5.2% and 6.5%) [16]. However, the resistance rates in Table 2 indicated that the two antibiotics were more suitable for the empiric therapy of the infection of ESC-resistant *S. sonnei* than that of ESC-resistant *S. flexneri* infection.

Fluoroquinolones are the popular antibiotics for the treatment of serious shigellosis in both adults and children. Research results from Gu et al. [23] showed the

resistance rate to CIP was 29.1% between 2007 and 2009 in the Asia-Africa area. Data from Henan Province, China displayed that 21% and 79% of *S. flexneri* strains showed high- or low-level resistance to CIP, respectively [6]. Yang et al. [16] reported that 27.9% and 9.7% *Shigella* were resistance to CIP and LEV, respectively, in Anhui province, China. In our study, a similar resistance rate to CIP (24.2%) and a higher resistance rate to LEV (16.0%) were exhibited (Table 2). Among the fluoroquinolone resistant isolates, 95.3% (82/86, resistance to CIP) and 89.5% (51/57, resistance to LEV) strains belonged to *S. flexneri*. The possible cause was that *S. flexneri* isolates often possessed plasmid-mediated quinolone resistance (PMQR) determinants or mutations in quinolone resistance-determining regions (QRDR) of gyrase and topoisomerase genes. This situation had been described in isolates from other regions of China by Zhang et al. [4], Zhu et al. [24] and Pu et al. [25].

So far, at least 109 variants of CTX-M enzymes (CTX-M-1 to 124) have been described. Of these CTX-Ms, 19 variants (CTX-M-15, 16, 19, 23, 25, 27, 32, 35, 37, 40, 42, 53, 54, 55, 57, 58, 62, 64, 82, 93) exhibit the increased hydrolysis activity against ceftazidime, and the others display a much higher rate of hydrolysis of cefotaxime than ceftazidime [26]. CTX-M-15 is the most usually detected CTX-M variant that hydrolyze ceftazidime at high level in enterobacteriaceae [26]. In this study, 28 *bla*_{CTX-M-15} positive ESC-resistant *Shigella* isolates were all resistant to ceftazidime (data not shown). No other CTX-M variant genes mediating high level ceftazidime resistance were found (Table 3). In the *bla*_{CTX-M} genes with higher catalytic efficiencies against cefotaxime than ceftazidime, *bla*_{CTX-M-14} was the most prevalent one (53.0%), and coincides with the data published worldwide in clinically important pathogens [26]. OXA-30 belongs to the class D oxacillinase group III, and mediates resistance to cefepime but not ceftazidime [27]. Unfortunately, 52 (26.3%) of our ESC-resistant *Shigella* isolates hosted *bla*_{OXA-30} and 12 of them carried *bla*_{CTX-M-15} concomitantly and conferred resistance to cefotaxime, ceftazidime and cefepime (Table 3). In the past decade, an emergence of ESBL-producing *Shigella* spp. carrying different types of ESBL genes has been described in different countries and regions [28]. However, only a few studies have reported in the world the existence of AmpC β -lactamases encoded by *bla*_{CMY-2} or *bla*_{DHA-1} in *Shigella* spp. [4,25,28-31]. In this study, we also found 2 AmpC β -lactamase producers with *bla*_{DHA-1} and *bla*_{CMY-2} in the three-dimensional extract test positive *Shigella* strains. The *bla*_{CMY-2} and the *bla*_{DHA-1} existed in 1 *S. flexneri* with *bla*_{CTX-M-14} and 1 *S. sonnei* with *bla*_{CTX-M-15} and *bla*_{OXA-30}, respectively (Table 3).

In this study, we detected several pathogenic genes (*ial*, *ipaH*, *set1*, *sen* and *virA*) for 198 ESC-resistant *Shigella* isolates (Table 4). It has been shown that *ial*

takes responsibility for penetration of epithelial cell by *Shigella* and *ipaH* also for spread from cell to cell [32,33]. All *Shigella* species studied were positive for the *ipaH* as expected because this gene exists in multiple copies on both the chromosome and the plasmid of *Shigella*. Conversely, the *ial* gene is exclusively located on the plasmid and was only detected in some *Shigella* isolates [8]. Indeed, less frequent examination of *ial* gene had been described by Luscher and Altwegg [34], Kingombe et al. [35] and Thong et al. [36]. However, this gene was found in all our ESC-resistant *Shigella* strains. Another virulence factor VirA is involved in the uptake, motility, and cell to cell transmission of *Shigella* within the human host. It is an essential virulence factor in *Shigella* disease pathogenesis [9]. The positive rate of *virA* implied all the isolates of our collection might have the ability (Table 4). The *set1* chromosomal gene encodes *Shigella* enterotoxin 1 (ShET-1, composed of one A and five B subunits), which is generated by *S. flexneri* (mainly in type 2a) and not found in other *Shigella* spp. [8,12,36-41]. The *sen* gene encoding *Shigella* enterotoxin 2 (ShET-2) is carried on a 140 MDa virulence plasmid. And the *sen* is present in all *Shigella* species. The both toxins are deemed to play a part in the clinical manifestation of shigellosis [8]. In our study, 79.3% of ESC-resistant *S. flexneri* strains were found to be *set1A* and *set1B* positive (62.1% isolates were serotype f2a, Table 4), and this is agreement with the previous results; however, 17.1% (24/140) ESC-resistant *S. sonnei* isolates also carried *set1A* and/or *set1B* genes (Table 4). The *set1A* and *set1B* genes are located on the *she* pathogenicity island (PAI), a chromosomal, laterally acquired, integrative element of *S. flexneri* [37]. Integrase-mediated excision can occur for the *she* PAI, and result in the formation of a circular excision product, which is a substrate for lateral transfer processes e.g. conjugation, packaging into phage particles and recombinase-mediated integration into the chromosome [42]. This may be the cause that the two determinants can be found in our ESC-resistant *S. sonnei* isolates studied. And the deficiency of *set1A* or *set1B*, or the existence of point mutations in the primer binding sites may be the possible explanation for the both genes not coexisting in some ESC-resistant *S. sonnei* isolates (Table 4). Furthermore, we found that the *S. flexneri* isolates with *set1* were more resistance to CIP, LEV ($p < 0.001$, each) and FEP ($p = 0.019$) than those without *set1*; for *S. sonnei*, the *set1* positive isolates were more likely resistance to SAM ($p < 0.001$), CIP, LEV ($p < 0.001$, each) and FEP ($p = 0.002$), and more likely sensitivity to CAZ ($p = 0.005$) than *set1* negative ones (data not shown). Nevertheless, we do not think there was correlation between *set1* and the antibiotics resistance, because none of reports have described currently that the invasion genetic elements carrying virulence genes known contain

resistance determinants simultaneously in *Shigella* spp.. These resistance differences may be only due to the spread of resistant plasmids among the different strains.

The results of ERIC-PCR typing showed that most of the cases of ESC-resistant *S. flexneri* and *S. sonnei* infections were caused by several identical strains, respectively [for *S. flexneri* isolates, the percentage of type A and type B accounting for 63.8% (37/58); for *S. sonnei*, type A, type B and type C encompassing 69.3% (97/140) of the isolates]. This indicates that clonal dissemination was likely to contribute most to the spread of ESC-resistant *S. flexneri* and *S. sonnei* within the region studied. Of 46 ESC-resistant *S. flexneri* isolates with the second largest number of the virulence gene composition (*ial* + *ipaH* + *virA* + *set1A* + *set1B* + *sen*) being resistant to 4 to 8 antibiotics, 60.3% belonged to type A (43.1%) and type B (17.2%). Of 114 ESC-resistant *S. sonnei* isolates with the first largest number of the virulence gene composition (*ial* + *ipaH* + *vir* + *sen*) being resistant to 3 to 6 antibiotics, 71.1% belonged to type A (57.9%) and type B (13.2%), but isolates that hosted the *set1* gene were more heterogeneous in ERIC-PCR pattern.

In the present study all the *Shigella* strains were isolated from intestinal clinic patients. No patients had been hospitalized or died following episodes of shigellosis. According to the clinical reports, early stage of shigellosis the patients infected by ESBL genes positive isolates were not more severe than those of the patients infected by ESBL genes negative isolates. However, most of these patients had longer course of treatment, because physicians used to treat diarrhea by prescribing cefotaxime or ceftriaxone (particularly for children) in the region studied. When treatment failed, other drugs (such as fluoroquinolone or β -lactamase inhibitors) would be used as a substitute for continuing treatment. In addition, all the patients with *Shigella* infection in the study were treated with antibiotics, therefore, we had no relevant data to compare the course of disease treated using antibiotics with that of disease treated without antibiotics.

Conclusions

This investigation found a striking difference from other regions of China with relatively low economic status in the distribution of species, from *S. flexneri* to *S. sonnei*, and a high prevalence of ESC-resistance among *Shigella* isolates. Antibiotic susceptibility testing displayed ESC-resistant *S. flexneri* isolates were much more resistant than ESC-resistant *S. sonnei* isolates. The analysis of extended-spectrum cephalosporin resistance genes showed the spread of the *Shigella* harboring a combination of genes *bla*_{CTX-M-15} and *bla*_{OXA-30}. This combination may be a serious threat to infection treatment because it is spread by conjugation and it displays high levels of resistance to cefotaxime, ceftazidime and cefepime. We also described the

prevalence of virulence genes in ESC-resistant *Shigella*, which may provide clues about the pathogenicity difference of this type of bacteria. ERIC-PCR typing supported the existence of specific clones responsible for the ESC-resistant *S. flexneri* and *S. sonnei* infection cases in the region studied. In China, especially in some undeveloped regions, there were limited and sporadic data on the epidemiology of *Shigella* spp. Therefore, we suggest the Ministry of Health National Antimicrobial Resistance Investigation Net (Mohnarin) to reinforce the surveillance for the development of antibiotic resistance and take measures to control shigellosis in China.

Abbreviations

ESC: Extended-spectrum cephalosporin; ESBL: Extended-spectrum beta-lactamase; *S. flexneri*: *Shigella flexneri*; *S. sonnei*: *Shigella sonnei*; CLSI: Clinical laboratory standards institute; PCR: Polymerase chain reaction; ERIC-PCR: Enterobacterial repetitive intergenic consensus sequence polymerase chain reaction; SPSS: Statistical package for social sciences.

Competing interests

The authors declare no conflict of interests.

Authors' contributions

CLZ performed the experiments; QZL designed the study, analyzed the clinical data and wrote this manuscript; JW, XC, LMS and YYG collected the clinical samples. All authors read and approved the final manuscript.

Acknowledgments

The authors thank Yunkui Zong for critical review of the manuscript. This work was supported by grants from Natural Science Foundation, Science and Technology Commission of Shanghai (No. 12ZR1425000), the National Natural Science Foundation of China (No. 81371872), and Xiaoshan District, Major Scientific and Technological Research Projects (No. 2013314).

Author details

¹Department of Clinical Laboratory, Zhejiang Xiaoshan Hospital, Zhejiang Province, China. ²Department of Clinical Laboratory, Shanghai First People's Hospital, School of Medicine, Shanghai Jiaotong University, 100 Haining Road, Shanghai 200080, People's Republic of China.

Received: 7 December 2013 Accepted: 8 May 2014

Published: 15 May 2014

References

1. Kuo CY, Su LH, Perera J, Carlos C, Tan BH, Kumarasinghe G, So T, Van PH, Chongthaleong A, Song JH, Chiu CH: **Antimicrobial susceptibility of *Shigella* isolates in eight Asian countries, 2001–2004.** *J Microbiol Immunol Infect* 2008, **41**:107–111.
2. Wang XY, Tao F, Xiao D, Lee H, Deen J, Gong J, Zhao Y, Zhou W, Li W, Shen B, Song Y, Ma J, Li ZM, Wang Z, Su PY, Chang N, Xu JH, Ouyang PY, von Seidlein L, Xu ZY, Clemens JD: **Trend and disease burden of bacillary dysentery in China (1991–2000).** *Bull World Health Organ* 2006, **84**:561–568.
3. Ji WJ, Xu XW, Dong F: **Genotype and drug resistance analysis of extended-spectrum beta-lactamases-producing *Shigella* in pediatric patients.** *Chin J Microbiol Immunol* 2010, **30**:472–476 (in Chinese).
4. Zhang W, Luo Y, Li J, Lin L, Ma Y, Hu C, Jin S, Ran L, Cui S: **Wide dissemination of multidrug-resistant *Shigella* isolates in China.** *J Antimicrob Chemother* 2011, **66**:2527–2535.
5. Zhang R, Zhou HW, Cai JC, Zhang J, Chen GX, Nasu M, Xie XY: **Serotypes and extended-spectrum β -lactamase types of clinical isolates of *Shigella* spp. from the Zhejiang province of China.** *Diagn Microbiol Infect Dis* 2011, **69**:98–104.
6. Xia S, Xu B, Huang L, Zhao JY, Ran L, Zhang J, Chen H, Pulsrikarn C, Pornruangwong S, Aarestrup FM, Hendriksen RS: **Prevalence and characterization of human *Shigella* infections in Henan Province, China, in 2006.** *J Clin Microbiol* 2011, **49**:232–242.
7. Xiong Z, Li T, Xu Y, Li J: **Detection of CTX-M-14 extended-spectrum beta-lactamase in *Shigella sonnei* isolates from China.** *J Infect* 2007, **55**:e125–e128.
8. Sousa MÃ, Mendes EN, Collares GB, Pêret-Filho LA, Penna FJ, Magalhães PP: ***Shigella* in Brazilian children with acute diarrhoea: prevalence, antimicrobial resistance and virulence genes.** *Mem Inst Oswaldo Cruz* 2013, **108**:30–35.
9. Yoshida S, Handa Y, Suzuki T, Ogawa M, Suzuki M, Tamai A, Abe A, Katayama E, Sasakawa C: **Microtubule-severing activity of *Shigella* is pivotal for intercellular spreading.** *Science* 2006, **314**:985–989.
10. Clinical and Laboratory Standards Institute: **Performance standards for antimicrobial susceptibility testing; Twenty-second Informational supplement, Document M100-S22.** Wayne, PA: CLSI; 2012.
11. Coudron PE, Moland ES, Thomson KS: **Occurrence and detection of AmpC beta-lactamases among *Escherichia coli*, *Klebsiella pneumoniae*, and *Proteus mirabilis* isolates at a Veterans Medical Center.** *J Clin Microbiol* 2000, **38**:1791–1796.
12. Vargas M, Gascon J, Jimenez De Anta MT, Vila J: **Prevalence of *Shigella* enterotoxins 1 and 2 among *Shigella* strains isolated from patients with traveler's diarrhea.** *J Clin Microbiol* 1999, **37**:3608–3611.
13. Villalobo E, Torres A: **PCR for detection of *Shigella* spp. in Mayonnaise.** *Appl Environ Microbiol* 1998, **64**:1242–1245.
14. Pérez-Pérez FJ, Hanson ND: **Detection of plasmid-mediated AmpC-lactamase genes in clinical isolates by using mitplex PCR.** *J Clin Microbiol* 2002, **40**:2153–2162.
15. Kumao T, Ba-Thein W, Hayashi H: **Molecular subtyping methods for detection of *Salmonella enterica* serovar Oranienburg outbreaks.** *J Clin Microbiol* 2002, **40**:2057–2061.
16. Yang H, Chen G, Zhu Y, Liu Y, Cheng J, Hu L, Ye Y, Li J: **Surveillance of antimicrobial susceptibility patterns among *Shigella* species isolated in China during the 7-year period of 2005–2011.** *Ann Lab Med* 2013, **33**:111–115.
17. Shen Y, Qian H, Gong J, Deng F, Dong C, Zhou L, Guo H: **High prevalence of antibiotic resistance and molecular characterization of integrons among *Shigella* isolates in Eastern China.** *Antimicrob Agents Chemother* 2013, **57**:1549–1551.
18. Chang Z, Lu S, Chen L, Jin Q, Yang J: **Causative species and serotypes of shigellosis in mainland China: systematic review and meta-analysis.** *PLoS ONE* 2012, **7**:e52515.
19. von Seidlein L, Kim DR, Ali M, Lee H, Wang X, Thiem VD, Canh do G, Chaicumpa W, Agtini MD, Hossain A, Bhutta ZA, Mason C, Sethabutr O, Talukder K, Nair GB, Deen JL, Kotloff K, Clemens J: **A multicentre study of *Shigella* diarrhoea in six Asian countries: disease burden, clinical manifestations, and microbiology.** *PLoS Med* 2006, **3**:e353.
20. Vinh H, Nhu NT, Nga TV, Duy PT, Campbell JJ, Hoang NV, Boni MF, My PV, Parry C, Nga TT, Van Minh P, Thuy CT, Diep TS, Phuong le T, Chinh MT, Loan HT, Tham NT, Lanh MN, Mong BL, Anh VT, Bay PV, Chau NV, Farrar J, Baker S: **A changing picture of shigellosis in southern Vietnam: shifting species dominance, antimicrobial susceptibility and clinical presentation.** *BMC Infect Dis* 2009, **9**:204.
21. Seol SY, Kim YT, Jeong YS, Oh JY, Kang HY, Moon DC, Kim J, Lee YC, Cho DT, Lee JC: **Molecular characterization of antimicrobial resistance in *Shigella sonnei* isolates in Korea.** *J Med Microbiol* 2006, **55**:871–877.
22. Wei HL, Wang YW, Li CC, Tung SK, Chiou CS: **Epidemiology and evolution of genotype and antimicrobial resistance of an imported *Shigella sonnei* clone circulating in central Taiwan.** *Diagn Microbiol Infect Dis* 2007, **58**:469–475.
23. Gu B, Cao Y, Pan S, Zhuang L, Yu R, Peng Z, Qian H, Wei Y, Zhao L, Liu G, Tong M: **Comparison of the prevalence and changing resistance to nalidixic acid and ciprofloxacin of *Shigella* between Europe-America and Asia-Africa from 1998 to 2009.** *Int J Antimicrob Agents* 2012, **40**:9–17.
24. Zhu YL, Yang HF, Liu YY, Hu LF, Cheng J, Ye Y, Li JB: **Detection of plasmid-mediated quinolone resistance determinants and the emergence of multidrug resistance in clinical isolates of *Shigella* in Sixian area, China.** *Diagn Microbiol Infect Dis* 2013, **75**:327–329.
25. Pu XY, Pan JC, Wang HQ, Zhang W, Huang ZC, Gu YM: **Characterization of fluoroquinolone-resistant *Shigella flexneri* in Hangzhou area of China.** *J Antimicrob Chemother* 2009, **63**:917–920.
26. Zhao WH, Hu ZQ: **Epidemiology and genetics of CTX-M extended-spectrum β -lactamases in Gram-negative bacteria.** *Crit Rev Microbiol* 2013, **39**:79–101.
27. Dubois V, Arpin C, Quentin C, Texier-Maugein J, Poirel L, Nordmann P: **Decreased susceptibility to cefepime in a clinical strain of *Escherichia coli* related to plasmid- and integron-encoded OXA-30 beta-lactamase.** *Antimicrob Agents Chemother* 2003, **47**:2380–2381.

28. Taneja N, Mewara A, Kumar A, Verma G, Sharma M: **Cephalosporin-resistant *Shigella flexneri* over 9 years (2001–09) in India.** *J Antimicrob Chemother* 2012, **67**:1347–1353.
29. Livermore DM, Mushtaq S, Nguyen T, Warner M: **Strategies to overcome extended-spectrum β -lactamases (ESBLs) and AmpC β -lactamases in *shigellae*.** *Int J Antimicrob Agents* 2011, **37**:405–409.
30. Huang IF, Chiu CH, Wang MH, Wu CY, Hsieh KS, Chiou CC: **Outbreak of dysentery associated with ceftriaxone-resistant *Shigella sonnei*: First report of plasmid-mediated CMY-2-type AmpC beta-lactamase resistance in *S. sonnei*.** *J Clin Microbiol* 2005, **43**:2608–2612.
31. Tajbakhsh M, García Migura L, Rahbar M, Svendsen CA, Mohammadzadeh M, Zali MR, Aarestrup FM, Hendriksen RS: **Antimicrobial-resistant *Shigella* infections from Iran: an overlooked problem?** *J Antimicrob Chemother* 2012, **67**:1128–1133.
32. Wei J, Goldberg MB, Burland V, Venkatesan MM, Deng W, Fournier G, Mayhew GF, Plunkett G 3rd, Rose DJ, Darling A, Mau B, Perna NT, Payne SM, Runyen-Janecky LJ, Zhou S, Schwartz DC, Blattner FR: **Complete genome sequence and comparative genomics of *Shigella flexneri* serotype 2a strain 2457T.** *Infect Immun* 2003, **71**:2775–2786.
33. Parsot C: ***Shigella* spp and enteroinvasive *Escherichia coli* pathogenicity factors.** *FEMS Microbiol Lett* 2005, **252**:11–18.
34. Lüscher D, Altwegg M: **Detection of *shigellae*, enteroinvasive and enterotoxigenic *Escherichia coli* using the polymerase chain reaction (PCR) in patients returning from tropical countries.** *Mol Cell Probes* 1994, **8**:285–290.
35. Kingombe CI, Cerqueira-Campos ML, Farber JM: **Molecular strategies for the detection, identification, and differentiation between enteroinvasive *Escherichia coli* and *Shigella* spp.** *J Food Prot* 2005, **68**:239–245.
36. Thong KL, Hoe SL, Puthucherry SD, Yasin R: **Detection of virulence genes in Malaysian *Shigella* species by multiplex PCR assay.** *BMC Infect Dis* 2005, **5**:8.
37. Niyogi SK, Vargas M, Vila J: **Prevalence of the *sat*, *set* and *sen* genes among diverse serotypes of *Shigella flexneri* strains isolated from patients with acute diarrhoea.** *Clin Microbiol Infect* 2004, **10**:574–576.
38. Noriega FR, Liao FM, Formal SB, Fasano A, Levine MM: **Prevalence of *Shigella* enterotoxin 1 among *Shigella* clinical isolates of diverse serotypes.** *J Infect Dis* 1995, **172**:1408–1410.
39. Yavzori M, Cohen D, Orr N: **Prevalence of the genes for *Shigella* enterotoxins 1 and 2 among clinical isolates of *Shigella* in Israel.** *Epidemiol Infect* 2002, **128**:533–535.
40. Roy S, Thanasekaran K, Dutta Roy AR, Sehgal SC: **Distribution of *Shigella* enterotoxin genes and secreted autotransporter toxin gene among diverse species and serotypes of *shigella* isolated from Andaman Islands, India.** *Trop Med Int Health* 2006, **11**:1694–1698.
41. Farfán MJ, Garay TA, Prado CA, Filliol I, Ulloa MT, Toro CS: **A new multiplex PCR for differential identification of *Shigella flexneri* and *Shigella sonnei* and detection of *Shigella* virulence determinants.** *Epidemiol Infect* 2010, **138**:525–533.
42. Sakellaris H, Luck SN, Al-Hasani K, Rajakumar K, Turner SA, Adler B: **Regulated site-specific recombination of the *she* pathogenicity island of *Shigella flexneri*.** *Mol Microbiol* 2004, **52**:1329–1336.

doi:10.1186/1471-2334-14-260

Cite this article as: Zhang et al.: Epidemic and virulence characteristic of *Shigella* spp. with extended-spectrum cephalosporin resistance in Xiaoshan District, Hangzhou, China. *BMC Infectious Diseases* 2014 **14**:260.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

