




Review

Coffee Chlorogenic Acids Incorporation for Bioactivity Enhancement of Foods: A Review

Alexis Rojas-González ^{1,2,†}, Claudia Yuritzí Figueroa-Hernández ^{3,†} , Oscar González-Ríos ¹,
Mirna Leonor Suárez-Quiroz ¹, Rosa María González-Amaro ⁴, Zorba Josué Hernández-Estrada ¹ 
and Patricia Rayas-Duarte ^{2,*} 

¹ Tecnológico Nacional de México/Instituto Tecnológico de Veracruz, M.A. de Quevedo 2779, Col. Formando Hogar, Veracruz 91897, Mexico; alexis.rojas@okstate.edu (A.R.-G.); oscar.gr@veracruz.tecnm.mx (O.G.-R.); mirna.sq@veracruz.tecnm.mx (M.L.S.-Q.); zorba.he@veracruz.tecnm.mx (Z.J.H.-E.)

² Robert M. Kerr Food & Agricultural Products Center, Oklahoma State University, 123 FAPC, Stillwater, OK 74078, USA

³ CONACYT-Tecnológico Nacional de México/Instituto Tecnológico de Veracruz, Unidad de Investigación y Desarrollo en Alimentos, M. A. de Quevedo 2779, Veracruz 91897, Mexico; claudia.fh@veracruz.tecnm.mx

⁴ CONACYT-Instituto de Ecología, A.C., Carretera Antigua a Coatepec 351, Col. El Haya, Xalapa, Veracruz 91073, Mexico; rosa.gonzalez@inecol.mx

* Correspondence: pat.rayas_duarte@okstate.edu; Tel.: +1-405-744-6468

† These authors contributed equally to this work.

Abstract: The demand of foods with high antioxidant capacity have increased and research on these foods continues to grow. This review is focused on chlorogenic acids (CGAs) from green coffee, which is the most abundant source. The main CGA in coffee is 5-O-caffeoylquinic acid (5-CQA). Coffee extracts are currently the most widely used source to enhance the antioxidant activity of foods. Due to the solubility of CGAs, their extraction is mainly performed with organic solvents. CGAs have been associated with health benefits, such as antioxidant, antiviral, antibacterial, anticancer, and anti-inflammatory activity, and others that reduce the risk of cardiovascular diseases, type 2 diabetes, and Alzheimer's disease. However, the biological activities depend on the stability of CGAs, which are sensitive to pH, temperature, and light. The anti-inflammatory activity of 5-CQA is attributed to reducing the proinflammatory activity of cytokines. 5-CQA can negatively affect colon microbiota. An increase in anthocyanins and antioxidant activity was observed when CGAs extracts were added to different food matrices such as dairy products, coffee drinks, chocolate, and bakery products. The fortification of foods with coffee CGAs has the potential to improve the functionality of foods.

Keywords: chlorogenic acids; coffee; 5CQA; functional foods; biological activity



Citation: Rojas-González, A.; Figueroa-Hernández, C.Y.; González-Ríos, O.; Suárez-Quiroz, M.L.; González-Amaro, R.M.; Hernández-Estrada, Z.J.; Rayas-Duarte, P. Coffee Chlorogenic Acids Incorporation for Bioactivity Enhancement of Foods: A Review. *Molecules* **2022**, *27*, 3400. <https://doi.org/10.3390/molecules27113400>

Academic Editor: Davide Bertelli

Received: 29 April 2022

Accepted: 20 May 2022

Published: 25 May 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The chlorogenic acids (CGAs) are a class of phenolic compounds widely distributed in various plants sources such as fruits, vegetables, coffee beans, tea, apples, and wine [1–3]. CGAs are esters of quinic acid (QA) and one trans-cinnamic acid residue such as caffeic acid (CA), *p*-coumaric acid (*p*-CoA), and ferulic acid (FA), which are known as caffeoylquinic acids (CQAs), *p*-coumaroylquinic acids (*p*-CoQAs) and feruloylquinic acid (FQAs) [1–5]. Caffeoylquinic acid may theoretically form four isomers, but only three are present in plants: 3-*O*-caffeoylquinic acid (3-CQA), neochlorogenic acid (5-*O*-caffeoylquinic acid, 5-CQA), or cryptochlorogenic acid (4-*O*-caffeoylquinic acid, 4-CQA). The most common isomer, 5-CQA, is an ester composed of caffeic acid and (–)-quinic acid and referred as chlorogenic acid [4,6]. According to the number of caffeoyl groups attached to the quinic acid, these CQAs can be classified into monophosphoylquinic acids (MCQAs), dicaffeoylquinic acids (DCQAs), tricaffeoylquinic acids (TCQAs), and tetracaffeoylquinic acids (tetra-CQAs) [3]. The chemical structures of the main CGAs are shown in Figure 1. The structural diversity and broad bioactivities of CGAs have increasingly attracted the attention of researchers [4,5,7–10]. It

has been demonstrated that these compounds are potent antioxidants and may also exert other physiological activities. For example, there is evidence that CGAs possess a wide variety of bioactivities, such as antiparasitic [11], antibacterial [12], anti-inflammatory [13], neuroprotective [14], anticancer [15], antiglycemic [16], and antiviral [17]. In addition, it has been demonstrated that CGAs have therapeutic effects in the prevention and treatment of some chronic and cardiovascular diseases [5,18,19]. This review aims to describe the main biological activities attributed to coffee CGAs, and their bioavailability and potential addition to different food matrices to obtain functional foods.




Figure 1. Chemical structures of main chlorogenic acids (CGAs) and isomers of caffeoylquinic acid.

2. Dietary Sources of Chlorogenic Acids (CGAs)

CGAs are a large family of esters of quinic acid and trans-cinnamic acids; up to date, at least 71 different chemical compounds are identified from different plants sources such as fruits, vegetables, coffee beans, tea, apples, artichoke, eggplant, and grapes [4,20,21]. However, those found in the highest concentration in plants are caffeoylquinic acids (CQAs), specifically mono- and di-CQAs, as well as the different isomeric forms of feruloylquinic acids (FQAs) [21,22]. Meinhart et al. [23] analyzed the CGAs concentration (CA, 3-CQA, 4-CQA, 5-CQA, 3,4-DQA, 3,5-DQA and 4,5-DQA) of 100 plants commonly used in Brazil as infusions. In their study, the highest concentrations of CGAs were yerba mate (*Ilex paraguariensis*), white and green tea (*Camellia sinensis*), and winter's bark (*Drimys winteri*). A study of the presence of CGAs in 53 vegetables consumed in Southern Brazil reported the highest concentrations of 3-CQA, 5-CQA, and 4-CQA in collard greens and chicory whereas the highest concentration of 3,4-DQA, 3,5-DQA, and 4,5-DQA were found in bay leaves and mustard [20]. At present, green coffee beans and yerba mate are recognized as the most important plant sources of CGAs, accounting for up to 6 to 12% in the case of green coffee and 9% for mate. 5-CQA is the most abundant CGA in green coffee beans, with a concentration of about 100 mg/g (dry basis), representing 76 to 84% of the total content of CGAs [4]. The main food sources of CGAs are shown in Table 1.

Table 1. Main sources of CGAs.

Source	Concentration (g/100 g) ¹ (dm)	CGA Composition	References
Artichoke	1–8	5-CQA, 1,5-DCQA, 3,4-DCQA and DCQA	[24]
Artichoke leaves	0.92	CA, 3-CQA, 4-CQA, 5-CQA, 3,4-DQA, 3,5-DQA and 4,5-DQA	[23]
Sweet potato leaves	-	3-CQA, 3,4-DCQA, 3,5-DCQA, 4,5-DCQA and 3,4,5-TCQA	[15]
White tea (<i>Camelia sinensis</i>) leaves	1.64	3-CQA, 4-CQA, 5-CQA, 3,4-DQA, 3,5-DQA and 4,5-DQA	[23]
Green tea (<i>Camelia sinensis</i>) leaves	1.32	3-CQA, 4-CQA, 5-CQA, 3,4-DQA, 3,5-DQA and 4,5-DQA	[23]
Yerba mate (<i>Ilex paraguariensis</i>) leaves and thalli	9.19	3-CQA, 4-CQA, 5-CQA, 3,4-DQA, 3,5-DQA and 4,5-DQA	[23]
Green coffee beans	4.10–11.3 ²	CQA, FQA and DCQA	[25]
Apples	0.38 0–0.2 g/L (juice)	3-CQA, 5-CQA, 4,5-DCQA	[26]
Pears	0.28 0–0.24 g/L (juice)	3-CQA, 5-CQA, 3,6-DCQA	[27]
Blueberries	2	5-CQA, 3-FQA	[28]
Grapes	0.15	5-CQA, CoQA	[29]
Spinach	0.2	<i>p</i> -CoQA	[30]
Beans and peas	0.12	<i>p</i> -CoQA	[31]
Stone fruits	0.01–0.6	<i>p</i> -CoQA, 5-CQA, FQA, 4,5-DCQA, 3,4-TCQA	[32]
Potato tubers	0.5–1.2	CQA; DCQA	[33]

¹ Units may have been changed for consistency and expressed in dry matter (dm). ² It depends on the variety and geographic origin of the coffee.

Coffee as a Source of CGAs

Coffee is one of the most widely consumed beverages in the world. This infusion contains several compounds that can exert beneficial biological activities for human health. Many beneficial effects have been investigated, mainly attributed to caffeine and other substances, such as polyphenols, mainly chlorogenic acids [1,34]. There are at least 30 different types of CGAs found in coffee, and this includes caffeoylquinic acids (CQAs), dicaffeoylquinic acids (DCQAs), tricaffeoylquinic acids (TCQAs), feruloylquinic acids (FQAs), and *p*-coumaroylquinic acids (*p*-CoQAs) [35]. One cup (200 mL) of coffee brew contains between 20–675 mg of CGAs depending on the variety of coffee and brewing method [36].

Whether green or roasted, the beneficial health effects of coffee have been attributed to the high content of CGAs and the antioxidant activity provided by the phenolic compounds in green coffee in addition to the those produced during the roasting process [37]. The concentration of active polyphenols inside green coffee depends on the variety of the bean and its geographical origin; in beverages, it also depends on the brewing process [38]. During the coffee roasting process, phenolic compounds undergo a series of intermolecular and intramolecular reactions and interactions [21]. Higher concentration of CGAs in lightly roasted coffee over dark roasted coffee has been established; however, the highest concentration of CGAs is found in green coffee beans [39]. In green coffee, CQAs alone account for up to 80% of the total CGAs and among CQAs, 5-CQA account for almost 60%.

Thus, 5-CQA is the most studied isomer of the CGAs and is responsible for the bitter and astringent taste in coffee [25].

3. Extraction of Chlorogenic Acids (CGAs) from Coffee

The extraction recovery of a wide variety of compounds from vegetal species is a critical step in the production of bioactive substances. The chemical properties of CGAs such as thermal stability, solubility, and oxidation-reduction reactions, need to be considered when combined with other substances [21]. Studies focused on the development of new extraction methods of CGA's have been made over the last couple of decades, mainly focusing on increasing mass transfer and extraction yields while minimizing the use of toxic organic solvents and energy consumption [40–46].

3.1. Organic Solvent Extraction

Madhava-Naidu et al. [46] extracted green coffee CGAs by sterilizing the beans at 120 °C for 20 min and Soxhlet extraction with hexane at different rates. The samples were then separated in glass columns and extracted with selected solvents at different ratios (80:20, 70:30, and 60:40) using mixtures of isopropanol and water. They obtained the best extraction yield (29.1%) and a CGAs content of 29.7% when they used a 60:40 isopropanol-water ratio in Robusta coffee, whereas in Arabica coffee the yield of extraction and CGAs content were 27.3 and 30.2%, respectively. Suárez-Quiroz et al. [40] compared four different methods of CGAs extraction using different solvents (water, aqueous methanol, aqueous isopropanol, and ethyl acetate). The extract yield values were not significantly different, demonstrating the high solubility of CGAs in organic solvents. CGA's extraction from green coffee (*C. arabica*) using water at 80 °C and activated carbon were not significantly different from the values of the previous investigation [47]. Thus, activated carbon is a suitable and more eco-friendly extraction method with a minimum of 97% CGAs purity in the extract was reported in this study [47].

Dibert et al. [48] tested the effect of different physicochemical parameters (temperature, particle diameter, and solvent-mass ratio) when extracting CGAs from green coffee beans with a methanol-water extraction (70:30 ratio) at three different temperatures (30, 40, and 50 °C). The highest yield of extraction of CGAs (18.1%) was obtained at 40 and 50 °C, when a mass-solvent ratio of 1:4 *w/v* was used. By increasing the mass ratio of green coffee beans an improvement in the yield of extraction of CGAs can be achieved.

3.2. Pulsed Electric Field Extraction

Bilge et al. [45] evaluated the effect of pulsed electric field on green and roasted *C. arabica* beans as a pretreatment by exposing them to monopolar pulses of 2 Hz with an interval of 0.5 s and generating an electric field of 28 kV/10 cm with water at 20 °C. The use of an electric field increased radical scavenging activity up to 31% and 11%, for green and roasted coffee beans, respectively, compared to untreated samples confirming that using electric pulses as a pretreatment before extraction can enhance the phenolic content extraction and reduce Maillard reaction products that occur at high temperatures of extraction and during the coffee roasting process. Phongsupa et al. [44] studied the extraction of CGAs by pulse electric field induction over *C. arabica*. The number of pulses and concentrations for this study was set to 1000 pulses at 5 kV and 62.7% methanol-water solution as solvent. The mass-solvent ratio with the most effective extraction was 0.75 g/mL and 30 s of blending which had the CGAs content of 9.8 µg/g. However, the results obtained in this study showed that an increase in the sample-solvent ratio leads to a higher concentration of CGAs [45].

4. Biological Activities of CGAs

Several studies have associated CGAs with beneficial health properties, such as antioxidant, antiviral, antibacterial, anticancer, and anti-inflammatory activity [4–6,49]. It has also been shown that it can modulate the gene expression of antioxidant enzymes

and reduce the risk of cardiovascular disease by suppressing the expression of P-selectin in platelets [49]. In addition, CGAs can reduce the relative risk of type 2 diabetes and Alzheimer's disease [6,50–54]. The main biological activities attributed to CGAs are shown in Figure 2.




Figure 2. Main biological activities attributed to CGAs.

Some of these properties are well recognized and demonstrated by *in vitro* and *in vivo* studies, such as antioxidant activity. However, other bioactivities of interest in recent years, although not yet well demonstrated, such as the potential anti-obesity [52,55–60] or prebiotic [61–63] properties of CGAs. In addition, it has also been shown that CGAs can modulate the activity of glucose-6-phosphatase, an enzyme involved in glucose metabolism, and therefore it may have a positive effect on diabetes management [64].

Furthermore, it is important to highlight that these biological activities are dependent on the CGA's stability. CGAs are particularly susceptible to environmental conditions, such as solvent type, pH, temperature, and light. These factors must be considered during the CGAs extraction. Moreover, the concentration of these compounds in plants is low. For this reason, the methodologies used for the CGAs extraction from plant sources must be efficient to guarantee the necessary concentration of CGAs to exert their biological activity.

4.1. Antioxidant Activity

There is a strong correlation between oxidative stress and the development of various degenerative diseases such as cancer and other aging-related diseases [65,66]. Extensive *in vitro* and *in vivo* studies have been performed to evaluate the antioxidant activity of CGAs [67]. As a result, CGAs are known to exhibit a radical scavenging effect similar to ascorbic acid [68]. In addition, CGAs can chelate transition metals such as Fe^{2+} to scavenge free radicals and disrupt chain reactions [21]. Studies have shown that CGAs may prevent the oxidation of low-density lipoproteins (LDL) induced by different oxidizing agents [69,70], as well as prevent DNA damage *in vitro* [71]. 5-CQA, which is the most important CGA in coffee, can scavenge 1,1-diphenyl-2-picrylhydrazyl radicals (DPPH), superoxide anions ($\text{O}_2^{\bullet-}$), hydroxyl radicals ($\bullet\text{OH}$), and peroxynitrite (ONOO^-) [72–74], and protect DNA from damage caused by oxidative stress in different studies [67,75].

Therefore, there is enough evidence to support that CGAs can inhibit the formation of reactive oxygen species and play a beneficial role in preventing oxidative and aging-related diseases [65,66]. However, studies indicate that these compounds may also act as potent pro-oxidants. Therefore, depending on their concentration, the presence of free transition metal ions, or their redox state, the antioxidant and pro-oxidant properties of CGAs can be modified [76–78].

4.2. Anti-Inflammatory Activity

Inflammation is a complex physiological process of tissue injury caused by exogenous or endogenous sources [67]. A prolonged unregulated inflammatory process can induce tissue damage and is the cause of many chronic pathologies, such as diabetes, alcoholic liver, chronic kidney disease, and cardiovascular and neurodegenerative diseases [79,80]. CGAs, mainly 5-CQA, have been shown anti-inflammatory activity by reducing pro-inflammatory cytokines, due to modulation of key transcription factors, such as tumor necrosis factor- α (TNF- α) and interleukins, such as IL-8 [67,81]. Another study performed in murine RAW264.7 macrophages showed that 5-CQA decreased lipopolysaccharide (LPS)-induced cyclooxygenase (COX-2) up-regulation at both the protein and mRNA levels, suggesting that 5-CQA might exert anti-inflammatory effects through inhibition of prostaglandin E2 (PGE2) production [82]. It has also been reported that CFA can enhance the wound healing process [67]. In a study with diabetic rats, oral administration of 5-CQA increased hydroxyproline concentrations and decreased malondialdehyde/nitric oxide levels in wound tissues. In addition, it allowed elevation of reduced glutathione [83,84]. Topical administration of 5-CQA-containing hydrogels to mouse skin wounds significantly reduced the size of the wound area in the inflammatory phase, improving the healing process [85].

4.3. Neuroprotective Activity

Alzheimer's disease is a neurodegenerative disease characterized by progressive deterioration of learning, memory, and other cognitive deficits, along with the extracellular deposition of β -amyloid peptides into the brain leading to neuroinflammation, synaptic loss and neuronal death [86,87]. According to Alzheimer Association [88], in 2050, the number of people aged 65 and older with Alzheimer's disease will reach 12.7 million. Several studies found an inverse relationship between coffee consumption and the development of Alzheimer's disease, suggesting its possible use in managing treatments [86,89–92]. The neuroprotective mechanisms of coffee are suggested to be related to the anti-inflammatory effects of caffeine and CGAs on A₁ and A₂ receptors. In addition, it reduces toxic deposits of β -amyloid peptides in the brain, which is a distinctive feature in Alzheimer's patients [6,90,92,93]. Furthermore, some coffee compounds could inhibit brain acetylcholinesterase and butyrylcholinesterase (causing a delay in the degradation of acetylcholine and butyrylcholine), resulting in the prevention of oxidative stress-induced neurodegeneration due to their high antioxidant activity [6,90,94].

On the other hand, murine model trials have shown a significant association between the consumption of CGAs and the prevention of the development of degenerative diseases and aging [6,95–98]. The effect of phenolic compounds from coffee on human cognitive function has not been well studied [99]. However, the number of in vitro studies concerning the neuroprotective effects of polyphenols is rapidly increasing. It has been demonstrated that intraperitoneal injections of 5-CQA reduced oxidative damage in the cerebellum of rats exposed to methotrexate, a drug with serious side effects used to treat some types of cancer, rheumatoid arthritis, and psoriasis [100]. In the same study, these researchers also observed that application of 5-CQA decreased lipopolysaccharide (LPS)-induced IL-1 β and (TNF- α) release in the substantia nigra, indicating neuroprotective effects of 5-CQA on neurodegenerative diseases caused by proinflammatory cytokines [100]. Taram et al. [101] studied the neuroprotective effects of 5-CQA and caffeic and ferulic acids on rat cerebellar granule neuron cultures. This research proposed that caffeic acid showed enhanced neuroprotection against a wide range of stressors compared to the other compounds evaluated. Thus, the authors suggest that caffeic acid could be a promising candidate in preclinical models of neurodegeneration [101].

4.4. Anticancer Activity

The antimutagenic properties of CGAs was demonstrated decades ago [102]. This activity is partially related to the antioxidant activity of these compounds since the overproduction of oxygen free radicals leads to oxidative DNA damage. This damage is leading

cause of the proliferation of several types of cancer, such as breast, colon, bladder, pancreatic, liver, skin, and prostate cancer [103]. Dietary polyphenols, including CGAs, can protect the initiation of tumor processes by inhibiting DNA lesions caused by both free radicals and carcinogens [104]. Indeed, some epidemiological studies demonstrated an inverse relationship between coffee consumption and the risk of certain types of cancer. This effect has been associated with the intake of CGAs [105–107]. Several mechanisms have suggested that CGAs may have a chemopreventive effect [80]. Among those, modulation of the expression of enzymes involved as endogenous antioxidant defenses, in DNA replication, as well as in cell differentiation and aging are prominent [104,108]. Moreover, metal chelation, inactivation of reactive compounds, and changes in metabolic pathways have been proposed to impact anticancer activity significantly [109]. Boettler et al. [110] demonstrated by *in vitro* and *in vivo* assays that coffee-derived CGAs can induce a cellular and tissue protection mechanism against carcinogenesis via the Nrf2/ARE pathway. This pathway regulates the expression of S-transferases (GST), γ -glutamate-cysteine ligase (γ GCL), NAD(P)H: quinone oxidoreductase 1 (NQO1), and heme oxygenase (H01). In another study by Feng et al. [108] using mouse epithelial JB6 cells, it was found that 5CQA had a protective effect against carcinogens. This effect was due to its ability to decrease the generation of free radicals and stimulate glutathione-S-transferase activity.

4.5. Antidiabetic Activity

According to International Diabetes Federation [111] diabetes (type 1 and 2) is one of the fastest-growing global health emergencies of the 21st century. It was estimated that 537 million adults aged 20–79 years are currently living with diabetes and type 2 diabetes mellitus (T2DM) is the most common type of diabetes, accounting for over 90% of all diabetes worldwide [111]. Several studies have demonstrated an association between moderate consumption of coffee and a lower risk of developing T2DM. This was observed in all sexes, obesity levels, and geographic locations [112–119]. This effect has been attributed to the bioactive compound 5-CQA. Through a meta-analysis, Huxley et al. [120] concluded that daily consumption of three to four cups of coffee decreased the risk of T2DM by 25%.

Furthermore, Bakuradze et al. [121] suggested that consumption of three to four cups of coffee per day could reduce oxidative damage, body fat mass, and energy/nutrient intake and that these effects were partially attributed to CGAs. Shearer et al. [122] studied the effects of regular and decaffeinated coffee (with CGAs) consumption for 28 days on insulin functions, *in vivo* using a rat model. They observed that the ingestion of decaffeinated coffee improves insulin-stimulated disposal in the high-fat-fed and insulin-resistant rats. Other suggested mechanisms of CGAs are related to the improvement of glucose and lipid metabolism by activating of AMP activated protein kinase (AMPK) [119], as shown in Figure 3. AMPK is a master sensor and regulator of cellular energy balance. This enzyme is activated by diverse pathological, metabolic, and pharmacological stressors such as hypoxia, exercise, thiazolidinediones, and metformin. This activation provokes the translocation of glucose transporter type 4 (GLUT4) from intracellular membranes to plasma and, therefore, the increase of glucose transport [119,123,124].

4.6. Cardiovascular Protection Activity

Currently, cardiovascular diseases (CVDs) comprise one of the leading causes of death and disability worldwide. The incidence of various chronic CVDs, including stroke, atherosclerosis, hypertension, ischemic heart disease, and heart failure, probably continues to increase [4]. Some risk factors, such as smoking, high blood pressure, hyperlipidemia, and hyperglycemia, have been reported to contribute, partially, to the development of CVDs [4]. According to the World Health Organization (WHO), ischemic heart disease is the leading cause of death worldwide, accounting for 16% of deaths worldwide (8.9 million people) [125]. Recently, many studies have shown that the consumption of CGAs-rich foods may be recommended to prevent CVDs [49,119,126–128]. The high antioxidant and anti-inflammatory activity of CGAs can improve endothelial dysfunction and reduce

insulin resistance which could be critical mechanisms to enhance the cardiovascular protection attributed to these compounds, as shown in a large number of in vitro and in vivo studies [67]. Taguchi et al. [129] observed that CGAs could improve endothelial function through by releasing of vasoactive molecules such as nitric oxide. This effect was studied in streptozotocin-treated diabetic rats. On the other hand, CGAs could decrease blood pressure by the following proposed mechanisms: (i) stimulation of nitric oxide production through the endothelium-dependent pathway [130], (ii) reduction of free radicals through decreased expression and activity of NADPH oxidase [131], and (iii) through inhibition of the angiotensin-converting enzyme (ACE) [67].




Figure 3. 5CQA-mediated regulation of glucose and lipid metabolism through activation of the AMPK pathway.

4.7. Antibacterial, Antifungal, and Antiviral Activity

The antimicrobial (bacteriostatic and bactericidal) effects of 5-CQA and coffee extracts on various types of detrimental microorganisms that may grow in different parts of the body, from oral bacteria causative of caries to harmful intestinal bacteria, are well known. Roasted *C. arabica* and *C. canephora* extracts and brews exhibited antibacterial activity against *Streptococcus mutans* and other oral types of bacteria [132,133]. Furthermore, 5-CQA can have a positive affect against the adverse microbiota present in the colon. Therefore, this chlorogenic acid can be used as a preservative and food additive [10]. For this reason, CGAs, mainly 5-CQA, could be potential natural antibacterial, antifungal and antiviral agents [2]. For example, 5-CQA exhibited a broad-spectrum antimicrobial activity against Gram-positive (*Streptococcus pneumoniae*, *Staphylococcus aureus*, and *Bacillus subtilis*) and Gram-negative (*Escherichia coli*, *Shigella dysenteriae*, and *Salmonella typhimurium*) pathogenic bacteria by increasing the membrane permeability, leading to plasma membrane barrier dysfunction, as well as leakage of nucleotide [134,135]. The suggested mechanism by which 5-CQA provokes the membrane disruption could involve the perturbation of the membrane lipid bilayer, resulting in cell leakage and dissipation of the membrane electrical potential [4,135].

In addition, Sung and Lee [136] studied the antifungal properties of 5-CQA against *Candida albicans*, a pathogenic yeast. They suggested that this compound could exert antifungal activity by disrupting the cell membrane structure and consequently, it can be used as an option for fungal treatment. In several studies, both caffeic acid and 5-CQA

have demonstrated antiviral activities against herpes simplex virus (HSV) types 1 and 2 [137], adenovirus [138], and HIV [139].

4.8. Other Bioactivities

4.8.1. Hepatoprotective Activity

The beneficial effects of coffee on liver diseases, in general, have been reported in several studies [140–142] for example, cirrhosis and hepatitis B and C [142]. Hepatic injury may be due to multiple factors, such as viral hepatitis, obesity, excessive alcohol consumption, and iron overload [67]. On the other hand, according to a meta-analysis of 16 human studies, coffee consumption (2 cups per day) decreased the risk of developing liver cancer by 40% compared to no coffee consumption [143,144]. The suggested mechanisms of hepatic protection were the prevention of cell apoptosis and oxidative stress damage due to the activation of natural antioxidant and anti-inflammatory systems [145,146]. These protective mechanisms have been mainly related to CGA [147] and caffeine [148], among other components of coffee.

4.8.2. Potential Prebiotic Activity

According to the International Scientific Association for Probiotics and Prebiotics (ISAPP), a prebiotic definition is “a substrate that is selectively utilized by host microorganisms conferring a health benefit” [149]. Usually, well-established prebiotics are carbohydrate-based, but other substances such as polyphenols and polyunsaturated fatty acids transformed into their respective conjugated fatty acids can potentially fit into this new prebiotic definition, provided there is sufficient evidence of their positive effect on the host [149]. The consumption of prebiotic foods or compounds selectively favors the growth of probiotic and other health-promoting microorganisms in the gut, especially *Bifidobacterium* and *Lactobacillus* [150–152]. Thus, indirectly, the health benefits of prebiotics are the following: (i) production of short-chain fatty acids that lower luminal pH, (ii) stimulation of the growth of beneficial intestinal bacteria and suppression of pathogenic bacteria [151,152], (iii) stimulation of the immune system [153,154], (iv) prevention of colon cancer [155], (v) decrease the prevalence to develop diabetes [156,157], and (vi) increased calcium absorption [158]. Furthermore, Kellow et al. [159] observed that dietary supplementation with prebiotics could reduce or delay the accumulation of advanced glycation end products (AGEs) formed through the Maillard reaction in individuals at high risk for type 2 diabetes and improve or restore the microbial balance within the gastrointestinal tract, potentially reducing AGE absorption.

Several studies have suggested that the non-absorbed part of 5-CQA and caffeic acid in the human gastrointestinal tract serves as a substrate for beneficial intestinal microbiota, thus stimulating their growth [160,161]. Whereas the bifidogenic effect of 5-CQA would seem consensus [61,62], the effect of 5-CQA on *Lactobacillus* growth remains debatable, as only selected strains can utilize it as a substrate [62,63]. Furthermore, Parkar et al. [61] reported an increase in short-chain fatty acids (butyric, acetic, and propionic acid) promoted by 5-CQA. Nevertheless, it has also been observed that 5-CQA promotes the growth of *Firmicutes* and *Bacteroides*, and *Clostridium*. Moreover, an inhibitory effect on the growth of *E. coli* has only been demonstrated in one study [135]. Therefore, more studies are needed to validate the effect of 5-CQA as a prebiotic.

5. Bioavailability of CGAs

Numerous studies have shown the potential health benefits of CGAs. Consequently, evidence of the absorption and bioavailability of CGAs is needed to evaluate these compounds' health benefits fully. However, the absorption and bioavailability of CGAs are controversial due to the significant interindividual differences regarding their utilization, metabolism, and excretion found in scientific and clinical studies.

5.1. Absorption of CGAs

Past studies have considered that 5-CQA, such as other phenolic compounds, could be poorly absorbed by the digestive system [162]. However, other studies have shown that a part of this compound can be absorbed intact in the stomach and/or small intestine [163,164]. It is now known that, on average, almost one third of the 5-CQA obtained from the diet is absorbed from the gastrointestinal tract into the bloodstream, although its absorption varies among humans [162,165–168]. For example, after coffee consumption, two plasma concentration peaks of CGAs corresponding to 5-CQA and DCQA were found at 0.5 to 1.0 and 1.5 to 4.0 h, respectively [163]. Furthermore, Mubarak et al. [169] reported a higher concentration of intact 5-CQA in plasma at 2.5 h in all healthy volunteers following intake of pure 5-CQA (400 mg, approximately corresponding to two cups of coffee). Therefore, it has been suggested that 5-CQA is absorbed through at least two pathways. One pathway may involve immediate absorption of intact 5-CQA in the stomach and/or upper gastrointestinal tract, whereas the other involves slow absorption of intact 5-CQA throughout the small intestine [4]. Additionally, Erk et al. [170] reported that a high intake of 5-CQA from coffee could modify gastrointestinal transport and influence its absorption and metabolism.

5.2. Metabolization of CGAs

The human metabolism of CGAs is somewhat complex but well defined. The main pathways of CGAs metabolism are as follows: (i) absorbed non-transformed, (ii) absorbed in the stomach or small intestine, with or without hydrolysis, and then conjugated (sulfate, glucuronide, or methyl) or otherwise metabolized (hydrogenated, α - or β -oxidized, conjugated with glycine), (iii) undergo gut microbiota-mediated metabolism, after which the microbial catabolites are absorbed without further change, or (iv) undergo metabolism via the intestinal microbiota, after which the microbial catabolites are absorbed and undergo mammalian phase II metabolism (conjugation with glucuronide, sulfate, methyl, or glycine) or are otherwise metabolized (hydrogenated, dehydrogenated, α - or β -oxidations) [171]. Thus, it has been found that 33% of the total intake of 5-CQA from the diet is absorbed intact, unhydrolyzed, in the stomach or upper intestine and subsequently passes into the bloodstream. About 7% of the total intake of 5-CQA is absorbed through the small intestine by hydrolysis to CA and QA. Furthermore, part of the metabolism of 5-CQA is mediated by the colonic microbiota. In some studies, traces of 5-CQA have been found in the urine (0.3–2.3%) after ingesting foods with a high content of this phenolic compound, indicating that the absorption of intact 5-CQA is intensively metabolized [163,165]. It is important to highlight that the absorbed part of 5-CQA and its metabolites can induce various physiological effects through the bloodstream, while unabsorbed 5-CQA can induce biological effects throughout the digestive tract, such as modification of the gut microbiota [62,165].

6. Incorporation of CGAs into Food Matrices

The growing demand for healthier foods and better lifestyles is relevant for consumers nowadays. Nutrition scientists and food scientists have established that the best way to enrich and fortify food products in overall nutrient intake with minimum side effects is by using extracts and compounds from natural sources (cereals, greens, fruits, etc.) [172]. Chlorogenic acids obtained from distinct vegetal species and their wastes can be used as natural ingredients for different food products as shown in Table 2. Plant foods such as vegetables and fruits are the main source of calories, carbohydrates, and other essential compounds for the human body and play an important role in human health, such as polyphenols [173]. To increase the intake of polyphenols and the level of acceptance among consumers around the globe, studies on food technology using polyphenols have been increasing over the last couple of decades. Polyphenols are a very significant source of phytochemicals which have been used by the pharmaceutical industry for many years in a wide variety of products [174].

Table 2. Principal incorporation of coffee extracts with CGAs into food products ^a.

Food Product	Technological Improvement	Extract Conditions	CGAs Content in Green Coffee Extract	Major Findings	Sensory Evaluation	References
Fried doughnuts	Dough stability	Heated at 110 °C for 15 min and Freeze-dried	25.5 g/100 g	Dough stability was not affected during mixing and GCA showed high stability increasing antioxidant activity	No significant difference up to 1% of GCE addition (Score 5–4.9)	[175]
Instant coffee	Fortification	Heated at high pressure at 180 °C for percolation and extraction	14.0 g/100 g	Enriched coffee with green coffee extract showed high antioxidant potential but decreased sensory score	No significant difference in <i>C. arabica</i> samples (Score 7.3–6.8)	[176]
Soymilk	Fortification	Heated aqueous extraction (1:10 <i>w/v</i>) at 100 °C for 1 h	N.A.	Phenolic compounds and antioxidant activity content increased significantly, and overall digestibility improved	No decrease in the acceptance level up to 0.25 mg/mL of CGA (Overall score 4.3–5.2)	[177]
Wheat bread	Dough stability and fortification	Heated aqueous extraction at 60, 70 and 80 °C for 1 h	37.3 g/100 g	GCE addition increased CGAs and antioxidant activity in bread, baking quality was not affected.	Maximum level of GCE without adverse effect was 1.5% flour basis (Overall score 64–60)	[178]
Liquid Khask	Enrichment	Heated aqueous extraction (1:10 <i>w/v</i>) at 100 °C for 30 min and encapsulated with water and oil emulsion	39.1 g/100 g	Encapsulated GCE protected color. pH remained unaffected and rheological properties were not affected and antioxidant activity highly increased	No significant difference up to 1% of encapsulated GCE addition (Score 4.7–4.9)	[179]
Dark chocolate	Enrichment	Heated aqueous extraction (1:5 <i>w/v</i>) at 80 °C for 30 and encapsulated	N.A.	Addition of CGAs (free and encapsulated) had no significant effect on dark chocolate color. However, the addition of free or encapsulated CGAs had a significant effect on chocolate flavor. This adverse effect of CGAs on chocolate flavor were lower in the case of encapsulated form addition	No significant difference in the bitterness up to 50.1 mg/5 kg of encapsulated CGAs (Score 1.5–2)	[180]

Table 2. Cont.

Food Product	Technological Improvement	Extract Conditions	CGAs Content in Green Coffee Extract	Major Findings	Sensory Evaluation	References
Yoghurt	Enrichment	Heated aqueous extraction (1:6 <i>w/v</i>) at 70 °C for 1 h. The extract was filtered and concentrated by evaporation (70 °C, 30 min) and spray drying	46.5 g/100 g	Green coffee-enriched yoghurt have desirable pH (4.7), acidity, color, and minimum syneresis. The flavor, texture and other sensory attributes of yoghurt were improved.	Higher score in overall acceptance up to 2% <i>w/v</i> of GCE	[181]

^a GCE, green coffee extract; GCA, green coffee addition; N.A., not available.

Corso et al. [176] studied the antioxidant properties of CGAs and enriching coffee itself. CGAs extract was obtained by a series of percolation stages with pressure water at 180 °C in the first and 100 °C at the final stages. The extract was freeze-dried and added to instant coffee formulations. It was added to obtain a concentration of 7% polyphenols in four different instant coffee formulations. The green coffee extract had 14% CQAs, and 5-CQA was the most abundant [25]. The formulation of instant coffee increased its 5-CQA content and showed 3.18 g/100 g compared to the control 1.20 g/100 g. Moreover, CGAs addition increases antioxidant activity in the instant coffee (evaluated by the ABTS and Folin methods); for enriched coffee, the antioxidant activity was in a range of 30.9–32.0 g of Trolox/100 g, whereas for the control, the content was 24.0–25.6 g of Trolox/100 g. Additionally, the authors reported that the antioxidant activity is not significantly affected by the roasting process. Since the polyphenols reduction (CGAs) is balanced by increasing melanoidin content, Vignoli et al. [182] had similar results. Moreover, no significant difference was found when the sensory evaluation was performed, and all formulations were accepted, obtaining average scores from 6.6 to 7.7 on a hedonic 10-point scale.

Bakery products are well-known sources of energy and nutrients such as carbohydrates, proteins, minerals, and vitamins. However, these also lack antioxidant-rich polyphenolic compounds, fiber, minerals, vitamin B6, thiamine, folate, vitamin E, and some phytochemicals, mostly because the bakery products are formulated with refined wheat flour [183,184]. A study to determine the functional and technological properties of GCE rich in hydroxycinnamic acids (CGAs) in wheat flour and bread was made by Mukkundur et al. [178]. Three levels of GCE obtained from defatted and decaffeinated *C. canephora* green coffee beans were added at 1, 1.5, and 2% on wheat flour. A decrease in total polyphenols (TTP), CGAs, and radical scavenging activity (RSA), 20.0, 36.2, and 93.1%, respectively, was observed due to the high temperature of extraction (80 °C). While the extract obtained at 60 °C was higher for TTP, CGAs, and RSA (21.4, 37.3, and 94.4%, respectively) due to the thermal sensibility of polyphenols [41]. In bread, CGAs addition was found to improve overall key parameters. First, the addition of 2% GCE improved the bread volume (565 cc) compared to the control (525 cc) due to polyphenols' interaction with gluten proteins and starch giving more tenacity and extensibility to the dough. GCE addition increased the greenness of bread crumb and reduced yellowness and lightness. This effect was expected because the green color indicates CGAs hydrolyzation and thermal degradation [41]. The texture of the bread was not significantly affected by GCE addition; however, the bread containing 2% GCE was softer (4.38 N) compared to the control (4.81 N). The content of TPP, RSA, and CGAs in bread significantly increased in all treatments; for TPP, the content was 0.16, 0.25, and 0.34% (for 1, 1.5, and 2% GCE, respectively) compared to the control (0.02%). RSA content increased from 12.7% for the control to 68.5% at the highest level of GCE addition. CGAs content increased from 0.28 to 0.54% with the treatments, whereas CGAs were not detected in control. Finally, for the sensory evaluation, the authors reported that despite the benefits of GCE addition in all three levels, the maximum level of enrichment without affecting the overall quality of the bread (especially taste) was 1.5% GCE.

The influence of addition of green coffee extract (GCE) prepared in aqueous extraction at 110 °C for 15 min with high content of chlorogenic acids in fried doughnuts was analyzed [175]. The GCE had a content of 25.5 g/100 g of polyphenols. The most abundant polyphenol was 5-CQA, two isomers of 5-CQA hydrolyzation (3-CQA and 4-CQA), and another ferulic compound (5-FQA). Results indicated a significant increase of antioxidant activity up to 37, 45, and 50% using three addition levels (0.25, 0.50, and 1%, respectively) compared to the control. The authors compared this enhancement with another extract from a different source of chlorogenic acids (green tea extract) GTE, and they found that GTE addition increased antioxidant activity by 22, 28, and 29% compared to the control. These data confirm that green coffee remains the vegetal species with the highest concentration of polyphenols, for example, CGAs. However, it is relevant to consider that the

frying process reduces the 5-CQA concentration at the final by hydrolyzing the diesters to monoesters.

Moreover, the addition of raw coffee beans in food products has been attempted. Another study on wheat bread properties was developed by Zain et al. [185]. The authors proposed to use grounded green coffee beans (GCB) instead of an extract to substitute the flour at three levels of addition (3, 5, and 7% GCB). After baking, there was a significant increase in the TPP content since the addition of 3, 5, and 7% GCB increased TPP in bread up to eight times at the highest level (1.61 mg GAE/g) compared to control (0.26 mg GAE/g) and even at the first level (3% GCB) TPP content was nearly twice as much compared to whole wheat bread. Since polyphenols were detected in control wheat bread, the authors concluded that it was the ferulic acid present naturally in wheat flour after the milling process and amino acids and smaller peptides formed in proteolysis of protein wheat flour during fermentation of bread. Antioxidant properties of the bread were also analyzed, and they found that with GCB addition, the RSA increased significantly (up to three times more) than the control. As explained by several authors, the increase in antioxidant activity is mainly attributed to the presence of phytochemicals in GCB (mainly chlorogenic acids). However, GCB addition did affect the sensory properties of bread. According to the sensory evaluation (shape, texture, attractiveness, color, chewability, odor, and taste) for control, the overall score was around 7.3–7.5, whereas for 3, 5, and 7% GCB bread, the scores were significantly lower (6.3, 5, and 4.2, respectively). Color and volume of bread were negatively affected; the more GCB was added, the color of the crumb and crust turned green and reduced volume. This suggested that GCB addition probably affects the stability of the gluten matrix, making a compact structure. It is well known that bread types depend on cultural and geographic requirements. Therefore, enriched food formulation must be designed according to the target market [186].

The fortification of dairy products has increased in the last couple of decades [187]. A study on the field of fortification of dairy products using encapsulated green coffee extract was made by Rahpeyma et al. [179]. The extract was obtained with boiling water at 110 °C for 30 min, cooled at room temperature and filtered. The authors used an emulsion microencapsulation technique using glycerol monostearate (GMS) and canola oil, and shaking at 4000 rpm and 70 °C. They used three levels of GCE addition (0.25, 0.5, and 1%) and three levels of encapsulated green coffee extract (EGCE) at 1.25, 2.5, and 5% EGCE. The TPP content and RSA of the extract were found to be 39 and 74%, respectively. Under the microencapsulation condition, the coating material greatly impacted the retention of phenolic compounds and antioxidant activity of the extract when added to kashk. The enrichment of kashk does not affect rheological properties such as viscosity across all treatments. Noteworthy, acidification by GCE, EGCE, and lactic fermentation caused a decrease in the negative electric charge of the micelles by degrading the calcium and inorganic phosphates.

Another study on dairy products was developed by Dönmez et al. [188]. They added green coffee powder (GCP) at two levels (1 and 2%) into homemade yogurt to analyze the polyphenols' activity and interaction with proteins from yogurt. The coffee addition reduced the serum release rate (syneresis) in yogurt. The serum separation was significantly restricted by half of the control rate with the highest addition level of GCP. Polyphenols are highly reactive to proteins since they can form protein-polyphenol complexes through multiple weak interactions (mainly hydrophobic, van der Waals, and hydrogen bridge-binding) formed between protein side chains and polyphenol aromatic rings [189]. However, interactions between GCP and proteins were strengthening the gel structure of yogurt and hence, affected its rheological behavior. The yogurt consistency was increased during the first 14 days of storage. There was no significant change in flow index of yogurt with the highest level of GCP during 21 days of storage. Color significantly changed with GCP addition; green color increased over storage time up to 40% of the greenness for day 21.

In dairy-free milk-based products, the effect of CGA addition was investigated by Seczyk et al. [177] with green coffee phenolics (CGAs) added to soymilk. The extract was

obtained by aqueous heat-assisted extraction using 10 g of green coffee, boiled in 100 mL of water at 110 °C, allowed to cool down at room temperature for 1 h with continuous orbital shaking and then filtered (Whatman No. 4). Six levels of addition were used in the study (0.0025, 0.05, 0.1, 0.25, 0.5, and 1 mg of phenolics (GAE, gallic acid equivalent) per 1 mL). Compared to the control, the content of phenolics increased up to 70% at the highest level of addition of CGAs. Antioxidant activity and reducing power were significantly affected, increasing the content up to 3.5–13.8 times more than the control.

Furthermore, CGAs addition also improved the digestibility of starch and proteins in soymilk, increasing digestibility up to 17.9% higher than the unfortified soymilk. CGE addition improved soymilk aroma and texture, with grass-lemon notes. The taste was positively affected by CGAs addition showing a score range between 5–5.3 whereas the control had a 4.7 score; however, at the highest level of fortification, the taste score significantly decreased (3.0) [177].

7. Future Perspectives

Coffee chlorogenic acids have multifunctional properties as phytochemical and nutraceutical. Because of these properties, 5-CQA received considerable attention for its potential functional effects. With the recent advances in food fortification and the growing research interest in CGAs from different sources, its use as a natural additive to increase the intake of phenolic compounds is attractive for the food industry. There are different methods to recover CGAs; however, these extraction methods are conditioned by their physical and chemical sensitivity. Extraction methods such as electric field pulse and activated carbon purification might be used to reduce the use of toxic solvents and high temperatures, which could contaminate and reduce CGAs content in the extract. However, more research on optimization for obtaining solvent-free coffee chlorogenic acid extracts is needed.

The potential to fortify foods with CGAs would represent an option for delivering antioxidant concentrations beneficial to sustain wellbeing and health in humans, pets, and farm animals. Due to CGA's susceptibility to temperature, pH, and light, some food processes are still challenging since they will reduce the content of CGAs in the final product. Several studies demonstrated increased antioxidant activity without affecting key quality parameters in baked products. CGAs are highly stable in yogurt and soymilk, enhancing the flavor and color while increasing their antioxidant activity. However, for products such as chocolate, more research is needed since CGAs addition significantly affects the flavor; for this reason, more research on the encapsulation of CGAs by different methods needs to be conducted to decrease the bitterness of foods enriched with CGAs is needed.

Author Contributions: Conceptualization, M.L.S.-Q., P.R.-D. and Z.J.H.-E.; formal analysis, investigation, writing—original draft preparation, A.R.-G. and C.Y.F.-H. (equal contribution); writing—review and editing, R.M.G.-A., O.G.-R., M.L.S.-Q., Z.J.H.-E. and P.R.-D.; funding acquisition, P.R.-D. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Hatch grant no. OKL03091 from the United States, Department of Agriculture (USDA), National Institute of Food and Agriculture and the Oklahoma Agricultural Experiment Station, Oklahoma State University.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Bułdak, R.J.; Hejmo, T.; Osowski, M.; Bułdak, Ł.; Kukla, M.; Polaniak, R.; Birkner, E. The impact of coffee and its selected bioactive compounds on the development and progression of colorectal cancer in vivo and in vitro. *Molecules* **2018**, *23*, 3309. [[CrossRef](#)] [[PubMed](#)]
2. Santana-Gálvez, J.; Cisneros-Zevallos, L.; Jacobo-Velázquez, D.A. Chlorogenic acid: Recent advances on its dual role as a food additive and a nutraceutical against metabolic syndrome. *Molecules* **2017**, *22*, 358. [[CrossRef](#)] [[PubMed](#)]
3. Liu, W.; Li, J.; Zhang, X.; Zu, Y.; Yang, Y.; Liu, W.; Xu, Z.; Gao, H.; Sun, X.; Jiang, X.; et al. Current advances in naturally occurring caffeoylquinic acids: Structure, bioactivity, and synthesis. *J. Agric. Food Chem.* **2020**, *68*, 10489–10516. [[CrossRef](#)]
4. Lu, H.; Tian, Z.; Cui, Y.; Liu, Z.; Ma, X. Chlorogenic acid: A comprehensive review of the dietary sources, processing effects, bioavailability, beneficial properties, mechanisms of action, and future directions. *Compr. Rev. Food Sci. Food Saf.* **2020**, *19*, 3130–3158. [[CrossRef](#)]
5. Li, L.; Su, C.; Chen, X.; Wang, Q.; Jiao, W.; Luo, H.; Tang, J.; Wang, W.; Li, S.; Guo, S. Chlorogenic acids in cardiovascular disease: A review of dietary consumption, pharmacology, and pharmacokinetics. *J. Agric. Food Chem.* **2020**, *68*, 6464–6484. [[CrossRef](#)]
6. Oboh, G.; Agunloye, O.M.; Akinyemi, A.J.; Ademiluyi, A.O.; Adefegha, S.A. Comparative study on the inhibitory effect of caffeic and chlorogenic acids on key enzymes linked to Alzheimer's disease and some pro-oxidant induced oxidative stress in rats' brain-in vitro. *Neurochem. Res.* **2013**, *38*, 413–419. [[CrossRef](#)]
7. Oboh, G.; Agunloye, O.M.; Adefegha, S.A.; Akinyemi, A.J.; Ademiluyi, A.O. Caffeic and chlorogenic acids inhibit key enzymes linked to type 2 Diabetes (in vitro): A comparative study. *J. Basic Clin. Physiol. Pharmacol.* **2015**, *26*, 165–170. [[CrossRef](#)]
8. Wang, L.N.; Wang, W.; Hattori, M.; Daneshmand, M.; Ma, C.M. Synthesis, Anti-HCV, antioxidant and reduction of intracellular reactive oxygen species generation of a chlorogenic acid analogue with an amide bond replacing the ester bond. *Molecules* **2016**, *21*, 737. [[CrossRef](#)]
9. Ye, X.; Li, J.; Gao, Z.; Wang, D.; Wang, H.; Wu, J. Chlorogenic acid inhibits lipid deposition by regulating the enterohepatic FXR-FGF15 pathway. *BioMed Res. Int.* **2022**, *2022*, 4919153. [[CrossRef](#)]
10. Naveed, M.; Hejazi, V.; Abbas, M.; Kamboh, A.A.; Khan, G.J.; Shumzaid, M.; Ahmad, F.; Babazadeh, D.; FangFang, X.; Modarresi-Ghazani, F.; et al. Chlorogenic Acid (CGA): A pharmacological review and call for further research. *Biomed. Pharmacother.* **2018**, *97*, 67–74. [[CrossRef](#)]
11. Smith, P.; Kiong Ho, C.; Takagi, Y.; Djaballah, H.; Shuman, S. Nanomolar inhibitors of *Trypanosoma brucei* RNA triphosphatase. *MBio* **2016**, *7*, e00058-16. [[CrossRef](#)] [[PubMed](#)]
12. Zhu, X.; Zhang, H.; Lo, R. Phenolic compounds from the leaf extract of artichoke (*Cynara scolymus* L.) and their antimicrobial activities. *J. Agric. Food Chem.* **2004**, *52*, 7272–7278. [[CrossRef](#)] [[PubMed](#)]
13. Zhao, Z.; Hee, S.S.; Satsu, H.; Totsuka, M.; Shimizu, M. 5-Caffeoylquinic acid and caffeic acid down-regulate the oxidative stress- and TNF- α -induced secretion of Interleukin-8 from Caco-2 Cells. *J. Agric. Food Chem.* **2008**, *56*, 3863–3868. [[CrossRef](#)] [[PubMed](#)]
14. Tian, X.; An, L.; Gao, L.Y.; Bai, J.P.; Wang, J.; Meng, W.H.; Ren, T.S.; Zhao, Q.C. Compound MQA, a caffeoylquinic acid derivative, protects against NMDA-induced neurotoxicity and potential mechanisms in vitro. *CNS Neurosci. Ther.* **2015**, *21*, 575–584. [[CrossRef](#)] [[PubMed](#)]
15. Kurata, R.; Adachi, M.; Yamakawa, O.; Yoshimoto, M. Growth suppression of human cancer cells by polyphenolics from sweetpotato (*Ipomoea batatas* L.) leaves. *J. Agric. Food Chem.* **2006**, *55*, 185–190. [[CrossRef](#)]
16. Pramanik, K.C.; Bhattacharya, P.; Biswas, R.; Bandyopadhyay, D.; Mishra, M.; Chatterjee, T.K. Hypoglycemic and antihyperglycemic activity of leaf extract of *Pluchea indica* Less. *Orient. Pharm. Exp. Med.* **2006**, *6*, 232–236. [[CrossRef](#)]
17. Mahmood, N.; Moore, P.S.; De Tommasi, N.; De Simone, F.; Colman, S.; Hay, A.J.; Pizza, C. Inhibition of HIV infection by caffeoylquinic acid derivatives. *Antivir. Chem. Chemother.* **2016**, *4*, 235–240. [[CrossRef](#)]
18. Hung, T.M.; Na, M.K.; Thuong, P.T.; Su, N.D.; Sok, D.E.; Song, K.S.; Seong, Y.H.; Bae, K.H. Antioxidant activity of Caffeoyl quinic acid derivatives from the roots of *Dipsacus asper* wall. *J. Ethnopharmacol.* **2006**, *108*, 188–192. [[CrossRef](#)]
19. El-Seedi, H.R.; El-Said, A.M.A.; Khalifa, S.A.M.; Göransson, U.; Bohlin, L.; Borg-Karlson, A.K.; Verpoorte, R. Biosynthesis, natural sources, dietary intake, pharmacokinetic properties, and biological activities of Hydroxycinnamic Acids. *J. Agric. Food Chem.* **2012**, *60*, 10877–10895. [[CrossRef](#)]
20. Meinhart, A.D.; Damin, F.M.; Caldeirão, L.; de Jesus Filho, M.; da Silva, L.C.; da Silva Constant, L.; Teixeira Filho, J.; Wagner, R.; Teixeira Godoy, H. Study of new sources of six chlorogenic acids and caffeic acid. *J. Food Compos. Anal.* **2019**, *82*, 103244. [[CrossRef](#)]
21. Upadhyay, R.; Mohan Rao, L.J. An outlook on chlorogenic acids—Occurrence, chemistry, technology, and biological activities. *Crit. Rev. Food Sci. Nutr.* **2013**, *53*, 968–984. [[CrossRef](#)] [[PubMed](#)]
22. Frosi, I.; Montagna, I.; Colombo, R.; Milanese, C.; Papetti, A. Recovery of chlorogenic acids from agri-food wastes: Updates on green extraction techniques. *Molecules* **2021**, *26*, 4515. [[CrossRef](#)] [[PubMed](#)]
23. Meinhart, A.D.; Damin, F.M.; Caldeirão, L.; da Silveira, T.F.F.; Filho, J.T.; Godoy, H.T. Chlorogenic acid isomer contents in 100 plants commercialized in Brazil. *Food Res. Int.* **2017**, *99*, 522–530. [[CrossRef](#)] [[PubMed](#)]
24. Lattanzio, V.; Cicco, N.; Linsalata, V. Antioxidant activities of artichoke phenolics. *Acta Hort.* **2005**, *681*, 421–428. [[CrossRef](#)]
25. Farah, A.; Donangelo, C.M. Phenolic compounds in Coffee. *Braz. J. Plant Physiol.* **2006**, *18*, 23–36. [[CrossRef](#)]
26. Lu, Y.; Foo, L.Y. Identification and quantification of major polyphenols in apple pomace. *Food Chem.* **1997**, *59*, 187–194. [[CrossRef](#)]
27. Wald, B.; Wray, V.; Galensa, R.; Herrmann, K. Malonated flavonol glycosides and 3,5-dicaffeoylquinic acid from pears. *Phytochemistry* **1989**, *28*, 663–664. [[CrossRef](#)]

28. Rodriguez-Mateos, A.; Cifuentes-Gomez, T.; Tabatabaee, S.; Lecras, C.; Spencer, J.P.E. Procyanidin, anthocyanin, and chlorogenic acid contents of highbush and lowbush blueberries. *J. Agric. Food Chem.* **2012**, *6*, 5772–5778. [[CrossRef](#)]
29. Somers, T.C.; Vérette, E.; Pockock, K.F. Hydroxycinnamate esters of *Vitis vinifera*: Changes during white vinification, and effects of exogenous enzymic hydrolysis. *J. Sci. Food Agric.* **1987**, *40*, 67–78. [[CrossRef](#)]
30. Mattila, P.; Hellström, J. Phenolic acids in potatoes, vegetables, and some of their products. *J. Food Compos. Anal.* **2007**, *20*, 152–160. [[CrossRef](#)]
31. Strack, D.; Hartfeld, F.; Austenfeld, F.A.; Grotjahn, L.; Wray, V. Coumaroyl-, caffeoyl- and feruloyltartronates and their accumulation in Mung bean. *Phytochemistry* **1985**, *24*, 147–150. [[CrossRef](#)]
32. Donovan, J.L.; Meyer, A.S.; Waterhouse, A.L. Phenolic composition and antioxidant activity of prunes and prune juice (*Prunus domestica*). *J. Agric. Food Chem.* **1998**, *46*, 1247–1252. [[CrossRef](#)]
33. Malmberg, A.G.; Theander, O. Determination of chlorogenic acid in potato tubers. *J. Agric. Food Chem.* **1985**, *33*, 549–551. [[CrossRef](#)]
34. Tajik, N.; Tajik, M.; Mack, I.; Enck, P. The potential effects of chlorogenic acid, the main phenolic components in coffee, on health: A comprehensive review of the literature. *Eur. J. Nutr.* **2017**, *56*, 2215–2244. [[CrossRef](#)]
35. Franca, A.S.; Oliveira, L.S. Coffee and its by-products as sources of bioactive compounds. In *Coffee: Production, Consumption and Health Benefits*, 1st ed.; Massey, J.L., Ed.; Nova Science Publishers: New York, NY, USA, 2016; pp. 1–28.
36. Jeszka-Skowron, M.; Sentkowska, A.; Pyrzyńska, K.; de Peña, M.P. Chlorogenic acids, caffeine content and antioxidant properties of green coffee extracts: Influence of green coffee bean preparation. *Eur. Food Res. Technol.* **2016**, *242*, 1403–1409. [[CrossRef](#)]
37. Perrone, D.; Farah, A.; Donangelo, C.M.; de Paulis, T.; Martin, P.R. Comprehensive analysis of major and minor chlorogenic acids and lactones in economically relevant brazilian coffee cultivars. *Food Chem.* **2008**, *106*, 859–867. [[CrossRef](#)]
38. Silva, C.T.; de Souza, M.C.; Machado, A.P.D.F.; do Nascimento, R.D.P.; da Cunha, D.T.; Bezerra, R.M.N.; Rostagno, M.A. Thermal stability and sensory evaluation of a bioactive extract from roasted coffee (*Coffea arabica*) beans added at increasing concentrations to conventional bread. *J. Food Process. Preserv.* **2021**, *45*, e15955. [[CrossRef](#)]
39. Somporn, C.; Kamtuo, A.; Theerakulpisut, P.; Siriamornpun, S. Effects of roasting degree on radical scavenging activity, phenolics and volatile compounds of arabica coffee beans (*Coffea arabica* L. cv. *Catimor*). *Int. J. Food Sci. Technol.* **2011**, *46*, 2287–2296. [[CrossRef](#)]
40. Suárez-Quiroz, M.; Alonso Campos, A.; Valerio Alfaro, G.; González-Ríos, O.; Villeneuve, P.; Figueroa-Espinoza, M. Isolation of green coffee chlorogenic acids using activated carbon. *J. Food Compos. Anal.* **2014**, *33*, 55–58. [[CrossRef](#)]
41. Upadhyay, R.; Ramalakshmi, K.; Jagan Mohan Rao, L. Microwave-assisted extraction of chlorogenic acids from green coffee beans. *Food Chem.* **2012**, *130*, 184–188. [[CrossRef](#)]
42. Yang, Z.; Tan, Z.; Li, F.; Li, X. An effective method for the extraction and purification of chlorogenic acid from ramie (*Boehmeria nivea* L.) leaves using acidic ionic liquids. *Ind. Crops Prod.* **2016**, *89*, 78–86. [[CrossRef](#)]
43. Budryn, G.; Nebesny, E.; Podsedek, A.; Zyzelewicz, D.; Materska, M.; Jankowski, S.; Janda, B. Effect of different extraction methods on the recovery of chlorogenic acids, caffeine and Maillard reaction products in coffee beans. *Eur. Food Res. Technol.* **2009**, *228*, 913–922. [[CrossRef](#)]
44. Phongsupa, J.; Yawootti, A.; Wattanuchariya, W. Chlorogenic acid extraction of local coffee beans by pulsed electric field. *AIP Conf. Proc.* **2021**, *2397*, 020004. [[CrossRef](#)]
45. Bilge, G.; Yurdakul, M.; Buzrul, S.; Bulut, O. Evaluation of the effect of pulsed electric field on coffee arabica beans. *Food Bioproc. Technol.* **2022**, *15*, 1073–1081. [[CrossRef](#)]
46. Madhava Naidu, M.; Sulochanamma, G.; Sampathu, S.R.; Srinivas, P. Studies on extraction and antioxidant potential of green coffee. *Food Chem.* **2008**, *107*, 377–384. [[CrossRef](#)]
47. Rakotomalala, J.J. Diversité Biochimique des Caféiers: Analyse des Acides Hydroxycinnamiques, Bases Puriques et Diterpènes Glycosidiques. Particularités des Caféiers Sauvages de la Région Malgache (*Mascarocoffea chev.*), 1993. Ph.D. Thesis, Université Montpellier II, Montpellier, France, 1992.
48. Dibert, K.; Cros, E.; Andrieu, J. Solvent extraction of oil and chlorogenic acid from green coffee. Part II: Kinetic data. *J. Food Eng.* **1989**, *10*, 199–214. [[CrossRef](#)]
49. Fuentes, E.; Caballero, J.; Alarcón, M.; Rojas, A.; Palomo, I. Chlorogenic acid inhibits human platelet activation and thrombus formation. *PLoS ONE* **2014**, *9*, e90699. [[CrossRef](#)]
50. Amato, A.; Terzo, S.; Mulè, F. Natural compounds as beneficial antioxidant agents in neurodegenerative disorders: A focus on Alzheimer’s disease. *Antioxidants* **2019**, *12*, 608. [[CrossRef](#)]
51. Heitman, E.; Ingram, D.K. Cognitive and neuroprotective effects of chlorogenic acid. *Nutr. Neurosci.* **2017**, *20*, 32–39. [[CrossRef](#)]
52. Bao, L.; Li, J.; Zha, D.; Zhang, L.; Gao, P.; Yao, T.; Wu, X. Chlorogenic acid prevents diabetic nephropathy by inhibiting oxidative stress and inflammation through modulation of the Nrf2/HO-1 and NF-κB pathways. *Int. Immunopharmacol.* **2018**, *54*, 245–253. [[CrossRef](#)]
53. Yan, Y.; Zhou, X.; Guo, K.; Zhou, F.; Yang, H. Use of chlorogenic acid against Diabetes Mellitus and its complications. *J. Immunol. Res.* **2020**, *2020*, 9680508. [[CrossRef](#)] [[PubMed](#)]
54. Ong, K.W.; Hsu, A.; Tan, B.K.H. Anti-diabetic and anti-lipidemic effects of chlorogenic acid are mediated by AMPK activation. *Biochem. Pharmacol.* **2013**, *85*, 1341–1351. [[CrossRef](#)] [[PubMed](#)]

55. Wang, Z.; Lam, K.L.; Hu, J.; Ge, S.; Zhou, A.; Zheng, B.; Zeng, S.; Lin, S. Chlorogenic acid alleviates obesity and modulates gut microbiota in high-fat-fed mice. *Food Sci. Nutr.* **2019**, *7*, 579–588. [[CrossRef](#)] [[PubMed](#)]
56. He, X.; Zheng, S.; Sheng, Y.; Miao, T.; Xu, J.; Xu, W.; Huang, K.; Zhao, C. Chlorogenic acid ameliorates obesity by preventing energy balance shift in high-fat diet induced obese mice. *J. Sci. Food Agric.* **2021**, *101*, 631–637. [[CrossRef](#)] [[PubMed](#)]
57. Ghadieh, H.E.; Smiley, Z.N.; Kopfman, M.W.; Najjar, M.G.; Hake, M.J.; Najjar, S.M. Chlorogenic acid/chromium supplement rescues diet-induced insulin resistance and obesity in mice. *Nutr. Metab.* **2015**, *12*, 19. [[CrossRef](#)] [[PubMed](#)]
58. Ye, X.; Liu, Y.; Hu, J.; Gao, Y.; Ma, Y.; Wen, D. Chlorogenic acid-induced gut microbiota improves metabolic endotoxemia. *Front. Endocrinol.* **2021**, *12*, 1717. [[CrossRef](#)]
59. Zheng, G.; Qiu, Y.; Zhang, Q.F.; Li, D. Chlorogenic acid and caffeine in combination inhibit fat accumulation by regulating hepatic lipid metabolism-related enzymes in mice. *Br. J. Nutr.* **2014**, *112*, 1034–1040. [[CrossRef](#)]
60. Zhong, Y.; Ding, Y.; Li, L.; Ge, M.; Ban, G.; Yang, H.; Dai, J.; Zhang, L. Effects and mechanism of chlorogenic acid on weight loss. *Curr. Pharm. Biotechnol.* **2020**, *21*, 1099–1106. [[CrossRef](#)]
61. Parkar, S.G.; Trower, T.M.; Stevenson, D.E. Fecal microbial metabolism of polyphenols and its effects on human gut microbiota. *Anaerobe* **2013**, *23*, 12–19. [[CrossRef](#)]
62. Mills, C.E.; Tzounis, X.; Oruna-Concha, M.J.; Mottram, D.S.; Gibson, G.R.; Spencer, J.P.E. In vitro colonic metabolism of coffee and chlorogenic acid results in selective changes in human faecal microbiota growth. *Br. J. Nutr.* **2015**, *113*, 1220–1227. [[CrossRef](#)]
63. Sales, A.L.; Depaula, J.; Mellinger Silva, C.; Cruz, A.; Lemos Miguel, M.A.; Farah, A. Effects of regular and decaffeinated roasted coffee (*Coffea arabica* and *Coffea canephora*) extracts and bioactive compounds on in vitro probiotic bacterial growth. *Food Funct.* **2020**, *11*, 1410–1424. [[CrossRef](#)] [[PubMed](#)]
64. Hemmerle, H.; Burger, H.J.; Below, P.; Schubert, G.; Rippel, R.; Schindler, P.W.; Paulus, E.; Herling, A.W. Chlorogenic acid and synthetic chlorogenic acid derivatives: Novel inhibitors of hepatic glucose-6-phosphate translocase. *J. Med. Chem.* **1997**, *40*, 137–145. [[CrossRef](#)] [[PubMed](#)]
65. Khansari, N.; Shakiba, Y.; Mahmoudi, M. Chronic inflammation and oxidative stress as a major cause of age-related diseases and cancer. *Recent Pat. Inflamm. Allergy Drug Discov.* **2009**, *3*, 73–80. [[CrossRef](#)] [[PubMed](#)]
66. Akash, M.S.H.; Rehman, K.; Chen, S. Effects of coffee on type 2 diabetes mellitus. *Nutrition* **2014**, *30*, 755–763. [[CrossRef](#)]
67. Liang, N.; Kitts, D.D. Role of chlorogenic acids in controlling oxidative and inflammatory stress conditions. *Nutrients* **2015**, *8*, 16. [[CrossRef](#)]
68. Nakatani, N.; Kayano, S.I.; Kikuzaki, H.; Sumino, K.; Katagiri, K.; Mitani, T. Identification, quantitative determination, and antioxidative activities of chlorogenic acid isomers in prune (*Prunus domestica* L.). *J. Agric. Food Chem.* **2000**, *48*, 5512–5516. [[CrossRef](#)]
69. Laranjinha, J.A.N.; Almeida, L.M.; Madeira, V.M.C. Reactivity of dietary phenolic acids with peroxy radicals: Antioxidant activity upon low density lipoprotein peroxidation. *Biochem. Pharmacol.* **1994**, *48*, 487–494. [[CrossRef](#)]
70. Gordon, M.H.; Wishart, K. Effects of chlorogenic acid and bovine serum albumin on the oxidative stability of low density lipoproteins in vitro. *J. Agric. Food Chem.* **2010**, *58*, 5828–5833. [[CrossRef](#)]
71. Cinkilic, N.; Cetintas, S.K.; Zorlu, T.; Vatan, O.; Yilmaz, D.; Cavas, T.; Tunc, S.; Ozkan, L.; Bilaloglu, R. Radioprotection by two phenolic compounds: Chlorogenic and quinic acid, on x-ray induced dna damage in human blood lymphocytes in vitro. *Food Chem. Toxicol.* **2013**, *53*, 359–363. [[CrossRef](#)]
72. Zang, L.Y.; Cosma, G.; Gardner, H.; Castranova, V.; Vallyathan, V. Effect of chlorogenic acid on hydroxyl radical. *Mol. Cell. Biochem.* **2003**, *247*, 205–210. [[CrossRef](#)]
73. Cha, J.W.; Piao, M.J.; Kim, K.C.; Yao, C.W.; Zheng, J.; Kim, S.M.; Hyun, C.L.; Ahn, Y.S.; Hyun, J.W. The polyphenol chlorogenic acid attenuates UVB-mediated oxidative stress in human HaCaT keratinocytes. *Biomol. Ther.* **2014**, *22*, 136–142. [[CrossRef](#)] [[PubMed](#)]
74. Kono, Y.; Kobayashi, K.; Tagawa, S.; Adachi, K.; Ueda, A.; Sawa, Y.; Shibata, H. Antioxidant activity of polyphenolics in diets. rate constants of reactions of chlorogenic acid and caffeic acid with reactive species of oxygen and nitrogen. *Biochim. Biophys. Acta Gen. Subj.* **1997**, *1335*, 335–342. [[CrossRef](#)]
75. Xu, J.G.; Hu, Q.P.; Liu, Y. Antioxidant and DNA-protective activities of chlorogenic acid isomers. *J. Agric. Food Chem.* **2012**, *60*, 11625–11630. [[CrossRef](#)] [[PubMed](#)]
76. Du, W.Y.; Xiao, Y.; Yao, J.J.; Hao, Z.; Zhao, Y.B. Involvement of NADPH oxidase in high-dose phenolic acid-induced pro-oxidant activity on rat mesenteric venules. *Exp. Ther. Med.* **2017**, *13*, 17–22. [[CrossRef](#)]
77. Kalinowska, M.; Bajko, E.; Matejczyk, M.; Kaczyński, P.; Łozowicka, B.; Lewandowski, W. The study of anti-/pro-oxidant, lipophilic, microbial and spectroscopic properties of new alkali metal salts of 5-O-caffeoylquinic acid. *Int. J. Mol. Sci.* **2018**, *19*, 463. [[CrossRef](#)]
78. Kalinowska, M.; Sienkiewicz-Gromiuk, J.; Świdorski, G.; Pietryczuk, A.; Cudowski, A.; Lewandowski, W. Zn(II) complex of plant phenolic chlorogenic acid: Antioxidant, antimicrobial and structural studies. *Materials* **2020**, *13*, 3745. [[CrossRef](#)]
79. Cachofeiro, V.; Goicochea, M.; de Vinuesa, S.G.; Oubia, P.; Lahera, V.; Llo, J. Oxidative stress and inflammation, a link between chronic kidney disease and cardiovascular disease. *Kidney Int.* **2008**, *74*, S4–S9. [[CrossRef](#)]
80. Farah, A.; de Paula Lima, J. Consumption of chlorogenic acids through coffee and health implications. *Beverages* **2019**, *5*, 11. [[CrossRef](#)]

81. Shin, H.S.; Satsu, H.; Bae, M.J.; Zhao, Z.; Ogiwara, H.; Totsuka, M.; Shimizu, M. Anti-inflammatory effect of chlorogenic acid on the IL-8 production in Caco-2 Cells and the dextran sulphate sodium-induced colitis symptoms in C57BL/6 Mice. *Food Chem.* **2015**, *168*, 167–175. [[CrossRef](#)]
82. Shan, J.; Fu, J.; Zhao, Z.; Kong, X.; Huang, H.; Luo, L.; Yin, Z. Chlorogenic acid inhibits lipopolysaccharide-induced Cyclooxygenase-2 expression in RAW264.7 cells through suppressing NF-KB and JNK/AP-1 activation. *Int. Immunopharmacol.* **2009**, *9*, 1042–1048. [[CrossRef](#)]
83. Bagdas, D.; Gul, N.Y.; Topal, A.; Tas, S.; Ozyigit, M.O.; Cinkilic, N.; Gul, Z.; Etoz, B.C.; Ziyankok, S.; Inan, S.; et al. Pharmacologic overview of systemic chlorogenic acid therapy on experimental wound healing. *Naunyn Schmiedebergs Arch. Pharmacol.* **2014**, *387*, 1101–1116. [[CrossRef](#)] [[PubMed](#)]
84. Bagdas, D.; Etoz, B.C.; Gul, Z.; Ziyankok, S.; Inan, S.; Turacozen, O.; Gul, N.Y.; Topal, A.; Cinkilic, N.; Tas, S.; et al. In vivo systemic chlorogenic acid therapy under diabetic conditions: Wound healing effects and cytotoxicity/genotoxicity profile. *Food Chem. Toxicol.* **2015**, *81*, 54–61. [[CrossRef](#)] [[PubMed](#)]
85. Affonso, R.C.L.; Voytena, A.P.L.; Fanan, S.; Pitz, H.; Coelho, D.S.; Horstmann, A.L.; Pereira, A.; Uarrota, V.G.; Hillmann, M.C.; Varela, L.A.C.; et al. Phytochemical composition, antioxidant activity, and the effect of the aqueous extract of coffee (*Coffea arabica* L.) bean residual press cake on the skin wound healing. *Oxidative Med. Cell. Longev.* **2016**, *2016*, 1923754. [[CrossRef](#)] [[PubMed](#)]
86. Gardener, S.L.; Rainey-Smith, S.R.; Villemagne, V.L.; Fripp, J.; Doré, V.; Bourgeat, P.; Taddei, K.; Fowler, C.; Masters, C.L.; Maruff, P.; et al. Higher coffee consumption is associated with slower cognitive decline and less cerebral A β -Amyloid accumulation over 126 months: Data from the Australian imaging, biomarkers, and lifestyle study. *Front. Aging Neurosci.* **2021**, *13*, 681. [[CrossRef](#)] [[PubMed](#)]
87. Villemagne, V.L.; Burnham, S.; Bourgeat, P.; Brown, B.; Ellis, K.A.; Salvado, O.; Szoeki, C.; Macaulay, S.L.; Martins, R.; Maruff, P.; et al. Amyloid β deposition, neurodegeneration, and cognitive decline in sporadic Alzheimer's Disease: A prospective cohort study. *Lancet Neurol.* **2013**, *12*, 357–367. [[CrossRef](#)]
88. Alzheimer Association. *2022 Alzheimer's Disease Facts and Figures*; John Wiley and Sons Inc.: Hoboken, NJ, USA, 2022. [[CrossRef](#)]
89. Basurto-Islas, G.; Blanchard, J.; Tung, Y.C.; Fernandez, J.R.; Voronkov, M.; Stock, M.; Zhang, S.; Stock, J.B.; Iqbal, K. Therapeutic benefits of a component of coffee in a rat model of Alzheimer's disease. *Neurobiol. Aging* **2014**, *35*, 2701–2712. [[CrossRef](#)] [[PubMed](#)]
90. Arendash, G.W.; Schleif, W.; Rezai-Zadeh, K.; Jackson, E.K.; Zacharia, L.C.; Cracchiolo, J.R.; Shippy, D.; Tan, J. Caffeine protects Alzheimer's mice against cognitive impairment and reduces brain β -Amyloid production. *Neuroscience* **2006**, *142*, 941–952. [[CrossRef](#)]
91. Wu, L.; Sun, D.; He, Y. Coffee intake and the incident risk of cognitive disorders: A dose–response meta-analysis of nine prospective cohort studies. *Clin. Nutr.* **2017**, *36*, 730–736. [[CrossRef](#)]
92. Dall'Igna, O.P.; Fett, P.; Gomes, M.W.; Souza, D.O.; Cunha, R.A.; Lara, D.R. Caffeine and Adenosine A2a receptor antagonists prevent β -Amyloid (25–35)-induced cognitive deficits in mice. *Exp. Neurol.* **2007**, *203*, 241–245. [[CrossRef](#)]
93. Pathak, L.; Agrawal, Y.; Dhir, A. Natural polyphenols in the management of major depression. *Expert Opin. Investig. Drugs* **2013**, *22*, 863–880. [[CrossRef](#)]
94. Rosso, A.; Mossey, J.; Lippa, C.F. Caffeine: Neuroprotective functions in cognition and Alzheimer's disease. *Am. J. Alzheimers Dis. Other Dement.* **2008**, *23*, 417–422. [[CrossRef](#)] [[PubMed](#)]
95. Hao, R.; Song, X.; Li, F.; Tan, X.; Sun-Waterhouse, D.; Li, D. Caffeic acid phenethyl ester reversed cadmium-induced cell death in hippocampus and cortex and subsequent cognitive disorders in mice: Involvements of AMPK/SIRT1 pathway and amyloid-tau-neuroinflammation axis. *Food Chem. Toxicol.* **2020**, *144*, 111636. [[CrossRef](#)] [[PubMed](#)]
96. Cho, E.S.; Jang, Y.J.; Hwang, M.K.; Kang, N.J.; Lee, K.W.; Lee, H.J. Attenuation of oxidative neuronal cell death by coffee phenolic phytochemicals. *Mutat. Res./Fundam. Mol. Mech. Mutagenesis* **2009**, *661*, 18–24. [[CrossRef](#)] [[PubMed](#)]
97. Zhang, Z.; Li, G.; Szeto, S.S.W.; Chong, C.M.; Quan, Q.; Huang, C.; Cui, W.; Guo, B.; Wang, Y.; Han, Y.; et al. Examining the neuroprotective effects of protocatechuic acid and chrysin on in vitro and in vivo models of Parkinson disease. *Free Radic. Biol. Med.* **2015**, *84*, 331–343. [[CrossRef](#)] [[PubMed](#)]
98. Sarroca, S.; Gatus, A.; Rodríguez-Farré, E.; Vilchez, D.; Pallàs, M.; Griñán-Ferré, C.; Sanfeliu, C.; Corpas, R. Resveratrol confers neuroprotection against high-fat diet in a mouse model of Alzheimer's disease via modulation of proteolytic mechanisms. *J. Nutr. Biochem.* **2021**, *89*, 108569. [[CrossRef](#)]
99. Lakey-Beitia, J.; Berrocal, R.; Rao, K.S.; Durant, A.A. Polyphenols as therapeutic molecules in Alzheimer's disease through modulating amyloid pathways. *Mol. Neurobiol.* **2014**, *51*, 466–479. [[CrossRef](#)]
100. Shen, W.; Qi, R.; Zhang, J.; Wang, Z.; Wang, H.; Hu, C.; Zhao, Y.; Bie, M.; Wang, Y.; Fu, Y.; et al. Chlorogenic acid inhibits LPS-induced microglial activation and improves survival of dopaminergic neurons. *Brain Res. Bull.* **2012**, *88*, 487–494. [[CrossRef](#)]
101. Taram, F.; Winter, A.N.; Linseman, D.A. Neuroprotection comparison of chlorogenic acid and its metabolites against mechanistically distinct cell death-inducing agents in cultured cerebellar granule neurons. *Brain Res.* **2016**, *1648*, 69–80. [[CrossRef](#)]
102. Mori, H.; Tanaka, T.; Shima, H.; Kuniyasu, T.; Takahashi, M. Inhibitory effect of chlorogenic acid on methylazoxymethanol acetate-induced carcinogenesis in large intestine and liver of hamsters. *Cancer Lett.* **1986**, *30*, 49–54. [[CrossRef](#)]
103. Toyokuni, S. Oxidative stress as an iceberg in carcinogenesis and cancer biology. *Arch. Biochem. Biophys.* **2016**, *595*, 46–49. [[CrossRef](#)]

104. Ramos, S. Cancer chemoprevention and chemotherapy: Dietary polyphenols and signalling pathways. *Mol. Nutr. Food Res.* **2008**, *52*, 507–526. [CrossRef] [PubMed]
105. Mishra, M.; Panta, R.; Miyares, M. Influence of coffee and its components on breast cancer: A review. *Asian Pac. J. Trop. Dis.* **2016**, *6*, 827–831. [CrossRef]
106. Li, Y.M.; Peng, J.; Li, L.Z. Coffee consumption associated with reduced risk of oral cancer: A meta-analysis. *Oral Surg. Oral Med. Oral Pathol. Oral Radiol.* **2016**, *121*, 381–389.e1. [CrossRef] [PubMed]
107. Nkondjock, A. Coffee consumption and the risk of cancer: An overview. *Cancer Lett.* **2009**, *277*, 121–125. [CrossRef]
108. Feng, R.; Lu, Y.; Bowman, L.L.; Qian, Y.; Castranova, V.; Ding, M. Inhibition of activator Protein-1, NF-KB, and MAPKs and induction of phase 2 detoxifying enzyme activity by chlorogenic acid. *J. Biol. Chem.* **2005**, *280*, 27888–27895. [CrossRef]
109. Kasai, H.; Fukada, S.; Yamaizumi, Z.; Sugie, S.; Mori, H. Action of chlorogenic acid in vegetables and fruits as an inhibitor of 8-hydroxydeoxyguanosine formation in vitro and in a rat carcinogenesis model. *Food Chem. Toxicol.* **2000**, *38*, 467–471. [CrossRef]
110. Boettler, U.; Volz, N.; Pahlke, G.; Teller, N.; Kotyczka, C.; Somoza, V.; Stiebitz, H.; Bytof, G.; Lantz, I.; Lang, R.; et al. Coffees rich in chlorogenic acid or N-methylpyridinium induce chemopreventive phase II-enzymes via the Nrf2/ARE pathway in vitro and in vivo. *Mol. Nutr. Food Res.* **2011**, *55*, 798–802. [CrossRef]
111. International Diabetes Federation. *IDF Diabetes Atlas 2021- 10th Edition* | *IDF Diabetes Atlas*; International Diabetes Federation: Brussels, Belgium, 2021.
112. Salazar-Martinez, E.; Willett, W.C.; Ascherio, A.; Manson, J.A.E.; Leitzmann, M.F.; Stampfer, M.J.; Hu, F.B. Coffee consumption and risk for type 2 Diabetes Mellitus. *Ann. Intern. Med.* **2004**, *140*, 1–8. [CrossRef]
113. Agardh, E.E.; Carlsson, S.; Ahlbom, A.; Efendic, S.; Grill, V.; Hammar, N.; Hilding, A.; Östenson, C.G. Coffee consumption, type 2 Diabetes and impaired glucose tolerance in Swedish men and women. *J. Intern. Med.* **2004**, *255*, 645–652. [CrossRef]
114. Lin, W.Y.; Xavier Pi-Sunyer, F.; Chen, C.C.; Davidson, L.E.; Liu, C.S.; Li, T.C.; Wu, M.F.; Li, C.I.; Chen, W.; Lin, C.C. Coffee consumption is inversely associated with type 2 Diabetes in Chinese. *Eur. J. Clin. Investig.* **2011**, *41*, 659–666. [CrossRef]
115. Pereira, M.A.; Parker, E.D.; Folsom, A.R. Coffee consumption and risk of type 2 Diabetes Mellitus: An 11-year prospective study of 28 812 postmenopausal women. *Arch. Intern. Med.* **2006**, *166*, 1311–1316. [CrossRef] [PubMed]
116. Tuomilehto, J.; Hu, G.; Bidel, S.; Lindström, J.; Jousilahti, P. Coffee consumption and risk of type 2 Diabetes Mellitus among middle-aged Finnish men and women. *J. Am. Med. Assoc.* **2004**, *291*, 1213–1219. [CrossRef] [PubMed]
117. van Dam, R.M. Coffee consumption and risk of type 2 Diabetes, cardiovascular diseases, and cancer. *Appl. Physiol. Nutr. Metab.* **2008**, *33*, 1269–1283. [CrossRef] [PubMed]
118. van Dam, R.M.; Feskens, E.J.M. Coffee consumption and risk of type 2 Diabetes Mellitus. *Lancet* **2002**, *360*, 1477–1478. [CrossRef]
119. Meng, S.; Cao, J.; Feng, Q.; Peng, J.; Hu, Y. Roles of chlorogenic acid on regulating glucose and lipids metabolism: A review. *Evid.-Based Complement. Altern.* **2013**, *11*, 801457. [CrossRef] [PubMed]
120. Huxley, R.; Man Ying Lee, C.; Barzi, F.; Timmermeister, L.; Czernichow, S.; Perkovic, V.; Grobbee, D.E.; Batty, D.; Woodward, M. Coffee, decaffeinated coffee, and tea consumption in relation to incident type 2 Diabetes Mellitus a systematic review with meta-analysis. *Arch. Intern. Med.* **2009**, *169*, 2053–2063. [CrossRef] [PubMed]
121. Bakuradze, T.; Boehm, N.; Janzowski, C.; Lang, R.; Hofmann, T.; Stockis, J.P.; Albert, F.W.; Stiebitz, H.; Bytof, G.; Lantz, I.; et al. Antioxidant-rich coffee reduces DNA damage, elevates glutathione status and contributes to weight control: Results from an intervention study. *Mol. Nutr. Food Res.* **2011**, *55*, 793–797. [CrossRef] [PubMed]
122. Shearer, J.; Sellars, E.A.; Farah, A.; Graham, T.E.; Wasserman, D.H. Effects of chronic coffee consumption on glucose kinetics in the conscious rat. *Can. J. Physiol. Pharmacol.* **2007**, *85*, 823–830. [CrossRef]
123. Kurth-Kraczek, E.J.; Hirshman, M.F.; Goodyear, L.J.; Winder, W.W. 5'AMP-activated protein kinase activation causes GLUT4 translocation in skeletal muscle. *Diabetes* **1999**, *48*, 1667–1671. [CrossRef]
124. Kahn, B.B.; Alquier, T.; Carling, D.; Hardie, D.G. AMP-activated protein kinase: Ancient energy gauge provides clues to modern understanding of metabolism. *Cell Metabol.* **2005**, *1*, 15–25. [CrossRef]
125. World Health Organization (WHO). The Top 10 Causes of Death. Available online: <https://www.who.int/en/news-room/fact-sheets/detail/the-top-10-causes-of-death> (accessed on 11 April 2022).
126. Mikami, Y.; Yamazawa, T. Chlorogenic acid, a polyphenol in coffee, protects neurons against glutamate neurotoxicity. *Life Sci.* **2015**, *139*, 69–74. [CrossRef] [PubMed]
127. Roshan, H.; Nikpayam, O.; Sedaghat, M.; Sohrab, G. Effects of green coffee extract supplementation on anthropometric indices, glycaemic control, blood pressure, lipid profile, insulin resistance and appetite in patients with the Metabolic Syndrome: A randomised clinical trial. *Br. J. Nutr.* **2018**, *119*, 250–258. [CrossRef] [PubMed]
128. Tsai, K.L.; Hung, C.H.; Chan, S.H.; Hsieh, P.L.; Ou, H.C.; Cheng, Y.H.; Chu, P.M. Chlorogenic acid protects against OxLDL-induced oxidative damage and mitochondrial dysfunction by modulating SIRT1 in endothelial cells. *Mol. Nutr. Food Res.* **2018**, *62*, 1700928. [CrossRef] [PubMed]
129. Taguchi, K.; Hida, M.; Matsumoto, T.; Ikeuchi-Takahashi, Y.; Onishi, H.; Kobayashi, T. Effect of short-term polyphenol treatment on endothelial dysfunction and thromboxane A2 levels in streptozotocin-induced diabetic mice. *Biol. Pharm. Bull.* **2014**, *37*, 1056–1061. [CrossRef]
130. Sato, Y.; Itagaki, S.; Kurokawa, T.; Ogura, J.; Kobayashi, M.; Hirano, T.; Sugawara, M.; Iseki, K. In vitro and in vivo antioxidant properties of chlorogenic acid and caffeic acid. *Int. J. Pharm.* **2011**, *403*, 136–138. [CrossRef]

131. Kanegae, M.P.P.; da Fonseca, L.M.; Brunetti, I.L.; de Oliveira Silva, S.; Ximenes, V.F. The reactivity of ortho-methoxy-substituted catechol radicals with sulfhydryl groups: Contribution for the comprehension of the mechanism of inhibition of NADPH oxidase by apocynin. *Biochem. Pharmacol.* **2007**, *74*, 457–464. [[CrossRef](#)]
132. Antonio, A.G.; Moraes, R.S.; Perrone, D.; Maia, L.C.; Santos, K.R.N.; Iório, N.L.P.; Farah, A. Species, roasting degree and decaffeination influence the antibacterial activity of coffee against *Streptococcus mutans*. *Food Chem.* **2010**, *118*, 782–788. [[CrossRef](#)]
133. Almeida, A.A.P.; Naghetini, C.C.; Santos, V.R.; Antonio, A.G.; Farah, A.; Glória, M.B.A. Influence of natural coffee compounds, coffee extracts and increased levels of caffeine on the inhibition of *Streptococcus mutans*. *Food Res. Int.* **2012**, *49*, 459–461. [[CrossRef](#)]
134. Li, G.; Wang, X.; Xu, Y.; Zhang, B.; Xia, X. Antimicrobial effect and mode of action of chlorogenic acid on *Staphylococcus aureus*. *Eur. Food Res. Technol.* **2014**, *238*, 589–596. [[CrossRef](#)]
135. Lou, Z.; Wang, H.; Zhu, S.; Ma, C.; Wang, Z. Antibacterial activity and mechanism of action of chlorogenic acid. *J. Food Sci.* **2011**, *76*, M398–M403. [[CrossRef](#)]
136. Sung, W.S.; Lee, D.G. Antifungal action of chlorogenic acid against pathogenic fungi, mediated by membrane disruption. *Pure Appl. Chem.* **2010**, *82*, 219–226. [[CrossRef](#)]
137. Khan, M.T.H.; Ather, A.; Thompson, K.D.; Gambari, R. Extracts and molecules from medicinal plants against herpes simplex viruses. *Antivir. Res.* **2005**, *67*, 107–119. [[CrossRef](#)] [[PubMed](#)]
138. Chiang, L.C.; Chiang, W.; Chang, M.Y.; Ng, L.T.; Lin, C.C. Antiviral activity of plantago major extracts and related compounds in vitro. *Antivir. Res.* **2002**, *55*, 53–62. [[CrossRef](#)]
139. Tamura, H.; Akioka, T.; Ueno, K.; Chujyo, T.; Okazaki, K.I.; King, P.J.; Robinson, W.E. Anti-Human immunodeficiency virus activity of 3,4,5-tricaffeoylquinic acid in cultured cells of lettuce leaves. *Mol. Nutr. Food Res.* **2006**, *50*, 396–400. [[CrossRef](#)]
140. Honjo, S.; Kono, S.; Coleman, M.P.; Shinchi, K.; Sakurai, Y.; Todoroki, I.; Umeda, T.; Wakabayashi, K.; Imanishi, K.; Nishikawa, H.; et al. Coffee consumption and serum aminotransferases in middle-aged Japanese men. *J. Clin. Epidemiol.* **2001**, *54*, 823–829. [[CrossRef](#)]
141. Wadhawan, M.; Anand, A.C. Coffee and liver disease. *J. Clin. Exp. Hepatol.* **2016**, *6*, 40–46. [[CrossRef](#)]
142. La Vecchia, C. Coffee, liver enzymes, cirrhosis and liver cancer. *J. Hepatol.* **2005**, *42*, 444–446. [[CrossRef](#)]
143. Bravi, F.; Bosetti, C.; Tavani, A.; Gallus, S.; La Vecchia, C. Coffee reduces risk for hepatocellular carcinoma: An updated meta-analysis. *Clin. Gastroenterol. Hepatol.* **2013**, *11*, 1413–1421.e1. [[CrossRef](#)]
144. Larsson, S.C.; Wolk, A. Coffee consumption and risk of liver cancer: A meta-analysis. *Gastroenterology* **2007**, *132*, 1740–1745. [[CrossRef](#)]
145. Yun, N.; Kang, J.W.; Lee, S.M. Protective effects of chlorogenic acid against ischemia/reperfusion injury in rat liver: Molecular evidence of its antioxidant and anti-inflammatory properties. *J. Nutr. Biochem.* **2012**, *23*, 1249–1255. [[CrossRef](#)]
146. Ji, L.; Jiang, P.; Lu, B.; Sheng, Y.; Wang, X.; Wang, Z. Chlorogenic acid, a dietary polyphenol, protects acetaminophen-induced liver injury and its mechanism. *J. Nutr. Biochem.* **2013**, *24*, 1911–1919. [[CrossRef](#)] [[PubMed](#)]
147. Shi, H.; Shi, A.; Dong, L.; Lu, X.; Wang, Y.; Zhao, J.; Dai, F.; Guo, X. Chlorogenic acid protects against liver fibrosis in vivo and in vitro through inhibition of oxidative stress. *Clin. Nutr.* **2016**, *35*, 1366–1373. [[CrossRef](#)]
148. Machado, S.R.; Parise, E.R.; de Carvalho, L. Coffee has hepatoprotective benefits in Brazilian patients with chronic Hepatitis C even in lower daily consumption than in American and European populations. *Braz. J. Infect. Dis.* **2014**, *18*, 170–176. [[CrossRef](#)] [[PubMed](#)]
149. Gibson, G.R.; Hutkins, R.; Sanders, M.E.; Prescott, S.L.; Reimer, R.A.; Salminen, S.J.; Scott, K.; Stanton, C.; Swanson, K.S.; Cani, P.D.; et al. Expert consensus document: The International Scientific Association for Probiotics and Prebiotics (ISAPP) consensus statement on the definition and scope of Prebiotics. *Nat. Rev. Gastroenterol. Hepatol.* **2017**, *14*, 491–502. [[CrossRef](#)] [[PubMed](#)]
150. Plamada, D.; Vodnar, D.C. Polyphenols—Gut microbiota interrelationship: A transition to a new generation of prebiotics. *Nutrients* **2022**, *14*, 137. [[CrossRef](#)]
151. Allsopp, P.; Possemiers, S.; Campbell, D.; Oyarzábal, I.S.; Gill, C.; Rowland, I. An exploratory study into the putative prebiotic activity of fructans isolated from *Agave angustifolia* and the associated anticancer activity. *Anaerobe* **2013**, *22*, 38–44. [[CrossRef](#)]
152. Saha, D.C.; Reimer, R.A. long-term intake of a high prebiotic fiber diet but not high protein reduces metabolic risk after a high fat challenge and uniquely alters gut microbiota and hepatic gene expression. *Nutr. Res.* **2014**, *34*, 789–796. [[CrossRef](#)]
153. Kumar, V.P.; Prashanth, K.V.H.; Venkatesh, Y.P. Structural analyses and immunomodulatory properties of fructo-oligosaccharides from onion (*Allium cepa*). *Carbohydr. Polym.* **2015**, *117*, 115–122. [[CrossRef](#)]
154. Kelly-Quagliana, K.A.; Nelson, P.D.; Buddington, R.K. Dietary oligofructose and inulin modulate immune functions in mice. *Nutr. Res.* **2003**, *23*, 257–267. [[CrossRef](#)]
155. Munjal, U.; Gleib, M.; Pool-Zobel, B.L.; Scharlau, D. Fermentation products of inulin-type fructans reduce proliferation and induce apoptosis in human colon tumour cells of different stages of carcinogenesis. *Br. J. Nutr.* **2009**, *102*, 663–671. [[CrossRef](#)]
156. Delzenne, N.M.; Cani, P.D.; Everard, A.; Neyrinck, A.M.; Bindels, L.B. Gut microorganisms as promising targets for the management of type 2 Diabetes. *Diabetologia* **2015**, *58*, 2206–2217. [[CrossRef](#)] [[PubMed](#)]
157. Cluny, N.L.; Eller, L.K.; Keenan, C.M.; Reimer, R.A.; Sharkey, K.A. Interactive effects of oligofructose and obesity predisposition on gut hormones and microbiota in diet-induced obese rats. *Obesity* **2015**, *23*, 769–778. [[CrossRef](#)] [[PubMed](#)]
158. Griffin, I.J.; Hicks, P.M.D.; Heaney, R.P.; Abrams, S.A. Enriched chicory inulin increases calcium absorption mainly in girls with lower calcium absorption. *Nutr. Res.* **2003**, *23*, 901–909. [[CrossRef](#)]

159. Kellow, N.J.; Coughlan, M.T.; Savige, G.S.; Reid, C.M. Effect of dietary prebiotic supplementation on advanced glycation, insulin resistance and inflammatory biomarkers in adults with pre-Diabetes: A study protocol for a double-blind placebo-controlled randomised crossover clinical trial. *BMC Endocr. Disord.* **2014**, *14*, 55. [[CrossRef](#)] [[PubMed](#)]
160. Rechner, A.R.; Spencer, J.P.E.; Kuhnle, G.; Hahn, U.; Rice-Evans, C.A. Novel biomarkers of the metabolism of caffeic acid derivatives in vivo. *Free Radic. Biol. Med.* **2001**, *30*, 1213–1222. [[CrossRef](#)]
161. Gonthier, M.P.; Remesy, C.; Scalbert, A.; Cheyrier, V.; Souquet, J.M.; Poutanen, K.; Aura, A.M. Microbial metabolism of caffeic acid and its esters chlorogenic and caftaric acids by human faecal microbiota in vitro. *Biomed. Pharmacother.* **2006**, *60*, 536–540. [[CrossRef](#)]
162. Farah, A.; de Paula Lima, J. Chlorogenic acids: Daily consumption through coffee, metabolism and potential health effects. In *Coffee: Consumption and Health Implications*, 1st ed.; Farah, A., Ed.; Royal Society of Chemistry: London, UK, 2019; pp. 364–415. [[CrossRef](#)]
163. Monteiro, M.; Farah, A.; Perrone, D.; Trugo, L.C.; Donangelo, C. Chlorogenic acid compounds from coffee are differentially absorbed and metabolized in humans. *J. Nutr.* **2007**, *137*, 2196–2201. [[CrossRef](#)]
164. Lafay, S.; Gil-Izquierdo, A.; Manach, C.; Morand, C.; Besson, C.; Scalbert, A. Chlorogenic acid is absorbed in its intact form in the stomach of rats. *J. Nutr.* **2006**, *136*, 1192–1197. [[CrossRef](#)]
165. Olthof, M.R.; Hollman, P.C.H.; Katan, M.B. Chlorogenic acid and caffeic acid are absorbed in humans. *J. Nutr.* **2001**, *131*, 66–71. [[CrossRef](#)]
166. Farah, A.; Duarte, G. Bioavailability and metabolism of chlorogenic acids from coffee. In *Coffee in Health and Disease Prevention*, 1st ed.; Preedy, V.R., Ed.; Academic Press: Oxford, UK, 2015; pp. 789–801.
167. Nabavi, S.F.; Tejada, S.; Setzer, W.N.; Gortzi, O.; Sureda, A.; Braidly, N.; Daglia, M.; Manayi, A.; Nabavi, S.M. Chlorogenic acid and mental diseases: From chemistry to medicine. *Curr. Neuropharmacol.* **2017**, *15*, 471. [[CrossRef](#)]
168. Farah, A.; Monteiro, M.; Donangelo, C.M.; Lafay, S. Chlorogenic acids from green coffee extract are highly bioavailable in humans. *J. Nutr.* **2008**, *138*, 2309–2315. [[CrossRef](#)] [[PubMed](#)]
169. Mubarak, A.; Bondonno, C.P.; Liu, A.H.; Considine, M.J.; Rich, L.; Mas, E.; Croft, K.D.; Hodgson, J.M. Acute effects of chlorogenic acid on nitric oxide status, endothelial function, and blood pressure in healthy volunteers: A randomized trial. *J. Agric. Food Chem.* **2012**, *60*, 9130–9136. [[CrossRef](#)] [[PubMed](#)]
170. Erk, T.; Williamson, G.; Renouf, M.; Marmet, C.; Steiling, H.; Dionisi, F.; Barron, D.; Melcher, R.; Richling, E. Dose-dependent absorption of chlorogenic acids in the small intestine assessed by coffee consumption in ileostomists. *Mol. Nutr. Food Res.* **2012**, *56*, 1488–1500. [[CrossRef](#)] [[PubMed](#)]
171. Clifford, M.N.; Kerimi, A.; Williamson, G. Bioavailability and metabolism of chlorogenic acids (Acyl-quinic acids) in humans. *Compr. Rev. Food Sci. Food Saf.* **2020**, *19*, 1299–1352. [[CrossRef](#)]
172. Vicentini, A.; Liberatore, L.; Mastrocola, D. Functional foods: Trends and development. *Ital. J. Food Sci.* **2016**, *28*, 338–352.
173. Yang, C.S.; Ho, C.T.; Zhang, J.; Wan, X.; Zhang, K.; Lim, J. Antioxidants: Differing meanings in food science and health science. *J. Agric. Food Chem.* **2018**, *66*, 3063–3068. [[CrossRef](#)]
174. Bagchi, D.; Moriyama, H.; Swaroop, A. *Green Coffee Bean Extract in Human Health*, 1st ed.; CRC-Press: Boca Raton, CA, USA, 2016; pp. 1–254. [[CrossRef](#)]
175. Budryn, G.; Zyzelewicz, D.; Nebesny, E.; Oracz, J.; Krysiak, W. Influence of addition of green tea and green coffee extracts on the properties of fine yeast pastry fried products. *Food Res. Int.* **2013**, *50*, 149–160. [[CrossRef](#)]
176. Corso, M.P.; Vignoli, J.A.; Benassi, M.D.T. Development of an instant coffee enriched with chlorogenic acids. *J. Food Sci. Technol.* **2016**, *53*, 1380–1388. [[CrossRef](#)]
177. Sęczyk, Ł.; Świeca, M.; Gawlik-Dziki, U. Soymilk enriched with green coffee phenolics—antioxidant and nutritional properties in the light of phenolics–food matrix interactions. *Food Chem.* **2017**, *223*, 1–7. [[CrossRef](#)]
178. Mukkundur Vasudevaiah, A.; Chaturvedi, A.; Kulathooran, R.; Dasappa, I. Effect of green coffee extract on rheological, physico-sensory and antioxidant properties of bread. *J. Food Sci. Technol.* **2017**, *54*, 1827–1836. [[CrossRef](#)]
179. Rahpeyma, E.; Sekhavatizadeh, S.S. Effects of encapsulated green coffee extract and canola oil on liquid Kashk quality. *Foods Raw Mater.* **2020**, *8*, 40–51. [[CrossRef](#)]
180. Zohreh, D. Properties of dark chocolate enriched with free and encapsulated chlorogenic acids extracted from green coffee. *Braz. J. Food Technol.* **2020**, *23*, e2019118. [[CrossRef](#)]
181. Pimpley, V.A.; Maity, S.; Murthy, P.S. Green coffee polyphenols in formulations of functional yoghurt and their quality attributes. *Int. J. Dairy Technol.* **2022**, *75*, 159–170. [[CrossRef](#)]
182. Vignoli, J.A.; Bassoli, D.G.; Benassi, M.T. Antioxidant activity, polyphenols, caffeine and melanoidins in soluble coffee: The influence of processing conditions and raw material. *Food Chem.* **2011**, *124*, 863–868. [[CrossRef](#)]
183. Yu, L.; Nanguet, A.L.; Beta, T. Comparison of antioxidant properties of refined and whole wheat flour and bread. *Antioxidants* **2013**, *2*, 370–383. [[CrossRef](#)]
184. Prückler, M.; Siebenhandl-Ehn, S.; Apprich, S.; Höltinger, S.; Haas, C.; Schmid, E.; Kneifel, W. Wheat bran-based biorefinery 1: Composition of wheat bran and strategies of functionalization. *LWT—Food Sci. Technol.* **2014**, *56*, 211–221. [[CrossRef](#)]
185. Zain, M.Z.M.; Baba, A.S.; Shori, A.B. Effect of polyphenols enriched from green coffee bean on antioxidant activity and sensory evaluation of bread. *J. King Saud Univ. Sci.* **2018**, *30*, 278–282. [[CrossRef](#)]

186. Dziki, D.; Gawlik-Dziki, U.; Pecio, Ł.; Różyło, R.; Świeca, M.; Krzykowski, A.; Rudy, S. Ground green coffee beans as a functional food supplement—Preliminary study. *LWT—Food Sci. Technol.* **2015**, *63*, 691–699. [[CrossRef](#)]
187. Hashemi Gahruie, H.; Eskandari, M.H.; Mesbahi, G.; Hanifpour, M.A. Scientific and technical aspects of yogurt fortification: A review. *Food Sci. Hum. Wellness* **2015**, *4*, 1–8. [[CrossRef](#)]
188. Dönmez, Ö.; Mogol, B.A.; Gökmen, V. Syneresis and rheological behaviors of set yogurt containing green tea and green coffee powders. *J. Dairy Sci.* **2017**, *100*, 901–907. [[CrossRef](#)]
189. Cheynier, V. Polyphenols in foods are more complex than often thought. *Am. J. Clin. Nutr.* **2005**, *81*, 223S–229S. [[CrossRef](#)] [[PubMed](#)]