

Extracellular mitochondria released from traumatized brains induced platelet procoagulant activity

Zilong Zhao,^{1,2} Yuan Zhou,^{1,2} Tristan Hilton,¹ Fanjian Li,² Cha Han,¹ Li Liu,² Hengjie Yuan,² Ying Li,² Xin Xu,¹ Xiaoping Wu,¹ Fangyi Zhang,³ Perumal Thiagarajan,⁴ Andrew Cap,⁵ Fu-Dong Shi,^{2,6} Jianning Zhang² and Jing-fei Dong^{1,7}

¹BloodWorks Research Institute, Seattle, WA, USA; ²Tianjin Institute of Neurology, Departments of Neurosurgery and Neurology, Tianjin Medical University General Hospital, Tianjin, China; ³Department of Neurosurgery, University of Washington School of Medicine, Seattle, WA, USA; ⁴Departments of Medicine and Pathology, Michael E. DeBakey VA Medical Center and Baylor College of Medicine, Houston, TX, USA; ⁵US Army Institute of Surgical Research, San Antonio, TX, USA; ⁶Department of Neurology, Barrow Neurological Institute, St. Joseph's Hospital and Medical Center, Phoenix, AZ, USA and ⁷Division of Hematology, Department of Medicine, University of Washington, School of Medicine, Seattle, WA, USA

ABSTRACT

Coagulopathy often develops soon after acute traumatic brain injury and its cause remains poorly understood. We have shown that injured brains release cellular microvesicles that disrupt the endothelial barrier and induce consumptive coagulopathy. Morphologically intact extracellular mitochondria accounted for 55.2% of these microvesicles, leading to the hypothesis that these extracellular mitochondria are metabolically active and serve as a source of oxidative stress that activates platelets and renders them procoagulant. In testing this hypothesis experimentally, we found that the extracellular mitochondria purified from brain trauma mice and those released from brains subjected to freeze-thaw injury remained metabolically active and produced reactive oxygen species. These extracellular mitochondria bound platelets through the phospholipid-CD36 interaction and induced α -granule secretion, microvesiculation, and procoagulant activity in an oxidant-dependent manner, but failed to induce aggregation. These results define an extracellular mitochondria-induced and redox-dependent intermediate phenotype of platelets that contribute to the pathogenesis of traumatic brain injury-induced coagulopathy and inflammation.

Introduction

Traumatic brain injury (TBI) induces coagulopathy that promotes secondary bleeding and propagates cerebral injury,¹ resulting in poor clinical outcomes of the patients.²⁻⁶ Laboratory findings suggest that coagulopathy results from a hypercoagulable state that rapidly develops into consumptive coagulopathy.^{1,7} This consumptive coagulopathy has been recapitulated in animal models of TBI with thrombotic and hemorrhagic manifestations in the pulmonary and cerebral microvasculature.^{8,9} Despite strong clinical and laboratory evidence of its presence and association with poor clinical outcomes, the pathogenesis of TBI-associated coagulopathy remains poorly understood.

We have recently shown in mouse models that cellular microvesicles released from traumatically injured brains disrupt endothelial cell junctions through a synergistic action with platelets.⁸ These brain-derived microvesicles are highly procoagulant due to the abundant expression of surface-exposed anionic phospholipids and tissue factor. The sudden and substantial release of brain-derived microvesicles into the circulation results in a consumptive coagulopathy that is characterized by progressive fibrinogen depletion from plasma and fibrin deposition in the vasculature.⁸



Haematologica 2020
Volume 105(1):209-217

Correspondence:

JING-FEI DONG
jfdong@bloodworksnw.org

JIANING ZHANG
jianningzhang@hotmail.com

Received: December 18, 2018.

Accepted: April 9, 2019.

Pre-published: April 11, 2019.

doi:10.3324/haematol.2018.214932

Check the online version for the most updated information on this article, online supplements, and information on authorship & disclosures: www.haematologica.org/content/105/1/207

©2020 Ferrata Storti Foundation

Material published in *Haematologica* is covered by copyright. All rights are reserved to the Ferrata Storti Foundation. Use of published material is allowed under the following terms and conditions:

<https://creativecommons.org/licenses/by-nc/4.0/legalcode>. Copies of published material are allowed for personal or internal use. Sharing published material for non-commercial purposes is subject to the following conditions: <https://creativecommons.org/licenses/by-nc/4.0/legalcode>, sect. 3. Reproducing and sharing published material for commercial purposes is not allowed without permission in writing from the publisher.



We also found that $55.2 \pm 12.6\%$ of annexin-V-binding microvesicles in peripheral blood of TBI mice were morphologically intact extracellular mitochondria (exMT).⁹ These exMT promoted coagulation through the surface-exposed anionic phospholipid cardiolipin. The findings raise an important question: are these exMT metabolically active and, if so, do they affect platelet function through redox-dependent mechanisms? The question is raised because exMT can be viable, transferred between cells, and influence the function of target cells.¹⁰⁻¹³ The influence of mitochondria on target cells has mostly been reported as protective,^{10,12,13} likely due to increased energy supply. However, reactive oxygen species (ROS) are generated during ATP production in mitochondria¹⁴ and are known to activate platelets.¹⁵⁻¹⁷ Several studies have shown that platelets in patients with TBI are activated but aggregate poorly.¹⁸ This platelet phenotype has also been reproduced in rat and swine models of TBI, in which platelets are activated,^{19,20} accumulate on the pia mater,²¹ and contribute to thrombosis in the lesion boundary zone.²² The underlying mechanism for this unique TBI-associated platelet phenotype remains poorly understood. Here we discuss results from a study that was designed to investigate the effect of exMT on platelet activation and procoagulant activity through *in vitro* experiments and in mouse models.

Methods

Mouse models

C57BL/6J male mice (12-20 weeks and 22-25 g), obtained from the Jackson Laboratory (Bar Harbor, ME, USA) were subjected to fluid percussion injury⁸ and blood samples were collected to quantify plasma exMT (*Online Supplementary Methods*).⁹ This mouse protocol was approved by the Institutional Animal Care and Use Committee of the BloodWorks Research Institute.

Flow cytometry

Platelet activation

We used an LSR II flow cytometer (Beckon Dickinson, San Jose, CA, USA) to detect platelet activation through several measurements: CD62p expression, the binding of PAC-1 antibody (BD Biosciences), which recognizes the active conformation of the fibrinogen receptor $\alpha IIb\beta 3$, annexin V binding,⁹ platelet-bound fibrinogen and coagulation factor V,^{25,26} and the formation of platelet-leukocyte complexes (*Online Supplementary Methods*).

Calcium influx

We used flow cytometry to measure the exMT-induced calcium influx in platelets labeled with 5 μM eFluor 514 Calcium Sensor Dye (Bioscience, San Diego, CA, USA) for 10 min at 37°C.⁹

Reactive oxygen species production

ExMT were suspended in Ca^{2+} -free HEPES buffered Tyrode solution (138 mM NaCl, 5.5 mM glucose, 10 mM HEPES, 12 mM NaHCO_3 , 2.9 mM KCl, 0.4 mM NaH_2PO_4 , 0.4 mM MgCl_2 , and 0.1% BSA; pH 7.2) and incubated for 30 min at 37°C with 10 μM of DCFH-DA dye (ThermoFisher).²⁴ As controls, labeled exMT were treated with either 50 μM N-acetylcysteine (NAC, Life Technologies, Grand Island, NY, USA), which is a precursor of the antioxidant glutathione, or 200 μM tert-butyl hydroperoxide (TBHP, Life Technologies), which increases intracellular ROS production.²⁷ ExMT fixed with 5% paraformaldehyde were also examined as a control.

Image flow cytometry for the extracellular mitochondria-platelet interaction

As we previously described,²⁸ platelet-rich plasma was incubated for 30 min at 37°C with exMT labeled with MitoTracker Green and an APC-conjugated CD41a antibody (BD Bioscience). The platelets were washed in phosphate-buffered saline and fixed in 1% paraformaldehyde. Images were acquired at 60 x magnification to collect $\geq 20,000$ cells from each sample using the Amnis ImageStream® X Mk II system (Amnis, Seattle, WA, USA). To distinguish exMT bound to the platelet surface from those internalized, platelets were digested with 1% trypsin for 10 min at 37°C after incubation with exMT. Mouse mitochondrial-specific DNA was then amplified from these exMT-treated and trypsinized platelets using a protocol modified from our previous study.⁹

CD36-extracellular mitochondria interaction

For *in vitro* experiments, platelet-rich plasma was incubated for 30 min at 37°C with a monoclonal CD36 antibody (Abcam, ab17044, Cambridge, MA, USA) or isotype IgG, followed by incubation with MitoTracker Green-labeled exMT and a PE-CD41a antibody for 30 min. For *in vivo* experiments, blood samples from non-injured mice infused with MitoTracker Green-labeled exMT were analyzed for exMT-bound platelets (MitoTracker Green⁺/CD41a⁺), platelet counts (CD41a⁺ platelet counts in 60 s), platelet CD62p expression, and platelet-leukocyte complexes (CD41a⁺/CD45⁺). The CD36 antibody used in this study recognizes both mouse and human CD36.^{29,30}

Statistical analysis

Categorical (frequency) variables were expressed as percentages and continuous variables were expressed as the mean \pm standard error of mean. The quantitative data were analyzed using SigmaPlot V. 11.2 (SYSTAT Software Inc., San Jose, CA, USA) for paired *t* tests, one way or repeated measures analysis of variance (ANOVA) as specified in each analysis. A *P* value of ≤ 0.05 was considered statistically significant.

Results

Extracellular mitochondria bound platelets *in vivo* and *in vitro*

Structurally intact (MitoTracker Green⁺) (Figure 1A) and cardiolipin-exposed (Figure 1B) exMT were detected in the peripheral blood of mice with acute TBI. The exMT formed complexes with approximately $12.3 \pm 5.8\%$ of circulating platelets that remained detectable for 6 h after TBI (Figure 1C). Low levels of exMT-platelet complexes were also detected in sham-treated mice and likely resulted from surgery-induced injuries. When co-incubated for 30 min at 37°C *in vitro*, purified exMT from brains subjected to freeze-thaw injury also formed complexes with platelets (Figure 1D) in a dose-dependent manner (Figure 1E). Transmission electron microscopy showed that exMT interacted with the body (Figure 1F) and filopodia (Figure 1G) of platelets.

The Amnis® imaging flow cytometer detected exMT on the surface of platelets (Figure 2A, top panel), being endocytosed by platelets (Figure 2A, middle panel) and fused with platelet membranes (Figure 2A, bottom panel). The internalization of exMT by platelets was further validated by co-incubation of human platelets with mouse exMT for 30 min at 37°C followed by the amplification of mouse mitochondria-derived DNA from exMT-treated platelets that were trypsinized to remove surface-bound exMT




Figure 1. Extracellular mitochondria were released from injured cells and bound platelets. (A, B) Extracellular mitochondria (exMT) from mice with traumatic brain injury (TBI) were detected by MitoTracker Green (# exMT/100 μ L plasma) (A) and anti-cardiolipin antibody (n=12, paired t test) (B). (C) Platelet-exMT complexes were detected in the blood of TBI mice using CD41a and TOM22 antibodies (n=8, paired t test). (D, E) After co-incubation for 30 min at 37 $^{\circ}$ C, exMT also bound platelets *in vitro*, as shown by a representative histogram (D) in a dose-dependent manner (n=30, one-way analysis of variance) (E). (F, G) Transmission electron microscopy (see *Online Supplementary Methods*) showed exMT binding to the body (F) and filopodia (G) of platelets (bar = 500 nm; arrow: mitochondria; *: platelet; representatives of 98 images reviewed).

(Figure 2B). Quantitatively, $13.7 \pm 9.1\%$ of platelets were exMT-bound after 30 min co-incubation (Figure 2C), which was consistent with the level of exMT-bound platelets found in TBI mice (Figure 1C). This exMT-platelet interaction was blocked by the phosphatidylserine-binding proteins annexin V and lactadherin (Figure 2C)^{28,31} and partially by a CD36 antibody (Figure 2D), which blocks the CD36-mediated binding of endothelial microvesicles to platelets,²⁹ suggesting that exMT bind platelets through cardiolipin exposed on the mitochondria and CD36 expressed on platelets. Furthermore, ATP production was dose-dependently increased in the exMT-treated platelets at levels that were significantly higher than in platelets activated with ADP (Figure 2E). We were unable to determine whether the ATP increase was caused by platelet-bound exMT or by exMT-induced ATP production of platelets. Finally, when added to platelets that had been treated with exMT, FITC-conjugated annexin V bound $16.2 \pm 6.3\%$ of platelets (Figure 2F).

Extracellular mitochondria were metabolically viable and activated platelets

We found that exMT from TBI mice stimulated platelets to express CD62p and this activity was partially blocked by the antioxidant glutathione (GSH) (Figure 3A). One pit-

fall of this experiment was that exMT purified from plasma of TBI mice were derived from multiple cells. It was technically challenging to separate brain-derived exMT from those of other cells, without damaging the viability of exMT. To address this concern, we tested exMT from brains subjected to freeze-thaw injury *in vitro*.⁹ These exMT produced ATP (Figure 3B) and ROS (Figure 3C). The ROS production was blocked by 50 μ M of the anti-oxidant NAC and enhanced by 200 μ M of the oxidant TBHP. The two agents have previously been shown to block and enhance ROS production of mammalian cells, respectively.^{32,27} This metabolic viability was further validated by the lack of ROS production from paraformaldehyde-fixed exMT (Figure 3C). After 30 min of incubation with exMT, both mouse (*Online Supplementary Figure S2*) and human platelets (Figure 3D) expressed CD62p on their surfaces. The exMT-induced CD62p expression was reduced by the antioxidants GSH (20 μ M), L-cysteine (0.5 mM), and NEM (2 mM) (Figure 3E). Testing the three thiol-modifying agents was necessary because GSH and L-cysteine reversibly interact with extracellular and intracellular oxidants, respectively, whereas NEM forms irreversible sulfur bonds with thiols. The exMT-treated platelets also increased calcium influx (*Online Supplementary Figure S3*), formed complexes with leukocytes (Figure 3F), and

released von Willebrand factor (Figure 3G). The exMT-induced CD62 expression and platelet-leukocyte aggregation were detected in 10-23% of platelets, which was consistent with the percentages of exMT-bound platelets found in TBI mice (Figure 1). We used platelets stimulated with collagen and thrombin as the control because this exMT-induced platelet phenotype resembled that of “coated platelets”.^{33,34} Together, these data suggested that exMT from TBI mice and those released from brains subjected to freeze-thaw injury *in vitro* were metabolically viable and activated platelets in an oxidant-dependent manner.

Using hopping probe ion conductance microscopy, we continuously monitored morphological changes of platelets adherent to fibrinogen in real-time (Figure 4A, top panel). After stimulation with exMT, adherent platelets underwent drastic membrane disintegration as exemplified in the middle panel of Figure 4A. This exMT-induced membrane disintegration was prevented by 20 μ M GSH (Figure 4A, bottom row). Consistent with the exMT-induced platelet disruption, CD41a⁺ platelet-derived membrane microvesicles were detected in the supernatant of exMT-treated platelets (Figure 4B) and

platelet counts were reduced after treatment with exMT (Figure 4C). The production of platelet microvesicles and the reduction of platelet counts were partially blocked by the anti-oxidant GSH.

In contrast to their induction of α -granule secretion, exMT at comparable doses failed to induce platelet aggregation (Figure 4D) and did not enhance the formation of platelet thrombosis on the collagen matrix under arterial shear stress (*Online Supplementary Figure S4A-C*). Furthermore, the exMT-treated platelets aggregated normally in response to collagen (Figure 4E) and ADP (Figure 4F), and were moderately primed for activation by sub-threshold concentrations of ADP and collagen (*Online Supplementary Figure S4D*). Neither the binding of PAC-1 antibody (*Online Supplementary Figure S5*), which recognizes the active conformation of integrin α IIb β 3, nor the surface density of the integrin α IIb β 3 (Figure 4G) was changed after exMT treatment, as compared to platelets stimulated with collagen.

Extracellular mitochondria-treated platelets were procoagulant

ExMT-treated platelets also expressed anionic phos-




Figure 2. Anionic phospholipid and CD36 mediated the extracellular mitochondria-platelet interaction. (A) Amnis® flow cytometric images show extracellular mitochondria (exMT) binding a platelet (top panel), endocytosed by a platelet (middle panel), and fused with platelet membrane (bottom panel) after 30 min co-incubation at 37 °C (representative images from 20,000 images randomly selected). (B) Platelets internalized exMT in a dose dependent manner, as determined by the presence of mouse mitochondrial DNA in mouse exMT-treated human platelets that were trypsinized to remove surface-bound exMT (n=20, one-way analysis of variance, ANOVA). (C, D) The formation of complexes between platelets and MitoTracker-Green⁺ exMT (1:1 ratio) was blocked by 20 μ g/mL of annexin V or 200 μ g/mL of lactadherin (n=56, one-way ANOVA) (C) and by an anti-CD36 antibody (n=20, one-way ANOVA) (D). (E) ATP production was increased in exMT-treated platelets (n=24, one-way ANOVA). (F) FITC-conjugated annexin V bound platelets that were pretreated with exMT (left: a representative flow cytometric histogram; right: a summary from 12 experiments; paired Student t test).

pholipids as indicated by annexin V binding (Figure 2F) and had elevated levels of fibrinogen (Figure 5A) and coagulation factor Va on their surfaces (Figure 5B), suggesting that exMT-treated platelets could promote coagulation.^{35,34} Consistent with this notion, exMT-treated platelets accelerated clot formation at a level comparable to that of collagen-activated platelets but lower than that of purified phosphatidylserine micelles (Figure 5C). When added to platelet-rich plasma, exMT-treated platelets significantly accelerated and enhanced thrombin-induced clot retraction (Figure 5D), which measures the integrin α IIb β 3-dependent retraction of platelet-bound fibrin fibers.^{35,36} The effect of exMT was comparable to that of collagen-stimulated platelets. This procoagulant activity was independent of tissue factor, which was not detected on the surface of exMT-treated platelets (Online Supplementary Figure S6).

Anti-CD36 antibody blocked extracellular mitochondria-induced platelet activation

To measure exMT-induced platelet activation *in vivo*, non-injured mice were infused with purified exMT. Thirty minutes after infusion, platelets became exMT-bound

(Figure 6A), expressed CD62p (Figure 6B), and formed complexes with leukocytes (Figure 6C). Mice infused with exMT also developed thrombocytopenia (Figure 6D). MitoTracker-Green-labeled exMT were detected in approximately 50% of platelet-leukocyte aggregates (Figure 6E). These phenotypic changes of platelets were prevented by infusing mice with exMT, together with a CD36 antibody but not isotype IgG.

Discussion

We investigated whether morphologically intact but cardiolipin-exposed exMT are metabolically active and, if so, whether they activate platelets through ROS. We have made several novel observations that define an exMT-induced and redox-dependent intermediate phenotype of platelets.

First, despite the cardiolipin translocation,⁹ exMT found in TBI mice or released from brains subjected to freeze-thaw injury produced ATP and ROS (Figure 3B, C). Because metabolic activity of mitochondria requires a functional membrane, exMT that interact with platelets




Figure 3. Extracellular mitochondria induced α -granule secretion of platelets through reactive oxygen species. (A) Extracellular mitochondria (exMT) purified from mice with traumatic brain injury stimulated platelets to express CD62 and this effect was blocked by 20 μ M of glutathione (GSH) (n=45, one-way analysis of variance, ANOVA). (B, C) ExMT released from brains subjected to freeze-thaw injury produced ATP (n=4, one-way ANOVA) (B) and reactive oxygen species (ROS) (2.5×10^9 /mouse of exMT; mean fluorescence intensity) at 37 °C *in vitro*. The ROS production was blocked by the antioxidant N-acetylcysteine, enhanced by the oxidant tert-butyl hydroperoxide, and not detected in paraformaldehyde-fixed exMT (n=18, one-way ANOVA) (C). (D) Platelets incubated with exMT at a ratio of 1:1 or 1:5 for 30 min at 37 °C expressed CD62p (n=24, one-way ANOVA). (E) The CD62p expression was reduced by 20 μ M of GSH, 200 μ M of L-cysteine, or 2 mM of NEM (n=30, one-way ANOVA). (F, G) ExMT-treated platelets formed complexes with leukocytes (n=24, one-way ANOVA) (F) and released von Willebrand factor (n=24, one-way ANOVA, Online Supplementary Methods) (G). For platelet activation, platelets stimulated with 5 μ g/mL of collagen or 50 nM of thrombin were used as controls to mimic "coated platelets".³³ NAC: N-acetylcysteine; TBHP: tert-butyl hydroperoxide; MFI: mean fluorescence intensity; NEM: N-ethylmaleimide; vWF: von Willebrand factor.




Figure 4. Extracellular mitochondria induced platelet disintegration, not aggregation. Platelets adherent to immobilized fibrinogen were treated with phosphate-buffered saline (PBS), extracellular mitochondria (exMT), or exMT + 20 μ M of glutathione (GSH) for 30 min at 37 °C and repeatedly scanned for up to 80 min by hopping probe ion conductance microscopy. (A) The images show adherent platelets treated with PBS (top panel), exMT at a 1:1 ratio with platelets (middle panel; arrow: membrane disintegration of an adherent platelet), and GSH-treated exMT (bottom panel). The images are representative of three to six independent experiments. (B) The supernatants were collected and stained for CD41a⁺ microvesicles by flow cytometry (n=18, one-way analysis of variance, ANOVA). (C) Platelet counts before and after exMT treatment either alone or with GSH (n=15, one-way ANOVA). (D-F) ExMT induced minimal platelet aggregation (D) (n=24, one-way ANOVA), but exMT-treated platelets aggregated normally when stimulated with collagen (E) or ADP (F) (n=54, repeated measures ANOVA). (G) CD61 expression on exMT-treated platelets was comparable to that on untreated platelets but increased upon stimulation with 5 μ g/mL of collagen (n=24, one-way ANOVA). MV: microvesicles; MFI: mean fluorescence intensity.

are likely those with an intact membrane. The metabolically active exMT formed complexes with platelets (Figures 1 and 2) which remained detectable in the circulation for at least 6 h after TBI (Figure 1C).⁹ Cardiolipin on exMT and CD36 on platelets mediated the exMT-platelet interaction because: (i) cardiolipin is the dominant anionic phospholipid expressed on exMT⁹ and (ii) the exMT-platelet interaction was blocked by the phospholipid binding proteins annexin V and lactadherin (Figure 2C) and by an antibody against CD36 (Figure 2D), which is a phospholipid receptor³⁷ that is expressed on platelets^{38,39} and promotes phospholipid-mediated endocytosis.⁴⁰

Second, exMT induced platelets to secrete their α -granule proteins (Figures 3 and 5) but failed to induce them to aggregate (Figure 4D) or promote platelet thrombus formation on the collagen matrix under arterial shear stress (Online Supplementary Figure S4A-C). There are several possible explanations for this apparent discrepancy. (i)

Platelet-bound exMT may have interfered with the fibrinogen coupling of platelets through steric hindrance. This is unlikely because exMT-bound platelets aggregated after stimulation with collagen or ADP at levels comparable to those of platelets that had not been treated with exMT (Figure 4E, F). (ii) ExMT only activated 10–25% of platelets, insufficient to induce platelet aggregation. A similar effect was observed with platelets simultaneously treated with two antibodies against the integrins α IIb and β 3,⁴¹ suggesting that α -granule secretion alone is insufficient to induce platelet aggregation, but may prime platelets for activation by other agonists. (iii) Some or all exMT-stimulated platelets underwent drastic membrane disintegration to produce microvesicles (Figure 4A, B), thereby becoming unavailable for aggregation. The third possibility is supported by the reduction of platelet counts after exMT treatment (Figure 4C). This exMT-induced intermediate platelet phenotype resembles “coat-




Figure 5. Extracellular mitochondria induced procoagulant activity of platelets and enhanced clot retraction. (A-C) Platelets incubated with extracellular mitochondria (exMT) at a ratio of 1:1 or 1:5 for 30 min at 37 °C expressed increased amounts of fibrinogen (A) (n=24, one-way analysis of variance, ANOVA) and coagulation factor Va (B) (n=24, one-way ANOVA), and had shortened clotting time (C) (n=24, one-way ANOVA). Platelets stimulated with 5 µg/mL of collagen or 50 nM of thrombin were again used as controls to mimic “coated platelets.” (D) Clot retraction in platelet-rich plasma was induced by 1 U/mL of thrombin in the presence of phosphate-buffered saline (PBS, left panel), exMT (1:1 ratio to platelets, middle panel), or 5 µg/mL of collagen (right panel). The graph in the right panel summarizes the results from multiple experiments (n=27, one-way ANOVA, *P<0.05 vs. PBS-treated). FVa: activated factor V; PS: phosphatidylserine.




Figure 6. CD36 antibody reduced extracellular mitochondria-induced platelet activation. Non-injured mice were infused with 2×10^8 /mouse of MitoTracker-Green-labeled extracellular mitochondria (exMT), together with CD36 antibody or isotype IgG. (A-F) Blood samples were collected 30 min after infusion to measure levels of exMT-bound platelets (A), CD62p⁺ platelets (B), platelet-leukocyte complexes (C), platelet counts (D), and platelet-leukocyte-exMT tri-complexes (E) by flow cytometry (n=16, one-way analysis of variance). (F) Schematic summary of the study. An injured brain releases metabolically active exMT that interact with platelets. These exMT induce α-granule secretion by platelets, but do not induce platelet aggregation. As a result, the exMT-bound platelets and platelet-derived microvesicles promote the systemic consumptive coagulopathy and inflammation that are consistently found during the acute phase of traumatic brain injury. WBC: white blood cell; ROS: reactive oxygen species; VWF: von Willebrand factor; FVa: activated factor V; MV: microvesicles.

ed platelets", which are defined as a subpopulation (12-30%) of platelets that express procoagulant activity upon stimulation by collagen and thrombin.^{25,42} This procoagulant activity was also detected in exMT-treated platelets (Figure 5).

Third, the metabolic viability of exMT also suggests a potential role of ROS in the development of this intermediate platelet phenotype because ROS are known to activate platelets.¹⁵⁻¹⁷ We have shown that ROS from exMT activated platelets (Figure 3) and induced platelet microvesiculation (Figure 4) in an oxidant-dependent manner. A physical contact between exMT and platelets appears to be required for the ROS-induced platelet secretion because ROS primarily affect exMT-bound platelets. This direct exMT-platelet interaction may concentrate ROS activity on exMT-bound cells or protect the oxidants from plasma antioxidants.

In summary, we have shown that exMT-bound platelets develop an intermediate phenotype using ROS as mediators. This intermediate phenotype is characterized by α -granule secretion, procoagulant activity and poor aggregation (Figure 6F). As a result, these procoagulant platelets

remain in the circulation to promote non-focal coagulation, similar to the consumptive coagulopathy found in TBI patients^{43,44} and mouse models.⁹ While previous studies have found protective effects of exMT on target cells,^{10,12,13} our study demonstrates that these exMT could also have detrimental effects by making platelets procoagulant and prothrombotic. The findings from this study could have much broader implications regarding exMT in other pathologies in which hypercoagulable and inflammatory states develop, such as severe infections, autoimmune diseases, and cancer.

Funding

This study was supported by NIH grants NS087296 and HL71895 (JFD), National Natural Science Foundation of China State Key Program grant 81330029 (JNZ), National Natural Science Foundation of China Major International Joint Research Project 81720108015 (JNZ), National Natural Science Foundation of China Research grants 81271364, 81271359 (JNZ) and 81601068 (ZLZ), Natural Science Foundation of Tianjin 17JQCQJC12100 (ZLZ) and a regional scientific grant 16PTSTJC00180 (JNZ).

References

- Cap AP, Spinella PC. Severity of head injury is associated with increased risk of coagulopathy in combat casualties. *J Trauma*. 2011;71(1 Suppl):S78-81.
- Sun Y, Wang J, Wu X, et al. Validating the incidence of coagulopathy and disseminated intravascular coagulation in patients with traumatic brain injury--analysis of 242 cases. *Br J Neurosurg*. 2011;25(3):363-368.
- Talving P, Benfield R, Hadjizacharia P, Inaba K, Chan LS, Demetriades D. Coagulopathy in severe traumatic brain injury: a prospective study. *J Trauma*. 2009;66(1):55-63.
- Stein SC, Young GS, Talucci RC, Greenbaum BH, Ross SE. Delayed brain injury after head trauma: significance of coagulopathy. *Neurosurgery*. 1992;30(2):160-165.
- Kaufman HH, Moake JL, Olson JD, et al. Delayed and recurrent intracranial hematomas related to disseminated intravascular clotting and fibrinolysis in head injury. *Neurosurgery*. 1980;7(5):445-449.
- Wafaisade A, Lefering R, Tjardes T, et al. Acute coagulopathy in isolated blunt traumatic brain injury. *Neurocrit Care*. 2010;12(2):211-219.
- Stein SC, Chen XH, Sinson GP, Smith DH. Intravascular coagulation: a major secondary insult in nonfatal traumatic brain injury. *J Neurosurg*. 2002;97(6):1373-1377.
- Tian Y, Salsbery B, Wang M, et al. Brain-derived microparticles induce systemic coagulation in a murine model of traumatic brain injury. *Blood*. 2015;125(13):2151-2159.
- Zhao Z, Wang M, Tian Y, et al. Cardiolipin-mediated procoagulant activity of mitochondria contributes to traumatic brain injury-associated coagulopathy in mice. *Blood*. 2016;127(22):2763-2772.
- Hayakawa K, Esposito E, Wang X, et al. Transfer of mitochondria from astrocytes to neurons after stroke. *Nature*. 2016;535(7613):551-555.
- Davis CH, Kim KY, Bushong EA, et al. Transcellular degradation of axonal mitochondria. *Proc Natl Acad Sci U S A*. 2014;111(26):9633-9638.
- Islam MN, Das SR, Emin MT, et al. Mitochondrial transfer from bone-marrow-derived stromal cells to pulmonary alveoli protects against acute lung injury. *Nat Med*. 2012;18(5):759-765.
- Moschoi R, Imbert V, Nebout M, et al. Protective mitochondrial transfer from bone marrow stromal cells to acute myeloid leukemic cells during chemotherapy. *Blood*. 2016;128(2):253-264.
- Schon EA, DiMauro S, Hirano M. Human mitochondrial DNA: roles of inherited and somatic mutations. *Nat Rev Genet*. 2012;13(12):878-890.
- Begonja AJ, Gambaryan S, Geiger J, et al. Platelet NAD(P)H-oxidase-generated ROS production regulates alphaIIb beta3-integrin activation independent of the NO/cGMP pathway. *Blood*. 2005;106(8):2757-2760.
- Arthur JF, Qiao J, Shen Y, et al. ITAM receptor-mediated generation of reactive oxygen species in human platelets occurs via Syk-dependent and Syk-independent pathways. *J Thromb Haemost*. 2012;10(6):1133-1141.
- Bosin TR, Schaltenbrand SL. Stimulation of platelet serotonin transport by substituted 1,4-naphthoquinone-induced oxidant stress. *Biochem Pharmacol*. 1991;41(6-7):967-974.
- Lorente L, Martin MM, Gonzalez-Rivero AF, et al. Serum soluble CD40 ligand levels are associated with severity and mortality of brain trauma injury patients. *Thromb Res*. 2014;134(4):832-836.
- Castellino FJ, Chapman MP, Donahue DL, et al. Traumatic brain injury causes platelet adenosine diphosphate and arachidonic acid receptor inhibition independent of hemorrhagic shock in humans and rats. *J Trauma Acute Care Surg*. 2014;76(5):1169-1176.
- Sillescu M, Johansson PI, Rasmussen LS, et al. Platelet activation and dysfunction in a large-animal model of traumatic brain injury and hemorrhage. *J Trauma Acute Care Surg*. 2013;74(5):1252-1259.
- Dietrich WD, Alonso O, Busto R, et al. Posttraumatic cerebral ischemia after fluid percussion brain injury: an autoradiographic and histopathological study in rats. *Neurosurgery*. 1998;43(3):585-593.
- Lu D, Mahmood A, Goussev A, Qu C, Zhang ZG, Chopp M. Delayed thrombosis after traumatic brain injury in rats. *J Neurotrauma*. 2004;21(12):1756-1766.
- Zhang JN, Bergeron AL, Yu Q, et al. Duration of exposure to high fluid shear stress is critical in shear-induced platelet activation-aggregation. *Thromb Haemost*. 2003;90(4):672-678.
- Yuan H, Houck KL, Tian Y, et al. Piperlongumine blocks JAK2-STAT3 to inhibit collagen-induced platelet reactivity independent of reactive oxygen species. *PLoS One*. 2015;10(12):e0143964.
- Alberio L, Safa O, Clemetson KJ, Esmon CT, Dale GL. Surface expression and functional characterization of alpha-granule factor V in human platelets: effects of ionophore A23187, thrombin, collagen, and convulxin. *Blood*. 2000;95(5):1694-1702.
- Tracy PB, Mann KG. Prothrombinase complex assembly on the platelet surface is mediated through the 74,000-dalton component of factor Va. *Proc Natl Acad Sci U S A*. 1983;80(8):2380-2384.
- Hierso R, Waltz X, Mora P, et al. Effects of oxidative stress on red blood cell rheology in sickle cell patients. *Br J Haematol*. 2014;166(4):601-606.
- Zhou Y, Cai W, Zhao Z, et al. Lactadherin promotes microvesicle clearance to prevent coagulopathy and improves survival of severe TBI mice. *Blood*. 2018;131(5):563-572.
- Ghosh A, Li W, Febbraio M, et al. Platelet CD36 mediates interactions with endothelial cell-derived microparticles and contributes to thrombosis in mice. *J Clin Invest*. 2008;118(5):1934-1943.

30. Pascual G, Avgustinova A, Mejetta S, et al. Targeting metastasis-initiating cells through the fatty acid receptor CD36. *Nature*. 2017;541(7635):41-45.
31. Saliba AE, Vonkova I, Gavin AC. The systematic analysis of protein-lipid interactions comes of age. *Nat Rev Mol Cell Biol*. 2015;16(12):753-761.
32. Samuni Y, Goldstein S, Dean OM, Berk M. The chemistry and biological activities of N-acetylcysteine. *Biochim Biophys Acta*. 2013;1830(8):4117-4129.
33. Dale GL, Friese P, Batar P, et al. Stimulated platelets use serotonin to enhance their retention of procoagulant proteins on the cell surface. *Nature*. 2002;415(6868):175-179.
34. Prodan CI, Vincent AS, Dale GL. Coated-platelet levels increase with number of injuries in patients with mild traumatic brain injury. *J Neurotrauma*. 2016;33(9):818-824.
35. Carr ME, Jr., Carr SL. Fibrin structure and concentration alter clot elastic modulus but do not alter platelet mediated force development. *Blood Coagul Fibrinolysis*. 1995;6(1):79-86.
36. Carr ME, Jr., Carr SL, Hantgan RR, Braaten J. Glycoprotein IIb/IIIa blockade inhibits platelet-mediated force development and reduces gel elastic modulus. *Thromb Haemost*. 1995;73(3):499-505.
37. Fadok VA, Bratton DL, Frasch SC, Warner ML, Henson PM. The role of phosphatidylserine in recognition of apoptotic cells by phagocytes. *Cell Death Differ*. 1998;5(7):551-562.
38. Rigotti A, Acton SL, Krieger M. The class B scavenger receptors SR-BI and CD36 are receptors for anionic phospholipids. *J Biol Chem*. 1995;270(27):16221-16224.
39. Podrez EA, Byzova TV, Febbraio M, et al. Platelet CD36 links hyperlipidemia, oxidant stress and a prothrombotic phenotype. *Nat Med*. 2007;13(9):1086-1095.
40. Tait JF, Smith C. Phosphatidylserine receptors: role of CD36 in binding of anionic phospholipid vesicles to monocytic cells. *J Biol Chem*. 1999;274(5):3048-3054.
41. Newman PJ, McEver RP, Doers MP, Kunicki TJ. Synergistic action of two murine monoclonal antibodies that inhibit ADP-induced platelet aggregation without blocking fibrinogen binding. *Blood*. 1987;69(2):668-676.
42. Hamilton SF, Miller MW, Thompson CA, Dale GL. Glycoprotein IIb/IIIa inhibitors increase COAT-platelet production in vitro. *J Lab Clin Med*. 2004;143(5):320-326.
43. Bulstrode H, Nicoll JA, Hudson G, Chinnery PF, Di Pietro V, Belli A. Mitochondrial DNA and traumatic brain injury. *Ann Neurol*. 2014;75(2):186-195.
44. Walko TD 3rd, Bola RA, Hong JD, et al. Cerebrospinal fluid mitochondrial DNA: a novel DAMP in pediatric traumatic brain injury. *Shock*. 2014;41(6):499-503.