

ORIGINAL ARTICLE

Positive effects of methylphenidate on hyperactivity are moderated by monoaminergic gene variants in children with autism spectrum disorders

JT McCracken¹, KK Badashova¹, DJ Posey², MG Aman³, L Scahill⁴, E Tierney⁵, LE Arnold³, B Vitiello⁶, F Whelan¹, SZ Chuang⁷, M Davies⁷, B Shah¹, CJ McDougle⁸ and EL Nurmi¹

Methylphenidate (MPH) reduces hyperactive-impulsive symptoms common in children with autism spectrum disorders (ASDs), however, response and tolerability varies widely. We hypothesized monoaminergic gene variants may moderate MPH effects in ASD, as in typically developing children with attention-deficit/hyperactivity disorder. Genotype data were available for 64 children with ASD and hyperactivity who were exposed to MPH during a 1-week safety/tolerability lead-in phase and 58 who went on to be randomized to placebo and three doses of MPH during a 4-week blinded, crossover study. Outcome measures included the Clinical Global Impression-Improvement (CGI-I) scale and the Aberrant Behavior Checklist (ABC-hyperactivity index). A total of 14 subjects discontinued the study because of MPH side effects. Subjects were genotyped for variants in *DRD1–DRD5*, *ADRA2A*, *SLC6A3*, *SLC6A4*, *MAOA* and *MAOB*, and *COMT*. Forty-nine percent of the sample met positive responder criteria. In this modest but relatively homogeneous sample, significant differences by *DRD1* ($P = 0.006$), *ADRA2A* ($P < 0.02$), *COMT* ($P < 0.04$), *DRD3* ($P < 0.05$), *DRD4* ($P < 0.05$), *SLC6A3* ($P < 0.05$) and *SLC6A4* ($P < 0.05$) genotypes were found for responders versus non-responders. Variants in *DRD2* ($P < 0.001$) and *DRD3* ($P < 0.04$) were associated with tolerability in the 14 subjects who discontinued the trial. For this first MPH pharmacogenetic study in children with ASD, multiple monoaminergic gene variants may help explain individual differences in MPH's efficacy and tolerability.

The Pharmacogenomics Journal (2014) **14**, 295–302; doi:10.1038/tpj.2013.23; published online 16 July 2013

Keywords: autism spectrum disorders; dopamine; genetics; hyperactivity; methylphenidate

INTRODUCTION

The autism spectrum disorders (ASDs) are a group of neurodevelopmental disorders including autistic disorder, Asperger's disorder and pervasive developmental disorder, not otherwise specified, which share impairment in social communication. Surveys estimate 1 out of 88 to 160 children as having an ASD.¹ Impairment from ASD varies considerably across its core dimensions, but also from associated maladaptive behaviors, such as symptoms resembling attention-deficit/hyperactivity disorder (ADHD).^{2–4} Moderate-to-severe hyperactive-impulsive behaviors overlap in one-third or more individuals with ASD. These behaviors are often targeted by pharmacologic and other treatment efforts,^{5,6} such as stimulants.

Despite common use of stimulants for ASD,^{7,8} variable effects and reduced tolerability was suggested from small early studies,^{9–12} but no moderators of response have been reported. In the largest stimulant trial in ASD, the Research Units on Pediatric Psychopharmacology (RUPP) Autism Network (2005) found methylphenidate (MPH) was superior to placebo in reducing ratings of hyperactivity,^{13,14} but only half (35/72; 49%) were considered clinical responders. In addition, 19% of subjects were withdrawn because of adverse effects; variability in optimal dose, adverse events and response was striking.¹⁵

Variability in response to stimulant treatment is well known in typically developing children,¹⁶ and there is interest in individual genetic variation moderating response in the treatment of ADHD.^{17,18} However, no reports examining the pharmacogenetics of stimulant response exist for children with ASD. In addition, although a small number of reports have described genetic associations with adverse events and stimulants, none in typical ADHD or subjects with ASD have examined possible genetic associations in those with 'intolerable' side effects leading to dropout.

Investigations in typically developing children have suggested that variants in catecholamine-related genes may be associated with MPH response in ADHD.¹⁷ The majority of these examined the variable number tandem repeat (VNTR) polymorphism located in the 3' untranslated region of the dopamine transporter gene (*SLC6A3*),^{17,19,20} the site of MPH action. A meta-analysis has suggested that the nine-repeat variant may confer reduced efficacy;²¹ however, recent reports showed enhanced response of 9/9 homozygotes with a dose-dependent effect.²² Investigations of stimulant response and the 48-base pair (bp) VNTR polymorphism in exon 3 of the dopamine D4 receptor (*DRD4*) gene¹⁷ have been mixed, with greater efficacy seen in subjects with one or two copies of the four-repeat VNTR genotype

¹Department of Psychiatry and Biobehavioral Sciences, University of California at Los Angeles, Semel Institute for Neuroscience, Los Angeles, CA, USA; ²Department of Psychiatry, Indiana University School of Medicine, Indianapolis, IN, USA; ³Departments of Psychiatry and Psychology, The Nisonger Center UCEDD, Ohio State University, Columbus, OH, USA; ⁴Child Study Center, Yale University, New Haven, CT, USA; ⁵Department of Psychiatry, Kennedy Krieger Institute, Johns Hopkins University, Baltimore, MD, USA; ⁶National Institute of Mental Health, National Institutes of Health, Bethesda, MD, USA; ⁷Department of Psychiatry, Columbia University, New York, NY, USA and ⁸Department of Psychiatry, Lurie Autism Center, Massachusetts General Hospital, Boston, MA, USA. Correspondence: Dr JT McCracken, Department of Psychiatry and Biobehavioral Sciences, UCLA NPI-Semel Institute, 760 Westwood Plaza, Los Angeles, CA 90024-1759, USA.

E-mail: jmcrcracken@mednet.ucla.edu

Received 9 December 2012; revised 21 April 2013; accepted 26 April 2013; published online 16 July 2013

(*DRD4A*),^{23–25} while others found opposite²⁶ or no effects.^{27–29} Variants such as *DAT1 9/10*, *DRD4.7* and the α -adrenergic receptor 2A (*ADRA2A*) influence the slope of dose-response relations.^{19,22} Few data are available regarding the impact of variation in other dopamine (DA)- receptors *DRD1*, *DRD2*, *DRD3* or *DRD5*. Although data are sparse and conflicting, some reports have associated MPH side effects with variants in *DAT1* and *DRD4* as well.^{19,25,30}

Awareness of pharmacogenetic interactions of other neurotransmitter genes with psychostimulant action has grown. Variants in *ADRA2A* influence response of inattentive symptoms to MPH.^{22,31,32} Interactions between DA and serotonin (5HT) have also been suggested as integral to MPH's behavioral and locomotor effects.³³ Two studies presented significant moderating effects of the 44-bp insertion–deletion polymorphism in the promoter region of the serotonin transporter gene (*SLC6A4*),^{19,34} although one report did not detect an effect.²⁹ Genetic variants in *COMT* also influence basal transmitter availability,³⁵ working memory³⁶ and cognitive response to acute stimulant administration.³⁷ In ADHD, three reports concur that the Val158Met *COMT* variant moderates MPH effects.^{19,38,39}

Variability in existing ADHD pharmacogenetic studies reflect the heterogeneity in methodology and outcomes. MPH dosing differs, there are few randomized, blinded, placebo-controlled studies, and only four assessed outcomes at multiple doses.^{19,20,25,31}

The purpose of this preliminary investigation was to thoroughly evaluate for the first time in subjects with ASD whether gene variants in candidate loci would be found to associate with the overall clinical response to MPH. Candidates were suggested as moderators by at least two prior pharmacogenetic studies of MPH (*DRD4*, *SLC6A3*, *SLC6A4*, *COMT* and *ADRA2A*), risk genes for ADHD, known targets of MPH action (*DRD1*, *DRD2* and *DRD5*)⁴⁰ or associated with psychomotor functioning and impulsivity (*DRD2* and *DRD3*).^{41,42} We hypothesized that gene variants suggested to exert significant effects on MPH response in typically developing children would also influence effects on response and tolerability in a homogeneous sample of children with ASD. However, we noted the multiple findings suggesting dysregulated DA function in ASD,^{43–45} which could alter the relations between gene variants and MPH response and tolerability.

MATERIALS AND METHODS

Subjects

The details of the recruitment and baseline characteristics of the sample have been described previously.⁴⁶ Briefly, subjects were recruited at the five centers that formed the RUPP Autism Network funded by the National Institute of Mental Health (NIMH). The study was approved by each local institutional review board and monitored by a central NIMH data and safety monitoring board. Written informed consent was obtained from a parent or guardian, and assent was given by the child when capable.

To be eligible for the study, subjects from 5 to 14 years were required to meet Diagnostic and Statistical Manual, version-IV criteria for autistic disorder, Asperger's disorder, or PDD-NOS using clinical examination and the Autism Diagnostic Interview-Revised,⁴⁷ with diagnoses following Diagnostic and Statistical Manual, version-IV criteria. All subjects had significant symptoms of ADHD (based on the Clinical Global Impression (CGI)-Severity and Swanson, Nolan, and Pelham-IV (SNAP-IV)), were medically healthy, and were not taking any concomitant psychotropic drugs during the treatment phase of the study.

Study design

Details of the study's design and rationale were published previously.⁴⁶ Briefly, it included a 1-week test dose phase of 1-day placebo, and 2 days each of the low, medium and high MPH doses used in the next phase to determine initial MPH tolerability, followed by a 4-week random-order, placebo-controlled, double-blind crossover phase to assess efficacy, and an 8-week open-label continuation phase for subjects showing a positive response to assess the maintenance of beneficial effects.

Subjects who were unable to tolerate MPH during the test dose phase as evidenced by a CGI rating of 'much worse' or 'very much worse' at a low or

medium dose of MPH were excluded before being randomized. Subjects who tolerated all three dose levels of MPH during the test dose phase entered into the 4-week double-blind crossover phase. Each subject received placebo and three different doses of MPH in random order, 1-week each.

Low, medium and high doses of MPH were assigned based on body weight to approximate 0.125, 0.25 and 0.5 mg kg⁻¹ per dose, administered morning and noon, with an additional half-dose given at 1600 hours. Subjects with moderate or greater side effects to the highest dose level of MPH in the test dose phase received the medium dose during 2 weeks of the double-blind, crossover phase. Clinicians, the patient and the caregiver were blind to treatment assignment during the crossover phase.

Positive response was defined as a rating of 'much improved' or 'very much improved' on the CGI completed by the masked study clinician and a significant reduction on the Aberrant Behavior Checklist (ABC)-hyperactivity subscale (defined as a 25% reduction by both parent and teacher or a 30% reduction by parent or teacher) at any given weekly assessment during the blinded phase, and completion of all weeks of the blinded phase. For the purpose of these responder analyses, subjects who dropped out at any point after entering the double-blind phase ($n = 8$) are grouped with non-responders. For the purpose of tolerability analyses, any subject who dropped out in either the test dose or double-blind phase secondary to adverse events formed the intolerant group ($n = 14$).

Genotyping

Ten candidate genes were selected for typing. A total of 36 variants were genotyped across the 10 loci. Common (>10%) locus variability was captured across the dopamine receptor genes *DRD1–DRD5*, *ADRA2A*, *MAOA* and *MAOB*, and *SLC6A4*. Several known functional variants were also genotyped, including the Val158Met *COMT* non-synonymous single-nucleotide polymorphism (SNP),¹⁹ and VNTRs in *SLC6A3* (480-bp VNTR in the 3' untranslated region),¹⁷ *DRD4* (exon 3 48-bp VNTR)⁴⁸ and *SLC6A4* (44-bp insertion–deletion in the promoter region (5-HTTLPR) with modifying SNP (rs25531) and intron 2 VNTR (STin2)).⁴⁹ To minimize multiple testing of non-independent genotypes, allele frequencies were computed and genotype classifications defined *a priori* to any outcome analyses.

The *SLC6A3* VNTR (*DAT1*) was genotyped using published methods and primers.⁵⁰ The *DRD4*-exon 3 VNTR was genotyped with the following modified primers: F—5'-CTACCTGCCCGCTCATG-3'; R—5'-CCGGTGATCTTG GCACGC-3' using previously described methods.⁵¹ The *SLC6A4* promoter region gene-linked polymorphic region (5-HTTLPR) short-long variant and intron 2 10/12 VNTR (STin2) were genotyped according to published protocols.⁵² For VNTR variants (5-HTTLPR, 5-HTT STin2, *DAT* and *DRD4*), PCR products were electrophoresed in 2% gold agarose (BMA, Rockland, ME, USA) gels in 1x TBE and imaged with ethidium bromide under fluorescent Kodak digital camera (Rochester, NY, USA). Alleles were determined by comparison with molecular weight standards and control individuals with previously determined genotypes. Samples were double-scored by two technicians.

Tag SNPs were selected using Haploview Software⁵³ publicly available from the HapMap Consortium to capture variability present at $\geq 10\%$ with an $r^2 \geq 0.8$. SNPs were genotyped using the Life Technologies (Life Technologies, Grand Island, NY, USA) TaqMan platform with Qiagen Type-it Fast SNP Probe PCR Kit (Qiagen, Valencia, CA, USA) according to the manufacturer's protocols. All genotypes were in Hardy–Weinberg Equilibrium ($P > 0.05$). Ten percent of the data set was genotyped in duplicate without any conflicting genotypes, and allele frequencies were checked for consistency with those reported by the HapMap Consortium.

Potential functional roles of associated non-coding variants were explored using the online UCSC Genome Browser⁵⁴ with ENCODE tracks⁵⁵ and the Broad Institute's Haploreg online resource.⁵⁶

Data analysis

All subjects randomized to the double-blind crossover phase were included in the responder analyses. Descriptive statistics are reported as the mean \pm s.d. Before examining allelic effects on response, a general linear model was constructed using the ABC-hyperactivity for each gene variant examined to test for the presence of a positive dose effect (for example, if a treatment effect existed). Dose (low, medium, high or placebo), genotype and the interaction dose by genotype were then entered as predictor variables of parent-rated ABC-hyperactivity scores. The primary analyses were response by genotype compared using χ^2 analyses or Fisher's exact tests. As the *DRD3* gly allele is the minor variant and some prior reports have shown biological differences between ser/ser and ser/gly versus gly/gly genotype groups, we restricted our comparison

accordingly.⁵⁷ As some investigators have noted significant moderation of stimulant effects of *DRD4* variants in typically developing children specifying genotypes by the presence of 0, 1 or 2 copies of the '7' allele,¹⁹ we also performed a similar three-way grouped analysis for this variant. Based on the fact that the previous literature on the *SLC6A3* VNTR has shown significant effects for 9/9 homozygotes, we analyzed our data according to this model.²² Statistical significance was set as $P < 0.05$, and $P < 0.002$ with correction for multiple testing.

RESULTS

Sixty-six children with autistic disorder ($n = 47$), Asperger's disorder ($n = 5$) or pervasive developmental disorder, not otherwise specified ($n = 14$) completed the test dose phase and entered the double-blind crossover phase. Six subjects discontinued during the test dose phase. Of the 72 subjects who entered the trial, 7 subjects were unable or refused to provide samples for DNA and data were missing for one subject, leaving 64 subjects for genotyping. Sixteen subjects received an additional week of medium dose during the crossover phase because of moderate or

greater side effects to the high dose in the test dose phase. Fifty-eight subjects were able to complete all 4 weeks of the crossover with the remainder leaving either because of adverse events ($n = 7$) or for other reasons ($n = 1$). The mean age of the subjects in this report was 6.90 years (± 2.2 , range 5.0–13.0); 75.9% were Caucasian, 87.9% were male; the mean intelligence quotient was 65.0 (± 33.3 range = 16–135). Neither age, ethnicity nor diagnosis was a significant covariate.

Table 1 shows parent and teacher rated SNAP-ADHD, SNAP-ODD and ABC-hyperactivity at baseline before entry into the double-blind phase. In general, subjects in the study had moderately severe symptoms of ADHD, and the optimal dose week demonstrated a moderate benefit associated with significantly reduced ABC-hyperactivity ratings and lower CGI-I scores. Analyses of ABC-hyperactivity scores across all four dose assessments showed a main effect for dose for all genotypes (all $P < 0.001$). Variability in individual dose-response trajectories and optimal dose was substantial (Figure 1).

After grouping subjects by *a priori* defined responder or non-responder categories, we analyzed the association of genotypes with responder status. Table 2 reports the results of χ^2 tests for genotype by response. Nominally significant responder by genotype effects (Figure 2) were noted for the dopamine receptors *DRD1* rs5326 ($P = 0.006$, Figure 2a) and rs4867798 ($P < 0.05$, Figure 2b), *DRD3* ser9gly rs6280 ($P < 0.05$, Figure 2c) and *DRD4* rs11246226 ($P < 0.04$, Figure 2d); the adrenergic receptor *ADRA2A* rs1800544 ($P < 0.02$, Figure 2e); the dopamine and serotonin transporters *SLC6A3* VNTR ($P < 0.05$, Figure 2f) and *SLC6A4* STin2 VNTR ($P < 0.05$, Figure 2g), respectively; and the catabolic enzyme *COMT* val/met rs4680 ($P < 0.05$, Figure 2h). No significant effects on response were evident for the *DRD2*, *DRD5* or *MAOA/B* variants tested.

In an effort to further reveal possible interactions between dose condition and genotype, in secondary analyses we tested the relationship between ABC-hyperactivity scores across the four treatment conditions for main effects of dose, genotype and dose by genotype interaction using analysis of covariance with baseline ABC-hyperactivity scores as the covariate. Results (Table 2) showed significant dose by genotype interactions for promoter SNPs in *DRD4* rs11246226 ($P < 0.02$) and *SLC6A4* rs12150214 ($P < 0.03$). In contrast to previous findings, *SLC6A3* VNTR analysis did not yield dose by genotype effects,²² however, our power to detect these effects was very low given the small number of subjects with the 9/9 genotype ($n = 3$), all of whom were responders.

The 14 subjects who dropped because of intolerance were similar to those completing the study (83% male, 83% Caucasian, mean age 8.6 years). Genetic comparison of those 14 subjects to trial completers revealed a protective effect for carriers of the minor allele at *DRD2* synonymous variant rs6275. Intolerance increased from 4% in minor allele carriers to 23% in those homozygous for the common allele ($P < 0.001$), a result surviving correction for multiple testing. The *DRD3* variant rs6280 (Ser9Gly) was nominally associated with tolerability, as intolerance increased from 3% in common allele homozygotes to 18% in carriers of the minor allele ($P = 0.031$), suggesting that the minor allele is a risk factor for stimulant intolerance.

DISCUSSION

In this first pharmacogenetic study of MPH in a modest but homogeneous (90% male, 90% stimulant naive) sample of children with ASD, improvement in hyperactive-impulsive symptoms associated with MPH administration was moderated significantly by several gene variants comprising elements of monoaminergic systems for dopamine, norepinephrine and serotonin, all of which have known regulatory effects on motor activity, and higher order cognitive functions.^{58–60} Of the 10 candidate genes queried based on prior research, our data supported a contribution of genetic variation in 7 genes (*SLC6A4*,

Table 1. ABC-hyperactivity ratings and SNAP ratings in genotyped children with ASD ($n = 58$) at entry into the double-blind phase

Outcome measure	N	Mean (\pm s.d.)	Item mean (\pm s.d.)
Parent SNAP-IV ADHD (18 items)	58	39.1 \pm 7.9	2.26 \pm 0.43
Parent SNAP-IV ODD (8 items)	58	9.67 \pm 6.17	1.21 \pm 0.78
Parent ABC-hyperactivity rating at baseline (16 items)	58	33.6 \pm 7.43	2.09 \pm 0.47
Teacher SNAP-IV ADHD (18 items)	54	35.74 \pm 8.23	2.05 \pm 0.46
Teacher SNAP-IV ODD (8 items)	54	8.48 \pm 4.64	1.07 \pm 0.59
Teacher ABC-hyperactivity rating (16 items)	54	31.46 \pm 8.68	1.94 \pm 0.55

Abbreviations: ABC, Aberrant Behavior Checklist; ADHD, attention-deficit/hyperactivity disorder; ASD, autism spectrum disorder; ODD, oppositional defiant disorder; SNAP-IV, Swanson, Nolan, and Pelham-IV. Varying sample size (N) because of missing teacher data on six subjects.




Figure 1. Individual parent-rated Aberrant Behavior Checklist (ABC)-hyperactivity scores during double-blind crossover trial. Top panel includes non-responders; bottom panel responders (see text for definition). Individual weekly ratings obtained at baseline and during random-order crossover weeks of methylphenidate treatment.

Table 2. MPH responder/non-responder number by genotypes from double-blind crossover trial

Gene	Variant	Alleles	Responder % (n)	Non-responder % (n)	Genotype × responder status P-value	Dose × genotype P-value
DRD1	rs4867798	C + TT	67% (22/33) 40% (10/25)	33% (11/33) 60% (15/25)	0.042	0.577
	rs5326	T + CC	82% (14/17) 44% (18/41)	18% (3/17) 56% (23/41)	0.006	0.215
DRD2	rs686				0.796	
	rs6277					
	rs6589377				0.714	
	rs4938019				0.292	
	rs7131056				0.505	
	rs1800498				0.714	
	rs2283265				0.566	
DRD3	rs6280	Ser/Ser	74% (14/19)	26% (5/19)	0.044	0.569
	(Ser9Gly)	Gly +	46% (18/39)	54% (21/39)		
	rs2134655				0.696	
	rs9880168				0.259	
	rs7633291				0.404	
	rs167771				0.795	
	rs3732790				0.397	
DRD4	rs11246226	AA C +	79% (11/14) 48% (21/44)	21% (3/14) 52% (23/44)	0.038	0.013
	rs3758653				0.271	
DRD5	Exon 3 VNTR				0.810	
	rs10033951				0.930	
SLC6A3	3'UTR VNTR	10 + 9/9	51% (27/53) 100% (3/3)	49% (26/53) 0% (0/3)	0.049	0.316
SLC6A4	rs12150214				0.078	0.026
	rs4251417				0.577	
	rs11080121				0.642	
	5HTT-LPR				0.881	
ADRA2A	STin2 VNTR	10 + 12/12	67% (22/33) 39% (9/23)	33% (11/33) 61% (14/23)	0.041	0.764
	rs1800544	CC G +	71% (20/28) 40% (12/30)	29% (8/28) 60% (18/30)	0.015	0.790
	rs12246561				0.319	
MAOA/B	rs3750625				0.546	
	rs1465108				0.501	
	rs3810709				0.416	
	rs3027399				0.073	
	rs10521432				0.308	
COMT	rs1799836				0.191	
	rs4680 (Val158Met)	Met + Val/Val	64% (25/38) 36% (7/19)	36% (13/38) 63% (12/19)	0.049	0.576

Abbreviation: MPH, methylphenidate.
Shaded values indicate significant *P*-values.

SLC6A3, *DRD1*, *DRD3*, *DRD4*, *COMT* and *ADRA2A*) to clinical MPH response. Although our sample size did not provide adequate power for multiple comparison correction, the number of positive associations observed is unlikely to occur by chance. An exact binomial calculation for the probability of 7 or more significant hits out of 36 variants, whereas not an exact *P*-value, suggests that many of these associations are in fact true positives ($P < 0.002$). These genetic effects may reveal some of the sources of marked variability in clinical outcomes observed in this trial and in clinical experience for this common treatment in ASD; in this trial only half of all subjects responded to moderate-dose MPH.

Our results add support to the previous suggestions of contributions of variants in several genes influencing MPH response in ADHD (*SLC6A4*, *SLC6A3*, *DRD4*, *COMT* and *ADRA2A*) but in some cases suggest new loci not previously implicated as MPH-response modifiers (*SLC6A4* STin2 VNTR and *DRD4* rs11246226). In addition, we also identified new associations with other genes not extensively examined thus far in the ADHD literature (*DRD1* rs5326 and rs4867798, *DRD3* ser9gly rs6280),

but rational candidates given their distribution and known cognitive and behavioral pharmacologic effects of agonists and antagonists at these targets. Our findings should encourage other investigations of these variants in larger samples.

Not surprisingly, DA transporter and DA receptor variants showed the strongest associations with treatment response. In contrast to earlier studies,^{19,20,61} our data agree with the recent report by Froehlich *et al.*,²² showing an enhanced MPH response in *SLC6A3* 9/9 homozygotes. *DRD1*, highly expressed in cortex and striatum, is important in cognitive and motor control,⁶² and may be a minor ADHD risk gene.^{63,64} *DRD1* was most strongly associated with clinical response with two SNPs (rs5326 and rs4867798; $P = 0.006$ and $P = 0.043$, respectively) in our sample. *In silico* exploration suggests that both *DRD1* SNPs (rs5326 and rs4867798) may be functional, as they map to areas of open chromatin in the 5' and 3' untranslated regions and appear to alter binding sites for the transcriptional repressor ZNF263 and the transcriptional activator POU2F1 respectively. The non-synonymous Ser9Gly variant in *DRD3* was also associated with




Figure 2. Genotype associations with methylphenidate clinical response rates. Positive response defined as Clinical Global Impression-Improvement (CGI-I) rating by blinded study clinician of 'much improved' or 'very much improved' and decrease in parent-rated Aberrant Behavior Checklist-Hyperactivity subscale of > 25% from baseline for any active drug week. $n = 58$ subjects. P -values reflect results of χ^2 tests. Panels a–h are plots of genotype group and their respective response rates for: 2a) *DRD1* rs5326; 2b) *DRD1* rs4867798; 2c) *DRD3* rs6280; 2d) *DRD4* rs11246226; 2e) *ADRA2A* rs1800544; 2f) *SLC6A3* VNTR; 2g) *SLC6A4* STin2; 2h) *COMT* rs4680.

MPH response. Ser/ser homozygotes demonstrated greater response than glycine allele carriers. Finally, we observed an association of a promoter SNP (rs11246226) in *DRD4* with both clinical response as well as a genotype by dose interaction. *DRD4* is well known as a minor risk gene for ADHD⁶⁵ and the exon 3 VNTR has been extensively studied for associations with cognitive phenotypes and drug response, but studies have not tested effects of other variants at this locus. It is difficult to ascertain the likely sources of differences between our data and some previous studies for effects of the *DRD4* exon 3 VNTR variant in particular, however, we note the literature is quite inconsistent with respect to its associations with response, even in reports from the same group.^{19,25} Consistent with other reports, *DRD2* was not found to associate with clinical response to MPH, surprising given the multiple functional variants and stimulant-induced displacement of *DRD2* binding following acute MPH.⁴⁰ Taken together, our findings of associations between *DRD1*, *DRD3* and *DRD4* and response measures suggest that common variants in the DA pathway do exert influences on the effects of MPH and moderate its DA-mediated impact on behavior.

Despite our modest sample size, the multiple (admittedly nominal) associations with several dopamine receptor subtypes (*DRD1*, *DRD3* and *DRD4*) lead us to speculate that the effects of gene variants in these targets may be amplified by ASD-related dopamine dysregulation. We posit that ASD-related differences in DA function may influence the interaction between gene variants and MPH, as evidence supports increased DA turnover (increased frontal uptake of ¹⁸F fluoro-3,4-dihydroxy-5-fluorophenylalanine and increased cerebrospinal fluid homovanillic acid) in ASD as well as increased dopamine transporter binding.^{44,45,66} Such ASD-related differences may shift dose-response, enhancing or distorting pharmacogenetic relationships seen in typically developing children. Indeed, the high rate of MPH intolerance, several-fold greater than reports from typically developing children, suggests greater sensitivity to MPH pro-dopaminergic effects in ASD. These data add modest support to the notion that although common gene variants influence psychostimulant response in both typically and atypically developing subjects with hyperactivity, there are some possible differences because of underlying disorder neurobiology.

Our findings also reinforce the emerging view that beneficial effects of stimulants are multifactorial in their mechanism,

reflecting drug effects on serotonergic and noradrenergic functions in addition to dopaminergic actions.⁵⁹ However, it is possible that identified non-DA gene effects may be also mediated via downstream effects on dopaminergic neurotransmission. For example, *SLC6A4* alleles associated with high expression (LPR I and particularly STin2 12 alleles), have been suggested to moderate mood effects of amphetamine in healthy subjects⁵² and in typically developing children with ADHD, and LPR III homozygous subjects showed less improvement in global functioning with MPH.³⁴ However, opposing effects of the two variants were seen on different outcome variables.¹⁹ Mice lacking *SLC6A3* still demonstrate reductions in activity from MPH, apparently mediated via 5HT.³³ The lower response rates in our 10/12 and 12/12 genotype subjects is in line with the known effects of the 12 allele on higher transporter expression. We also noted a tagging SNP in the first intron of *SLA6A4*, just upstream of the translational start, was associated with differential response at high dose MPH. This SNP, while intronic, is in a linkage disequilibrium block showing multiple enhancer histone marks suggesting an area of active transcriptional regulation.

Similarly, these data also contribute to the growing support for the involvement of the noradrenergic system in stimulant response in ADHD, consistent with direct effects of stimulant-induced NE release⁵⁸ and the reciprocal relationship of noradrenergic–dopaminergic pathways.⁶⁷ Our data did identify a moderating effect of the *ADRA2A* promoter SNP rs1800544 and clinical response, extending prior observations.^{31,32} Online database mining supports a functional effect of the associated variant, as this SNP maps to a site of open chromatin in the *ADRA2A* promoter with histone marks and binding sites for the transcription factors RAD21 and CTCF, which have a role in promoter methylation, and ultimately expression.

Relevant to dopaminergic and noradrenergic transmission, we found support for earlier observations^{19,38,39} that variants in *COMT* moderated clinical benefit from MPH. However, contrary to studies in healthy subjects³⁷ and typical ADHD, in our data set *met*-carriers showed a greater reduction in hyperactivity with MPH treatment than *val/val* homozygotes. Indeed, *met/met* subjects had a 75% (9/12) response rate versus 37% (7/19) for *val/val* homozygotes, a twofold differential. If ASD is associated with reduced tonic dopamine (consequences of increased DAT and increased presynaptic uptake), the resulting hypothesized shift in dose-response

could favor *met/met* individuals, as *val/val* subjects may lack sufficient dopamine availability to overcome reduced DA tone despite MPH releasing effects.

Our report is novel in our analyses of possible genetic underpinnings of stimulant intolerance, not heretofore researched in the ADHD pharmacogenetic literature. A *DRD2* variant, rs6275, not associated with response, was significantly associated with MPH intolerance and remained significant after correction. Homozygotes for the common allele at rs6275 showed a sixfold greater rate of intolerance. The non-synonymous Ser9Gly (rs6280) variant in *DRD3* was associated with MPH response and tolerability, as gly allele carriers were highly likely to either be intolerant of MPH and/or be less responsive than ser/ser homozygotes. These variants merit additional investigation in relationship to a range of stimulant-induced adverse effects in much larger samples.

Strengths of our findings include the thorough genetic analyses, use of a rigorous study design including comparisons of placebo versus multiple fixed-dose treatment conditions, blinded evaluations by multiple informants (parents, teachers and study clinicians), application of a careful, multi-informant based and clinically relevant definition of responder status, and recruitment of a largely male, stimulant-naïve, homogenous sample. Relative to the only other small ($n \leq 13$), controlled studies of stimulants in ASD, our sample is more homogenous with respect to sex, ethnicity, treatment history and diagnosis.^{11,12} Capitalizing on the strengths of our design, we were able to examine genetic effects observed under optimal treatment conditions, as determined by blinded assessments and multiple raters.

Limitations of this work include the short duration of MPH treatment observation periods, the modest sample size and the restrictions of genotyping to selected candidate genes. Our observations could be associated with an increased risk of type I statistical error.

In summary, the reduction of hyperactivity by MPH in individuals with ASD may be moderated at the level of individual genetic variability, particularly at loci that influence monoaminergic signaling. ADHD symptoms are common in ASD, and MPH remains the best empirically validated treatment for this target, although variation in response and tolerability is large. Additional studies replicating our observations and identifying other potential individual predictors of response to MPH in individuals with ASD are strongly needed. Improving overall outcomes from long-term intervention efforts for ASD remains a major clinical and public health challenge.

CONFLICT OF INTEREST

Dr Aman has received consulting fees from Bristol-Myers Squibb, BioMarin, Roche and Supernus. Dr Aman also reports research support from Bristol-Myers Squibb and Johnson and Johnson. Dr Arnold has received research funding from Curemark, Shire and Lilly, and has consulted on advisory boards for AstraZeneca, Biomarin, Novartis, Noven, Seaside Therapeutics and Shire. Dr McCracken reports receiving consulting fees from BioMarin, Novartis and PharmaNet; he also reports research support from Bristol-Myers Squibb, Roche and Seaside Therapeutics. Dr McDougale reports having received consultant fees from Bristol-Myers Squibb, Hoffman-LaRoche and Forest Research Institute; he has also received research support and is on the speakers' bureau of Bristol-Myers Squibb. Dr Scahill reports receiving consultant fees from Brackett, Pfizer, Hoffman, BioMarin; he has also received research support from Pfizer, Shire and Hoffman. The remaining authors declare no conflict of interest.

ACKNOWLEDGEMENTS

The RUPP Autism Network comprises the following investigators listed by role and study site: Ohio State University: principal investigator Michael G Aman, PhD, co-investigators L Eugene Arnold, MEd, MD, Yaser Ramadan, MD, Andrea Witwer, MA, Ronald Lindsay, MD and Patricia Nash, MD; University of California at Los Angeles: principal investigator James T McCracken, MD, co-investigators Bhavik Shah, MD, James McGough, MD, Pegeen Cronin, PhD, Lisa Lee, BA; Indiana University: principal

investigator Christopher J McDougale, MD, co-investigators David Posey, MD, Naomi Swiezy, PhD, Arlene Kohn, BA; Yale University: principal investigator Lawrence Scahill, MSN, PhD, co-investigators Andres Martin, MD, Kathleen Koenig, MSN, Fred Volkmar, MD, Deirdre Carroll, MSN, Allison Lancor, BS; Kennedy Krieger Institute: principal investigator Elaine Tierney, MD, co-investigators Jaswinder Ghuman, MD, Nilda Gonzalez, MD, Marco Grados, MD; National Institute of Mental Health: principal investigator Benedetto Vitiello, MD, co-investigator Louise Ritz, MBA; statisticians: Shirley Z Chuang, MS, Mark Davies, MPH, of Columbia University; data management: James Robinson, MEd, Don McMahon, MS, Nathan Kline Institute. Research supported by NIMH contracts N01 MH-70070 (principal investigator: Dr McCracken), N01 MH-70009 (principal investigator: Dr Scahill), N01 MH-70001 (principal investigator: Dr McDougale), and N01 MH 80011 (principal investigator: Dr Aman); by NIH Division of Research Resources General Clinical Research Center grants M01 RR-00750 (to Indiana University), M01 RR-00052 (to John Hopkins University), M01 RR-00034 (to Ohio University) and M01 RR-06022 (to Yale University); by NIMH grants MH-01805 (to Dr McCracken) NIMH grants MH094613 and T32MH073517 (Dr Nurmi) and MH-68627 (to Dr Posey); and by funding from the Korczak Foundation (to Dr Scahill). The opinions and assertions contained in this report are the private views of the authors and are not to be construed as official or as reflecting the views of the National Institute of Mental Health, the National Institutes of Health, or the Department of Health and Human Services.

REFERENCES

- Centers for Disease Control (2007). Prevalence of autism spectrum disorders, 14 Sites, United States, 2002. *Morbidity Mortality Weekly Rep.* pp 2007; **56**: 12–28.
- Lecavalier L. Behavioral and emotional problems in young people with pervasive developmental disorders: relative prevalence, effects of subject characteristics, and empirical classification. *J Autism Dev Disord* 2006; **36**: 1101–1114.
- McCracken JT. Disruptive behaviors in children with autism spectrum disorders. In: Coyle JHE, Klevzon A (eds) *Textbook of Autism Spectrum Disorders*. American Psychiatric Publishing: New York, NY, USA, 2011, pp 159–167.
- Simonoff E, Pickles A, Charman T, Chandler S, Loucas T, Baird G. Psychiatric disorders in children with autism spectrum disorders: prevalence, comorbidity, and associated factors in a population-derived sample. *J Am Acad Child Adolesc Psychiatry* 2008; **47**: 921–929.
- Frazier JA, Biederman J, Bellordre CA, Garfield SB, Geller DA, Coffey BJ *et al*. Should the diagnosis of attention-deficit/hyperactivity disorder be considered in children with pervasive developmental disorder? *J Attention Disord* 2001; **4**: 203–211.
- Goldstein S, Schwabach AJ. The comorbidity of pervasive developmental disorder and attention deficit hyperactivity disorder: results of a retrospective chart review. *J Autism Dev Disord* 2004; **34**: 329–339.
- Aman MG, Lam KS, Van Bourgondien ME. Medication patterns in patients with autism: temporal, regional, and demographic influences. *J Child Adolesc Psychopharmacol* 2005; **15**: 116–126.
- Esbensen AJ, Greenberg JS, Seltzer MM, Aman MG. A longitudinal investigation of psychotropic and non-psychotropic medication use among adolescents and adults with autism spectrum disorders. *J Autism Dev Disord* 2009; **39**: 1339–1349.
- Campbell M, Fish B, David R, Shapiro T, Collins P, Koh C. Response to triiodothyronine and dextroamphetamine: a study of preschool schizophrenic children. *J Autism Child Schizophr* 1972; **2**: 343–358.
- Campbell M, Small AM, Collins PJ, Friedman E, David R, Genieser N. Levodopa and levoamphetamine: a crossover study in young schizophrenic children. *Curr Ther Res Clin Exp* 1976; **19**: 70–86.
- Quintana H, Birmaher B, Stedje D, Lennon S, Freed J, Bridge J *et al*. Use of methylphenidate in the treatment of children with autistic disorder. *J Autism Dev Disord* 1995; **25**: 283–294.
- Handen BL, Johnson CR, Lubetsky M. Efficacy of methylphenidate among children with autism and symptoms of attention-deficit hyperactivity disorder. *J Autism Dev Disord* 2000; **30**: 245–255.
- Aman MG, Singh NN, Stewart AW, Field CJ. The aberrant behavior checklist: a behavior rating scale for the assessment of treatment effects. *Am J Ment Defic* 1985; **89**: 485–491.
- Aman MG, Singh NN, Stewart AW, Field CJ. Psychometric characteristics of the aberrant behavior checklist. *Am J Ment Defic* 1985; **89**: 492–502.
- Posey DJ, Aman MG, McCracken JT, Scahill L, Tierney E, Arnold LE *et al*. Positive effects of methylphenidate on inattention and hyperactivity in pervasive developmental disorders: an analysis of secondary measures. *Biol Psychiatry* 2007; **61**: 538–544.
- Greenhill LL, Abikoff HB, Arnold LE, Cantwell DP, Conners CK, Elliott G *et al*. Medication treatment strategies in the MTA Study: relevance to clinicians and researchers. *J Am Acad Child Adolesc Psychiatry* 1996; **35**: 1304–1313.

- 17 Stein MA, McGough JJ. The pharmacogenomic era: promise for personalizing attention deficit hyperactivity disorder therapy. *Child Adolesc Psychiatr Clin N Am* 2008; **17**: 475–490, xi-xii.
- 18 Nurmi EL, McCracken JT. Genetics in child psychiatry. In Sadock BJ, Sadock VA, Ruiz P, Kaplan HI, Ovid Technologies I (eds) *Kaplan & Sadock's Comprehensive Textbook of Psychiatry, 10th edn*. Wolters Kluwer Health/Lippincott Williams & Wilkins: Philadelphia, PA, USA, 2009.
- 19 McGough JJ, McCracken JT, Loo SK, Manganiello M, Leung MC, Tietjens JR *et al*. A candidate gene analysis of methylphenidate response in attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry* 2009; **48**: 1155–1164.
- 20 Stein MA, Waldman ID, Sarampote CS, Seymour KE, Robb AS, Conlon C *et al*. Dopamine transporter genotype and methylphenidate dose response in children with ADHD. *Neuropsychopharmacology* 2005; **30**: 1374–1382.
- 21 Purper-Ouakil D, Wohl M, Orejarena S, Cortese S, Boni C, Asch M *et al*. Pharmacogenetics of methylphenidate response in attention deficit/hyperactivity disorder: association with the dopamine transporter gene (SLC6A3). *Am J Med Genet B Neuropsychiatr Genet* 2008; **147B**: 1425–1430.
- 22 Froehlich TE, Epstein JN, Nick TG, Melguizo Castro MS, Stein MA, Brinkman WB *et al*. Pharmacogenetic predictors of methylphenidate dose-response in attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry* 2011; **50**: 1129–1139 e1122.
- 23 Cheon KA, Kim BN, Cho SC. Association of 4-repeat allele of the dopamine D4 receptor gene exon III polymorphism and response to methylphenidate treatment in Korean ADHD children. *Neuropsychopharmacology* 2007; **32**: 1377–1383.
- 24 Hamarman S, Fossella J, Ulger C, Brimacombe M, Dermody J. Dopamine receptor 4 (DRD4) 7-repeat allele predicts methylphenidate dose response in children with attention deficit hyperactivity disorder: a pharmacogenetic study. *J Child Adolesc Psychopharmacol* 2004; **14**: 564–574.
- 25 McGough J, McCracken J, Swanson J, Riddle M, Kollins S, Greenhill L *et al*. Pharmacogenetics of methylphenidate response in preschoolers with ADHD. *J Am Acad Child Adolesc Psychiatry* 2006; **45**: 1314–1322.
- 26 Tahir E, Yazgan Y, Cirakoglu B, Ozbay F, Waldman I, Asherson PJ. Association and linkage of DRD4 and DRD5 with attention deficit hyperactivity disorder (ADHD) in a sample of Turkish children. *Mol Psychiatry* 2000; **5**: 396–404.
- 27 Kooij JS, Boonstra AM, Vermeulen SH, Heister AG, Burger H, Buitelaar JK *et al*. Response to methylphenidate in adults with ADHD is associated with a polymorphism in SLC6A3 (DAT1). *Am J Med Genet B Neuropsychiatr Genet* 2008; **147B**: 201–208.
- 28 van der Meulen EM, Bakker SC, Pauls DL, Oteman N, Kruitwagen CL, Pearson PL *et al*. High sibling correlation on methylphenidate response but no association with DAT1-10R homozygosity in Dutch sibpairs with ADHD. *J Child Psychol Psychiatry* 2005; **46**: 1074–1080.
- 29 Zeni CP, Guimaraes AP, Polanczyk GV, Genro JP, Roman T, Hutz MH *et al*. No significant association between response to methylphenidate and genes of the dopaminergic and serotonergic systems in a sample of Brazilian children with attention-deficit/hyperactivity disorder. *Am J Med Genet B Neuropsychiatr Genet* 2007; **144B**: 391–394.
- 30 Gruber R, Joobor R, Grizenko N, Leventhal BL, Cook Jr EH, Stein MA. Dopamine transporter genotype and stimulant side effect factors in youth diagnosed with attention-deficit/hyperactivity disorder. *J Child Adolesc Psychopharmacol* 2009; **19**: 233–239.
- 31 Polanczyk G, Zeni C, Genro JP, Guimaraes AP, Roman T, Hutz MH *et al*. Association of the adrenergic alpha2A receptor gene with methylphenidate improvement of inattentive symptoms in children and adolescents with attention-deficit/hyperactivity disorder. *Arch Gen Psychiatry* 2007; **64**: 218–224.
- 32 da Silva TL, Pianca TG, Roman T, Hutz MH, Faraone SV, Schmitz M *et al*. Adrenergic alpha2A receptor gene and response to methylphenidate in attention-deficit/hyperactivity disorder-predominantly inattentive type. *J Neural Transm* 2008; **115**: 341–345.
- 33 Gainetdinov RR, Wetsel WC, Jones SR, Levin ED, Jaber M, Caron MG. Role of serotonin in the paradoxical calming effect of psychostimulants on hyperactivity. *Science* 1999; **283**: 397–401.
- 34 Seeger G, Schloss P, Schmidt MH. Marker gene polymorphisms in hyperkinetic disorder—predictors of clinical response to treatment with methylphenidate? *Neurosci Lett* 2001; **313**: 45–48.
- 35 Alfimova MV, Golimbet VE, Gritsenko IK, Lezheiko TV, Abramova LI, Strel'tsova MA *et al*. Interaction of dopamine system genes and cognitive functions in patients with schizophrenia and their relatives and in healthy subjects from the general population. *Neurosci Behav Physiol* 2007; **37**: 643–650.
- 36 de Frias CM, Annerbrink K, Westberg L, Eriksson E, Adolfsson R, Nilsson LG. Catechol O-methyltransferase Val158Met polymorphism is associated with cognitive performance in nondemented adults. *J Cogn Neurosci* 2005; **17**: 1018–1025.
- 37 Mattay VS, Goldberg TE, Fera F, Hariri AR, Tessitore A, Egan MF *et al*. Catechol O-methyltransferase val158-met genotype and individual variation in the brain response to amphetamine. *Proc Natl Acad Sci USA* 2003; **100**: 6186–6191.
- 38 Cheon KA, Jun JY, Cho DY. Association of the catechol-O-methyltransferase polymorphism with methylphenidate response in a classroom setting in children with attention-deficit hyperactivity disorder. *Int Clin Psychopharmacol* 2008; **23**: 291–298.
- 39 Kereszturi E, Tarnok Z, Bognar E, Lakatos K, Farkas L, Gadoros J *et al*. Catechol-O-methyltransferase Val158Met polymorphism is associated with methylphenidate response in ADHD children. *Am J Med Genet B Neuropsychiatr Genet* 2008; **147B**: 1431–1435.
- 40 Volkow ND, Wang GJ, Fowler JS, Ding YS. Imaging the effects of methylphenidate on brain dopamine: new model on its therapeutic actions for attention-deficit/hyperactivity disorder. *Biol Psychiatry* 2005; **57**: 1410–1415.
- 41 Kollins SH, Anastopoulos AD, Lachiewicz AM, FitzGerald D, Morrissey-Kane E, Garrett ME *et al*. SNPs in dopamine D2 receptor gene (DRD2) and norepinephrine transporter gene (NET) are associated with continuous performance task (CPT) phenotypes in ADHD children and their families. *Am J Med Genet B Neuropsychiatr Genet* 2008; **147B**: 1580–1588.
- 42 Lee B, London ED, Poldrack RA, Farahi J, Nacca A, Monterosso JR *et al*. Striatal dopamine d2/d3 receptor availability is reduced in methamphetamine dependence and is linked to impulsivity. *J Neurosci* 2009; **29**: 14734–14740.
- 43 Nieminen-von Wendt TS, Metsahonkala L, Kulomaki TA, Aalto S, Autti TH, Vanhala R *et al*. Increased presynaptic dopamine function in Asperger syndrome. *Neuroreport* 2004; **15**: 757–760.
- 44 Makkonen I, Riikonen R, Kokki H, Airaksinen MM, Kuikka JT. Serotonin and dopamine transporter binding in children with autism determined by SPECT. *Dev Med Child Neurol* 2008; **50**: 593–597.
- 45 Nakamura K, Sekine Y, Ouchi Y, Tsujii M, Yoshikawa E, Futatsubashi M *et al*. Brain serotonin and dopamine transporter bindings in adults with high-functioning autism. *Arch Gen Psychiatry* 2010; **67**: 59–68.
- 46 RUPP Autism Network. Randomized, controlled, crossover trial of methylphenidate in pervasive developmental disorders with hyperactivity. *Arch Gen Psychiatry* 2005; **62**: 1266–1274.
- 47 Lord C, Rutter M, Le Couteur A. Autism diagnostic interview-revised: a revised version of a diagnostic interview for caregivers of individuals with possible pervasive developmental disorders. *J Autism Dev Disord* 1994; **24**: 659–685.
- 48 Faraone SV, Perlis RH, Doyle AE, Smoller JW, Goralnick JJ, Holmgren MA *et al*. Molecular genetics of attention-deficit/hyperactivity disorder. *Biol Psychiatry* 2005; **57**: 1313–1323.
- 49 Wendland JR, Martin BJ, Kruse MR, Lesch KP, Murphy DL. Simultaneous genotyping of four functional loci of human SLC6A4, with a reappraisal of 5-HTTLPR and rs25531. *Mol Psychiatry* 2006; **11**: 224–226.
- 50 Cook Jr. EH, Stein MA, Krasowski MD, Cox NJ, Olkon DM, Kieffer JE *et al*. Association of attention-deficit disorder and the dopamine transporter gene. *Am J Hum Genet* 1995; **56**: 993–998.
- 51 Shaikh S, Collier D, Kerwin RW, Pilowsky LS, Gill M, Xu WM *et al*. Dopamine D4 receptor subtypes and response to clozapine. *Lancet* 1993; **341**: 116.
- 52 Lott DC, Kim SJ, Cook EH, de Wit H. Serotonin transporter genotype and acute subjective response to amphetamine. *Am J Addict* 2006; **15**: 327–335.
- 53 Barrett JC, Fry B, Maller J, Daly MJ. Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics* 2005; **21**: 263–265.
- 54 Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM *et al*. The human genome browser at UCSC. *Genome Res* 2002; **12**: 996–1006.
- 55 Rosenbloom KR, Dreszer TR, Long JC, Malladi VS, Sloan CA, Raney BJ *et al*. ENCODE whole-genome data in the UCSC Genome Browser: update 2012. *Nucleic Acids Res* 2012; **40**(Database issue): D912–D917.
- 56 Ward LD, Kellis M. HaploReg: a resource for exploring chromatin states, conservation, and regulatory motif alterations within sets of genetically linked variants. *Nucleic Acids Res* 2012; **40**(Database issue): D930–D934.
- 57 Bakker PR, van Harten PN, van Os J. Antipsychotic-induced tardive dyskinesia and the Ser9Gly polymorphism in the DRD3 gene: a meta analysis. *Schizophr Res* 2006; **83**: 185–192.
- 58 Arnsten AF. Catecholamine influences on dorsolateral prefrontal cortical networks. *Biol Psychiatry* 2011; **69**: e89–e99.
- 59 Dalley JW, Cardinal RN, Robbins TW. Prefrontal executive and cognitive functions in rodents: neural and neurochemical substrates. *Neurosci Biobehav Rev* 2004; **28**: 771–784.
- 60 Oades RD, Sadile AG, Sagvolden T, Viggiano D, Zuddas A, Devoto P *et al*. The control of responsiveness in ADHD by catecholamines: evidence for dopaminergic, noradrenergic and interactive roles. *Dev Sci* 2005; **8**: 122–131.
- 61 Purper-Ouakil D. [Use of psychotropic drugs in children]. *Arch Pediatr* 2008; **15**: 1834–1836.
- 62 Williams GV, Goldman-Rakic PS. Modulation of memory fields by dopamine D1 receptors in prefrontal cortex. *Nature* 1995; **376**: 572–575.
- 63 Misener VL, Luca P, Azeke O, Crosbie J, Waldman I, Tannock R *et al*. Linkage of the dopamine receptor D1 gene to attention-deficit/hyperactivity disorder. *Mol Psychiatry* 2004; **9**: 500–509.

- 64 Bobb AJ, Addington AM, Sidransky E, Gornick MC, Lerch JP, Greenstein DK *et al*. Support for association between ADHD and two candidate genes: NET1 and DRD1. *Am J Med Genet B Neuropsychiatr Genet* 2005; **134B**: 67–72.
- 65 Bobb AJ, Castellanos FX, Addington AM, Rapoport JL. Molecular genetic studies of ADHD: 1991 to 2004. *Am J Med Genet B Neuropsychiatr Genet* 2005; **132B**: 109–125.
- 66 Gillberg C, Svennerholm L. CSF monoamines in autistic syndromes and other pervasive developmental disorders of early childhood. *Br J Psychiatry* 1987; **151**: 89–94.
- 67 Guiard BP, El Mansari M, Blier P. Cross-talk between dopaminergic and norenergic systems in the rat ventral tegmental area, locus ceruleus, and dorsal hippocampus. *Mol Pharmacol* 2008; **74**: 1463–1475.



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/3.0/>