

Photoplethysmography rhythm interpretation: an essential skill in an era of novel technologies

Ying X. Gue^{1,2}, Wahbi K. El-Bouri ^{1,2}, and Gregory Y.H. Lip^{1,2,3*}

¹Liverpool Centre for Cardiovascular Science, University of Liverpool and Liverpool Heart & Chest Hospital, Liverpool L7 8TX, UK; ²Department of Cardiovascular and Metabolic Medicine, University of Liverpool, Liverpool L69 3BX, UK; and ³Aalborg Thrombosis Research Unit, Department of Clinical Medicine, Aalborg University, 9000 Aalborg, Denmark

Received 28 July 2021; accepted 30 July 2021; online publish-ahead-of-print 3 August 2021

This editorial refers to ‘The photoplethysmography dictionary: practical guidance on signal interpretation and clinical scenarios from TeleCheck-AF’, by R.M.J. van der Velden et al. doi: 10.1093/ehjdh/ztab050.

Introduction

Innovations in smart wearables and mHealth have provided avenues in screening and monitoring of cardiovascular health at a population wide level. With the disruption to conventional healthcare during the coronavirus-19 (COVID-19) pandemic, the role of remote monitoring and management of different health conditions have accelerated and have taken a huge step forward, providing clinicians with many novel devices to support these endeavours.

The TeleCheck-AF study is a multicentre study utilizing photoplethysmography (PPG) technology through a built-in camera allowing semi-continuous heart rate and rhythm monitoring for patients with atrial fibrillation (AF).¹ Despite the widespread use of PPG-based technology, clinicians and health care providers have not been introduced to it formally and in a structured way like in the case of electrocardiograms (ECGs).

In this issue of the journal, the TeleCheck-AF investigators provided, based on their experience, a structured stepwise practical guide on evaluating and interpreting PPG signals. This comprises of a simple five-step approach: (i) checking the quality of tracing, (ii) checking the output of the PPG FibriCheck© algorithm, (iii) checking for regularity, (iv) checking medical history to increase likelihood of arrhythmia and increase pre-test probability, and (v) proceeding with further diagnostic testing and therapy as indicated.¹

Apart from this, the investigators also shared some common clinical scenarios to help clinicians better understand the application in real life, including the important limitations of PPG such as the inability to differentiate between regular tachycardias and the requirements for ECG documentation for the final diagnosis.

This is a pertinent topic, with the current rapid advancements and adoption of novel devices into routine clinical practice. It certainly appears that the integration of practical guides of interpretation of PPG into the medical curriculum, similar to that of ECG interpretation, may not be too far in the future of medical education.

Where do we go from here?

Artificial intelligence (AI) algorithms were used in TeleCheck-AF to determine which patients were clearly in AF. Those where the diagnosis was uncertain were then checked for regularity. Nonetheless, clinicians should be provided with the knowledge to interpret the results of these devices not only to double-check the interpretation of the algorithm but also to unravel the enigma surrounding new technology. Also, the future of medical education goes beyond the skill of PPG interpretation and, whilst not explicitly tackled in van der Velden et al.,¹ is a current theme that runs throughout.

The use of AI to assist with interpretation should not be a deterrent for healthcare professionals to learn its interpretation. Indeed, there has been an exponential growth in AI use the diagnosis and risk assessment of patients with AF.² Indeed, AI has also been applied to help improve stroke risk stratification, accounting for dynamic changes in stroke risks.^{3,4}

The bottom line is that AI and *in silico* medical technologies, along with robotics and medical devices, are developing at a rapid pace. Undoubtedly, these new technologies will play a vital role in the future of medicine, as will the ability to understand their use and limitations. Hence, medical education must expand to incorporate the use and assessment of novel technologies in clinical settings. While attitudes to new technologies are on the whole positive, uptake of these technologies remains poor, primarily due to a lack of knowledge of the technology and discomfort when using them.^{5,6}

The TeleCheck-AF project is therefore an important step towards bridging that knowledge gap and developing familiarity with PPG

The opinions expressed in this article are not necessarily those of the Editors of the *European Heart Journal – Digital Health* or of the European Society of Cardiology.

*Corresponding author. Tel: 0151 7949020, Email: gregory.lip@liverpool.ac.uk

© The Author(s) 2021. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com




Figure 1 Schematic demonstrating four necessary considerations for the successful adoption of novel technology.

rhythm interpretation. Without these initiatives to enhance adoption, the technology itself is ineffectual, and as such this is a crucial step in the improvement of digital health provision.

For the successful adoption of novel technologies, it is not only sufficient that there is robust evidence to support their clinical use (Figure 1). The successful application of mobile health (mHealth) technologies to improve patient care in AF has been in the mAFA-II trial, which was a cluster randomized trial comparing the ABC (Atrial fibrillation Better Care) pathway against usual care, resulting in a significant reduction in stroke/thromboembolism, mortality and bleeding with the mAFA App-based intervention.⁷ Importantly, the long-term extension cohort showed a high adherence (>70%) and persistence (>90%) of use.⁸ This is important given that ABC pathway compliance has major implications for reducing mortality and morbidity from stroke, major bleeding and hospitalizations.^{9,10}

Nonetheless, healthcare professionals must also understand the use of the technology and its limitations. The limitations are particularly important as this will create clear bounds of use. Beyond an understanding of the technology and the limitations of its use, data security, ownership, and access are crucial aspects of technology integration that are needed to build trust. Again, this is an area where technology can help, through the use of blockchain for example.¹¹ Care must also be taken to avoid the exacerbation of existing health inequalities through the adoption of digital healthcare solutions. It is sobering to remember that just under half the world's population has no access to the internet¹² and that lack of internet access is associated with poorer health.¹³

Taken together, strong evidence, strong data security, a workforce that is familiar and understands the technology, along with a focus on healthcare equity and equality, all combine to accelerate the successful adoption of these novel healthcare technologies. Time will tell.

Conflict of interest: Consultant and speaker for BMS/Pfizer, Boehringer Ingelheim and Daiichi-Sankyo. No fees are received personally. WKE and YXG has no conflicts of interests to declare.

References

- van der Velden RMJ, Verhaert DVM, Hermans ANL, Duncker D, Manninger M, Betz K, Gawalko M, Desteghe L, Pisters R, Hemels M, Pison L, Sohaib A, Sultan A, Steven D, Wijtvlit P, Gupta D, Svennberg E, Luermans JCLM, Chaldoupi M, Vernooij K, den Uijl D, Lodzinski P, Jansen WPJ, Eckstein J, Bollmann A, Vandervoort P, Crijns HJGM, Tieleman R, Heidbuchel H, Pluymaekers NAHA, Hendriks JM, Linz D; TeleCheck-AF Investigators. The photoplethysmography dictionary: practical guidance on signal interpretation and clinical scenarios from TeleCheck-AF. *Eur Heart J Digit Health* 2021;doi:10.1093/ehjdh/ztob050.
- Olier I, Ortega-Martorell S, Pieroni M, Lip GYH. How machine learning is impacting research in atrial fibrillation: implications for risk prediction and future management. *Cardiovasc Res* 2021;117:1700–1717.
- Lip GYH, Tran G, Genaidy A, Marroquin P, Estes C, Landsheftl J. Improving dynamic stroke risk prediction in non-anticoagulated patients with and without atrial fibrillation: Comparing common clinical risk scores and machine learning algorithms. *Eur Heart J Qual Care Clin Outcomes* 2021.
- Lip GYH, Genaidy A, Tran G, Marroquin P, Estes C, Sloop S. Improving stroke risk prediction in the general population: a comparative assessment of common clinical rules, a new multimorbidity index, and machine-learning-based algorithms. *Thromb Haemost* 2021.
- Ayatollahi H, Sarabi FZP, Langarizadeh M. Clinicians' knowledge and perception of telemedicine technology. *Perspect Health Inf Manag* 2015;12:1c.
- Oh S, Kim JH, Choi S-W, Lee HJ, Hong J, Kwon SH. Physician confidence in artificial intelligence: an online mobile survey. *J Med Internet Res* 2019;21:e12422.
- Guo Y, Lane DA, Wang L, Zhang H, Wang H, Zhang W, Wen J, Xing Y, Wu F, Xia Y, Liu T, Wu F, Liang Z, Liu F, Zhao Y, Li R, Li X, Zhang L, Guo J, Burnside G, Chen Y, Lip GYH; mAFA-App II Trial Investigators. Mobile health technology to improve care for patients with atrial fibrillation. *J Am Coll Cardiol* 2020;75:1523–1534.
- Guo Y, Guo J, Shi X, Yao Y, Sun Y, Xia Y, Yu B, Liu T, Chen Y, Lip G; mAFA-App II Trial investigators. Mobile health technology-supported atrial fibrillation screening and integrated care: a report from the mAFA-II trial Long-term Extension Cohort. *Eur J Intern Med* 2020;82:105–111.
- Romiti GF, Pastori D, Rivera-Caravaca JM, Ding WY, Gue YX, Menicelli D, Gumprecht J, Koziel M, Yang P-S, Guo Y, Lip GY, Proietti M. Adherence to the 'Atrial Fibrillation Better Care' (ABC) pathway in patients with atrial fibrillation. *Thromb Haemost* 2021.
- Yoon M, Yang P-S, Jang E, Yu HT, Kim T-H, Uhm J-S, Kim J-Y, Sung J-H, Pak H-N, Lee M-H, Joung B, Lip GYH. Improved population-based clinical outcomes of patients with atrial fibrillation by compliance with the simple ABC (Atrial Fibrillation Better Care) pathway for integrated care management: a nationwide cohort study. *Thromb Haemost* 2019;19:1695–1703.
- El-Bouri WK, Gue Y, Lip GYH. 'Rise of the machines': the next frontier in individualized medicine. *Cardiovasc Res* 2021.
- Makri A. Bridging the digital divide in health care. *Lancet Digit Health* 2019;1:e204–e205.
- Sieck CJ, Sheon A, Ancker JS, Castek J, Callahan B, Siefert A. Digital inclusion as a social determinant of health. *NPJ Digit Med* 2021;4:52.