

SCIENTIFIC REPORTS



OPEN

Litter input decreased the response of soil organic matter decomposition to warming in two subtropical forest soils

Received: 18 April 2016
Accepted: 02 September 2016
Published: 20 September 2016

Qingkui Wang^{1,2}, Tongxin He¹ & Jing Liu^{1,3}

Interaction effect of temperature and litter input on SOM decomposition is poor understood, restricting accurate prediction of the dynamics and stocks of soil organic carbon under global warming. To address this knowledge gap, we conducted an incubation experiment by adding ¹³C labeled leaf-litter into a coniferous forest (CF) soil and a broadleaved forest (BF) soil. In this experiment, response of the temperature sensitivity (Q_{10}) of SOM decomposition to the increase in litter input was investigated. The temperature dependences of priming effect (PE) and soil microbial community were analyzed. The Q_{10} for CF soil significantly decreased from 2.41 in no-litter treatment to 2.05 in litter-added treatment and for BF soil from 2.14 to 1.82, suggesting that litter addition decreases the Q_{10} . PE in the CF soil was 24.9% at 20 °C and 6.2% at 30 °C, and in the BF soil the PE was 8.8% at 20 °C and -7.0% at 30 °C, suggesting that PE decreases with increasing temperature. Relative PE was positively related to the concentrations of Gram-negative bacterial and fungal PLFAs. This study moves a step forward in understanding warming effect on forest carbon cycling by highlighting interaction effect of litter input and warming on soil carbon cycling.

Soil in terrestrial ecosystems contains 1550 Pg organic C¹, which is a result of the balance between input from leaves and output from decomposition of soil organic matter (SOM)². SOM decomposition is not only controlled by temperature and moisture² but also affected by litter input³. The global increase of 1.1–6.4 °C in temperature by the end of the 21st century is likely to alter the input of litter to soil systems^{4,5}. This alteration of exogenous substrate input will change soil organic C (SOC) stocks through priming effect (PE)⁶. In addition, changes in the temperature sensitivity (Q_{10}) of SOM decomposition induced by increasing substrate input also determine the response of terrestrial C balance to global warming⁷. Therefore, improving prediction of SOC dynamics under global warming is important to understand the combined effect of temperature and litter input on SOM decomposition.

Unfortunately, a consensus has not yet been reached on the magnitude of Q_{10} of SOM decomposition because of a large variation on Q_{10} , although considerable efforts have been exerted to study Q_{10} values^{2,8,9}. This inconsistency may stem from confounding factors such as substrate availability and microbial community structure^{10,11}. Increased litter input to soil systems is likely to alter substrate availability by releasing labile C during litter decomposition. Recently, a number of empirical studies have identified the control of labile substrate availability to Q_{10} of microbial respiration^{11–13}; these studies demonstrated that the addition of readily available C substrate significantly increased Q_{10} values. Despite its importance, substrate availability is probably the least studied factor that affects the Q_{10} of SOM decomposition². The abovementioned studies generally used glucose as sources of labile substrate. However, as an important source of labile substrate in actual forest ecosystems, how litter addition affects Q_{10} still remains poor understood.

Another important process that affects SOC cycling is PE, being defined as the changes in the decomposition rate of SOM after input of exogenous substrates¹⁴. Several studies on PE have been conducted, but the direction and intensity of PE in various experiments are large uncertain, which range from -95.1% to 1207%^{15,16}. This

¹Key Laboratory of Forest Ecology and Management, Institute of Applied Ecology, Chinese Academy of Sciences, Shenyang 110164, P. R. China. ²Huitong Experimental Station of Forest Ecology, Chinese Academy of Sciences, Huitong 418307, P.R. China. ³University of Chinese Academy of Sciences, Beijing 100049, P.R. China. Correspondence and requests for materials should be addressed to Q.W. (email: wqkui@163.com)

		Common bacteria	GP	GN	GP/GN	Fungi	B/F	Actinomycete
20 °C	CF	9.25 ± 0.88aA	11.55 ± 1.09aA	5.97 ± 0.30aA	1.93 ± 0.12aA	3.88 ± 0.42aA	6.93 ± 0.39bA	2.94 ± 0.22aA
	CFL	11.51 ± 0.23bA	12.61 ± 0.41aA	7.34 ± 0.32bA	1.72 ± 0.13aA	6.00 ± 0.23bB	5.25 ± 0.21aA	3.48 ± 0.19aA
	BF	19.92 ± 0.13aA	18.27 ± 0.83aA	14.60 ± 1.42aA	1.26 ± 0.12aA	9.74 ± 0.39aA	5.42 ± 0.19bA	5.10 ± 0.29aA
	BFL	22.12 ± 0.76bA	17.37 ± 1.01aA	15.32 ± 0.54aA	1.13 ± 0.08aA	12.33 ± 0.51bA	4.45 ± 0.21aA	5.30 ± 0.26aA
30 °C	CF	9.25 ± 0.27aA	10.41 ± 0.34aA	5.74 ± 0.34aA	1.82 ± 0.11aA	3.28 ± 0.19aA	7.74 ± 0.29bA	2.98 ± 0.18aA
	CFL	11.73 ± 0.68bA	11.91 ± 0.46bA	6.75 ± 0.28bA	1.76 ± 0.12aA	4.51 ± 0.24bA	6.74 ± 0.32aB	3.33 ± 0.17aA
	BF	22.63 ± 0.95aB	19.08 ± 0.92aA	17.05 ± 0.84aA	1.12 ± 0.09aA	9.54 ± 0.55aA	6.17 ± 0.28aB	5.18 ± 0.20aA
	BFL	26.12 ± 0.45bB	21.76 ± 0.57bB	19.63 ± 1.27aB	1.11 ± 0.10aA	11.25 ± 0.61bA	6.01 ± 0.23aB	5.98 ± 0.55aA

Table 1. Changes in the concentrations (nmol g⁻¹ soil) of PLFAs and two PLFA ratios in soils with needle addition at 20 and 30 °C at the end of 42-day incubation period. Different lower case letters after data denote significant effect of litter addition on soil microbial community for same soils; different capital letters after data denote significant effect of temperature on soil microbial community for same soils. GP and GN represent Gram-positive and Gram-negative bacteria.

uncertainty is attributed to the fact that PE is dependent on many factors, such as incubation temperature^{17,18}, soil nutrient availability^{19–21}, and characteristics of soil microbial community⁹. For example, Kuzyakov predicted that incubation temperature may mediate PE because increasing temperature can accelerate most enzyme activities¹⁷; however, direct evidence is lacking on the influence of temperature on the PE in forest ecosystems, although some studies reported the impacts of temperature on rhizosphere PE in arable soils^{22,23}. Therefore, it remains unclear whether SOC losses through PE would change after temperature increase.

The important role of biotic factors (e.g., microbial community structure) in SOM decomposition is now being recognized^{17,20,24}. The alteration of soil microbial community structure and activity by exogenous substrate addition has been demonstrated^{20,25}, which may affect Q_{10} ¹³ and PE^{26,27}. Some studies found that the level of microbial community response affected Q_{10} ⁹. In addition, a shift from C limitation for soil microorganisms to nutrient limitation induced by exogenous C addition may result in microbial mining additional SOM to acquire nutrients, and therefore positive PE^{20,28}. However, how the response of soil microbial community structure to temperature affects Q_{10} and the PE of SOM decomposition remains unclear.

In the present study, we presented the result of an incubation experiment that combined short-term manipulations of temperature and substrate (¹³C-labeled leaf litter) addition in two different soils from a coniferous forest (CF) and an evergreen broadleaved forest (BF). The ¹³C isotopic differences between litter and native SOC allowed the quantification of CO₂-C sources from decomposition of litter and SOM. These data were used to determine Q_{10} and PE in different soils. The present study aimed to investigate the interactive effect of exogenous substrate input and warming on SOM decomposition in subtropical forests. Based on our knowledge, we hypothesized that litter input would decrease Q_{10} , and warming would decrease PE intensity.

Results

Soil microbial community composition. Litter addition increased the common bacterial concentration by 24.4% and 26.8% in CF soils and 11.0% and 15.4% in BF soils at 20 and 30 °C, respectively (Table 1). Fungal concentration was increased by 54.6% and 37.5% in CF soils and 26.6% and 17.9% in BF soils at 20 and 30 °C, respectively. However, litter addition increased Gram-positive bacterial concentrations averagely by 14.2% in two soils only at 30 °C and increased Gram-negative bacterial concentrations only in the CF soils by 22.9% and 17.6% at 20 and 30 °C, respectively. Litter addition decreased the ratio of bacteria to fungi with an average of 18.4%, with the exception of BF soil at 30 °C. Increasing temperature increased bacterial concentrations averagely by 15.8% and the ratio of bacteria to fungi averagely by 22.4% in BF soils. Compared with CF soils, BF soils had higher microbial concentrations measured by phospholipid fatty acids (PLFAs) but lower than the ratios of Gram-positive to Gram-negative bacteria and bacteria to fungi at the same temperature.

Litter-C fate. Expressed as percentage of the amount of added litter, CO₂ released rate from litter differed among treatments (Fig. 1). However, the pattern of litter decomposition was similar among all treatments. Increasing temperature promoted litter decomposition averagely by 52.4%. However, litter decomposition between in CF and BF soils did not differ. At the end of incubation, approximately 10% of the added litter-C was respired as CO₂ at 20 °C and 12.7% at 30 °C for both soils.

The amount of litter-derived C incorporated into PLFAs was affected by temperature and soil type (Fig. 2). Over the 42-day incubation, the litter-C incorporated into total PLFAs was 1.98 mg C kg⁻¹ and 1.48 mg C kg⁻¹ soil for CF soil at 20 °C and 30 °C, respectively, and 1.53 mg C kg⁻¹ and 1.24 mg C kg⁻¹ for BF soil. Elevating temperature decreased the amount of litter-C incorporated into fungal PLFAs in the both soils, as well as the common and Gram-negative bacterial PLFAs in the CF soil.

Temperature sensitivity of SOM decomposition. The temperature sensitivity of SOM decomposition in the treatments without litter addition was significantly different in the CF and BF soils (Fig. 3). CF soils (2.41 ± 0.10) had slightly higher Q_{10} value than BF soils (2.14 ± 0.13) at the end of the 42-day incubation. A significant decline in Q_{10} was detected after litter addition, and Q_{10} decreased to 2.05 in the CF soils and 1.82 in the BF soils, indicating that litter addition decreased the response of SOM decomposition to warming. The Q_{10}




Figure 1. The cumulative fraction of added litter C released as CO₂ during the 42-day incubation period.




Figure 2. Concentration of litter-derived C within different PLFAs 42 days after treatments. Significant differences ($P < 0.05$) are labeled by asterisks between treatment pairs (t-test). Data are means with SD ($n = 3$).




Figure 3. Temperature sensitivity (Q_{10}) of SOM and litter decomposition in the control (no litter) and treated (added litter) soils from a coniferous forest (CF) and a broadleaved forest (BF) during the 42-day incubation period.

dynamics in the CF soils without litter remained constant, but Q_{10} in the BF soils without litter initially increased slightly and then decreased significantly. By contrast, litter addition led to a significant increase in the Q_{10} with time, indicating that litter addition modifies the dynamics of the temperature sensitivity of SOM decomposition.




Figure 4. Cumulative C respired as CO₂ from SOM in the control soils and litter-amended soils at 20 and 30 °C from a CF and a BF during the 42-day incubation period.




Figure 5. The relative priming effect of litter addition on SOM decomposition in soils from a CF and a BF at 20 and 30 °C.

	Bacteria	Fungi	Bacteria:fungi	GP	GN	GP:GN	Actinomycete
CF	0.451	0.939**	-0.923**	0.630	0.781*	-0.492	0.377
BF	-0.940**	0.839*	-0.955**	-0.923**	0.898**	0.334	-0.648

Table 2. Relationship between priming effect and concentrations of different microbial groups in CF and BF soils. *, **Denote significant difference at $P < 0.05$ and 0.01 . GP and GN represent Gram-positive and Gram-negative bacteria.

Priming effect. The cumulative CO₂-C released from SOM in the control at 30 °C was 140% higher for CF soils and 113% higher for BF soils than their counterparts at 20 °C (Fig. 4). After adding litters, the CO₂-C that respired from SOM at 30 °C was 104% and 82% higher for CF and BF soils than that at the 20 °C. These results indicate that litter addition decreases the response of SOM decomposition to temperature elevation. In the control, the SOM decomposition of BF soils was 30% greater than that of CF soils at 20 °C, and 16% higher at 30 °C, whereas in the litter-added treatments, the SOM decomposition of BF soils was 14% greater than that of CF soils at 20 °C, and 2% higher at 30 °C.

Litter addition primed SOM decomposition, but relative PE differed in CF and BF soils (Fig. 5). PE was higher in the CF soils (24.9%) than in the BF soils (8.8%) at 20 °C. Stimulation effect (positive PE = 6.2%) was observed in the CF soils at 30 °C, but inhibition effect (negative PE = -7.0%) was found in the BF soils. This result indicates that elevating temperature decreases PE in both soils at the same degree. A significantly positive correlation was observed between relative PE and the concentrations of fungi and Gram-negative bacteria PLFAs, and a negative correlation was found between relative PE and the ratio of bacteria to fungi in both forest soils (Table 2).

Discussion

To the best of our knowledge, this study is the first to investigate the effect of litter input on the temperature sensitivity of SOM decomposition in forest ecosystems. The finding that litter addition decreased the temperature sensitivity of SOM decomposition supported our first hypothesis; however, this finding contradicts the observations

	CF soil	BF soil
SOC (g·kg ⁻¹)	26.2 ± 2.3a	51.7 ± 1.2b
Total N (g·kg ⁻¹)	1.72 ± 0.20a	3.85 ± 0.13b
C:N ratio	15.2 ± 1.1b	13.4 ± 0.4a
Total P (g·kg ⁻¹)	0.20 ± 0.01a	0.35 ± 0.09b
NH ₄ ⁺ -N (mg·kg ⁻¹)	4.77 ± 1.01a	8.46 ± 2.00b
Available P (mg·kg ⁻¹)	1.99 ± 0.45a	2.11 ± 0.59a
Sand content (%)	8.70 ± 1.91a	8.11 ± 1.73a
Silt content (%)	42.76 ± 1.30a	43.95 ± 1.29a
Clay content (%)	48.54 ± 1.95a	47.94 ± 2.01a

Table 3. Properties of surface mineral soils in a coniferous forest (CF) and a broadleaved forest (BF). Different letters followed data in the same row denote significant difference at $P < 0.05$.

of some studies that suggested that the addition of readily available carbon substrate increased $Q_{10}^{10,12,13}$. This different response may be attributed to the difference in the quality of substrates added to the soils. The litter used in our experiment was recalcitrant to soil microorganisms, whereas in other studies sucrose or glucose was used. Another potential explanation is the fact that the water-soluble N in the litter entered the soils and thereby increased soil N availability, resulting in a low response of SOM decomposition to elevated temperature. Decline in the Q_{10} of SOM decomposition after litter addition suggests that the stimulation effect of warming on SOM decomposition will be decreased by the increase in litter input under atmospheric CO₂ enrichment and can partly abate C losses in this subtropical region. Therefore, this will result in a less positive feedback to climate change than previously expected for subtropical forest ecosystems.

Soil microbial community is generally limited by C²⁹, and some studies documented that an increase in substrate availability positively affects temperature sensitivity^{10,13}. In the present study, the significant increase in the Q_{10} of soils with litters during incubation was attributed to the gradual release of dissolved organic substrate in litters to the soil, which provided energy for soil microorganisms. In the BF soils without litter, a decrease in Q_{10} with incubation time was also attributed to the decrease of substrate availability because of the depletion of liable substrate in soils.

Regardless of the litter addition, higher Q_{10} was observed in the CF soils than in the BF soils. This result may be explained by the higher C:N ratio of the CF soil than that of the BF soil (Table 3) because low-quality SOM with high C:N ratio has higher temperature sensitivity than high-quality SOM with low C:N ratio³⁰. Some studies also found that SOM quality influences the temperature sensitivity of SOM decomposition^{31,32}. CF soils also had a lower N content than BF soils (Table 3), which could result in significant N depletion with increasing temperature and potentially high SOM decomposition. In the present study, CF soils had higher ratios of bacteria:fungi and Gram-positive:Gram-negative bacteria than BF soils (Table 1). The pine litter used in the present study was more compositionally similar to the litter from CF than BF. Thus, in comparison with BF soil, the microbes that decompose pine litter would be more likely to use CF soil as C source under elevated temperature in this forest ecosystem, which increased SOM mined by microbes to gain additional N in CF soil.

An important finding in our study is the strong decrease in PE with increasing soil temperature in the subtropical forest ecosystem. This finding indicates that the magnitude of PE depends on environmental temperature. Our findings confirmed the prediction of Kuzyakov using experimental data¹⁷. In a meta-analysis literature, Zhang *et al.* also found that PE intensity below 20 °C (38.4%) was significantly higher than that above 20 °C (20.2%)¹⁶. One potential mechanism for this phenomenon is the fact that fast growth of microbes at high temperature could lead to more N immobilization³³. Another potential explanation is the high SOM decomposition level in soils without litter at high temperature (Fig. 4). A similar observation was also reported by Thiessen *et al.*⁹. Our findings indicate that global warming will likely decrease the effect of PE induced by litter input on SOC storage in the subtropical forests. The response of PE to temperature change is crucial in predicting the response of SOM decomposition in future climate change. Therefore, additional experiments on this topic should be widely conducted in a range of forest ecosystems in future research.

A higher PE was observed in CF soils than in BF soils (Fig. 5), indicating that external C input will result in higher native SOC loss in CF soils than in BF soils. According to discussions of Zhang *et al.*¹⁶, differences in soil properties (e.g., NH₄⁺-N content, C:N ratio) can partially explain PE variations. Two mechanisms may explain these results. The first is that low N availability stimulates soil microbes to mine SOM and search for nutrients to sustain their growth^{14,34}. In the present study, BF soils had richer amounts of nutrients (e.g., N, P) and lower C:N ratio than CF soils (Table 3), which result in lower PE in BF soils than that in CF soils. In a temperate forest, Wang *et al.* demonstrated that soils with high C:N ratio had high PE after soluble C addition³⁵. The second mechanism is the concept of the “nutrient bank” mechanism proposed by Fontaine and Barot³⁶, wherein microbial N mining regulates SOM decomposition and nutrient release. This mechanism is supported by the negative relationship between PE and nutrient availability^{37,38}.

PE also depends on the characteristics of soil microbial community^{9,17,28}. In the present study, soil microbial growth and changes in community structure occurred after litter input (Table 1), which were partly responsible for PEs. Recently, some studies also suggested that PE may be related to soil microbial community because PEs are accompanied by changes in the composition of soil microbial community^{20,39,40}; and Bell *et al.* showed that PE was related to the size of biomass and its composition, particularly bacteria:fungi ratio²⁶. In the present study, significantly positive relationships were observed between fungi, Gram-negative bacteria and PE (Table 1),

suggesting that fungi and Gram-negative bacteria play an important role in PE. This result also confirms previous observations that Gram-negative bacteria or fungi may be particularly important in metabolizing soil SOM^{37,39,41}.

In conclusion, litter input decreases the temperature sensitivity of SOM decomposition. This finding suggests that, in the context of global warming, stimulation of increasing temperature on SOM decomposition is likely to be decreased by increased litter input under elevated atmospheric CO₂ in the subtropical China. Litter addition induces PE, but PE magnitude decreases with elevated temperature. Thus, acceleration of global warming on SOM decomposition may be partly compensated by the decrease in PE with increasing temperature in subtropical forest ecosystems. Significant correlations between PE and the concentrations of Gram-negative bacterial and fungal PLFAs indicate that Gram-negative bacteria and fungi may have important functions in metabolizing SOM and PE. To better understand and model influence of global change on SOC cycle, an accurate evaluation of the sources of primed C from recent and old SOM pools is required in future.

Material and Methods

Study sites and soil sample preparation. The soil samples were collected from a CF and an evergreen BF at Huitong National Research Station of Forest Ecosystem (26°40'N, 109°26'E) in Huitong County, Hunan Province, China. The mean annual temperature (1998 to 2014) was 16.5 °C and the mean annual rainfall was 1200 mm to 1400 mm. The soil derived from shale is classified as Ultisols according to the second edition of U.S. Soil Taxonomy. The CF is a monoculture plantation of *Cunninghamia lanceolata* (about 26 years old), and the BF is dominated by *Cyclobalanopsis glauca*, *Machilus pauhoi*, and *Liquidam bar formosana*. Mineral soil was randomly sampled from 0 cm to 10 cm depth after removing organic matter layer. Fresh soil was immediately transported to the laboratory and sieved through a 2-mm mesh sieve. The remaining organic residues and stones were carefully removed by hand from the soils. The physicochemical properties of the two soils are shown in Table 3.

Experiment design and CO₂ measurement. All soil samples were pre-incubated for one week before incubation, and then divided into two temperature groups (20 and 30 °C). A trace technique was used to label tree seedling in a growth chamber^{42,43}. The seedlings were continuously labeled for three months with ¹³CO₂ gas with an abundance of 99.9 atom%. In this incubation experiment, we used the needles that newly emerged during labeling to insure uniform ¹³C distribution among the structural and metabolic litter components. In order to test whether the needles were uniformly labeled, ground needles were extracted with hot-water (80 °C) for 16 h with three replicates, and then the δ¹³C of extract and residues were measured. The δ¹³C of the extract and residues were 1330 ± 14‰ (mean ± SD) and 1313 ± 17‰, respectively. The δ¹³C of labeled needles was 1318‰. The C, N, P, and lignin concentrations in the needles were 477.1, 19.2, 1.51, and 312.6 g·kg⁻¹, respectively. Each temperature group was split into four amendment treatments: (1) CF soil without litter (CF), (2) BF soil without litter (BF), (3) CF soil with litter (CFL), and (4) BF soil with litter (BFL). In the treatments with litter, the amount of added litter C was equal to 5% of the SOC content. Ground litter (<250 μm) was homogeneously mixed with soil. The soil samples were incubated in 500 mL Mason jars (250 g fresh soil in each Mason jar) at two temperatures under aerobic conditions. Each treatment had three replicates (i.e., three Mason jars). Meanwhile, three Mason jars without soil samples were also incubated at each temperature. Vials (50 mL) without lids that contain 20 mL of 0.2 M NaOH solution were placed to trap respired CO₂ in each Mason jar. Vials containing NaOH solution were taken after 1, 2, 6, 13, 21, 30, and 42 days and replaced by new NaOH vials. At the same time Mason jars were flushed with compressed air without CO₂ for 20 min to replenish O₂. Soil water content was measured by weighing each Mason jar every three days during the first 14 days and once a week after that, and deionized water was added to maintain moisture at 60% of field capacity.

To determine the δ¹³C of released CO₂, 10 mL of NaOH solution was collected from the glass vial containing 20 mL of NaOH solution on each collection date. The δ¹³C of CO₂ in the NaOH solution was measured using a stable isotope-ratio mass spectrometer⁴⁴. The remaining 10 mL of NaOH solution was used to determine the amount of released CO₂ using alkali-trapping techniques. The amount of CO₂ was determined by titration with 0.1 M HCl to pH 8.3 after precipitation of carbonates with BaCl₂, and calculated as the difference of the evolved CO₂ from the Mason jars with and without soil.

Partitioning CO₂-C from litter and SOM sources. The amount of CO₂-C measured by titration is the sum of litter and SOM decomposition. The application of ¹³C labeled litter allowed the partitioning of the total respired C into CO₂ derived from litter and SOM. We assumed that isotopic fractionation of litter was negligible during litter decomposing. A mass balance equation was used to separate the amount of CO₂-C derived from litter and SOM decomposition⁴⁰:

$$C_L = C_T \times (\delta_T - \delta_S) / (\delta_L - \delta_S) \quad (1)$$

$$C_S = C_T \times (\delta_L - \delta_T) / (\delta_L - \delta_S) \quad (2)$$

In Equations (1) and (2), C_T (C_T = C_L + C_S) is the total amount of CO₂-C during the considered time period, and δ_T is the corresponding value of δ¹³C. C_L is the amount of C derived from the added litter, and δ_L is δ¹³C of the litter. C_S is the amount of C derived from SOM, and δ_S is δ¹³C of SOM.

Calculations of temperature sensitivity and priming effect. The temperature sensitivity (Q₁₀) of litter and SOM decomposition was calculated by the following equation⁹:

$$Q_{10} = R_{30} / R_{20} \quad (3)$$

where R₂₀ and R₃₀ are respired CO₂ rate at 20 and 30 °C, respectively.

The relative priming effect (PE, %) induced by the added litter was calculated by comparing the amount of CO₂ derived from native SOM in the litter-containing soil samples with the amount of CO₂ derived from native SOM in no-litter added soil samples³⁴:

$$PE(\%) = (CO_{2\text{treatment}} - CO_{2\text{control}})/CO_{2\text{control}} \times 100 \quad (4)$$

where CO_{2treatment} is the accumulated amount of CO₂ derived from SOM in the treatments with litter addition, and CO_{2control} is the amount of CO₂ derived from SOM without litter addition.

PLFA determination. Soil microbial community composition was determined by PLFAs as biomarkers for different microbial groups. At the end of incubation, part of the soil was taken and immediately freeze-dried for PLFA analysis. Lipid extraction and PLFA analyses were performed based on the method described by White and Ringelberg⁴⁴. Freeze-dried soil (5 g) was extracted using chloroform:methanol:phosphate buffer (1:2:0.8). The extracted PLFAs were purified on silica columns with chloroform, acetone, and methanol, amended with methyl-nonadecanoate as internal standard for quantifying the PLFAs, and converted to fatty acid methyl esters (FAMES) through alkaline methanolysis. The concentration and isotopic composition of individual FAMES was analyzed by gas chromatography–tandem mass spectrometry (Thermo Fisher). Qualitative standard mixes (37 Comp. FAME Mix and Bacterial Acid Methyl Esters CP Mix, Sigma-Aldrich) were used to identify the peak. Total bacterial content was calculated by summing i15:0, a15:0, 15:0, i16:0, 16:1ω7c, 16:1ω9c, 16:0, a17:0, i17:0, cy17:0, 17:0, 18:0, cy19:0, and 20:0 PLFAs⁴⁵. Short- or odd-chain saturated PLFA (15:0, 16:0, 17:0, 18:0, and 20:0) were considered as common bacteria⁴⁶, and PLFAs (i15:0, a15:0, i16:0, i17:0, and a17:0) were used as markers for the Gram-positive bacteria; PLFAs (16:1ω7c, 16:1ω9c, cy17:0, and cy19:0) were used as markers for the Gram-negative bacteria²⁵. The 18:1ω9c, 18:1ω7c, and 18:2ω9,12c PLFAs were used as markers for fungi⁴⁵. Mid-chain branched saturated PLFAs (10Me16:0, 10Me17:0, 10Me18:0) were associated with actinomycetes⁴⁶.

The δ¹³C values of individual PLFAs were determined using isotope ratio mass spectrometry, as described by Moore–Kucera and Dick²⁵. The concentration (mg C kg⁻¹ soil) of litter-derived C within PLFAs was calculated using the equations described by Nottingham *et al.*⁴¹

$$Pi = (\delta^{13}C_i - \delta^{13}C_c)/(\delta^{13}C_i - \delta^{13}C_l) \quad (5)$$

where δ¹³C_i is the δ¹³C enrichments (‰) of individual PLFA in the soils with leaf-litter at the end of incubation; δ¹³C_c is the δ¹³C enrichments (‰) of individual PLFA in the control soils; and δ¹³C_l is the δ¹³C of the labeled leaf litters (‰). The total labeled leaf-litter-derived C in each PLFA was calculated by multiplying each Pi by individual PLFA abundances.

Statistical analysis. The data were statistically analyzed using SPSS version 19.0 for Windows. The effect of litter addition on soil PLFA concentrations and the effect of temperature on the concentration of litter-derived C within different PLFA groups were tested by t-test after ANOVA. Significant differences in litter decomposition, Q₁₀ values of SOM decomposition, and cumulative CO₂-C from SOM among different treatments were tested by Tukey's HSD after repeated measure ANOVA. ANOVA followed by Tukey HSD was performed to test significant differences in the concentration of litter-derived C within microbial groups and PE. Pearson correlation was performed to test the relationship between the PE and concentrations of microbial groups. *P* < 0.05 was considered significant.

References

- Lal, R. Carbon sequestration. *Philosophical Transactions of the Royal Society B* **363**, 815–830 (2008).
- Davidson, E. A. & Janssens, I. A. Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature* **440**, 165–173 (2006).
- Wang, Q., He, T., Wang, S. & Liu, L. Carbon input manipulation affects soil respiration and microbial community composition in a subtropical coniferous forest. *Agricultural and Forest Meteorology* **178–179**, 152–160 (2013).
- IPCC. Climate Change 2007: The Physical Science Basis, Contribution of Working Group-I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change Cambridge University Press, Cambridge (2007).
- Bond-Lamberty, B. & Thompson, A. Temperature-associated increase in the global soil respiration record. *Nature* **464**, 579–582 (2010).
- Qiao, N. *et al.* Soil Biology & Biochemistry Labile carbon retention compensates for CO₂ released by priming in forest soils. *Global Change Biology* **20**, 1943–1954 (2014).
- Cox, P. M., Betts, R. A., Jones, C. D., Spall, S. A. & Totterdell, I. J. Acceleration of global warming due to carbon-cycle feedbacks in a coupled climate model. *Nature* **408**, 184–187 (2000).
- Knorr, W., Prentice, I. C., House, J. I. & Holland, E. A. Long-term sensitivity of soil carbon turnover to warming. *Nature* **433**, 298–301 (2005).
- Thiessen, S., Gleixner, G., Wutzler, T. & Reichstein, M. Both priming and temperature sensitivity of soil organic matter decomposition depend on microbial biomass—An incubation study. *Soil Biology & Biochemistry* **57**, 739–748 (2013).
- Gershenson, A., Bader, N. E. & Cheng, W. Effects of substrate availability on the temperature sensitivity of soil organic matter decomposition. *Global Change Biology* **15**, 176–183 (2009).
- Fissore, C., Giardina, C. P. & Kolka, R. K. Reduced substrate supply limits the temperature response of soil organic carbon decomposition. *Soil Biology & Biochemistry* **67**, 306–311 (2013).
- Zhu, B. & Cheng, W. X. Rhizosphere priming effect increases the temperature sensitivity of soil organic matter decomposition. *Global Change Biology* **17**, 2172–2183 (2011).
- Pang, X., Zhu, B., Lu, X. & Cheng, W. Labile substrate availability controls temperature sensitivity of organic carbon decomposition at different soil depths. *Biogeochemistry* **126**, 85–98 (2015).
- Kuzyakov, Y., Friedel, J. K. & Stahr, K. Review of mechanisms and quantification of priming effects. *Soil Biology & Biochemistry* **32**, 1485–1498 (2000).
- García-Pausas, J. & Paterson, E. Microbial community abundance and structure are determinants of soil organic matter mineralisation in the presence of labile carbon. *Soil Biology & Biochemistry* **43**, 1705–1713 (2011).
- Zhang, W., Wang, X. & Wang, S. Addition of external organic carbon and native soil organic carbon decomposition: a meta-analysis. *Plos One* **8**, e54779 (2013).

17. Kuzyakov, Y. Priming effects: Interactions between living and dead organic matter. *Soil Biology & Biochemistry* **42**, 1363–1371 (2010).
18. Fang, C. M. Similar response of labile and resistant soil organic matter pools to changes in temperature. *Nature* **433**, 57–59 (2005).
19. Chen, R. *et al.* Soil C and N availability determine the priming effect: microbial N mining and stoichiometric decomposition theories. *Global Change Biology* **20**, 2356–2367 (2014).
20. Wang, Q., Wang, Y., Wang, S., He, T. & Liu, L. Fresh carbon and nitrogen inputs alter organic carbon mineralization and microbial community in forest deep soil layers. *Soil Biology & Biochemistry* **72**, 145–151 (2014).
21. Nottingham, A. T., Turner, B. L., Stoot, A. W. & Tanner, E. V. J. Nitrogen and phosphorus constrain labile and stable carbon turnover in lowland tropical forest soils. *Soil Biology & Biochemistry* **80**, 26–33 (2015).
22. Bader, N. E. & Cheng, W. X. Rhizosphere priming effect of *Populus fremontii* obscures the temperature sensitivity of soil organic carbon respiration. *Soil Biology & Biochemistry* **39**, 600–606 (2007).
23. Kuzyakov, Y., Hill, P. W. & Jones, D. L. Root exudate components change litter decomposition in a simulated rhizosphere depending on temperature. *Plant and Soil* **290**, 293–305 (2007).
24. Karhu, K. *et al.* Temperature sensitivity of soil respiration rates enhanced by microbial community response. *Nature* **513**, 81–84 (2014).
25. Moore-Kucera, J. & Dick, R. P. Application of ¹³C-labeled litter and root materials for *in situ* decomposition studies using phospholipid fatty acids. *Soil Biology & Biochemistry* **40**, 2485–2493 (2008).
26. Bell, J. M., Smith, J. L., Bailey, V. L. & Bolton, H. Priming effect and C storage in semi-arid no-till spring crop rotations. *Biology and Fertility of Soils* **37**, 237–244 (2003).
27. Streit, K. *et al.* Soil warming alters microbial substrate use in alpine soils. *Global Change Biology* **20**, 1327–1338 (2014).
28. Blagodatskaya, E. V. & Kuzyakov, Y. Mechanisms of real and apparent priming effects and their dependence on soil microbial biomass and community structure: critical review. *Biology and Fertility of Soils* **45**, 115–131 (2008).
29. Cleveland, C., Nemergut, D., Schmidt, S. & Townsend, A. Increases in soil respiration following labile carbon additions linked to rapid shifts in soil microbial community composition. *Biogeochemistry* **82**(3), 229–240 (2007).
30. Moore-Kucera, J. & Dick, R. P. A pulse-chase method to ¹³Carbon-label Douglas-fir seedlings for decomposition studies. *Soil Science* **173**, 46–53 (2008).
31. Lin, J., Zhu, B. & Cheng, W. Decadally cycling soil carbon is more sensitive to warming than faster-cycling soil carbon. *Global Change Biology* **21**, 4602–4612 (2015).
32. Wetterstedt, J. Å., Persson, T. & Ågren, G. I. Temperature sensitivity and substrate quality in soil organic matter decomposition: results of an incubation study with three substrates. *Global Change Biology* **16**, 1806–1819 (2010).
33. Giardina, C. P. & Ryan, M. G. Evidence that decomposition rates of organic carbon in mineral soil do not vary with temperature. *Nature* **404**, 858–861 (2000).
34. Hamdi, S., Moyano, F., Sall, S., Bernoux, M. & Chevallier, T. Synthesis analysis of the temperature sensitivity of soil respiration from laboratory studies in relation to incubation methods and soil conditions. *Soil Biology & Biochemistry* **58**, 115–126 (2013).
35. Melillo, J. M. *et al.* Soil Warming and Carbon-Cycle Feedbacks to the Climate System. *Science* **298**, 2173–2176 (2002).
36. Hamer, U. & Marschner, B. Priming effects in different soil types induced by fructose, alanine, oxalic acid and catechol addition. *Soil Biology & Biochemistry* **37**, 445–454 (2005).
37. Wang, H. *et al.* The priming effect of soluble carbon inputs in organic and mineral soils from a temperate forest. *Oecologia* **178**, 1239–1250 (2015).
38. Fontaine, S., Barot, S., Fontaine, S. & Barot, S. Size and functional diversity of microbe populations control plant persistence and long-term soil carbon accumulation. *Ecology Letters* **8**, 1075–1087 (2005).
39. Fontaine, S. *et al.* Fungi mediate long term sequestration of carbon and nitrogen in soil through their priming effect. *Soil Biology & Biochemistry* **43**, 86–96 (2011).
40. Dimassi, B. *et al.* Effect of nutrients availability and long-term tillage on priming effect and soil C mineralization. *Soil Biology & Biochemistry* **78**, 332–329 (2014).
41. Kramer, C. & Gleixner, G. Variable use of plant- and soil-derived carbon by microorganisms in agricultural soils. *Soil Biology & Biochemistry* **38**, 3267–3278 (2006).
42. Blagodatskaya, E., Yuyukina, T., Blagodatsky, S. & Kuzyakov, Y. Three-source-partitioning of microbial biomass and of CO₂ efflux from soil to evaluate mechanisms of priming effects. *Soil Biology & Biochemistry* **43**, 778–786 (2011).
43. Nottingham, A. T., Griffiths, H., Chamberlain, P. M., Stoot, A. W. & Tanner, E. V. J. Soil priming by sugar and leaf-litter substrates: A link to microbial groups. *Applied Soil Ecology* **42**, 183–190 (2009).
44. White, D. C. & Ringelberg, D. B. Signature Lipid Biomarker Analysis. In Burlage, R. S., Atlas, R., Stahl, D., Geesey, G. & Sayler, G. (Eds.) *Techniques in Microbial Ecology* Oxford University Press, New York, pp. 255–272 (1998).
45. Hill, G. T. *et al.* Methods for assessing the composition and diversity of soil microbial communities. *Applied Soil Ecology* **15**, 25–36 (2000).
46. Zelles, L. Fatty acid patterns of phospholipids and lipopolysaccharides in the characterisation of microbial communities in soil: a review. *Biology and Fertility of Soils* **29**, 111–129 (1999).

Acknowledgements

This research was supported by the National Key Research and Development Program of China (Grant no. 2016YFA0600801), and the National Natural Science Foundation of China (grant no. 31570466). We are grateful to Xiuyong Zhang and Ke Huang for helping preparation of incubation soil samples.

Author Contributions

Q.W. conceived and designed this experiment and wrote this manuscript. T.H. and J.L. performed the experiment and measured all soil variables.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Wang, Q. *et al.* Litter input decreased the response of soil organic matter decomposition to warming in two subtropical forest soils. *Sci. Rep.* **6**, 33814; doi: 10.1038/srep33814 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016