

RESEARCH ARTICLE

Open Access



Common variants in *LEPR*, *IL6*, *AMD1*, and *NAMPT* do not associate with risk of juvenile and childhood obesity in Danes: a case–control study

Mette Hollensted^{1†}, Tarunveer S Ahluwalia^{1,2,3*†}, Christian Theil Have¹, Niels Grarup¹, Cilius Esmann Fonvig^{1,4}, Tenna Ruest Haarmark Nielsen^{1,4}, Cæcilie Trier^{1,4}, Lavinia Paternoster⁵, Oluf Pedersen¹, Jens-Christian Holm⁵, Thorkild I A Sørensen^{1,5,6} and Torben Hansen¹

Abstract

Background: Childhood obesity is a highly heritable disorder, for which the underlying genetic architecture is largely unknown. Four common variants involved in inflammatory-adipokine triggering (*IL6* rs2069845, *LEPR* rs1137100, *NAMPT* rs3801266, and *AMD1* rs2796749) have recently been associated with obesity and related traits in Indian children. The current study aimed to examine the effect of these variants on risk of childhood/juvenile onset obesity and on obesity-related quantitative traits in two Danish cohorts.

Methods: Genotype information was obtained for 1461 young Caucasian men from the Genetics of Overweight Young Adults (GOYA) study (overweight/obese: 739 and normal weight: 722) and the Danish Childhood Obesity Biobank (TDCOB; overweight/obese: 1022 and normal weight: 650).

Overweight/obesity was defined as having a body mass index (BMI) ≥ 25 kg/m²; among children and youths, this cut-off was defined using age and sex-specific cut-offs corresponding to an adult body mass index ≥ 25 kg/m². Risk of obesity was assessed using a logistic regression model whereas obesity-related quantitative measures were analyzed using a general linear model (based on z-scores) stratifying on the case status and adjusting for age and gender. Meta-analyses were performed using the fixed effects model.

Results: No statistically significant association with childhood/juvenile obesity was found for any of the four gene variants among the individual or combined analyses (rs2069845 OR: 0.94 CI: 0.85–1.04; rs1137100 OR: 1.01 CI: 0.90–1.14; rs3801266: 0.96 CI: 0.84–1.10; rs2796749 OR: 1.02 CI: 0.90–1.15; $p > 0.05$). However, among normal weight children and juvenile men, the *LEPR* rs1137100 A-allele significantly associated with lower BMI ($\beta = -0.12$, $p = 0.0026$).

(Continued on next page)

* Correspondence: veertarun@gmail.com

†Equal contributors

¹The Novo Nordisk Foundation Center for Basic Metabolic Research, Section of Metabolic Genetics, Faculty of Health and Medical Sciences, University of Copenhagen, DIKU Building, Universitetsparken 1, 2100 Copenhagen, Denmark

²COPSAC, Copenhagen Prospective Studies on Asthma in Childhood, Herlev and Gentofte Hospital, University of Copenhagen, Ledreborg Allé 34, DK-2820 Copenhagen, Denmark

Full list of author information is available at the end of the article

(Continued from previous page)

Conclusions: The *IL6*, *LEPR*, *NAMPT*, and *AMD1* gene variants previously found to associate among Indian children did not associate with risk of obesity or obesity-related quantitative measures among Caucasian children and juvenile men from Denmark.

Keywords: Childhood obesity, Juvenile obesity, Single nucleotide polymorphism, Case-control study, Body mass index, BMI, Obesity

Background

The burden of childhood obesity has increased dramatically within the past two decades and as of today one in every three children (6–9 years old) within the European Union is overweight or obese [1]. Childhood obesity is associated with numerous related complications and serious diseases later in life, including cardiovascular disease [2], type 2 diabetes, several cancers and rheumatologic disorders, as well as social stigmatization and psychological health problems [3, 4]. Although the recent increase in the prevalence of childhood obesity is most likely a result of environmental changes, a genetic component contributing to the risk of obesity is also evident from twin studies [5, 6]. To date, genome-wide association studies (GWAS) have identified well over 100 adiposity susceptibility loci [7, 8], however, only a limited number of these have been implicated in childhood obesity [9–11].

Recent studies suggest that white adipose tissue inflammation has an emerging role in obesity pathophysiology [12]. White adipose tissue plays an important role in the production of bioactive adipokines that possess immune-regulatory properties in addition to regulation of systemic metabolism. A recent study explored the role of genes encoding adipokines and inflammatory markers (*interleukin 6*, *IL6*; *leptin receptor*, *LEPR*, and *nicotinamide phosphoribosyltransferase*, *NAMPT*) in 2,261 normal weight and 907 overweight/obese children of Indian origin (age range: 11–17 years). Overweight/obesity was defined using age and sex-specific cut-offs corresponding to a BMI ≥ 25 kg/m² in adults [13]. When employing a two stage study design, all three variants (*IL6*, *LEPR*, and *NAMPT*) were found to associate with an increased prevalence of childhood obesity (OR_{combined} >1.35) and related quantitative measures (BMI, weight, waist and hip circumference, $p \leq 2.0 \times 10^{-3}$) [14].

A follow-up study in the Indian cohort reported the *AMD1* gene (*adenosylmethionine decarboxylase 1*) single nucleotide polymorphism (SNP) rs2796749 G allele to be associated with childhood obesity (OR = 1.35; $p = 1.9 \times 10^{-6}$) and related quantitative measures (BMI, weight, height, waist and hip circumference, as well as plasma leptin levels ($p \leq 6.7 \times 10^{-4}$) [15]. *AMD1* encodes an important intermediate enzyme involved in the homocysteine pathway.

Previous studies examining the effect of *IL6*, *LEPR*, *NAMPT*, and *AMD1* variants on risk of childhood obesity in Caucasians are very limited and have generally been small and statistically underpowered [16, 17]. Thus, the aim of the present study was to replicate the findings by Tabassum et al. [14, 15] by examining the effect of four SNPs: *IL6* rs2069845, *LEPR* rs1137100, *NAMPT* rs3801266, and *AMD1* rs2796749 on the risk of childhood/juvenile obesity and anthropometric traits (BMI, height, and weight) in the two Danish study populations comprising cases of childhood or juvenile obesity and normal weight controls.

Methods

The present study was conducted in accordance with the Helsinki declaration and approved by the Danish Data Protection Agency and by the regional scientific ethics committees. Prior to this study, all participants aged 18 or parents of younger children signed an informed written consent.

Study participants

GOYA male

In the recruitment of these participants, a case-cohort sampling design was employed in which cases were defined as the individuals with the upper-most extreme BMI in the cohort, and controls were randomly selected from the same cohort [18].

During 1943–1977, 362200 Caucasian men with a mean age of ~20 years were examined at draft boards in the area of Copenhagen, Denmark. At the draft board examinations, standing height (without shoes) and weight (in underwear only) were measured. Obesity was defined as 35 % overweight relative to a local standard in use at the time (mid 1970's). This definition corresponds to a BMI ≥ 31.0 kg/m², which proved to be above the 99th percentile. All obese men ($n = 1930$) and a randomly selected control group ($n = 3601$), comprising one percent of all participants, were sampled on the basis of the records from the draft board examinations.

In 1991–1994, at the mean age of 46 years, all obese men and 50 % of the randomly selected young men from the study population were invited for a follow-up survey (753 obese and 879 control men attended), at which time non-fasting venous blood samples were obtained

[19]. By using a sampling fraction of 0.5 % (50 % of 1 %), the included controls effectively represent approximately 158000 men among whom the case group was the most obese. Criteria for invitation to follow-up surveys and participation have been described elsewhere [20].

In the current study, while replicating the original studies by Tabassum et al. [14, 15], we adopted the BMI cut-off corresponding to overweight/obesity defined as BMI ≥ 25 kg/m², and 1461 ($n_{\text{overweight/obese}}$: 739 and $n_{\text{normal weight}}$: 722) men had complete genetic and phenotypic information representing the GOYA (male) cohort.

The Danish Childhood Obesity Biobank (TDCOB)

All children were recruited through The Children's Obesity Clinic, Copenhagen University Hospital Holbæk, Denmark. Obese children, defined by a BMI above the 90th percentile for age and sex according to Danish standard BMI charts [21], aged 3 to 18 years were treated using the Children's Obesity Clinic's Treatment Protocol [22]. In addition, an age and gender matched control group was recruited from schools and high schools in the surrounding area.

Height and weight were measured to the nearest 0.1 cm and 0.1 kg, respectively, using an integrated calibrated weight and stadiometer (ADE, Modell MZ10023, Germany) while wearing light indoor clothing. Danish standard BMI charts [21] were used to calculate BMI (kg/m²) into BMI z-scores. Prior to analysis, children with syndromes (e.g. Turner syndrome, Down's syndrome) or other clinical diagnoses known to associate with obesity were excluded.

A total of 1672 children ($n_{\text{overweight/obese}}$: 1022 and $n_{\text{normal weight}}$: 650), from the TDCOB cohort participated in the current study.

Clinical characteristics for both study populations are listed in Table 1.

Genotyping

GOYA (Male)

Genotyping on the Illumina Human 610 k quad bead chip was performed at the Centre National de Génotypage, Evry, France. Genotypes were called using the Gen Call algorithm from Genome Studio package. In brief, the data quality control included 1) exclusion of SNPs with a minor allele frequency <1% and >5 % missing genotypes, 2) exclusion of those failing Hardy Weinberg equilibrium (HWE, $p < 10^{-7}$), and 3) including only European individuals (clustering with the HapMap CEU population (Utah residents with Northern and western European ancestry)). Further checks on heterozygosity, sample duplicates, individual relatedness, or with sex discrepancy were made. Imputation was carried out to HapMap release 22 (CEU individuals) using Mach 1.0, Markov Chain Haplotyping [23].

Table 1 Clinical characteristics of participants from the GOYA (male) and the TDCOB study

Characteristics	GOYA	TDCOB
N_{total} (%men)	1461 (100)	1672 (41.9)
$N_{\text{overweight/obese}}/N_{\text{normal weight}}$	739/722	1022/650
Age _{overweight/obese}	19.9 ± 1.9	11.7 ± 3.2
Age _{normal weight}	19.9 ± 1.7	12.7 ± 3.3
BMI _{overweight/obese}	32.6 ± 3.0	27.2 ± 5.8
zBMI _{overweight/obese}	6.4 ± 1.6	2.8 ± 0.70
BMI _{normal weight}	21.1 ± 1.7	18.4 ± 2.7
zBMI _{normal weight}	0.04 ± 1.0	0.11 ± 0.89
Height _{overweight/obese}	177.3 ± 6.1	153.3 ± 16.3
zHeight _{overweight/obese}	0.10 ± 0.9	0.96 ± 0.98
Height _{normal weight}	176.7 ± 6.6	155.4 ± 16.3
zHeight _{normal weight}	0.007 ± 1.0	0.66 ± 0.99
Weight _{overweight/obese}	102.7 ± 11.5	66.5 ± 25.6
zWeight _{overweight/obese}	5.28 ± 1.6	4.0 ± 0.52
Weight _{normal weight}	66.1 ± 6.9	45.9 ± 14.5
zWeight _{normal weight}	0.011 ± 1.0	3.86 ± 0.51

Data are means ± standard deviations. Individuals were classified as being overweight/obese or normal weight using age and sex-specific cut-offs corresponding to a BMI ≥ 25 kg/m² in adults

Finally, we extracted genotypes for two SNPs (*LEPR* rs1137100 and *AMD1* rs2796749) available on the SNP array, whereas the other two (*IL6* rs2069845, *NAMPT* rs3801266) were imputed with an $r^2 \geq 0.8$.

TDCOB

The SNPs were genotyped on Illumina Infinium Human-CoreExome Beadchip platform at the Novo Nordisk Foundation Centre for Basic Metabolic Research's laboratory at Symbion, Copenhagen, Denmark. Genotypes were called with Illumina standard pipeline in Genome Studio. The data quality control involved 1) HWE filter: variants with $p_{\text{HWE}} > 0.005$ were retained, 2) ethnicity checks using principal component analyses were performed, and individuals identified as ethnic outliers or with extreme positive or negative inbreeding coefficients were removed. Checks on heterozygosity, sample duplicates, or with sex discrepancy were made. Additional genotypes were imputed into 1000 genomes phase 1 [24] using impute2 [25]. All variants examined in the current study were imputed with high quality (info >0.95). In case of siblings, only the sibling with the best overall genotype call-rate was retained in the study (regardless of gender).

Statistical analyses

The quantitative measures were transformed to z-scores (standard deviations (sd) from the study reference group (controls) mean) using the equation z (z-score) = $(x - \mu)/\sigma$ where x is the raw measure, μ is the mean value for the

study reference group, and σ is the sd for that reference group. Logistic regression was used to compare allele frequencies in the dichotomized case–control groups whereas a general linear model was applied to test quantitative traits using an additive genetic model while adjusting for age and gender. For these analyses, we adopted both the age and gender dependent BMI cut-off corresponding to an adult BMI ≥ 25 kg/m² provided by Cole et al. [13] as per the studies by Tabassum et al. [14, 15], as well as a BMI cut-off of 31 kg/m², taking into account the original GOYA study design. No transformation of traits was performed as they followed normal distributions within case or control groups. Linear regression analyses were performed separately in the case and cohort groups owing to the trait distribution as per study design. A *p*-value of less than 0.05 was considered statistically significant. Association tests were conducted on the imputed genotype likelihoods using the frequentist score method of the SNPtest tool (version 2.5) with a standard additive model among TDCOB children and using R statistical software (version 2.12.1) [26] in the GOYA males.

We combined the individual study results (obesity risk and quantitative trait (zBMI, zWeight, zHeight) association results) to perform a fixed effects (inverse variance) meta-analysis where weights are proportional to the squared standard error of the effect estimates, as implemented in the METAL software [27]. The heterogeneity in effect sizes across the two participating cohorts was evaluated using the chi-square test for heterogeneity (I^2) also through METAL.

External data look-up

External data from published GWAS meta-analyses on BMI was contributed by GIANT consortia investigators and was freely accessible [28].

Results

Association with obesity

We found no association between the four genetic loci (*IL6* rs2069845, *LEPR* rs1137100, *NAMPT* rs3801266, and *AMD1* rs2796749) and prevalence of obesity (corresponding to an adult BMI cut-off of 25 kg/m²) among Danish young men or children (Table 2). We also checked for associations taking into account the original GOYA study design (corresponding to a BMI cut-off of 31 kg/m²) but the results remained insignificant ($p > 0.05$; data not shown).

Furthermore, no statistically significant association with obesity was found for either of the variants in the combined analyses (rs2069845: OR [95 % CI] = 0.94 [0.85–1.04], $p_{\text{combined}} = 0.24$; rs1137100: OR [95 % CI] = 1.01 [0.90–1.14], $p_{\text{combined}} = 0.46$; rs3801266: OR [95 % CI] = 0.96 [0.84–1.10], $p_{\text{combined}} = 0.58$; and rs2796749: OR [95 % CI] = 1.02 [0.90–1.15], $p_{\text{combined}} = 0.68$; Table 2).

Association with obesity-related quantitative traits

We then examined the four SNPs for association with obesity-related traits (zBMI, zHeight, and zWeight) in the normal weight and overweight/obese groups separately.

Among the normal weight juvenile men, the *LEPR* rs1137100 A-allele significantly associated with zBMI ($\beta = -0.17$, $p_{\text{GOYA}} = 0.0036$), and this association remained significant in the combined analyses ($\beta = -0.12$, $p_{\text{combined}} = 0.0026$; Table 3).

Furthermore, we observed some cohort specific associations that did not reach statistical significance in the combined analyses. Among GOYA, the *LEPR* SNP associated with reduced zWeight ($\beta = -0.12$, $p_{\text{GOYA}} = 0.046$) whereas the *AMD1* rs2796749 associated with zHeight ($\beta = -0.14$, $p_{\text{GOYA}} = 0.017$; Table 3). While taking the original study design for GOYA into account (BMI cut-off of 31 kg/m²), we found similar associations for *LEPR* rs1137100 with zBMI and zWeight ($\beta_{\text{GOYA_zBMI}} = -0.14$, $p_{\text{GOYA_zBMI}} = 0.014$; and $\beta_{\text{GOYA_zWeight}} = -0.11$, $p_{\text{GOYA_zWeight}} = 0.045$) whereas non-significant for *AMD1* rs2796749 and zHeight ($p > 0.05$) (data not shown).

In TDCOB, *AMD1* rs2796749 was found to associate with zBMI ($\beta = -0.16$, $p_{\text{TDCOB}} = 0.0077$) while *IL6* rs2069845 modestly associated with zHeight ($\beta = -0.05$, $p = 0.047$; Table 3).

Among the overweight/obese individuals, no significant association with zBMI, zWeight or zHeight was identified in either of the examined cohorts or in the combined analysis (Table 4). A *p*-value corresponding to each chi-square test for heterogeneity has also been indicated (P_{HET}) in the results tables (Tables 2, 3, and 4).

Discussion

Common variations in four genes involved in inflammatory and homocysteine pathways (*IL6*, *LEPR*, *NAMPT*, and *AMD1*) have recently been shown to associate with childhood obesity among children of Indian origin [14, 15]. We aimed to replicate these findings, however, we found none of the variants to exert a strong effect on risk of obesity among Danish children and juveniles (OR ≤ 1.15 , as based on 95 % CI, Table 2). We did, however, observe a modest association between the *LEPR* variant and lowered BMI among non-obese Danish children and young men, albeit the directionality of these associations was opposite to what was observed among Indian children [14]. In the studies by Tabassum et al., *IL6*, *LEPR*, and *NAMPT* variants also associated with hip and waist circumference [14, 15], but unfortunately, information on these traits were not available in the present study.

LEPR is involved in the regulation of fat metabolism, and although mutations in *LEPR* robustly associate with monogenic forms of obesity [29, 30], studies focusing on complex forms of obesity report conflicting results for several variants, including Q223R, K109R and K656N, in

Table 2 Association with obesity: Case-control analysis in GOYA (male) and TDCOB study participants

SNP	Locus	Alleles (effect/other)	GOYA (n = 1461)				TDCOB (n = 1672)				Combined (n = 3133)			
			n (cases/controls)	EAf (cases/controls)	p	OR (95 % CI)	n (cases/controls)	EAf (cases/controls)	p	OR (95 % CI)	n (cases/controls)	p	OR (95 % CI)	I ² (p _{HET})
rs2069845	IL6	G/A	739/722	0.49/0.49	0.88	0.98 (0.85–1.14)	1022/650	0.47/0.49	0.13	0.88 (0.76–1.01)	1761/1372	0.24	0.94 (0.85–1.04)	0 (0.35)
rs1137100	LEPR	A/G	739/722	0.74/0.74	0.69	0.96 (0.81–1.14)	1022/650	0.75/0.74	0.42	1.05 (0.90–1.23)	1761/1372	0.46	1.01 (0.90–1.14)	0 (0.40)
rs3801266	NAMPT	T/C	739/722	0.82/0.83	0.96	0.99 (0.82–1.20)	1022/650	0.83/0.84	0.47	0.92 (0.76–1.11)	1761/1372	0.58	0.96 (0.84–1.10)	0 (0.63)
rs2796749	AMD1	G/C	739/722	0.78/0.78	0.94	1.0 (0.84–1.19)	1022/650	0.79/0.78	0.59	1.03 (0.87–1.22)	1761/1372	0.68	1.02 (0.90–1.15)	7 (0.73)

Cases and controls were defined using age and sex-specific cut-offs corresponding to a BMI ≥ 25 kg/m² in adults. p-values are for an additive genetic model and are adjusted for age
 EAf Effect allele frequency, BMI body mass index, CI confidence interval, HET heterogeneity, OR Odds Ratio, SNP single nucleotide polymorphism

Table 3 Association with quantitative traits in 1372 normal weight individuals

SNP	Locus	Alleles (effect/other)	Effect allele frequency	GOYA (n = 722)			TDCOB (n = 650)			Combined (n = 1372)			I^2 (p_{HET})
				β	SE	p	β	SE	p	β	SE	p	
zBMI													
rs2069845	<i>IL6</i>	G/A	0.49	0.04	0.05	0.34	-0.02	0.05	0.62	0.01	0.04	0.76	3 (0.31)
rs1137100	<i>LEPR</i>	A/G	0.74	-0.17	0.06	0.0036	-0.08	0.06	0.15	-0.12	0.04	0.0026	20 (0.26)
rs3801266	<i>NAMPT</i>	T/C	0.83	-0.03	0.07	0.58	0.04	0.07	0.51	0.01	0.05	0.93	0 (0.39)
rs2796749	<i>AMD1</i>	G/C	0.78	0.03	0.06	0.60	-0.16	0.06	0.0077	-0.06	0.04	0.12	80 (0.03)
zHeight													
rs2069845	<i>IL6</i>	G/A	0.49	0.02	0.05	0.65	-0.07	0.06	0.19	-0.02	0.04	0.56	36 (0.20)
rs1137100	<i>LEPR</i>	A/G	0.74	0.01	0.06	0.63	-0.08	0.06	0.17	-0.04	0.04	0.36	5 (0.30)
rs3801266	<i>NAMPT</i>	T/C	0.83	0.07	0.07	0.32	-0.12	0.08	0.12	-0.02	0.05	0.75	69 (0.07)
rs2796749	<i>AMD1</i>	G/C	0.78	-0.12	0.06	0.046	0.10	0.07	0.15	-0.02	0.05	0.64	82 (0.01)
zWeight													
rs2069845	<i>IL6</i>	G/A	0.49	0.06	0.05	0.29	-0.05	0.03	0.047	-0.03	0.02	0.19	69 (0.06)
rs1137100	<i>LEPR</i>	A/G	0.74	-0.14	0.06	0.017	-0.02	0.03	0.44	-0.05	0.03	0.08	68 (0.07)
rs3801266	<i>NAMPT</i>	T/C	0.83	0.02	0.07	0.82	-0.01	0.04	0.79	-0.01	0.03	0.89	0 (0.75)
rs2796749	<i>AMD1</i>	G/C	0.78	-0.07	0.06	0.29	-0.01	0.03	0.95	-0.01	0.03	0.66	0 (0.33)

Effect sizes (β), standard error (SE) and p -values are for an additive genetic model based on z -scores of each trait and are adjusted for age. Effect allele frequency is the mean frequency observed for the GOYA and TDCOB cohort control individuals. $P_{add} < 0.05$ is significant and is shown in bold. Values of I^2 are percentages BMI body mass index, CI confidence interval, HET heterogeneity, SE standard error, SNP single nucleotide polymorphism

adults [31–33]. Reported *LEPR* associations with obesity among children and adolescents have also been inconsistent: Mexican ($n = 128$) and Polish ($n = 142$) studies found no association between *LEPR* rs1137100 [K109R] and obesity [16, 34], whereas a small study of 136 Japanese

children found the rs1137100 A-allele to associate with increased risk of being overweight [35]. In the current study, the rs1137100 A-allele associated significantly with a lowered BMI but not with obesity. However, this result was primarily driven by the normal weight juvenile men and

Table 4 Association with quantitative traits in 1761 overweight/obese individuals

SNP	Locus	Alleles (Effect/other)	Effect allele frequency	GOYA (n = 739)			TDCOB (n = 1022)			Combined (n = 1761)			I^2 (p_{HET})
				β	SE	p	β	SE	p	β	SE	p	
zBMI													
rs2069845	<i>IL6</i>	G/A	0.48	0.02	0.09	0.78	0.01	0.03	0.90	0.01	0.03	0.84	0 (0.83)
rs1137100	<i>LEPR</i>	A/G	0.74	0.11	0.10	0.28	-0.01	0.03	0.71	0.01	0.03	0.99	20 (0.26)
rs3801266	<i>NAMPT</i>	T/C	0.83	-0.09	0.12	0.45	-0.05	0.04	0.19	-0.06	0.04	0.13	0 (0.77)
rs2796749	<i>AMD1</i>	G/C	0.78	-0.11	0.11	0.31	-0.06	0.04	0.08	-0.07	0.04	0.05	0 (0.69)
zHeight													
rs2069845	<i>IL6</i>	G/A	0.48	0.04	0.05	0.39	-0.01	0.04	0.79	0.01	0.03	0.70	0 (0.41)
rs1137100	<i>LEPR</i>	A/G	0.74	0.05	0.05	0.35	-0.05	0.05	0.28	-0.01	0.04	0.89	49 (0.16)
rs3801266	<i>NAMPT</i>	T/C	0.83	-0.04	0.06	0.54	-0.02	0.06	0.70	-0.03	0.04	0.48	0 (0.85)
rs2796749	<i>AMD1</i>	G/C	0.78	0.05	0.06	0.43	-0.04	0.05	0.51	0.01	0.04	0.94	2 (0.31)
zWeight													
rs2069845	<i>IL6</i>	G/A	0.48	0.07	0.09	0.41	-0.02	0.02	0.46	-0.01	0.02	0.61	0 (0.33)
rs1137100	<i>LEPR</i>	A/G	0.74	0.13	0.10	0.19	0.02	0.03	0.45	0.03	0.03	0.28	11 (0.28)
rs3801266	<i>NAMPT</i>	T/C	0.83	-0.11	0.11	0.33	0.03	0.03	0.38	0.02	0.03	0.54	24 (0.25)
rs2796749	<i>AMD1</i>	G/C	0.78	-0.03	0.11	0.75	0.02	0.03	0.56	0.01	0.03	0.63	0 (0.65)

Effect sizes (β), standard error (SE) and p -values are for an additive genetic model based on z -scores of each trait and is adjusted for age. Effect allele frequency is the mean frequency observed for the GOYA and TDCOB cohort obese individuals. $P_{add} < 0.05$ is significant. Values of I^2 are percentages Add additive, BMI body mass index, CI confidence interval, HET heterogeneity, SE standard error, SNP single nucleotide polymorphism

not by the children. These inconsistent findings could imply that the effects of rs1137100 may get modified by additional genetic or environmental factors which may vary between populations. Recent studies in Europeans have suggested obesity related measures to be influenced by interactions between *LEPR* polymorphisms and a polymorphism of the tyrosine phosphatase 1B [36] and ponderal index [37] and future studies focusing on such “gene-gene” and “gene-environment” interactions are thus warranted.

In the current study, no association with obesity was identified for the *IL6* variant in the individual cohorts or in the combined analysis. Previous studies examining the putative association between variations in the promoter region of *IL6* and risk of obesity in adults have yielded contradictory results [32, 38–40], and studies among children are limited. Reports on Mexican adolescents (age range: 14–19 years) found *IL6* variants (rs1800795, rs1800796, and rs1800797) to associate with quantitative obesity-related traits and prevalence of obesity [41]. Similarly, a study on Greek primary school children ($n = 184$) reported rs1800795 to associate with obesity-related traits [42]. In the HapMap database, rs1800795 and rs1800797 (for CEU population) are both in relatively high linkage disequilibrium (LD) with rs2069845 ($r^2 > 0.83$, $D' > 0.93$) which was examined in the present study. The inconsistency between study results may be due to differences in age or gender stratification in the examined study populations, and/or it may relate to statistical power issues.

Variations in *NAMPT* have in recent studies in German and Brazilian children and adolescents not associated with risk of obesity [17, 43] and the similar was observed in the current study involving Danes. However, a French study including children and adults reported a rare *NAMPT* variant (rs10487818) associating with decreased risk of obesity [44]. Since the rare variant (rs10487818) examined among French [44] was unavailable in our study and in the Exome variant server [45], it may represent a population-specific signal.

Albeit we found no association between *AMD1* rs2796749 and risk of obesity, the *AMD1* (rs2796749) G-allele strongly correlated with a lower zBMI among the TDCOB non-obese children and borderline with the TDCOB obese children but not among the GOYA men, suggesting some protective effect of *AMD1* SNP on BMI among children but not young men. Again, it is noteworthy that the *AMD1* G-allele associated with a lower zBMI among Danish TDCOB children whereas directionality of effect among Indian children was opposite. As the first study to report an association between an *AMD1* variant and childhood obesity, Tabassum et al. identified a relatively high OR (1.35, $p = 1.9 \times 10^{-6}$) for *AMD1* rs2796749 [11], perhaps reflecting a case of “winner’s curse”, or it may indicate that the effect of *AMD1* rs2796749 is population-specific.

All of the four SNPs examined in the present study were included in both a large meta-analysis ($n = 249796$) of European-ancestry-based GWAS [46], and a more recent combined GWAS and Metabochip meta-analysis ($n = 339224$) [8]. Both studies were performed in adults, and no statistically significant association with BMI was observed for any of the variants [8, 46], suggesting that none of the four variants significantly influence the risk of obesity in adulthood and that putative effects may be exerted only in childhood. Although the GOYA study participants were sampled at ~20 years of age, the group of overweight/obese individuals already had a much increased BMI at age seven years, compared to the population sample and this deviation increased as the children grew older [47]. However, BMI data from age seven years were only available in a subset ($n \sim 172$ in each group) compared to available data at mean age 20 years ($n \sim 722$ in each group). For that reason we define the individuals with data at age 20 years as juvenile onset obese/overweight cases and controls, in the current study. Moreover, as our statistical power to detect the large ORs reported by Tabassum and colleagues was very high (>90 % for variants with risk allele frequencies >0.40, data not shown), our study design should have enabled us to identify a large effect of these variants on childhood/juvenile obesity, if it had been exerted among Danish individuals.

The identified differences between the current study results and those reported among Indian children [14, 15] may be due to the following reasons: firstly, the observed phenotype at a given BMI cutoff is known to differ between various ethnic populations, and risk factors for cardiovascular disease can be identified at a lower BMI in Asian and Indian populations compared to European [48]. This suggests that applying the same BMI cut-off of 25 kg/m² in both, the Indian and Danish populations may confound the association. Furthermore, genotype frequencies and LD patterns may differ between the populations, and environmental factors such as lifestyle may also modify the effects exerted by genetic variants. Finally, population stratification may also play a role. Genetic marker based clustering analyses have therefore been applied to exclude ethnic outliers in the current study as well by Tabassum and colleagues [14, 15] ruling out the possibility that major population stratification plays a role for the observed differences between the two studies.

Conclusions

The current study illustrates that four variants (*IL6* rs2069845, *LEPR* rs1137100, *NAMPT* rs3801266, and *AMD1* rs2796749) implicated in childhood obesity among Indians are not strongly associated with obesity among Danish children and juveniles.

Abbreviations

Add: additive; AMD1: adenosylmethionine decarboxylase 1; BMI: body mass index; CEU: Utah residents with Northern and western European ancestry; CI: confidence interval; GOYA: Genetics of Overweight Young Adults; GWAS: genome-wide association study; HET: heterogeneity; HWE: Hardy Weinberg equilibrium; IL6: interleukin 6; LD: linkage disequilibrium; LEPR: leptin receptor; MAF: minor allele frequency; NAMPT: nicotinamide phosphoribosyltransferase; OR: odds ratio; SD: standard deviation; SE: standard error; SNP: single nucleotide polymorphism; TDCOB: The Danish Childhood Obesity Biobank.

Competing interests

MH holds shares in Novozymes A/S, and TH and OP hold shares in Novo Nordisk A/S. All other authors declare that there is no competing interest associated with this manuscript.

Authors' contributions

The concept and idea regarding the epidemiological studies underlying the study populations were conceived by TIAS and JCH, and the collection and genotyping of study participants was planned and performed by TIAS, JCH, OBP, TH, MH, CEF, TRHN, and CT. SNP calling and quality control of genotype data for the GOYA cohort was performed by LP. The original hypothesis regarding the study was conceived by MH, TSA, and TH. Detailed planning of analyses and study design was performed by MH, TSA, NG, CTH, and TH. MH, TSA, OBP, TH, JCH, and TIAS contributed to the establishment of study population databases specific for this study. Statistical analyses in association studies were performed by MH, TSA, CTH, and NG. The initial draft was written by MH, and the final draft was prepared by TSA, MH, and TH. All authors revised the manuscript and contributed to the discussion. The final manuscript was read and approved by all authors.

Acknowledgements

The authors wish to thank the staff of the Copenhagen City Heart study as well as the staff at The Children's Obesity Clinic for recruitment and examination of participants. Furthermore, we wish to thank B. Andersen, G. J. Klavsen and M. Møller for laboratory assistance, G. Lademann for secretarial support, A. Nielsen, P. Sandbeck and M. Appleyard for data management, and M. Kristensen for grant management.

MH received PhD scholarship funding from TARGET (The impact of our genomes on individual treatment response in obese children, <http://metabol.ku.dk/research/research-projectsites/target/>), The Danish PhD School of Molecular Metabolism (<http://metabolism-endocrinology.dk/>) and the Copenhagen Graduate School of Health Sciences and acknowledges all grant providers. Tarunveer S. Ahluwalia received his Post-Doctoral Research grant from the Gene-diet Interactions in Obesity project (GENDINO, www.gendinob.dk) and acknowledges the same.

This study was conducted as part of the activities of the Novo Nordisk Foundation Centre for Basic Metabolic Research (NNF CBMR, <http://metabol.ku.dk/research/section-for-metabolic-genetics/>) also supported by the Danish Diabetes Academy (<http://www.danishdiabetesacademy.dk/>). Furthermore, this study was part of the research activities of the Danish Childhood Obesity Biobank (ClinicalTrials.gov ID-no.: NCT00928473) as well as related to TARGET, BIOCHILD (Genetics and systems biology of childhood obesity in India and Denmark, <http://biochild.ku.dk/>). We also thank the MRC centre for Causal Analyses in Translational Epidemiology (MRC CAITE) for the genotype data quality control carried out there. Finally, we thank all the study participants.

Author details

¹The Novo Nordisk Foundation Center for Basic Metabolic Research, Section of Metabolic Genetics, Faculty of Health and Medical Sciences, University of Copenhagen, DIKU Building, Universitetsparken 1, 2100 Copenhagen, Denmark. ²COPSAC, Copenhagen Prospective Studies on Asthma in Childhood, Herlev and Gentofte Hospital, University of Copenhagen, Ledreborg Allé 34, DK-2820 Copenhagen, Denmark. ³Steno Diabetes Center, Gentofte, Denmark. ⁴The Children's Obesity Clinic, Department of Pediatrics, Copenhagen University Hospital Holbæk, Holbæk, Denmark. ⁵MRC Integrative Epidemiology Unit, School of Social and Community Medicine, University of Bristol, Bristol, UK. ⁶Institute of Preventive Medicine, Bispebjerg and Frederiksberg Hospital, The Capital Region, Copenhagen, Denmark.

Received: 7 April 2015 Accepted: 9 November 2015

Published online: 11 November 2015

References

- Wijnhoven TM, van Raaij JM, Spinelli A, Starc G, Hassapidou M, Spiroski I, et al. WHO european childhood obesity surveillance initiative: body mass index and level of overweight among 6-9-year-old children from school year 2007/2008 to school year 2009/2010. *BMC Public Health*. 2014;14:806.
- Baker JL, Olsen LW, Sørensen TIA. Childhood body-mass index and the risk of coronary heart disease in adulthood. *N Engl J Med*. 2007;357(23):2329-37.
- World Health Organization: Why does childhood overweight and obesity matter? 2015 [http://www.who.int/dietphysicalactivity/childhood_consequences/en/]
- Organization WH. Why does childhood overweight and obesity matter? 2014.
- Elks CE, den Hoed M, Zhao JH, Sharp SJ, Wareham NJ, Loos RJ, et al. Variability in the heritability of body mass index: a systematic review and meta-regression. *Front Endocrinol (Lausanne)*. 2012;3:29.
- Silventoinen K, Rokholm B, Kaprio J, Sørensen TIA. The genetic and environmental influences on childhood obesity: a systematic review of twin and adoption studies. *Int J Obes (Lond)*. 2010;34(1):29-40.
- Garur N, Sandholt CH, Hansen T, Pedersen O. Genetic susceptibility to type 2 diabetes and obesity: from genome-wide association studies to rare variants and beyond. *Diabetologia*. 2014;57(8):1528-41.
- Locke AE, Kahali B, Berndt SI, Justice AE, Pers TH, Day FR, et al. Genetic studies of body mass index yield new insights for obesity biology. *Nature*. 2015;518(7538):197-206.
- Fernandez JR, Klimentidis YC, Dulin-Keita A, Casazza K. Genetic influences in childhood obesity: recent progress and recommendations for experimental designs. *Int J Obes (Lond)*. 2012;36(4):479-84.
- Bradfield JP, Taal HR, Timpson NJ, Scherag A, Lecoeur C, Warrington NM, et al. A genome-wide association meta-analysis identifies new childhood obesity loci. *Nat Genet*. 2012;44(5):526-31.
- den Hoed M, Ekelund U, Brage S, Grøntved A, Zhao JH, Sharp SJ, et al. Genetic susceptibility to obesity and related traits in childhood and adolescence: influence of loci identified by genome-wide association studies. *Diabetes*. 2010;59(11):2980-8.
- Exley M, Hand LE, O'Shea D, Lynch L. The interplay between the immune system and adipose in obesity. *J Endocrinol*. 2014;223(2):R41-8.
- Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ*. 2000;320(7244):1240-3.
- Tabassum R, Mahendran Y, Dwivedi OP, Chauhan G, Ghosh S, Marwaha RK, et al. Common variants of IL6, LEPR, and PBEF1 are associated with obesity in Indian children. *Diabetes*. 2012;61(3):626-31.
- Tabassum R, Jaiswal A, Chauhan G, Dwivedi OP, Ghosh S, Marwaha RK, et al. Genetic variant of AMD1 is associated with obesity in urban Indian children. *PLoS One*. 2012;7(4):e33162.
- Pyrzak B, Wisniewska A, Kucharska A, Wasik M, Demkow U. No association of LEPR Gln223Arg polymorphism with leptin, obesity or metabolic disturbances in children. *Eur J Med Res*. 2009;14 Suppl 4:201-4.
- Körner A, Böttcher Y, Enigk B, Kiess W, Stumvoll M, Kovacs P. Effects of genetic variation in the visfatin gene (PBEF1) on obesity, glucose metabolism, and blood pressure in children. *Metabolism*. 2007;56(6):772-7.
- Paternoster L, Evans DM, Aagaard Nohr E, Holst C, Gaborieau V, Brennan P, et al. Genome-Wide Population-Based Association Study of Extremely Overweight Young Adults - The GOYA Study. *PLoS One*. 2011;6(9):e24303.
- Schnohr P, Jensen G, Lange P, Scharling H, Appleyard M. The Copenhagen City Heart Study - introduction. *Eur Heart J Supplements*, Vol. 3 (Suppl H) July 2001.
- Sonne-Holm S, Sørensen TIA, Jensen G, Schnohr P. Independent effects of weight change and attained body weight on prevalence of arterial hypertension in obese and non-obese men. *BMJ*. 1989;299(6702):767-70.
- Nysom K, Molgaard C, Hutchings B, Michaelsen KF. Body mass index of 0 to 45-y-old Danes: reference values and comparison with published European reference values. *Int J Obes Relat Metab Disord*. 2001;25(2):177-84.
- Holm JC, Gamborg M, Bille DS, Grønbaek HN, Ward LC, Faerk J. Chronic care treatment of obese children and adolescents. *Int J Pediatr Obes*. 2011;6(3-4):188-96.
- Li Y, Abecasis GR. Mach 1.0: rapid haplotype reconstruction and missing genotype inference. *Am J Hum Genet*. 2006;S79:416.
- Delaneau O, Marchini J, 1000GP-Consortium. Integrating sequence and array data to create an improved 1000 Genomes Project haplotype reference panel. *Nat Commun*. 2014;5:3934.

25. Marchini J, Howie B, Myers S, McVean G, Donnelly P. A new multipoint method for genome-wide association studies by imputation of genotypes. *Nat Genet.* 2007;39(7):906–13.
26. The R project for Statistical Computing [<http://www.r-project.org/>]
27. METAL [<http://www.sph.umich.edu/csg/abecasis/Metal/>]
28. The GIANT Consortium: Genetic Investigation of ANthropometric Traits. 2015. [http://www.broadinstitute.org/collaboration/giant/index.php/GIANT_consortium]
29. Farooqi IS, Wangensteen T, Collins S, Kimber W, Matarese G, Keogh JM, et al. Clinical and molecular genetic spectrum of congenital deficiency of the leptin receptor. *N Engl J Med.* 2007;356(3):237–47.
30. Dubern B, Clement K. Leptin and leptin receptor-related monogenic obesity. *Biochimie.* 2012;94(10):2111–5.
31. Bender N, Allemann N, Marek D, Vollenweider P, Waeber G, Mooser V, et al. Association between variants of the leptin receptor gene (LEPR) and overweight: a systematic review and an analysis of the CoLaus study. *PLoS One.* 2011;6(10):e26157.
32. Yu Z, Han S, Cao X, Zhu C, Wang X, Guo X. Genetic polymorphisms in adipokine genes and the risk of obesity: a systematic review and meta-analysis. *Obesity.* 2012;20(2):396–406.
33. Heo M, Leibel RL, Fontaine KR, Boyer BB, Chung WK, Koulu M, et al. A meta-analytic investigation of linkage and association of common leptin receptor (LEPR) polymorphisms with body mass index and waist circumference. *Int J Obes Relat Metab Disord.* 2002;26(5):640–6.
34. Angel-Chavez LI, Tene-Perez CE, Castro E. Leptin receptor gene K656N polymorphism is associated with low body fat levels and elevated high-density cholesterol levels in Mexican children and adolescents. *Endocr Res.* 2012;37(3):124–34.
35. Okada T, Ohzeki T, Nakagawa Y, Sugihara S, Arisaka O. Impact of leptin and leptin-receptor gene polymorphisms on serum lipids in Japanese obese children. *Acta Paediatr.* 2010;99(8):1213–7.
36. Santaniemi M, Ukkola O, Kesaniemi YA. Tyrosine phosphatase 1B and leptin receptor genes and their interaction in type 2 diabetes. *J Intern Med.* 2004; 256(1):48–55.
37. Labayen I, Ruiz JR, Moreno LA, Ortega FB, Beghin L, DeHenauw S, et al. The effect of ponderal index at birth on the relationships between common LEP and LEPR polymorphisms and adiposity in adolescents. *Obesity.* 2011;19(10):2038–45.
38. Huth C, Illig T, Herder C, Gieger C, Grallert H, Vollmert C, et al. Joint analysis of individual participants' data from 17 studies on the association of the IL6 variant -174G > C with circulating glucose levels, interleukin-6 levels, and body mass index. *Ann Med.* 2009;41(2):128–38.
39. Grallert H, Huth C, Kolz M, Meisinger C, Herder C, Strassburger K, et al. IL-6 promoter polymorphisms and quantitative traits related to the metabolic syndrome in KORA S4. *Exp Gerontol.* 2006;41(8):737–45.
40. Qi L, Zhang C, van Dam RM, Hu FB. Interleukin-6 genetic variability and adiposity: associations in two prospective cohorts and systematic review in 26,944 individuals. *J Clin Endocrinol Metab.* 2007;92(9):3618–25.
41. Ramírez-López G, Portilla-de Buen E, Sánchez-Corona J, Salmerón-Castro J, Mendoza-Carrera F. Interleukin-6 polymorphisms are associated with obesity and hyperglycemia in Mexican adolescents. *Arch Med Res.* 2013;44(1):62–8.
42. Dedoussis GV, Manios Y, Choumerianou DM, Yiannakouris N, Panagiotakos DB, Skenderi K, et al. The IL-6 gene G-174C polymorphism related to health indices in Greek primary school children. *Obes Res.* 2004;12(7):1037–41.
43. Belo VA, Luizon MR, Lacchini R, Miranda JA, Lanna CM, Souza-Costa DC, Tanus-Santos JE. The effects of NAMPT haplotypes and metabolic risk factors on circulating visfatin/NAMPT levels in childhood obesity. *Int J Obes (Lond).* 2013.
44. Blakemore AI, Meyre D, Delplanque J, Vatin V, Lecoœur C, Marre M, et al. A rare variant in the visfatin gene (NAMPT/PBEF1) is associated with protection from obesity. *Obesity.* 2009;17(8):1549–53.
45. NHLBI Exome Sequencing Project (ESP). The Exome Variant Server. 2015 [<http://evs.gs.washington.edu/EVS/>]
46. Speliotes EK, Willer CJ, Berndt SI, Monda KL, Thorleifsson G, Jackson AU, et al. Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. *Nat Genet.* 2010;42(11):937–48.
47. Sørensen TIA, Sonne-Holm S. Risk in childhood of development of severe adult obesity: retrospective, population-based case-cohort study. *Am J Epidemiol.* 1988;127(1):104–13.
48. WHO Consultation. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet.* 2004; 363(9403):157–63.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

