




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Study of SEIR epidemic model and scenario analysis of COVID-19 pandemic

Subrata Paul^a, Animesh Mahata^{b,*}, Uttam Ghosh^c, Banamali Roy^d

^a Department of Mathematics, Arambagh Government Polytechnic, Arambagh, West Bengal, India

^b Mahadevnagar High School, Maheshitala, Kolkata- 700141, West Bengal, India

^c Department of Applied Mathematics, University of Calcutta, Kolkata-700009, India

^d Department of Mathematics, Bangabasi Evening College, Kolkata-700009, West Bengal, India

ARTICLE INFO

Keywords:

COVID-19

Epidemic

SEIR model

Stability analysis

Case study: India and Brazil

ABSTRACT

In recent times, the Coronavirus disease (caused by COVID-19) is evidently observed to be the extremely contagious one with high fatality rate worldwide. In March 2020, the disease was declared a “global pandemic” by the World Health Organization (WHO). So far, there is no known/effective vaccine or medicine. In this paper, we propose and analyze an SEIR compartment model. We also compare and analyze the case study of India and Brazil. The model system is discussed by using MATLAB (2018a) software and the numerical results are verified graphically.

1. Introduction

Coronavirus disease 2019 (COVID-19), that has generated a pandemic, is primarily a respiratory illness, caused by a novel coronavirus that spreads from person to person, poses a serious public health risk with its high contagion rate. In most cases, the infected people perceive mild respiratory symptoms that usually disappear on their own, but some people develop severe illness, like pneumonia [1]. The virus is transmitted through contact with an infected person or via respiratory droplets when an infected person coughs or sneezes.

Since December 2019, an outbreak of a new coronavirus named SARS-COV-2 has been reported in many countries and has infected thousands of people all over the world with a high mortality rate. The World Health Organization has declared COVID-19 to be a pandemic. First it was identified in Wuhan city, Hubei Province of China on December 31, 2019 [2]. The symptoms of COVID-19 appearing within 2–14 days after infection include fever, cough, a running nose and difficulty in breathing. There is a higher risk of infection if one has been in a contaminated area or if one has been in close contact with a person infected with the new coronavirus. There is also a higher risk if one suffers from co-morbidities. In recent times, most of the countries are in the clutches of COVID-19 and a large number of people are infected. COVID-19, has seriously affected the lives of people in multiple ways and incurred huge losses to the economy.

Coronavirus is common among animals. However, animal to human transmission is reportedly rare. It causes damage to the respiratory tract

ranging from the common cold to severe conditions like SARS [3]. The outbreak of Coronavirus (officially known as Covid-19), which started in China, has so far killed 567,657 people and infected 12,844,410 people across the globe (as on July 12, 2020) [4]. In India, 849,553 confirmed cases and 22,674 deaths have been reported so far according to official figures released by the Union Ministry of Health and Family Welfare (MoHFW) [5].

Mathematical models can be viable tools in analyzing the spread and control of infectious disease; for instance, the epidemic model developed by Kermack and Mckendrick in 1927 [6]. In epidemiology there are different models to predict and explain the dynamics of an epidemic. The data of the Covid-19 outbreak can also be studied through various mathematical models such as SIR, SEIR (Susceptible, Exposed, Infected and Recovery), SIQR (Susceptible, Infectious, Quarantined and Recovered) and so on [7–12]. Tang, Wang, Li and Bragazzi [13] presented a compartmental deterministic model that would integrate the clinical development of the disease, the epidemiological status of the patient and the measures for intervention. In this situation, several studies have been conducted using real time data of the affected countries and distinct characteristics of the outbreak [14,15] have been examined. Kamrujjaman, Ghosh and Islam [16] developed the Susceptible Exposed Infectious Recovered model (SEIR) to clarify the dynamics of COVID-19 case. The SEIR model divides the population into four parts, namely the susceptible $S(t)$, the exposed $E(t)$, the infected $I(t)$ and the recovered $R(t)$ at time t . Several researchers have worked on mathematical modeling of the novel coronavirus. It exposes that those SIS, SIR and SEIR models can

* Corresponding author.

E-mail address: animeshmahata8@gmail.com (A. Mahata).

<https://doi.org/10.1016/j.egg.2021.100087>

Received 30 July 2020; Received in revised form 4 May 2021; Accepted 21 May 2021

Available online 31 May 2021

2405-9854/© 2021 Elsevier Inc. All rights reserved.




Fig. 1. The Diagram of the SEIR model.




Fig. 2. Confirmed cases, Recovered cases and Deceased cases with respect to time (Days).

Table 1
Model parameters and their descriptions.

Notation	Interpretations
β	Effective contact rate
λ	Birth rate of susceptible
μ	Mortality rate
k	Progression rate exposed to infected
γ	Recovery rate

Table 2
Parameter values of proposed model of system (i).

Parameter	Value for India	Value for Brazil	Reference
β	0.476	0.4417	Estimated
λ	0.0182	0.01867	Estimated
μ	0.0073	0.00626	Estimated
k	0.071	0.77751938	[5]
γ	0.286	0.07143	[5]
R_0	3.67	2.8421	Estimated

reflect the dynamics of various epidemics well. Concurrently, different models have been used to model the COVID-19 [17–24]. Its one variant, SEIR [25] is considered to be the most appropriate modeling technique for COVID-19. It may be observed that the specifications of this virus require more complex models in the days to come.

Some of the frequently asked questions that need to be answered are as follows. How many people exactly recover from COVID-19 over a certain period of time? How many people are infected from the disease? How many people died from it? Recently, several mathematical models suitable for the study of the dynamics and evolution of this pandemic

have been published as an attempt to answer the above questions.

In the present work we introduce and analyze the most basic transmission model for a directly transmitted infectious disease caused by bacteria, viruses or fungi. Direct transmission occurs through individual-to-individual contact, through a sneeze or cough or through skin-skin contact.

The purpose of the present work is:

1. Study of the dynamical behavior of the model and its stability.
2. Confirmation of the results by numerical simulation to control the spread of COVID-19.
3. Comparison and analysis of the scenarios of two countries, namely India and Brazil.
4. Validation and discussion of the model in COVID-19 cases of a state named West Bengal in India.

Brazil is the fifth largest country in the world with India at the seventh position. It has a much lower population density than India. However, Brazil shows higher number of confirmed cases and a higher mortality rate than India. The above observations motivate us to conduct a comparative study of COVID -19 cases between India and Brazil.

The article is organized as follows:

In Section 2 we describe the SEIR Model and study the equilibrium points. Section 3 is devoted to the discussion of stability analysis and stability criterion of the model. The case studies of India and Brazil are presented in Section 4. In Section 5 we perform numerical simulation using MATLAB. We discuss the comparison between India and Brazil in Section 6. In Section 7, we exhibit the validation of the model. Finally, Section 8 includes conclusion of the paper.

2. Model formulation

Consider a Susceptible –Exposed –Infected – Recovered (SEIR) model for the present epidemic. The SEIR disease transmission model is based on several strong assumptions [26]. The population (N) is divided into four classes: the susceptible individuals (S), the exposed individuals (E), the infected individuals (I) and the recovered individuals (R) at any time $t \geq 0$ (see Fig. 1). Therefore, we have

$$N(t) = S(t) + E(t) + I(t) + R(t)$$

$$\left. \begin{aligned} \frac{dS}{dt} &= \lambda - \beta SI - \mu S, \\ \frac{dE}{dt} &= \beta SI - (\mu + k) E, \\ \frac{dI}{dt} &= kE - (\mu + \gamma) I, \\ \frac{dR}{dt} &= \gamma I - \mu R, \end{aligned} \right\} \quad (i)$$

with initial condition $S(0) = S_0 > 0$, $E(0) = E_0 \geq 0$, $I(0) = I_0 > 0$ and $R(0) = R_0 \geq 0$.




Fig. 3. New cases, Cumulative cases and New death, Cumulative death with respect to time (Days).




Fig. 4. New cases, Cumulative cases and New death, Cumulative death with respect to time (Days).

2.1. Positivity and boundedness of solutions

Theorem. All the variables are non-negative for all $t \geq 0$.

The closed region $\Omega = \left\{ (S, E, I, R) \in \mathbb{R}^4 : 0 < N \leq \frac{\lambda}{\mu} \right\}$ is positive invariant for the system (i).

Proof. From the equation (i), we get

$$\frac{dS}{dt} = \lambda - \beta SI - \mu S \geq -(\beta I + \mu) S$$

$$\text{We have, } S(t) \geq S(0) \exp \left(- \int_0^t (\beta I + \mu) dp \right) > 0.$$

$$\text{Now } \frac{dE}{dt} = \beta SI - (\mu + k) E \geq -(\mu + k) E.$$

$$\text{We have, } E(t) \geq E(0) \exp \left(- \int_0^t (\mu + k) dp \right) > 0.$$




Fig. 5. Confirmed cases, Recovered cases and Death cases with respect to time (Days).




Fig. 6. Time series plot of equation (i) with parameter values for India given in Table 2.

$$\text{Also } \frac{dI}{dt} = kE - (\mu + \gamma)I \geq -(\mu + \gamma)I.$$

$$\text{We have, } I(t) \geq I(0) \exp\left(-\int_0^t (\gamma + \mu) dp\right) > 0.$$

$$\text{Now } \frac{dR}{dt} = \gamma I - \mu R \geq -\mu R.$$

$$\text{We have, } R(t) \geq R(0) \exp\left(-\int_0^t (\mu) dp\right) > 0.$$

$$\text{Again } \frac{d(S+E+I+T)}{dt} = \lambda - \mu(S+E+I+T).$$

Therefore,

$$\frac{dN}{dt} = \lambda - \mu N \dots$$

$$\text{If } \lambda - \mu N < 0 \text{ then } \frac{dN}{dt} < 0.$$

Therefore, the equation (ii) is bounded by $\frac{\lambda}{\mu}$.

Then we get S, E, I and R as positive function.

2.2. Basic reproduction number, disease-free equilibrium state and epidemic equilibrium state

The basic reproduction number R_0 , is the number of secondary infections that one infected person would produce in a fully susceptible population through the entire duration of the infectious period. R_0 provides a threshold condition for the stability of the disease-free equilibrium point (for most models).

The disease-free equilibrium point is locally asymptotically stable

(ii)




Fig. 7. Time series plot of equation (i) with parameter values for Brazil given in Table 2.





Fig. 8. Shows that global stability of E_1 in S-E-I plane using different initial values $((1,0,1), (2,1,2)$ and $(3,1.5,3)$) and the values of parameters for (a) India (b) Brazil.

when $R_0 < 1$ i.e., the disease dies out. The disease-free equilibrium point is unstable when $R_0 > 1$ i.e., the disease establishes itself in the population or an epidemic occurs. Since, the considered model has disease-free equilibrium at $(\frac{\lambda}{\mu}, 0, 0, 0)$, the basic reproduction number can be found analytically.

The basic reproduction number (R_0) for the COVID-19 model can be obtained from the leading eigen value of the matrix FV^{-1} [27] where,




Fig. 9. Confirm Cases with respect to time (Days).




Fig. 10. Recovery Cases with respect to time (Days).




Fig. 11. Death Cases with respect to time (Days).




Fig. 12. Infected region.




Fig. 13. Infected cases.

$$F = \begin{bmatrix} \frac{\beta\lambda}{\mu} & 0 \\ 0 & 0 \end{bmatrix} \text{ and } V = \begin{bmatrix} 0 & k + \mu \\ \gamma + \mu & -k \end{bmatrix}.$$

Therefore, the reproduction number

$$(R_0) = \frac{k\beta\lambda}{\mu(k + \mu)(\gamma + \mu)} \dots \quad (\text{iii})$$

3. Stability analysis

The equilibrium points are obtained by equating the right hand side of the equations in system (i) to zero.

$$\frac{dS}{dt} = \frac{dE}{dt} = \frac{dI}{dt} = \frac{dR}{dt} = 0.$$

The two equilibrium points are given by $E_0 = (\frac{\lambda}{\mu}, 0, 0, 0)$ which is the disease-free equilibrium point and $E_1 = (S^*, E^*, I^*, R^*)$ which is the

unique epidemic point of the system (i), where.

$$S^* = \frac{\lambda - (\mu + k)E}{\mu},$$

$$E^* = \frac{\{\mu(\mu + k)(\mu + \gamma)\}(R_0 - 1)}{\beta k(\mu + k)},$$

$$I^* = \frac{kE}{\mu + \gamma},$$

$$R^* = \frac{\lambda k E}{\mu(\mu + \gamma)}.$$

With R_0 given by equation (iii). In case of an epidemic, E^* will exist only when $R_0 > 1$.

3.1. Theorem

The disease-free equilibrium of the system is locally stable if $R_0 < 1$ and unstable if $R_0 > 1$.

Proof: From equation (i) we consider




Fig. 14. Infected cases.

$$\lambda - \beta SI - \mu S = F_1,$$

$$\beta SI - (\mu + k)E = F_2,$$

$$kE - (\mu + \gamma)I = F_3,$$

$$\gamma I - \mu R = F_4.$$

The Jacobian matrix is,

$$J = \begin{bmatrix} -\beta I - \mu & 0 & -\beta S & 0 \\ \beta I & -(\mu + k) & \beta S & 0 \\ 0 & k & -(\mu + \gamma) & 0 \\ 0 & 0 & \gamma & -\mu \end{bmatrix}$$

At the equilibrium point $E_0 = (\frac{\lambda}{\mu}, 0, 0, 0)$, the Jacobian matrix becomes,

$$J(E_0) = \begin{bmatrix} -\mu & 0 & -\frac{\beta\lambda}{\mu} & 0 \\ 0 & -(\mu + k) & \frac{\beta\lambda}{\mu} & 0 \\ 0 & k & -(\mu + \gamma) & 0 \\ 0 & 0 & \gamma & -\mu \end{bmatrix}$$

Therefore, its characteristic equation is

$$\begin{vmatrix} -\mu - x & 0 & -\frac{\beta\lambda}{\mu} & 0 \\ 0 & -(\mu + k) - x & \frac{\beta\lambda}{\mu} & 0 \\ 0 & k & -(\mu + \gamma) - x & 0 \\ 0 & 0 & \gamma & -\mu - x \end{vmatrix} = 0$$

The characteristic roots are $-\mu$, $-\mu$, $-(\mu + \lambda)$ and $(\mu + k)(R_0 - 1)$.

The first three roots are negative and the last one is will be negative if $R_0 < 1$ and positive if $R_0 > 1$.

Hence the equilibrium point E_0 is locally asymptotically stable if $R_0 < 1$ and unstable if $R_0 > 1$.

3.2. Theorem

If $R_0 > 1$, the epidemic equilibrium E_1 is locally asymptotically stable.

Proof: We consider the equations

$$F_1 = \lambda - \beta SI - \mu S,$$

$$F_2 = \beta SI - (\mu + k)E,$$

$$F_3 = kE - (\mu + \gamma)I,$$

$$F_4 = \gamma I - \mu R.$$

The Jacobian matrix.

$$J = \begin{bmatrix} -\beta I - \mu & 0 & -\beta S & 0 \\ \beta I & -(\mu + k) & \beta S & 0 \\ 0 & k & -(\mu + \gamma) & 0 \\ 0 & 0 & \gamma & -\mu \end{bmatrix}.$$

At the equilibrium point $E_1 = (S^*, E^*, I^*, R^*)$, the Jacobian matrix becomes,

$$J(E_1) = \begin{bmatrix} -\beta I^* - \mu & 0 & -\beta S^* & 0 \\ \beta I^* & -(\mu + k) & \beta S^* & 0 \\ 0 & k & -(\mu + \gamma) & 0 \\ 0 & 0 & \gamma & -\mu \end{bmatrix}.$$

Therefore, its characteristic equation is

$$\begin{vmatrix} -\beta I^* - \mu - x & 0 & -\beta S^* & 0 \\ \beta I^* & -(\mu + k) - x & \beta S^* & 0 \\ 0 & k & -(\mu + \gamma) - x & 0 \\ 0 & 0 & \gamma & -\mu - x \end{vmatrix} = 0.$$

Or, $(-\mu - x)(x^3 + ax^2 + bx + c) = 0$, where

$$a = \beta I^* + 3\mu + k + \gamma$$

$$b = (\beta I^* + \mu)(2\mu + k + \gamma) + (\mu + k)(\mu + \gamma)$$

$$c = (\beta I^* + \mu)(\mu + k)(\mu + \gamma) - \mu\beta k S^*$$

By Routh-Hurwitz Criterion, the system (i) is locally asymptotically stable if $a > 0$, $b > 0$, $ab > c$.

Thus, E_1 is a locally asymptotically stable equilibrium point.

3.3. Theorem

The disease-free equilibrium of the system (i) is globally asymptotically stable if $R_0 < 1$.

Proof: Considering the following linear Lyapunov function

$$L = B_1 E + B_2 I$$

With Lyapunov derivative (where a dot represents differentiation with respect to time)

$$\dot{L} = B_1 \dot{E} + B_2 \dot{I}$$

Substituting the expression for \dot{E} and \dot{I} from (i), we have

$$\frac{dL}{dt} = B_1 [\beta SI - (\mu + k)E] + B_2 [kE - (\mu + \gamma)I] \dots \quad (iv)$$

$$\dot{V} = \left(\lambda - \beta SI - \mu S - \frac{S^*(\lambda - \beta SI - \mu S)}{S} \right) + \left((\beta SI - (\mu + k)E) - \frac{E^*(\beta SI - (\mu + k)E)}{E} \right) + Q \left((kE - (\mu + \gamma)I) - \frac{I^*(kE - (\mu + \gamma)I)}{I} \right) \dots \quad (vii)$$

Little perturbation from equation (iv) with the reproduction number (iii) gives:

$$B_1 = \lambda k, \quad B_2 = \mu(\mu + k) \dots \quad (v)$$

Substituting the expression of B_1, B_2 obtained from equation (v) we have:

$$\begin{aligned} \frac{dL}{dt} &= \beta SI \lambda k - (\mu + \gamma) \mu(\mu + k) I \\ &= I [\beta S \lambda k - (\mu + \gamma) \mu(\mu + k)] \\ &= I [(\mu + \gamma) \mu(\mu + k)] \left[\frac{\beta S \lambda k}{(\mu + \gamma) \mu(\mu + k)} - 1 \right] \end{aligned}$$

Since, $S = \frac{\lambda}{\mu} \leq N$, it follows that

$$\dot{V} = \left(\beta S^* I^* + \mu S^* - \mu S - \frac{S^*(\beta S^* I^* + \mu S^* - \beta SI - \mu S)}{S} \right) + \left((\beta SI - (\mu + k)E) - \frac{E^*(\beta SI - (\mu + k)E)}{E} \right) + Q \left((kE - (\mu + \gamma)I) - \frac{I^*(kE - (\mu + \gamma)I)}{I} \right) \dots \quad (x)$$

$$\frac{dL}{dt} \leq I [(\mu + \gamma) \mu(\mu + k)] \left[\frac{\beta \lambda k}{(\mu + \gamma) \mu(\mu + k)} - 1 \right]$$

$$= > \frac{dL}{dt} \leq I [(\mu + \gamma) \mu(\mu + k)] [R_0 - 1]$$

Hence if $R_0 < 1$, then $\frac{dL}{dt} < 0$. Hence, by LaSalle's extension to Lyapunov's principle [28,29], the disease free equilibrium points is globally asymptotically stable.

3.4. Theorem

If $R_0 > 1$, the epidemic equilibrium E_1 is globally asymptotically stable.

Proof: Consider the model (i) and $R_0 > 1$, so that the epidemic equilibrium E_1 of model exists.

We consider the following non-linear Lyapunov function of Goh-Volterra type:

$$V = \left(S - S^* - \log \frac{S}{S^*} \right) + \left(E - E^* - \log \frac{E}{E^*} \right) + Q \left(I - I^* - \log \frac{I}{I^*} \right)$$

With Lyapunov derivative (where a dot represents differentiation with respect to time)

$$\dot{V} = \left(\dot{S} - \frac{S^* \dot{S}}{S} \right) + \left(\dot{E} - \frac{E^* \dot{E}}{E} \right) + Q \left(\dot{I} - \frac{I^* \dot{I}}{I} \right) \dots \quad (vi)$$

Substituting the value $\dot{S}, \dot{E}, \dot{I}$ from (i) into (vi), we have

At steady state from equation (i) we have:

$$\lambda = \beta S^* I^* + \mu S^* \dots \quad (viii)$$

Substituting equation (viii) into (vii) gives:

$$\begin{aligned} \dot{V} &= \left(\beta S^* I^* + \mu S^* - \beta SI - \mu S - \frac{S^*(\beta S^* I^* + \mu S^* - \beta SI - \mu S)}{S} \right) + \left((\beta SI - (\mu + k)E) - \frac{E^*(\beta SI - (\mu + k)E)}{E} \right) \\ &+ Q \left((kE - (\mu + \gamma)I) - \frac{I^*(kE - (\mu + \gamma)I)}{I} \right) \dots \quad (ix) \end{aligned}$$

Further simplification gives:

Collecting all infected class without single star (*) from (x) and equating to zero:

$$S^* \beta I - (\mu + k)E + Q(kE - (\mu + \gamma)I) = 0 \dots \quad (xi)$$

A little perturbation of steady state from (i) and (xi) resulted into:

$$Q = \frac{S^* \beta}{(\mu + \gamma)} \cdot (\mu + k) = \frac{I^* S^* \beta}{E^*}, \quad k = \frac{(\mu + \gamma) I^*}{E^*} \dots \quad (xii)$$

Substituting the expression from (xii) into (x) gives:

$$\begin{aligned} \dot{V} &= \left(\beta S^* I^* + \mu S^* - \mu S - \frac{S^*(\beta S^* I^* + \mu S^* - \mu S)}{S} \right) + \\ &\left(-\frac{E^* \beta SI}{E} + I^* S^* \beta \right) + \left(-\frac{I^* S^* E \beta I}{I E^*} + \beta S^* I^* \right) \dots \quad (xiii) \end{aligned}$$

Finally, since the arithmetic mean exceeds the geometric mean, we

have

$$\left(2 - \frac{s}{S^*} - \frac{S^*}{S}\right) \leq 0, \quad \left(3 - \frac{S^*}{S} - \frac{I^* E}{IE^*} - \frac{SE^* I}{E}\right) \leq 0$$

Thus, $\dot{V} \leq 0$ for $R_0 > 1$.

Hence, V is a Lyapunov function, by LaSalle's Invariance Principle [29], the epidemic equilibrium E_1 is globally asymptotically stable.

4. Case study

In this section we observe the scenario of COVID-19 pandemic in India and Brazil.

4.1. India

We have considered the number of confirmed cases, death cases and recovered cases of India as given in Ref. [4]. The first case of COVID-19 was reported on January 30, 2020. During the month of February, the number of cases reported was only 3 and remained the same during the whole month. In March it was noted that the number of cases started increasing. However, an explosion in the number happened ultimately in May 2020. To describe the spread of COVID-19 using SEIR model, few considerations and assumptions were made due to limited availability of data [4,13,30,31]. We have considered the number of infected, recovered and deceased cases from 2nd June to July 2, 2020 [32] which is depicted in Fig. 2. The estimated values of the parameters given in Table 1 are as follows: effective contact rate $\beta \approx 0.476$, birth rate of susceptible $\lambda \approx 0.0182$ and mortality rate $\mu \approx 0.0073$. We obtain the recovery rate $\gamma \approx 0.286$.

In Fig. 3, the bar diagram shows the number of new cases as well as cumulative new cases and new death along with cumulative death from 30th January to 3rd July 2020 in India due to COVID-19 [32]. Therefore, we obtain an estimation of the value of R_0 from the equation (iii) as 3.67.

In this study, we have only predicted the number of confirmed cases. We have used only time series data for confirmed cases and death cases. An increase in active cases is an alarming situation because as discussed earlier, India is one of the largest populations in the whole world.

4.2. Brazil

The first case in Brazil was a 61-year-old man who had returned from Lombardy (Italy) and tested positive for the virus. The study of evolution of COVID 19 and its prediction in Brazil is based on the data provided by Ref. [5] during the period of 26th February to July 3, 2020. The estimated values of the parameters given in Table 1 are as follows: effective contact rate $\beta \approx 0.4417$, birth rate of susceptible $\lambda \approx 0.01867$ and mortality rate $\mu \approx 0.00626$. We obtain the recovery rate $\gamma \approx 0.07143$. Therefore, we obtain an estimation of the value of R_0 from the equation (iii) as 2.8421.

Fig. 4 shows the number of new cases as well as cumulative cases and new death as well as cumulative death from 26th February to July 3, 2020 of Brazil due to COVID-19.

Fig. 5 is a diagrammatic representation of confirmed cases, death cases and recovered cases during the period of 2nd March to July 3, 2020 in Brazil. It is observed that the confirmed cases are gradually increasing with respect to death cases from 2nd March to July 3, 2020.

5. Numerical simulation

In this section, we implement meticulous numerical validation of the results obtained analytically. We have used mathematical software MATLAB (2018a) to numerically approximate the solution of our model system (i) [14,16].

The estimated values of the parameters in the case of COVID-19 in

India and Brazil are as follows:

6. Comparison between India and Brazil

Fig. 9 is the graphical representation of comparison of confirmed cases during 2nd march to 3rd July 2020 between India and Brazil, according to the data provided in Ref. [5] and Ref. [32]. Although the number of infected cases gradually increases during this period, Brazil shows a much higher rate of increase than India.

Fig. 10 and Fig. 11 are the graphical representation of comparison of recovered cases and death cases respectively during 2nd march to 3rd July 2020 between India and Brazil, according to the data provided in Ref. [5] and Ref. [32]. There is a gradual increase in the recovery rate as well as death rate for both the countries during this period. The representation also indicates a greater recovery rate and death rate for Brazil.

Fig. 12 shows that the geographical distribution of infected cases on 3rd July 2020 of India and Brazil. It should be noted that the average testing rate (per million) in both the countries are way behind the testing rate in some of the developed countries of the world. It is expected that with an increase in the testing rate there will be an increase in the infected regions in both the countries.

7. Model validation and discussion

In this section, the infection rate is taken into consideration for validation of the solution obtained from Equation (i) with real time data. For k , μ and γ as given in Table 2 and if the population does not follow lockdown (i.e., $E \approx I$), we have from the equation (i),

$$\frac{dI}{dt} = kE - (\mu + \gamma) I.$$

$$\text{Then, } \frac{dI}{dt} = [k - (\mu + \gamma)] I.$$

Therefore,

$$I(t) = I_0 \exp [k - (\mu + \gamma)] t \dots \quad (\text{xiv})$$

I_0 being the initial number of infected cases.

Fig. 13 shows the goodness of fit of Eq. (xiv) with the real time data of total infected cases in India as reported from 2nd March to July 3, 2020. Fitted parameters are $\{k - (\mu + \gamma)\} = 0.2171$ with 95% confidence interval as (0.1952, 0.239) and $I_0 = 368.5$ with 95% confidence interval as (228.8, 508.2).

Fig. 14 shows the goodness of fit of Eq. (xiv) with the real time data of total infected cases in Brazil as reported from 2nd March to July 3, 2020. Fitted parameters are $\{k - (\mu + \gamma)\} = 0.2166$, with 95% confidence interval as (0.173, 0.2603) and $I_0 = 618.3$ with 95% confidence interval as (151.3, 1085).

8. Conclusion

The Susceptible-Exposed-Infection-Recovered (SEIR) model is used in this paper to study the dynamical behavior and stability of the model characterizing the spread of COVID-19. Graphical representations of comparison of the spread of the disease and its mortality rates are extensively discussed for India and Brazil. The essential parameters namely effective contact rate (β), birth rate of susceptible (λ), mortality rate (μ) and recovery rate (γ) are estimated for both countries using the current data [5,32] and are shown in Table 2. The reproduction number (R_0) given by equation (iii) is 3.67 for India and 2.8421 for Brazil which gives a clear indication that India will be encountering an increased number of infected cases in near future. Fig. 6 and Fig. 7 indicates a time series plot of the SIER model for given R_0 in case of India and Brazil respectively. It may be observed that the COVID-19 scenario will prevail for a much longer period of time than expected. Fig. 8 shows that the infected cases exhibit a rapid ascent and reaches its peak for different increasing values of contact rate (β).

We have compared the analytical results of infected cases obtained in

equation (iv) with the numerical data in Figs. 13 and 14. The graphs are in good agreement with the numerical simulations performed with the help of MATLAB.

From our study of the present situation it is recommended that the rate of disease transmission needs to be controlled, otherwise a huge proportion of population will be affected within an extremely short period of time. Among the popular preventive measures are enforcement of lockdown, curfews and specification of containment zones. Mainly, the government's strategy must be to restrict the contamination by reducing unnecessary mobilization and restricting unwanted social interactions.

We wish to extend our study towards a modified SEIR compartmental model accounting for infection from undiagnosed individuals and for different levels of population isolation in order to evaluate effects of contact reduction in the epidemic temporal dynamics.

CRedit authorship contribution statement

Subrata Paul: Conceptualization, Methodology. **Animesh Mahata:** Data curation, Software, Investigation. **Uttam Ghosh:** Software, Validation. **Banamali Roy:** Supervision, Writing – review & editing.

Acknowledgement

The authors are grateful to the reviewers for their valuable comments and suggestions.

References

- [1] D.D. Richman, R.J. Whitley, F.G. Hayden, *Clinical Virology*, John Wiley & Sons, 2016.
- [2] C. Wang, P.W. Horby, F.G. Hayden, G.F. Gao, A novel coronavirus outbreak of global health concern, *Lancet* 395 (10223) (2020) 470–473.
- [3] Centers for disease control and prevention, Novel coronavirus, 2020, <https://www.cdc.gov/coronavirus/2019-ncov>, 2019.
- [4] World Health Organization, Coronavirus world health organization, Available: <https://www.who.int/health-topics/coronavirus>, 2020, 19.
- [5] <https://www.worldometers.info/coronavirus/>.
- [6] N.O. Kermack, A.G. Mackendrick, Contribution to mathematical theory of epidemics, *P. Roy. Soc. Lond. A. Math. Phys. Eng. Sci.* 115 (270) (1927) 700–721.
- [7] C. Ji, D. Jiang, N. Shi, Multigroup SIR epidemic model with stochastic perturbation, *Phys. Stat. Mech. Appl.* 390 (10) (2011) 1747–1762.
- [8] M. Liu, C. Bai, K. Wang, Asymptotic stability of a two group stochastic SEIR model with infinite delays, *Commun. Nonlinear Sci. Numer. Simulat.* 19 (10) (2014) 3444–3453.
- [9] R. Sun, J. Shi, Global stability of multi-group epidemic model with group mixing and nonlinear incidence rates, *Appl. Math. Comput.* 218 (2) (2011) 280–286.
- [10] Q. Yang, X. Mao, Extinction and recurrence of multi-group SEIR epidemic models with stochastic perturbations, *Nonlinear Anal. R. World Appl.* 14 (3) (2013) 1434–1456.
- [11] J. Yu, D. Jiang, N. Shi, Global stability of two-group SIR model with random perturbation, *J. Math. Anal. Appl.* 360 (1) (2009) 235–244.
- [12] C. Yuan, D. Jiang, D. O'Regan, R.P. Agarwal, Stochastically asymptotically stability of the multi-group SEIR and SIR models with random perturbation, *Commun. Nonlinear Sci. Numer. Simulat.* 17 (6) (2012) 2501–2516.
- [13] B. Tang, X. Wang, Q. Li, N.L. Bragazzi, S. Tang, Y. Xiao, J. Wu, Estimation of the transmission risk of the 2019-nCoV and its implication for public health interventions, *J. Clin. Med.* 9 (2) (2020) 462.
- [14] A. Tiwari, Modeling and Analysis of COVID-19 Epidemic in India, 2020 medRxiv preprint.
- [15] H.B. Fredj, F. Cherif, Novel Corona Virus Disease Infection in Tunisia: Mathematical Model and the Impact of the Quarantine Strategy, ELSEVIER, 2020.
- [16] Md Kamrujjaman, U. Ghosh, Md. S. Islam, Pandemic and the dynamics of SEIR model, Case COVID-19 (2020).
- [17] Sk. S. Nadim, I. Ghosh, J. Chattopadhyay, Short-term Predictions and Prevention Strategies for Covid-2019: A Model Based Study, 2020 arXiv preprint arXiv: 2003.08150.
- [18] A. Kucharski, T. Russell, C. Diamond, Y. Liu, J. Edmunds, S. Funk, R. Eggo, Early dynamics of transmission and control of COVID-19: a mathematical modelling study, *Lancet Infect. Dis.* (2020), [https://doi.org/10.1016/S1473-3099\(20\)30144-4](https://doi.org/10.1016/S1473-3099(20)30144-4).
- [19] Quilty, J. Billy, S. Clifford, et al., Effectiveness of airport screening at detecting travellers infected with novel coronavirus (2019-ncov), *Euro Surveill.* 25 (5) (2020).
- [20] H. Wang, Z. Wang, Y. Dong, R. Chang, et al., Phase-adjusted estimation of the number of coronavirus disease 2019 cases in wuhan, China, *Cell. Discov.* 6 (10) (2020), <https://doi.org/10.1038/s41421-020-0148-0>.
- [21] L. Ferretti, C. Wymant, M. Kendall, et al., Quantifying SARS-CoV-2 transmission suggests epidemic control with digital contact tracing, *Science* (2020), <https://doi.org/10.1126/science.abb6936>.
- [22] S. Zhao, S. Musa, Q. Lin, J. Ran, et al., Estimating the unreported number of novel coronavirus (2019-nCoV) cases in China in the first half of January 2020: a data-driven modelling analysis of the early outbreak, *J. Clin. Med.* 9 (2020) 388.
- [23] S. Zhao, Q. Lin, J. Ran, et al., Preliminary estimation of the basic reproduction number of novel coronavirus (2019-nCoV) in China, from 2019 to 2020: a data-driven analysis in the early phase of the outbreak, *Int. J. Infect. Dis.* 92 (2020) 214–217.
- [24] Joseph T. Wu, Kathy Leung, Gabriel M. Leung, Now casting and forecasting the potential domestic and international spread of the 2019-ncov outbreak originating in wuhan, China: a modelling study, *Lancet* 395 (10225) (2020) 689–697.
- [25] R.M. Anderson, R.M. May, *Infectious Diseases of Humans*, Oxford University Press, Oxford, 1991.
- [26] J. Murray, *Mathematical Biology I*, third ed., Springer-Verlag, Heidelberg, 2002.
- [27] O. Diekmann, J.A.P. Heesterbeek, M.G. Roberts, The construction of next-generation matrices for compartmental epidemic models, *J. R. Soc. Interface* 7 (47) (2009) 873–885.
- [28] L. Perko, *Differential Equations and Dynamical Systems*, Springer, 2000.
- [29] M.Y. Li, H.L. Smith, L. Wang, Global dynamics of an SEIR epidemic model with vertical transmission, *SIAM J. Appl. Math.* 62 (2001) 58.
- [30] A.O. Victor, Mathematical Predictions for COVID-19 as a Global Pandemic, 2020 medRxiv preprint.
- [31] M. Martcheva, *An Introduction to Mathematical Epidemiology*, Springer, New York, 2015.
- [32] India COVID-19 Tracker, Retrieved, <https://www.covid19india.org/2020>, 2020, 07-13.