



# Leaf coordination between petiole vascular development and water demand in response to elevated CO<sub>2</sub> in tomato plants

Itay Cohen<sup>1</sup> | Juliana Espada Lichston<sup>1,3</sup>  | Cristiane Elizabeth Costa de Macêdo<sup>3</sup> | Shimon Rachmilevitch<sup>2</sup> 

<sup>1</sup>Jacob Blaustein Center for Scientific Cooperation, The Blaustein Institutes for Desert Research, Ben-Gurion University of the Negev, Beersheba, Israel

<sup>2</sup>The French Associates Institute for Agriculture and Biotechnology of Drylands, The Blaustein Institutes for Desert Research, Ben-Gurion University of the Negev, Beersheba, Israel

<sup>3</sup>Federal University of Rio Grande do Norte, Natal, Brazil

## Correspondence

Shimon Rachmilevitch, The French Associates Institute for Agriculture and Biotechnology of Drylands, The Blaustein Institutes for Desert Research, Ben-Gurion University of the Negev, Beersheba 8499000, Israel.  
Email: rshimon@bgu.ac.il

## Funding information

Israeli Ministry of Agriculture and Rural Development (Eugene Kandel Knowledge Centers)

## Abstract

The rise in atmospheric CO<sub>2</sub> has a profound impact on plants physiology and performance. Stomatal gas exchange such as reduction in water loss via transpiration and higher photosynthetic rates are among the key plant physiological traits altered by the increase of CO<sub>2</sub>. Water acquired in plant roots is transported via the xylem vessels to the shoots. Under conditions of elevated CO<sub>2</sub>, water flux decreases due to higher water use efficiency and a decline in stomatal conductance. However, the mechanism by which the shoot vascular development is affected under elevated CO<sub>2</sub> is still largely unclear in herbaceous crops. In the current study, tomato plants were exposed to either 400 or 800 ppm of CO<sub>2</sub> and were analyzed for growth, leaf area, gas exchange rate, and petiole anatomy. Elevated CO<sub>2</sub> caused a reduction in metaxylem vessel diameter, which in turn, decreased leaf theatrical conductivity by 400% as compared with plants grown under ambient CO<sub>2</sub>. This work links anatomical changes in the petioles to the rise in atmospheric CO<sub>2</sub> and water use. Plant water demand declined under elevated CO<sub>2</sub>, while photosynthesis increased. Thus, the decrease in leaf specific conductivity was attributed to lower water consumption in leaf gas exchange and, by extension, to higher leaf water use efficiency. As the global climate changes and water scarcity becomes more common, such anatomical alterations caused by elevated CO<sub>2</sub> may affect plant response to water limitation. Further research on petiole anatomical alterations under conditions of combined climate change factors such as drought and heat with elevated CO<sub>2</sub> may assist in clarifying the responses expected by future climate scenarios.

## KEYWORDS

carbon assimilation (A), elevated CO<sub>2</sub> (eCO<sub>2</sub>), leaf specific conductivity (K<sub>L</sub>), tomato (*Solanum lycopersicum*), transpiration rates (Tr), water use efficiency (WUE)

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *Plant Direct* published by American Society of Plant Biologists and the Society for Experimental Biology and John Wiley & Sons Ltd.

## 1 | INTRODUCTION

Since the beginning of the Industrial Revolution (mid-18th century), the use of fossil fuels and deforestation have caused a continuous increase in the atmospheric CO<sub>2</sub> concentration as compared with previous centuries. The current concentration is ~410 μmol mol<sup>-1</sup>, while projections predict an increase to 530–970 μmol mol<sup>-1</sup> by the end of the twenty-first century (Kant et al., 2012). This increase in atmospheric CO<sub>2</sub> levels has substantial effects on plant photosynthesis and growth in both managed and natural environments (Bloom et al., 2012).

Photosynthesis involves leaf gas exchange of CO<sub>2</sub> and water through the stomata; therefore, when the gradient of CO<sub>2</sub> between leaf intercellular spaces and the atmosphere increases, the rise in atmospheric CO<sub>2</sub> concentration helps plants to lower stomatal water loss per unit of CO<sub>2</sub> gained (Paudel et al., 2018).

Elevated CO<sub>2</sub> (eCO<sub>2</sub>) simultaneously increases carbon assimilation (A) and decreases the transpiration rate (T) on leaf, plant, and ecosystem scales (Mohamed et al., 2013). Dividing A by T (A/T) generates the widely used term water use efficiency (WUE), which describes how efficiently carbon is assimilated per unit of water evaporated on the leaf scale (Blum, 2009; Medrano et al., 2015). Numerous studies have reported that eCO<sub>2</sub> increases WUE in C3 plants (Ainsworth & Long, 2005; Franzaring et al., 2011; Wu et al., 2004; Wullschleger et al., 2002). This higher WUE is attributed to the increase in frequent stomatal closure events due to higher leaf intercellular CO<sub>2</sub> concentrations (Leakey et al., 2009). Because eCO<sub>2</sub> increases net carbon gain, it was proposed that an increase in WUE under eCO<sub>2</sub> is beneficial for plants under limited water availability, particularly in dry lands (Wu et al., 2004). Heat and drought stresses are key climate change factors negatively impacting photosynthesis and crop productivity (Cohen, Zandalinas, Fritschi, & Mittler, 2020; Cohen, Zandalinas, Huck, et al., 2020). This positive impact of CO<sub>2</sub> fertilization on photosynthesis may partly off-set the negative impacts of climate change factors on plants in both managed and natural environments.

Water acquired by the root system is carried by the transpiration stream in the vascular tissue to the shoot, mainly for stomatal gas exchange (Leakey et al., 2009). Xylem is a vascular tissue found in all vascular plants and functions as a rapid conduit of water and a mechanical support for vertical growth (Růžička et al., 2015). Recently, it was shown that the transport of water through the xylem vessels is over 10<sup>6</sup> times more efficient than transport via the parenchyma cells or plasmodesmata (Qaderi et al., 2019).

A previous study observed that increasing the CO<sub>2</sub> levels caused alterations in the tomato root vascular system's ability to conduct water, especially when nitrogen availability is low (Cohen et al., 2019) or when nitrogen is applied as ammonium (Cohen et al., 2018). Water conductivity in xylem vessel obeys Hagen–Poiseuille's law, and the flow rate in the xylem is mostly determined by the sum of the diameters of each individual xylem vessel (Hochberg et al., 2015; Tyree et al., 1994). Accordingly, for any given leaf, an increase in the leaf petiole xylem vessel size results in higher water conductivity and greater leaf specific conductivity (K<sub>L</sub> hydraulic conductivity leaf

area<sup>-1</sup>) (Rodríguez-Gamir et al., 2010). However, an increase in the stem and petiole xylem vessel diameter also raises the risk of cavitation (Cochard, 2002) and decreases embolism resistance. Thus, xylem size development is a trade-off between water conductivity and embolism resistance (Gleason et al., 2016; Wagner et al., 1998). It was suggested that crop plants are sensitive to drought-induced embolism because of the low cost of xylem tissue construction (Liu et al., 2020).

Moreover, genotypes with lower axial conductivity can maintain a more favorable hydration status under water limitation than high conductivity genotypes (Strock et al., 2020). In oaks (*Quercus robur* L.), it was shown that trees with larger xylem vessels displayed higher mortality under prolonged droughts (Qaderi et al., 2019).

Recently, eCO<sub>2</sub> was found to alter root xylem development (Cohen et al., 2019), and other studies showed that stem xylem was also affected by eCO<sub>2</sub> (Kostiainen et al., 2004; Liu et al., 2020). However, the effects of eCO<sub>2</sub> on the size of petiole vessels and leaf specific conductivity (K<sub>L</sub>) remain uncharacterized.

The current study hypothesized that the decrease in petiole metaxylem vessel size and K<sub>L</sub> was due to a reduction in WUE caused by eCO<sub>2</sub>.

The present study investigated the effects of eCO<sub>2</sub> on the development of xylem vessels in the petioles and on the leaf specific conductivity of tomato plants.

## 2 | MATERIALS AND METHODS

### 2.1 | Plant material and growth conditions

Tomato plants (*Solanum lycopersicum* cv. Brigade, Hishtil Co.; Ashkelon, Israel) were planted in an environmentally controlled greenhouse. Throughout the experiment, the plants were grown at day/night temperatures of 25/17°C, respectively, supplied with either ambient (400 μmol mol<sup>-1</sup>) or elevated (800 μmol mol<sup>-1</sup>) CO<sub>2</sub> concentrations and relative humidity of ~45%. The plants were grown during April–May of 2017, with a day length of about 13 h under natural sunlight.

CO<sub>2</sub> levels in the two greenhouses were continuously monitored with an infrared gas analyzer sensor (Gascard NG; Edinburgh Sensors), and CO<sub>2</sub> was automatically resupplied on demand.

The nutrient solution for each treatment contained 6.5, 1.5, 2.6, 4, 5, and 4 mM of N, P, Mg, Ca, K, and S, respectively, and micro-nutrients (Cohen et al., 2019).

### 2.2 | Gas exchange measurements

Gas exchange measurements were conducted on young fully mature leaves prior to plant harvest. Measurements were taken during the mid-morning (10:00–12:00), and the net CO<sub>2</sub> assimilation rates (A), transpiration (T), and stomatal conductance (g<sub>s</sub>) of the young fully mature leaves of each plant were measured using a portable Li-6400XT Infrared Gas Analyzer (Li-Cor Biosciences Inc.; NE, USA). The



open photosynthetic system was equipped with an external carbon dioxide source, in order to maintain the leaf chamber at a CO<sub>2</sub> concentration of either 400 or 800 μmol mol<sup>-1</sup>. The chamber temperature was adjusted to 24°C, and the photosynthetic active radiation (PAR) rate was 1,000 μmol photons m<sup>-2</sup> s<sup>-1</sup> (Cohen et al., 2019).

Leaf instantaneous WUE was calculated from gas exchange measurements as the ratio between net carbon assimilation (A) and transpiration rate (T) (Medrano et al., 2015).

### 2.2.1 | Biomass measurement

At the end of the experiment, the aboveground portion of the plants was harvested and collected for fresh weight biomass analysis.

## 2.3 | Leaf area measurements

The leaf area measurements of whole young fully mature tomato leaves were taken by digital image analysis, using Easy Leaf Area software (Easlon & Bloom, 2014). The leaves were collected at the end of study, and the same leaves were later subjected to anatomical analysis. Photographs were taken with an iPhone 5 mobile phone and calibrated using a 4-cm<sup>2</sup> red calibration area, according to the software guidelines.

## 2.4 | Anatomy

At the end of the study, the harvested petioles of young fully mature leaves and preserved them in 70% ethanol till further processing. The samples were analyzed with cross sections made freehand. Sections were depigmented with commercial sodium hypochlorite (8%) in proportions of 1:1 hypochlorite and distilled water, which in a few seconds, removed the natural plant pigments. Subsequently, the samples were washed in distilled water and stained with aqueous solutions of Alcian Blue (1% w/v) and Safranin 1% (w/v) for 3 min each. Cross sections were placed on slides with glycerin 50% in distilled water, examined through an Axio ImagerA1 microscope with dark-field illumination (Zeiss; Goettingen, Germany), and photographed with an AxioCam HRC camera (Zeiss). The total number of xylem vessels in each petiole was counted, and the areas of individual metaxylem vessels and total xylem tissue were analyzed using image analysis software; we did not measure the area of individual protoxylem vessels (Digimizer version 4.2.5; Ostend, Belgium) as described by Kraus and Arduin (1997).

## 2.5 | Theoretical conductivity

Leaf theoretical conductivity was calculated as described in Shtein et al. (2016). Using the modified Hagen–Poiseuille's law, the current study divided the leaf area into the sum of the calculated theoretical

conductivity =  $\frac{\pi \rho}{128 \eta} \times (d^4)$  of each individual xylem vessel, and the collective conductivity of all individual vessels in a given petiole was calculated ( $d$  is the vessel diameter,  $\rho$  is the fluid density [assumed to be 1,000 kg m<sup>-3</sup>], and  $\eta$  is the viscosity [assumed to be 1 × 10<sup>-9</sup> MPa × s]).

## 2.6 | Experimental design and statistical analyses

The experiment was designed in a random plot with eight replications, with CO<sub>2</sub> treatments as a single factor.


A statistical analysis was performed using Student's *t* test, with an  $\alpha$  level of .05 using JMP 10 (SAS).

## 3 | RESULTS

eCO<sub>2</sub> significantly increased the aboveground biomass accumulation of tomato plants. The dry weights of the plants were 150 and 170 g for the ambient and eCO<sub>2</sub>-grown plants, respectively (Figure 1a). This difference in dry biomass corresponds to a 14% increase in eCO<sub>2</sub>-grown plants. Average leaf area slightly increased in response to CO<sub>2</sub> enrichment, displaying areas of 150 and 158 cm<sup>2</sup> for ambient and eCO<sub>2</sub>-grown plants, respectively (Figure 1b).

The photosynthetic rates were 13.5 and 17 μmol CO<sub>2</sub> s<sup>-1</sup> m<sup>-2</sup> for the ambient and eCO<sub>2</sub>-grown tomato plants, respectively. This significant difference in the leaf-based photosynthetic rate constituted a 26% increase in eCO<sub>2</sub>-grown plants (Figure 2a). In addition to photosynthesis, transpiration also differed significantly between CO<sub>2</sub> treatments. Tomato plants exhibited transpiration rates of 0.62 and 0.5 mmol H<sub>2</sub>O s<sup>-1</sup> m<sup>-2</sup> for the ambient and eCO<sub>2</sub> treatments, respectively. This difference corresponds to a 20% decrease in the transpiration of eCO<sub>2</sub>-grown plants (Figure 2b). Stomatal conductance was also found to be significantly affected by eCO<sub>2</sub>; tomato plants displayed stomatal conductance rates of 0.2 and 0.12 mmol s<sup>-1</sup> m<sup>-2</sup> in the ambient and eCO<sub>2</sub>-grown plants, respectively (Figure 2c). This difference corresponds to a 40% decrease in the displayed stomatal conductance of eCO<sub>2</sub>-grown plants. As a result of the described alterations in leaf gas exchange rates, leaf instantaneous WUE was accordingly changed. Tomato plants displayed WUE rates of 2.8 and 3.8 for ambient and eCO<sub>2</sub>-grown plants, respectively. This difference corresponds to a 35% increase in WUE for the eCO<sub>2</sub>-grown plants (Figure 2d).

The total area of individual petioles' xylem tissue was not significantly different between the two CO<sub>2</sub> treatments (as seen in Figure 3a). Similarly, the number of metaxylem vessels in each petiole was found to be similar in both treatments, ranging from 90–120 vessels per petiole (Figure 3b). In contrast, under eCO<sub>2</sub>, the average diameter of an individual metaxylem vessel decreased by an average of 40% in comparison with that under ambient CO<sub>2</sub> conditions (Figure 3c). These differences in individual metaxylem vessel diameters are shown in Figure 3d,e. The distribution of vessel size diameter is presented in Figure 4a,b. The median vessel diameter observed was



**FIGURE 1** Tomato plant biomass (a) and average leaf area (b) as affected by CO<sub>2</sub> treatment ( $n = 8$ ). Results are expressed as means  $\pm$  standard error. Bars labeled with an asterisk are significantly different ( $p < .05$ )

17.6 and 13.4  $\mu\text{m}$  for ambient and eCO<sub>2</sub>-grown plants, respectively. While xylem vessels demonstrated different responses to eCO<sub>2</sub>, the average parenchyma cell diameter in the petiole was not different under eCO<sub>2</sub> conditions (Figure 3f).

Theoretical conductivity was calculated from the metaxylem vessel size and leaf area data.  $K_L$  significantly decreased in plants under eCO<sub>2</sub>, from 0.042 to 0.01 ( $[\text{kg s}^{-1} \text{m}^{-1} \text{MPa}^{-1}] \times 10^{-4}$ ); this response represents an over 400% decrease in theoretical conductivity for plants grown under eCO<sub>2</sub> compared with plants grown under ambient CO<sub>2</sub> (Figure 5).


## 4 | DISCUSSION

The current study demonstrates that tomato petiole vascular anatomy displays phenotypic plasticity under eCO<sub>2</sub> as compared with growth under ambient CO<sub>2</sub> conditions. This alteration coincides with the decrease in leaf transpiration rate and leaf specific conductivity. Vascular tissue development is balanced between the need to rapidly conduct water and the need to provide cavitation resistance (Hacke & Sperry, 2001). Previous studies showed a tight correlation between leaf stomatal conductance and petiole xylem water transport capacity (Brocius & Hacke, 2016). While other studies presented changes in plant vascular anatomy under eCO<sub>2</sub>, these reports either focused on stress (Domec et al., 2015; Hao et al., 2018) or described changes in the phloem (Ainsworth & Lemonnier, 2018). Leaf area did not change

(Figure 2e), leaf thickness increased and specific leaf area decreased (Figure 2d) under eCO<sub>2</sub>, which is an adaptation and acclimation mechanism to the increase of CO<sub>2</sub>. Similar responses were shown in other C3 crops under eCO<sub>2</sub> (Gillespie et al., 2012), and as a consequence, this alteration could explain the reduction in stomatal conductance as well.

When plants are grown under eCO<sub>2</sub>, their WUE increases since, for each carbon molecule obtained, plants evaporate fewer water molecules. In tomato plants, the current study showed that doubling the CO<sub>2</sub> concentrations resulted in an approximately 35% increase in WUE (Figure 2d), which was the result of a simultaneous increase (~20%) in CO<sub>2</sub> assimilation and a decrease (~30%) in transpiration. Similarly, due to their capacity to concentrate CO<sub>2</sub> in the bundle-sheath cells, C4 plants function as if they were under constant eCO<sub>2</sub> conditions (Chaves & Oliveira, 2004; Rachmilevitch et al., 1999). Accordingly, as a result of their higher WUE, it was previously suggested that C4 plants display smaller vessel diameters and lower  $K_L$  than C3 plants grown in similar habitats, as shown by Kocacinar and Sage (2004).

Moreover, the current study indicates that eCO<sub>2</sub> did not affect the number of vessels per petiole, the total xylem tissue area, or the size of parenchyma cells. In contrast, the individual metaxylem vessels were significantly smaller in the eCO<sub>2</sub>-grown plants, demonstrating that the metaxylem vessel size, not the actual number of vessels, was affected by CO<sub>2</sub> concentrations. In *Pinus radiata*, eCO<sub>2</sub> was shown to decrease the lignin content and the area occupied by xylem vessels,




**FIGURE 2** Tomato leaf carbon assimilation (a), transpiration rate (b), stomatal conductance (c), water use efficiency (d), and leaf vapor pressure deficit (VPD) (e) as affected by CO<sub>2</sub> treatment. Results are expressed as means ± standard error. Bars labeled with an asterisk are significantly different ( $p < .05$ )


indicating an increase in sapwood density at eCO<sub>2</sub> due to thicker tracheid walls (Atwell et al., 2003). In addition, eCO<sub>2</sub> reduced stem hydraulic conductivity in oaks and eucalyptus, which was attributed to a decrease in vessel diameter (Atwell et al., 2007; Heath et al., 1997).

The current study results suggest that the reduction in leaf water requirement under eCO<sub>2</sub> led to a decrease in the petiole metaxylem vessel diameter, similarly to what occurs under other environmental conditions that limit water conductivity in the foliage, such as drought and salinity (Baum et al., 2000; Hochberg et al., 2015; Jerszurki et al., 2020; Lynch et al., 2014). Since stomatal closure and water conductivity are coupled, it is likely that changes in one will also affect the other (Qaderi et al., 2019). When water consumption is low, there is less need to produce larger vessels, which are prone to embolism. Previous studies in different plant species suggested that eCO<sub>2</sub> either increased (Liu et al., 2020), decreased (Medeiros & Ward, 2013; Rico et al., 2013), or had minimal impact on xylem vessel diameter (Kostiainen et al., 2004). Additionally, it was suggested that under mild drought, smaller vessels can be beneficial in improving drought tolerance, not just due to lower embolism risk, but also because of the lower axial water transportation, which may create a more favorable

soil water status (Liu et al., 2020; Medeiros & Ward, 2013). Due to the intense use of fossil fuels, atmospheric CO<sub>2</sub> levels are steadily rising, and drought and heat episodes are projected to increase in frequency in the near future due to climatic changes (Lobell & Gourdji, 2012). The current study results are supported by the previous work of Kocacinar et al. (2008), which showed that plants operated in elevated-CO<sub>2</sub>-like conditions (C4 plants) developed smaller metaxylem vessels than C3 plants from the same habitats. It was also demonstrated that increasing CO<sub>2</sub> from subambient to ambient results in higher (~20%) cavitation resistance (Rico et al., 2013). This suggests that a rise in atmospheric CO<sub>2</sub> may partially mitigate the negative effects of drought on plants, both directly due to reduced water losses and indirectly by increasing the resistance to cavitation due to small xylem size. The decrease in theoretical conductivity under eCO<sub>2</sub> indicates coordination between vascular plasticity to environmental factors that affect water consumption, such as the rise in atmospheric CO<sub>2</sub>, and the way in which plants balance water conductivity with the ability to resist cavitation. In addition, stomatal conductance is sensitive to changes in leaf VPD (Dai et al., 1992). In our study, leaf VPD differed among the CO<sub>2</sub> treatments; therefore, the reduction in




**FIGURE 3** Average number of xylem vessels in petioles (a), xylem tissue area (b) ( $n = 8$ ), average area of individual metaxylem vessel (c) ( $n = 100-110$ ), transverse sections of plant petioles under ambient CO<sub>2</sub> (d) and elevated CO<sub>2</sub> (e), and average parenchyma cell diameter (f). Arrows show metaxylem vessels as affected by CO<sub>2</sub> treatment. Results are expressed as means  $\pm$  standard error. Bars labeled with an asterisk are significantly different ( $p < .05$ )



**FIGURE 4** Distribution of vessel diameters in petioles of tomato plants grown at either ambient (a) or elevated (b) CO<sub>2</sub>





**FIGURE 5** Leaf theoretical conductivity as affected by CO<sub>2</sub> treatment ( $n = 8$ ). Results are expressed as means  $\pm$  standard error. Bars labeled with an asterisk are significantly different ( $p < .05$ )

stomatal conductance and transpiration is probably due to both a reduction in metaxylem diameter and an increase in leaf VPD. Thus, understanding the inverse impacts of drought and eCO<sub>2</sub> on plant growth and development may be more significant than ever, especially since, globally, C<sub>3</sub> plants constitute 90% of plant species (Ibrahim et al., 2008). This study demonstrated that the rise in CO<sub>2</sub> will also induce the development of smaller xylem vessels in petioles that potentially have a great impact on sensitivity to embolism.

## 5 | CONCLUSIONS

Plants plastically in response to climate change factors occur at all levels, and in our particular case, the vascular system. Our data indicate that tomato, a C<sub>3</sub> model crop, displayed a plastic response in xylem vessel size and theoretical conductivity under eCO<sub>2</sub>, and this reduction in xylem vessel size may be associated with lower leaf water consumption under eCO<sub>2</sub>. The impact of combined climate change factors (heat, drought, and eCO<sub>2</sub>) on the development of the plant's vascular system needs further attention, especially in forest trees, where massive cavitation negatively affects trees' ability to recover and increase the chances of wildfires.

### ACKNOWLEDGMENTS

The authors acknowledge the valuable contributions of Liron Summerfield for all her great help.

### FUNDING INFORMATION

This research was partially supported by the Israeli Ministry of Agriculture and Rural Development (Eugene Kandel Knowledge Centers) as part of the program The Root of the Matter: The root zone knowledge center for leveraging modern agriculture.

### CONFLICT OF INTEREST

The authors declare no conflict of interest associated with the work described in this manuscript.

### ORCID

Juliana Espada Lichston <https://orcid.org/0000-0002-4042-4524>

Shimon Rachmilevitch <https://orcid.org/0000-0003-3600-5949>

### REFERENCES

- Ainsworth, E. A., & Lemmonier, P. (2018). ScienceDirect phloem function: A key to understanding and manipulating plant responses to rising atmospheric [CO<sub>2</sub>]? *Current Opinion in Plant Biology*, 43, 50–56. <https://doi.org/10.1016/j.pbi.2017.12.003>
- Ainsworth, E. A., & Long, S. P. (2005). What have we learned from 15 years of free air CO<sub>2</sub> enrichment (FACE)? A meta-analytic review of the response of photosynthesis, canopy properties and plant production to rising CO<sub>2</sub>. *The New Phytologist*, 179(2), 351–372. <https://doi.org/10.1111/j.1469-8137.2004.01224.x>
- Atwell, B. J., Henery, M. L., Rogers, G. S., et al. (2007). Canopy development and hydraulic function in eucalyptus tereticornis grown in drought in CO<sub>2</sub>-enriched atmospheres. *Functional Plant Biology*, 34(12), 1137–1149.
- Atwell, B. J., Henery, M. L., & Whitehead, D. (2003). Sapwood development in *Pinus radiata* trees grown for three years at ambient and elevated carbon dioxide partial pressures. *Tree Physiology*, 23, 13–21. <https://doi.org/10.1093/treephys/23.1.13>
- Baum, S. F., Tran, P. N., & Silk, W. K. (2000). Effects of salinity on xylem structure and water use in growing leaves of sorghum. *The New Phytologist*, 146, 119–127. <https://doi.org/10.1046/j.1469-8137.2000.00625.x>
- Bloom, A. J., Asensio, J. S. R., Randall, L., Rachmilevitch, S., Cousins, A. B., & Carlisle, E. A. (2012). CO<sub>2</sub> enrichment inhibits shoot nitrate assimilation in C<sub>3</sub> but not C<sub>4</sub> plants and slows growth under nitrate in C<sub>3</sub> plants. *Ecology*, 93, 355–367. <https://doi.org/10.1890/11-0485.1>
- Blum, A. (2009). Effective use of water (EUW) and not water-use efficiency (WUE) is the target of crop yield improvement under drought stress. *F Crop Res*, 112, 119–123. <https://doi.org/10.1016/j.fcr.2009.03.009>
- Brocius, C. A., & Hacke, U. G. (2016). Stomatal conductance scales with petiole xylem traits in *Populus* genotypes. *Functional Plant Biology*, 43(6), 553–562.
- Chaves, M. M., & Oliveira, M. M. (2004). Mechanisms underlying plant resilience to water deficits: Prospects for water-saving agriculture. *Journal of Experimental Botany*, 55, 2365–2384. <https://doi.org/10.1093/jxb/erh269>
- Cochard, H. (2002). Xylem embolism and drought-induced stomatal closure in maize. *Planta*, 215, 466–471. <https://doi.org/10.1007/s00425-002-0766-9>
- Cohen, I., Halpern, M., Yermiyahu, U., Bar-Tal, A., Gendler, T., & Rachmilevitch, S. (2019). CO<sub>2</sub> and nitrogen interaction alters root anatomy, morphology, nitrogen partitioning and photosynthetic acclimation of tomato plants. *Planta*, 250, 1423–1432. <https://doi.org/10.1007/s00425-019-03232-0>
- Cohen, I., Rapaport, T., Berger, R. T., & Rachmilevitch, S. (2018). The effects of elevated CO<sub>2</sub> and nitrogen nutrition on root dynamics. *Plant Science*, 272, 294–300. <https://doi.org/10.1016/j.plantsci.2018.03.034>
- Cohen, I., Zandalinas, S. I., Fritschi, F. B., & Mittler, R. (2020). The impact of water deficit and heat stress combination on the molecular response, physiology, and seed production of soybean. *Physiologia Plantarum*, 172(1–12), 41–52. <https://doi.org/10.1111/ppl.13269>

- Cohen, I., Zandalinas, S. I., Huck, C., Fritschi, F. B., & Mittler, R. (2020). Meta-analysis of drought and heat stress combination impact on crop yield and yield components. *Physiologia Plantarum*, 1–11, 66–76. <https://doi.org/10.1111/ppl.13203>
- Domec, J., Smith, D. D., & McCulloh, K. (2015). A synthesis of the effects of atmospheric carbon dioxide enrichment on plant hydraulics: implications for whole-plant water use efficiency and resistance to drought. *Plant, Cell & Environment*, 40, 921–937. <https://doi.org/10.1111/pce.12843>
- Easlon, H. M., & Bloom, A. J. (2014). Easy leaf area: Automated digital image analysis for rapid and accurate measurement of leaf area. *Applications in Plant Sciences*, 2, 1400033. <https://doi.org/10.3732/apps.1400033>
- Franzaring, J., Weller, S., Schmid, I., & Fangmeier, A. (2011). Growth, senescence and water use efficiency of spring oilseed rape (*Brassica napus* L. cv. Mozart) grown in a factorial combination of nitrogen supply and elevated CO<sub>2</sub>. *Environmental and Experimental Botany*, 72, 284–296. <https://doi.org/10.1016/j.envexpbot.2011.04.003>
- Gillespie, K. M., Xu, F., Richter, K. T., et al. (2012). Greater antioxidant and respiratory metabolism in field-grown soybean exposed to elevated O<sub>3</sub> under both ambient and elevated CO<sub>2</sub>. *Plant, Cell and Environment*, 35, 169–184. <https://doi.org/10.1111/j.1365-3040.2011.02427.x>
- Gleason, S. M., Westoby, M., Jansen, S., Choat, B., Hacke, U. G., Pratt, R. B., & Zanne, A. E. (2016). Weak tradeoff between xylem safety and xylem-specific hydraulic efficiency across the world's woody plant species. *The New Phytologist*, 209, 123–136. <https://doi.org/10.1111/nph.13646>
- Hacke, U. G., & Sperry, J. S. (2001). Functional and ecological xylem anatomy. *Perspect Plant Ecol Evol Syst*, 4, 97–115. <https://doi.org/10.1007/978-3-319-15783-2>
- Hao, G.-Y., Holbrook, N. M., Zwieniecki, M. A., Gutschick, V. P., & BassiriRad, H. (2018). Coordinated responses of plant hydraulic architecture with the reduction of stomatal conductance under elevated CO<sub>2</sub> concentration. *Tree Physiology*, 2, 1041–1052. <https://doi.org/10.1093/treephys/tpy001>
- Heath, J., Kerstiens, G., & Tyree, M. T. (1997). Stem hydraulic conductance of European beech (*Fagus sylvatica* L.) and pedunculate oak (*Quercus robur* L.) grown in elevated CO<sub>2</sub>. *Journal of Experimental Botany*, 48(7), 1487–1489.
- Hochberg, U., Degu, A., Gendler, T., Fait, A., & Rachmilevitch, S. (2015). The variability in the xylem architecture of grapevine petiole and its contribution to hydraulic differences. *Functional Plant Biology*, 42, 357–365. <https://doi.org/10.1071/FP14167>
- Ibrahim, D. G., Gilbert, M. E., Ripley, B. S., & Osborne, C. P. (2008). Seasonal differences in photosynthesis between the C3 and C4 subspecies of *Alloteropsis semialata* are offset by frost and drought. 31, 1038–1050. <https://doi.org/10.1111/j.1365-3040.2008.01815.x>
- Jerszurki, D., Sperling, O., Parthasarathi, T., Lichston, J. E., Yaaran, A., Moshelion, M., Rachmilevitch, S., & Lazarovitch, N. (2020). Wide vessels sustain marginal transpiration flux and do not optimize inefficient gas exchange activity under impaired hydraulic control and salinity. *Physiologia Plantarum*, 170, 60–74. <https://doi.org/10.1111/ppl.13107>
- Kant, S., Seneweera, S., Rodin, J., Materne, M., Burch, D., Rothstein, S. J., & Spangenberg, G. (2012). Improving yield potential in crops under elevated CO<sub>2</sub>: Integrating the photosynthetic and nitrogen utilization efficiencies. *Frontiers in Plant Science*, 3, 1–9. <https://doi.org/10.3389/fpls.2012.00162>
- Kocacinar, F., & Sage, R. F. (2004). Photosynthetic pathway alters hydraulic structure and function in woody plants. *Oecologia*, 139, 214–223. <https://doi.org/10.1007/s00442-004-1517-3>
- Kostiainen, K., Kaakinen, S., Saranpää, P., Sigurdsson, B. D., Linder, S., & Vapaavuori, E. (2004). Effect of elevated [CO<sub>2</sub>] on stem wood properties of mature Norway spruce grown at different soil nutrient availability. *Global Change Biology*, 10, 1526–1538. <https://doi.org/10.1111/j.1365-2486.2004.00821.x>
- Kraus, J.E.; Arduin M (1997) *Manual basico de metodos em morfologia vegetal*. EDUR.
- Leakey, A. D. B., Ainsworth, E. A., Bernacchi, C. J., Rogers, A., Long, S. P., & Ort, D. R. (2009) Elevated CO<sub>2</sub> effects on plant carbon, nitrogen, and water relations: Six important lessons from FACE. *Journal of Experimental Botany* 60:2859–2876. <https://doi.org/10.1093/jxb/erp096>
- Liu, J., Kang, S., Davies, W. J., & Ding, R. (2020). Elevated [CO<sub>2</sub>] alleviates the impacts of water deficit on xylem anatomy and hydraulic properties of maize stems. *Plant, Cell & Environment*, 43, 563–578. <https://doi.org/10.1111/pce.13677>
- Lobell, D. B., & Gourdji, S. M. (2012). The influence of climate change on global crop productivity. *Plant Physiology*, 160, 1686–1697. <https://doi.org/10.1104/pp.112.208298>
- Lynch, J. P., Chimungu, J. G., & Brown, K. M. (2014). Root anatomical phenes associated with water acquisition from drying soil: Targets for crop improvement. *Journal of Experimental Botany*, 65, 6155–6166. <https://doi.org/10.1093/jxb/eru162>
- Medeiros, J. S., & Ward, J. K. (2013). Increasing atmospheric [CO<sub>2</sub>] from glacial to future concentrations affects drought tolerance via impacts on leaves, xylem and their integrated function. *New Phytologist*, 199(3), 738–748.
- Medrano, H., Tomás, M., Martorell, S., Flexas, J., Hernández, E., Rosselló, J., Pou, A., Escalona, J. M., & Bota, J. (2015). From leaf to whole-plant water use efficiency (WUE) in complex canopies: Limitations of leaf WUE as a selection target. *Crop J*, 3, 220–228. <https://doi.org/10.1016/j.cj.2015.04.002>
- Mohamed, S. J., Jellings, A. J., & Fuller, M. P. (2013). Positive effects of elevated CO<sub>2</sub> and its interaction with nitrogen on safflower physiology and growth. *Agronomy for Sustainable Development*, 33, 497–505. <https://doi.org/10.1007/s13593-013-0137-x>
- Paudel, I., Halpern, M., Wagner, Y., Raveh, E., Yermiyahu, U., Hoch, G., & Klein, T. (2018). Elevated CO<sub>2</sub> compensates for drought effects in lemon saplings via stomatal downregulation, increased soil moisture, and increased wood carbon storage. *Environmental and Experimental Botany*, 148, 117–127. <https://doi.org/10.1016/j.envexpbot.2018.01.004>
- Qaderi, M. M., Martel, A. B., & Dixon, S. L. (2019) Environmental factors influence plant vascular system and water regulation. *Plants*, 8, 1–23. <https://doi.org/10.3390/plants8030065>
- Rachmilevitch, S., Reuveni, J., Percy, R. W., & Gale, J. (1999). A high level of atmospheric oxygen, as occurred toward the end of the Cretaceous period, increases leaf diffusion conductance. *Journal of Experimental Botany*, 50, 869–872. <https://doi.org/10.1093/jxb/50.335.869>
- Rico, C., Pittermann, J., Polley, H. W., et al. (2013). The effect of subambient to elevated atmospheric CO<sub>2</sub> concentration on vascular function in *Helianthus annuus*: Implications for plant response to climate change. *New Phytologist*, 199(4), 956–965.
- Rodríguez-Gamir, J., Intrigliolo, D. S., Primo-Millo, E., & Forner-Giner, M. A. (2010). Relationships between xylem anatomy, root hydraulic conductivity, leaf/root ratio and transpiration in citrus trees on different rootstocks. *Physiologia Plantarum*, 139, 159–169. <https://doi.org/10.1111/j.1399-3054.2010.01351.x>
- Růžička, K., Ursache, R., Hejálko, J., & Helariutta, Y. (2015). Xylem development - from the cradle to the grave. *The New Phytologist*, 207, 519–535. <https://doi.org/10.1111/nph.13383>
- Shtein, I., Hayat, Y., Munitz, S., Harcavi, E., Akerman, M., Drori, E., Schwartz, A., & Netzer, Y. (2016). From structural constraints to hydraulic function in three *Vitis* rootstocks. *Trees*, 1–11, 851–861. <https://doi.org/10.1007/s00468-016-1510-6>
- Strock, C. F., Burridge, J. D., Niemiec, M. D., Brown, K. M., & Lynch, J. P. (2020). Root metaxylem and architecture phenotypes integrate to





- regulate water use under drought stress. *Plant, Cell & Environment*, 44(1), 49–67. <https://doi.org/10.1111/pce.13875>
- Tyree, M. T., Davis, S. D., & Cochard, H. (1994). Biophysical perspectives of xylem evolution: Is there a tradeoff of hydraulic efficiency for vulnerability to dysfunction? *IAWA Journal*, 15, 335–360. <https://doi.org/10.1163/22941932-90001369>
- Wagner, K. R., Ewers, F. W., & Davis, S. D. (1998). Tradeoffs between hydraulic efficiency and mechanical strength in the stems of four co-occurring species of chaparral shrubs. *Oecologia*, 117, 53–62. <https://doi.org/10.1007/s004420050631>
- Wu, D.-X., Wang, G.-X., Bai, Y.-F., & Liao, J.-X. (2004). Effects of elevated CO<sub>2</sub> concentration on growth, water use, yield and grain quality of wheat under two soil water levels. *Agriculture, Ecosystems and Environment*, 104, 493–507. <https://doi.org/10.1016/j.agee.2004.01.018>
- Wullschleger, S. D., Tschaplinski, T. J., & Norby, R. J. (2002). Plant water relations at elevated CO<sub>2</sub>—Implications for water-limited environments. *Plant, Cell and Environment*, 25, 319–331. <https://doi.org/10.1046/j.1365-3040.2002.00796.x>

**How to cite this article:** Cohen, I., Lichston, J. E., de Macêdo, C. E. C., & Rachmilevitch, S. (2022). Leaf coordination between petiole vascular development and water demand in response to elevated CO<sub>2</sub> in tomato plants. *Plant Direct*, 6(1), e371. <https://doi.org/10.1002/pld3.371>