

RESEARCH ARTICLE

Personalized expression of bitter ‘taste’ receptors in human skin

Lauren Shaw¹, Corrine Mansfield¹, Lauren Colquitt¹, Cailu Lin¹, Jaime Ferreira², Jaime Emmetsberger², Danielle R. Reed^{1*}

1 Monell Chemical Senses Center, Philadelphia, PA, United States of America, **2** Estee Lauder Companies, Inc., Melville, NY, United States of America

* reed@monell.org



OPEN ACCESS

Citation: Shaw L, Mansfield C, Colquitt L, Lin C, Ferreira J, Emmetsberger J, et al. (2018) Personalized expression of bitter ‘taste’ receptors in human skin. PLoS ONE 13(10): e0205322. <https://doi.org/10.1371/journal.pone.0205322>

Editor: Maik Behrens, Leibniz-Institute for Food Systems Biology at the TU Munich, GERMANY

Received: July 7, 2018

Accepted: September 24, 2018

Published: October 17, 2018

Copyright: © 2018 Shaw et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All original data are available from Center for Open Science (osf.io/6azgj) and all data from GTEx is available through dbGAP (accession number phs000424.v7.p2).

Funding: This research was funded in part by the Penn Skin Biology and Diseases Resource-based Center with grant funding from the National Institute of Arthritis and Musculoskeletal and Skin Diseases (P30-AR069589), Monell Institutional Funds, and a grant from Estée Lauder (<https://www.elcompanies.com/>). The funders provided feedback on study design, data collection and

Abstract

The integumentary (i.e., skin) and gustatory systems both function to protect the human body and are a first point of contact with poisons and pathogens. These systems may share a similar protective mechanism because, as we show here, both human taste and skin cells express mRNA for bitter ‘taste’ receptors (*TAS2Rs*). We used gene-specific methods to measure mRNA from all known bitter receptor genes in adult human skin from freshly biopsied samples and from samples collected at autopsy from the Genotype-Tissue Expression project. Human skin expressed some but not all *TAS2Rs*, and for those that were expressed, the relative amounts differed markedly among individuals. For some *TAS2Rs*, mRNA abundance was related to presumed sun exposure based on the location from which the skin sample was collected (*TAS2R14*, *TAS2R30*, *TAS2R42*, and *TAS2R60*), sex (*TAS2R3*, *TAS2R4*, *TAS2R8*, *TAS2R9*, *TAS2R14*, and *TAS2R60*), and age (*TAS2R5*), although these effects were not large. These findings contribute to our understanding of extraoral expression of chemosensory receptors.

Introduction

Humans have at least five widely accepted types of taste receptors: salty, sour, sweet, bitter, and umami. The bitter receptors, called taste receptor type 2 (*TAS2R*), are G protein-coupled receptors that protect humans from ingesting toxins [1]. In the gustatory pathway when bitter compounds bind to a *TAS2R* protein on a taste cell, a conformational change of the protein elicits a signaling cascade. This indirectly induces the release of intracellular calcium, which leads to depolarization and neurotransmitter release, thereby activating sensory neurons that send signals to the central nervous system for bitter perception [2]. Humans have 25 bitter receptors, the *TAS2R* proteins, that are encoded by the *TAS2R* genes located on chromosomes 5, 7, and 12 (Fig 1).

Recently, scientists have identified bitter receptors in locations of the body other than the taste cells. This expression and activation of extragustatory *TAS2Rs* will not lead to taste perception, but instead will elicit distinct cell-type-specific physiological responses. The results of several studies have demonstrated that the extraoral expression of *TAS2Rs* is involved in or

analysis, provided advice on the decision to publish, and helped prepare the manuscript.

Competing interests: The nature of our funding source, in this case, Estée Lauder, does not alter our adherence to PLOS ONE policies on sharing data and materials.

regulate important biological processes germane to the nature of the tissue in which they reside. Bitter receptors have been implicated in the relaxation of smooth muscle, vasoconstriction, gut motility, bronchodilation, nutrient sensing, insulin release, and the release of the antimicrobial peptide, β -defensin [3–7]. As an example, studies performed by Lee *et al.* demonstrated that susceptibility to upper respiratory infection depends on an inborn genotype within one of these bitter receptor genes (*TAS2R38*). Gram-negative bacteria secrete a quorum-sensing molecule that is an agonist of the *TAS2R38* receptor. People with non-functional alleles of this receptor are more susceptible to sinonasal infection because of impairments in this bactericidal pathway [8]. The broader implications of this result are that bitter receptors expressed in extraoral areas may be involved in innate immunity.

Building on this observation, we conducted a study to assess the gene expression patterns of all 25 *TAS2R* genes in skin, since it is a barrier organ and a first line of defense against invading pathogens, presenting both innate and adaptive immune functions. In addition, at least one cell type in human skin (keratinocytes) expresses olfactory receptors, which are similar to bitter taste receptors [9]. Other investigators have measured mRNA expression of a few *TAS2R* genes in skin with conflicting results, perhaps owing to lack of appropriate controls against genomic DNA contamination [10, 11]. Here, we combine results from a smaller biopsy study using quantitative PCR (qPCR) and appropriate controls with a larger autopsy study using an RNA-seq method to get a more complete understanding of *TAS2R* mRNA expression patterns in human skin.

Results

Sample integrity

RNA and DNA were extracted from 15 whole skin samples provided by the University of Pennsylvania Department of Dermatology (Tables 1 and 2) and from one fungiform taste papilla (FP) biopsy obtained from a separate donor as a representative of taste tissue. One sample (004) did not produce viable RNA (RNA integrity number equivalents = 1.0) and was eliminated from the study. Using the remaining RNA samples, cDNA was synthesized and tested for the presence of unwanted genomic DNA using the Abelson 1 (*ABL1*) gene [12]. This is a necessary step since the *TAS2R* protein-coding sequences are within single exons, and *TAS2R* primers cannot be designed to differentiate between genomic DNA and cDNA. Based on the results, three of the samples (005, 006, and 007) were unlikely to contain cDNA because they did not express this gene, and two (009 and 014) had residual genomic DNA after a second DNase treatment (S1 Fig). These five samples were eliminated from the study.

Samples obtained from Penn Dermatology after Mohs surgery vary in size because the procedure requires surgeons to continue removing tissue until all cancerous cells are gone and only healthy tissue remains. The depth of the surgery therefore varies by individual. The samples obtained in this study consist of healthy skin that was removed to properly close the wound at the end of the procedure. Thus, each biopsy sample is unique [13]. To characterize the skin layers and cell types represented in the biopsy samples from Penn Dermatology, qPCR was performed for seven skin-layer- and cell-type-specific markers, standardized to *GAPDH*. As expected, biopsy samples differed in the relative abundance of cell-layer markers (Fig 2)[14, 15].

PCR amplification

To investigate whether bitter taste receptor mRNA is expressed in human skin, PCR experiments were performed with two technical replicates for each of the 25 *TAS2R* genes (S2–S26 Figs), which were compared against two positive controls: (a) genomic DNA from skin and (b)




Fig 1. Bitter receptor locations in the human genome. The location of *TAS2R* genes on human chromosomes 5, 7, and 12 marked by red bars.

<https://doi.org/10.1371/journal.pone.0205322.g001>

fungiform papillae cDNA. Of the 25 *TAS2R* genes, only three showed no expression (*TAS2R1*, 7, and 8), 19 showed variable expression (*TAS2R3*, 4, 5, 9, 13, 14, 16, 20, 31, 38–43, 45, 46, 50,

Table 1. Penn Dermatology subject characteristics.

Sample No.	Age	Gender	Site
001	64	F	Face
002	81	M	Cheek
003	52	M	Scalp
004 [#]	75	M	Neck
005 [#]	62	M	Left temple
006 [#]	63	F	Right cheek
007 [#]	46	M	Left temple
008	70	M	Right cheek
009 [#]	54	M	Nose
010	87	M	Leg
011	58	F	Left posterior thigh
012	56	M	Right cheek
013	52	M	Right leg
014 [#]	55	F	Left supraclavicular
015	55	M	Eyebrow

Individual information for skin samples obtained from Penn Dermatology.

[#]Samples that did not pass our sample integrity tests.

<https://doi.org/10.1371/journal.pone.0205322.t001>

Table 2. Combined subject characteristics.

Characteristic	Group	N subjects Penn Dermatology (N = 9)	N subjects GTEx (N = 914)
Age	20–29	0	68
	30–39	0	70
	40–49	0	150
	50–59	5	300
	60+	4	326
Sex	F	2	311
	M	7	603
Sun Exposure	Yes	9	508
	No	0	406

Summary of demographics of nine viable skin biopsies obtained from Penn Dermatology and of 914 skin biopsies from the GTEx data set. All samples from Penn Dermatology were presumed to be sun-exposed based on the physical location of the sample, e.g., cheek; the GTEx consortium identified the samples from the lower leg as sun-exposed whereas those from the skin of the suprapubic region are identified as not sun-exposed. However, we caution that skin location is a proxy measure of long-term sun-exposure and may not be accurate in all cases.

<https://doi.org/10.1371/journal.pone.0205322.t002>

and 60), and three showed universal expression (*TAS2R10*, 19, and 30) (Fig 3). The genomic DNA positive controls were amplified in every case; however, there was variability in *TAS2R* expression in the FP, suggesting that *TAS2Rs* are expressed at low levels even in taste tissue. This low abundance may explain the variability of expression between technical replicates, as shown in S2–S26 Figs and summarized by the yellow cells in Fig 3. PCR experiments were also performed for *GNAT3*, a gene encoding for the α -subunit of the taste-associated G protein gustducin, and keratin 10 (*KRT10*), a positive epithelial marker (Fig 3 and S27 and S28 Figs). *GNAT3* was detected in taste tissue, as expected, and in four skin samples (002, 003, 008, and 015), perhaps suggesting some similarity between the pathway elicited in skin and the initial steps of the gustatory pathway. As anticipated, *KRT10* was detected in FP and all skin samples. All primers are listed in Table 3.


Quantitative PCR analysis

To quantify mRNA abundance, qPCR was performed on each of the 25 *TAS2R* genes standardized to the housekeeping gene *GAPDH* (Fig 4). The results show variable expression of the *TAS2R* genes across samples, which was expected based on the results of the PCR amplification experiments. The taste tissue sample showed variable expression across receptor type. We also confirmed some expression of *GNAT3* in samples using qPCR standardized to the housekeeping gene *GAPDH* (Fig 4).

Taste-related genes are minimally expressed even in taste tissue, and our agarose gel PCR and qPCR results do not correlate exactly as expected, as is the case when expression of mRNA is near the level of detection [16]. It is possible that individual variation seen here may be due in some part to technical aspects of the biopsy procedure. Despite these limitations, these results suggested that a study of *TAS2R* mRNA expression with a larger sample size was warranted. To do so, we turned to a large and publicly available RNA-seq data set.

GTEx data analysis

After appropriate approvals, we obtained RNA-seq expression data from the Genotype-Tissue Expression project (GTEx; #12732: Bitter receptor gene expression: patterns across tissues). The data were measured at the gene level in RPKM units (reads per kilobase of transcript per million mapped reads) and we extracted the expression data for 25 bitter receptor genes. The



Location	Cell Type:	Enriched in layer:	Marker
Epidermis	Corneocyte	Stratum corneum	CDSN
	Keratinocyte	Stratum granulosum	FLG
	Keratinocyte	Stratum spinosum	DKMN
	Keratinocyte	All but basal layer	KRT10
	Melanocyte	Stratum basale	TYR
Dermis	Collagen	Dermis papillary	COL4A1
	Collagen	Dermis reticulum	COL1A1

Fig 2. Quantification of skin-specific gene expression—qPCR results from cDNA of FP and skin samples. Data are from amplification of skin-specific markers characterized in the table [14, 15]. Data from all markers are represented in order of skin layer for each individual biopsy, with the top of the epidermis (*CDSN*) as the lightest bar section and the bottom of the dermis (*COL1A1*) as the darkest bar section. Results were standardized to the housekeeping gene *GAPDH* and expressed as $2^{-\Delta\Delta C_t}$.

<https://doi.org/10.1371/journal.pone.0205322.g002>

data analyzed consisted of 914 skin samples that varied in location (presumed to be sun-exposed from lower leg or not-sun-exposed from suprapubic region), sex, and age (Tables 2 and 4).

This data set was used because RNA-seq provides more accurate detection of low-abundance transcripts and because it provided a large sample size of tissue collected following a standard procedure. There was heterogeneity of variance between the *TAS2R* genes, but the most highly expressed bitter receptor genes were *TAS2R5*, 14, 20, and 4 (Fig 5). For statistical analysis of sun-exposure, we considered only subjects who had donated both sun-exposed and not sun-exposed tissue (n = 299) and performed Kruskal-Wallis tests to detect differences in the distribution of gene expression levels based on sun-exposure. Results based on tissue type indicated significantly lower expression levels in sun-exposed skin for *TAS2R14* (Median diff. = 0.084,




Fig 3. Results from two rounds of PCR. Each column is labeled by a gene, with members of the *TAS2R* family in the order of location on human chromosomes. Each row is labeled by a sample ID, where 'gDNA' represents genomic DNA (positive control), 'FP' represents taste tissue, and 'W' represents water (a negative control). Green box, bands in both experiments; yellow box, bands in one experiment; red box, no bands. * indicates that there was only one PCR experiment for that gene.

<https://doi.org/10.1371/journal.pone.0205322.g003>

$p < 0.05$), *TAS2R30* (Median diff. = 0.009, $p < 0.01$), and *TAS2R42* (Median diff = 0, $p < 0.05$), but significantly higher expression levels in sun-exposed skin for *TAS2R60* (Median diff = 0.046, $p < 0.0001$) (Fig 6 and S1 Table). We also observed a small sex difference in mRNA expression. In skin from the suprapubic area, females' expression was significantly higher for *TAS2R3* (Median diff. = 0.034, $p < 0.01$), *TAS2R4* (Median diff. = 0.126, $p < 0.01$), and *TAS2R8* (Median diff = 0, $p < 0.05$) (Fig 7, S2 Table). In skin from the lower leg, females' expression was significantly lower for *TAS2R3* (Median diff. = 0.023, $p < 0.05$), *TAS2R9* (Median diff = 0, $p < 0.01$), and *TAS2R14* (Median diff = 0.080, $p < 0.01$) (Fig 7, S3 Table). Finally, there was a positive correlation between increasing age and expression of *TAS2R5* gene but only in not-sun-exposed skin ($p = 0.001$) (Fig 8).

Discussion

Previous studies have shown bitter taste receptor expression in many tissues, including the airway, gastrointestinal tract, and testes [2]. Here, we provide a comprehensive analysis of bitter taste receptor expression in skin using two types of skin samples and three methods of analysis. We show a large-scale assessment of expression of all 25 *TAS2R* genes in adult human skin. By compounding our data, it was discovered that bitter receptors can be detected in the skin, though not ubiquitously. The pattern of results suggests an association between *TAS2R* expression and chromosomal location. For instance, there is no expression of the *TAS2R* gene on chromosome 5 and little to no expression of the first few *TAS2Rs* on chromosome 12. We found that some bitter receptors are not expressed at all, some are variably expressed among people, and some are expressed in almost all skin samples we tested. Variability in more highly expressed receptors is correlated with skin location (presumed-sun-exposed vs. non-exposed), sex, and age. Expression of the taste-related gene *GNAT3* suggests that these receptors are functional in the skin and that the pathway may be G protein-dependent. Although we may speculate on the significance of these findings, we are not yet able to determine causation between the factors stated above and expression level. We must further clarify that sun

Table 3. Primer sequences.

Primer	Sequences (5'-3')	DNA (bp)	RNA (bp)
TAS2R1	F: TGTGGTGGTGAATGGCATTG R: CAGCACTTACTGTGGAGGAGAAC	813	813
TAS2R3	F: ACACATGATTCAGGGATAAATAATGCAAA R: TTAGCCATCTTGGTTTTGGTAGGAAATT	575	575
TAS2R4	F: TACAGTGGTCAATTGCAAACTTGG R: AATGTCTCGGAGAGTAAAGGGTGG	749	749
TAS2R5	F: TGGTCCTCATATAACCTCATTATCCTGG R: CTGCCATGAGTGTCTCCA	667	667
TAS2R7	F: TGTTTTATATTGGTGCTATATCCAGATGTCTATGC R: GGATAAATGAATGACTTGAGGGGTAGATTAGAG	658	658
TAS2R8	F: CAATTTAGTTATCGCCAGAATTTGTTTGATC R: TTATTTAAACAATTAATAAGTGAGTGACCCAAGG	723	723
TAS2R9	F: TGAATTGACCATAGGGATTTGGG R: ATAATTAGAATGAATGAATGGCTTGATGG	807	807
TAS2R10	F: GACTTGTAACACTGCATTGACTGTGCC R: AAAGAGGCTTGCTTTAGCTTGCTG	783	783
TAS2R13	F: GGGTCAGTAAAAGAGAGCTGTCTCTC R: ATCAGAAGAAAGGAGTGGCTTGAAG	742	742
TAS2R14	F: GCTTTGGCAATCTCTCGAATTAGC R: CTCTAAATCTTTGTGACCTGAGGGC	796	796
TAS2R16	F: CCTGGGAATTTTTAATATCCTTACATTCTGGT R: GAAGCGCGCTTTCATGCTT	419	419
TAS2R19	F: GGTTTACTCTGGGTCATGTTATTC R: TTTGCTCTGCTGTGCTCCTAAG	606	606
TAS2R20	F: GCACTGATAAATTTTCATTGCCTGG R: TTGTTCCCCCAAATCAGAATGAAT	770	770
TAS2R30	F: GGTGTTATTAATACTACATTGGTATGCAACTC R: AAGACAGGTTGCTTTTCCAGC	603	603
TAS2R31	F: CATTGGTAAATTCATTGAGC R: GATATCATTATGGACAGAAAGTAAAC	661	661
TAS2R38	F: ACAGTGATTGTGTGCTGCTG R: GCTCTCCTCAACTTGGCATT	766	766
TAS2R39	F: TGTCGCCATTTCTCATCACCTTA R: ATGAGTGGCTGGCAGGGTAG	841	841
TAS2R40	F: AGAGTGCATCACTGGCATCCTT R: GAGGATGAGAAAGTAGCTGGTGGC	685	685
TAS2R41	F: GGTGCTGCCCTTGGATATGA R: TGAAGATGAGGATGAAGGGATGG	738	738
TAS2R42	F: ATGGCCACCGAATTGGACA R: GCTTGCTGTTTCCCAGAATGAG	871	871
TAS2R43	F: GGTCTCCAGAGTTGGTTTTGC R: TCTTGTTCCTCCAAATCAGG	698	698
TAS2R45	F: CTCCTTTGCTGACCAAATGTGC R: GAACGGGTGGGCTGAAGAAC	709	709
TAS2R46	F: GAGTTGAATCCAGCTTTTAAC R: ATAGCTGAATGCAATAGCTTC	606	606
TAS2R50	F: GGTAATTTTCATTGACTGGGTGAAGAG R: CCTTGCTAACCATGACAACCTGGG	710	710
TAS2R60	F: CAGGCAATGGCTTCATCACTG R: TCCCACACCCAGAATTTAAAGTCC	748	748
ABL1	F: AGCATCTGACTTTGAGCC R: CCCATTGTGATTATAGCCTAAGAC	793	193
KRT10	F: CCTTCGAAATGTGTCCACTGG R: CAGGGATTGTTTCAAGGCCA	—	290

(Continued)

Table 3. (Continued)

Primer	Sequences (5'-3')	DNA (bp)	RNA (bp)
GNAT3	F: TCTGGGTATGTGCCAAATGA R: GGCCAGTGTATTCTGGAAA	—	386

The oligonucleotide sequences and the corresponding amplicon sizes are given for genomic DNA and cDNA. F, Forward; R, reverse; bp, base pairs.

<https://doi.org/10.1371/journal.pone.0205322.t003>

exposure has acute and long-term effects and our characterization of sun exposure based on location refers only to presumed long-term exposure. Future studies should focus on determining how any of these factors including sex might directly alter *TAS2R* expression in the skin.

The role of bitter receptors in the skin may become apparent after exploring the most highly expressed receptors and their known agonists. Some *TAS2R* proteins are promiscuous and bind to a wide variety of substances, whereas others have more specificity and bind to one or a few known substances. The protein products of *TAS2R5* and *TAS2R20*, two of the most highly expressed genes in the GTEx data set, are narrowly tuned and recognize one to three of 104 known bitter compounds [10]. *TAS2R4*, the product of *TAS2R4*, another highly expressed gene in this study, is intermediate and binds to 6–16 known bitter compounds. Finally, the




Fig 4. Quantification of bitter taste-related gene expression—qPCR results from cDNA of skin samples after amplification for genes of interest. cDNA was amplified with primers for *GNAT3*, *KRT10*, and the 25 *TAS2R* genes. Data were standardized to the housekeeping gene *GAPDH*, and $2^{-\Delta\Delta Ct}$ was calculated. Results were plotted with individual values in gray and mean across all subjects in red ($n = 9$). Data points for the FP sample are in blue.

<https://doi.org/10.1371/journal.pone.0205322.g004>

Table 4. GTEEx subject characteristics.

<i>Skin sample type</i>	<i>N subjects</i>
Sun-exposed (lower leg)	209
Not sun-exposed (suprapubic area)	107
Both	299
Total	914

Information about post-mortem tissue samples donated by individuals from the Genotype-Tissue Expression project (GTEEx) data set.

<https://doi.org/10.1371/journal.pone.0205322.t004>

TAS2R14 product, *TAS2R14* is broadly tuned and binds to 33 known bitter substances, including synthetic medicinal compounds [17, 18]. Interestingly, *TAS2R38*, the gene for the bitter receptor that enhances innate immunity of the upper respiratory system by recognizing bacteria [8], is rarely or never expressed in skin. We do not know whether the agonists for bitter receptors in skin are endogenous compounds, a pathogen product, or some other exogenous ligand. Further experiments should investigate the cellular response in skin when exposed to compounds similar to known agonists of these bitter receptors to learn more about their potential functions.

Determining the cellular expression of *TAS2R* proteins in skin is an important next step. Bitter receptors are typically expressed in cells known to have chemosensory functions and these cell types are typically sparsely distributed (nose, gut, and tongue). Although we do not know which cell type in human skin expresses *TAS2R* mRNA, previous studies suggest that they may be in the epidermis, and potentially expressed by keratinocytes [10, 11]. There may also be previously uncharacterized cell types in human skin similar to solitary chemosensory cells that express bitter receptors [19], where we speculate that they may function in innate immunity, wound healing, and/or differentiation. Future studies should attempt to determine the localization of *TAS2Rs* in skin potentially through immunocytochemistry, which would require validating human *TAS2R* antibodies, or *in situ* mRNA hybridization.

Materials and methods

Sample collection and DNA/RNA extraction

Staff at the University of Pennsylvania Department of Dermatology collected healthy skin from 15 Mohs surgery patients for this study ($n = 4$ female/11 male; mean age, 62 ± 11.24 years). The Mohs procedure is used to remove cancerous skin and requires removal of additional healthy skin to facilitate proper closure of the wound [13]. We received this additional healthy skin on the day of its removal. The information obtained about each subject was provided by the department and is summarized in Tables 1 and 2. Removal location was provided and based on that information as well as the proximity to cancerous skin we presumed that all samples should be considered sun-exposed. We also obtained one FP biopsy from the tongue of a separate donor as a positive control for *TAS2R* expression. FP were removed from the surface of the tongue using curved spring micro-scissors [20]. The papillae and skin tissue (0.5 mg) were mechanically homogenized and DNA and RNA was extracted using the Zymo Duet DNA/RNA MiniPrep Plus kit, following the protocol for solid tissue. DNA and RNA was quantified with the Thermo Fisher Scientific NanoDrop 1000 Spectrophotometer and measured RNA degradation through RNA integrity number equivalents generated by the Agilent TapeStation and High Sensitivity ScreenTape Assay. The RNA underwent an extra DNase treatment using the Thermo Fisher TURBO DNA-free Kit; RNA (100ng) in water (5 μ L) was




Fig 5. Expression levels of *TAS2R* genes from RNA-seq obtained from the GTEx database. Data are plotted with individual RPKM values in gray points and mean across all samples in red lines ($N = 914$).

<https://doi.org/10.1371/journal.pone.0205322.g005>

then reverse transcribed into cDNA using the NuGEN Ovation RNA Amplification System V2 protocol, purified with the QIAquick PCR Purification Kit, and again quantified. The Institutional Review Board at the University of Pennsylvania approved the collection of skin biopsies for this use.

Primers and PCR amplification

Primer sets for *KRT10* and *GNAT3* were designed using the NCBI Primer-BLAST tool. The *ABL1* primers are designed to span introns, leading to expected bands at 793 base pairs for genomic DNA and 193 base pairs for cDNA [12]. Primer sets for all 25 *TAS2R* genes have been previously published [11]. PCR reactions using primers listed in Table 3 (Invitrogen, Carlsbad, CA, USA) were performed according to the Invitrogen Platinum Taq Green Hot Start DNA Polymerase protocol with a 1 μ L template. The total amount of genomic DNA from each sample was 10 ng, and the total amount of cDNA from each sample was 50 ng. A StepOne Thermocycler was used according to the following profile: one cycle of 4 min at 94°C; 40 cycles of 1 min at 94°C, 1 min at 55°C, 2 min at 72°C; one cycle of a final hold at 4°C. Fragments were




Fig 6. Effect of sun exposure on *TAS2R* expression from the GTEx data. Expression levels of bitter receptor genes from RNA-seq obtained from the GTEx database are separated based on sun exposure. Data are plotted as mean and SD across subjects that donated both skin sample types (N = 299 for each sample type). * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$; **** $p < 0.0001$.

<https://doi.org/10.1371/journal.pone.0205322.g006>

detected by staining with SYBR Green Safe. The PCR products were electrophoresed on a 1.0% gel in TAE buffer.

Real-time qPCR

Real Time qPCR reactions were performed in 10 μ L of water in a 384-well plate according to the TaqMan Fast Advanced Master Mix protocol with 1 μ L template and run in triplicate. The total amount of cDNA from each sample was 50 ng. Primers for skin-specific markers, *TAS2Rs*, and a pre-developed endogenous control, *GAPDH* were used. PCR reactions were performed with the QuantStudio 12K Flex Real-Time PCR machine and amplification was evaluated by comparative analysis based on cycle threshold [21]. Graphs were generated using GraphPad Prism 7 (La Jolla, CA, USA).

GTEx database analysis

RNA-seq data from 914 post-mortem tissue samples were provided by the GTEx project (Tables 2 and 4), with information about each sample, including the age and sex of the tissue donor, and tissue type (sun-exposed skin from lower leg or sun-unexposed skin from suprapubic region). Details on skin sample removal, sectioning, and preservation can be found in the GTEx Tissue Harvesting Work Instruction (<https://biospecimens.cancer.gov>). For the 25 bitter receptor genes from 914 samples, the gene expression RPKM values were normalized for all samples of the same tissue type. Due to the heterogeneity of variance between the genes, we used the non-parametric Kruskal-Wallis test to detect differences in the distribution of expression levels based on effects of sun exposure and of sex within each tissue type (S1–S3 Tables). For analysis of effects of sun exposure, only data from the 299 subjects that donated both types




Fig 7. Effect of sex on TAS2R expression from the GTEx data. Expression levels of bitter receptor genes from RNA-seq obtained from the GTEx database are separated based on sex and presumed sun exposure. Data are plotted as mean and SD across males (N = 603) and females (N = 311) that donated both skin sample types. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$; **** $p < 0.0001$.

<https://doi.org/10.1371/journal.pone.0205322.g007>

of samples were included. For effects of sex in skin from the lower leg, all 508 tissue samples were included, and from the suprapubic area, all 406 tissue samples were included. Data for sun exposure effects and sex were analyzed in R version 3.4.2, and graphs were generated in GraphPad Prism 7. Effects of age were analyzed via correlation and plotted in R (version 3.4.2) and R-studio (version 1.0.136). We deposited a data analysis script based in R on Github (<https://github.com/DanielleReed/TAS2R38>).




Fig 8. Correlation plots of TAS2R expression against age from the GTEx data. Individual RPKM data are plotted separated by sun exposure and in order of increasing age of the subject for each receptor. R values and p values are given on the corresponding plot.

<https://doi.org/10.1371/journal.pone.0205322.g008>

Supporting information

S1 Fig. Gene expression of *ABLI1*. PCR was performed with genomic DNA from skin (gDNA), a mixture of genomic DNA and cDNA from skin (Mix), cDNA from fungiform papillae (FP), and cDNA from 14 skin samples (001–015). Water was used as a no-template control. The larger band at 793 base pairs (bp) includes introns, and the smaller band at 293 bp does not contain introns. Genomic DNA was used as a positive control for the larger band size. A mix was used as a positive control for both bands. The smear at FP is likely caused by nonspecific binding.

(JPG)

S2 Fig. Gene expression of *TAS2R1*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 813 bp. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.

(JPG)

S3 Fig. Gene expression of *TAS2R3*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 575 bp. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.

(JPG)

S4 Fig. Gene expression of *TAS2R4*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 749 bp. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.

(JPG)

S5 Fig. Gene expression of *TAS2R5*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 667 bp. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.

(JPG)

S6 Fig. Gene expression of *TAS2R7*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 658 bp. The experiment was replicated

(bottom panel) because taste receptors are not abundant and can have variable results.
(JPG)

S7 Fig. Gene expression of *TAS2R8*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 723 bp. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.
(JPG)

S8 Fig. Gene expression of *TAS2R9*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 807 bp. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.
(JPG)

S9 Fig. Gene expression of *TAS2R10*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 783 bp. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.
(JPG)

S10 Fig. Gene expression of *TAS2R13*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 742 bp. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.
(JPG)

S11 Fig. Gene expression of *TAS2R14*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 796 bp. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.
(JPG)

S12 Fig. Gene expression of *TAS2R16*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 419 bp. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.
(JPG)

S13 Fig. Gene expression of *TAS2R19*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 606 bp. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.
(JPG)

S14 Fig. Gene expression of *TAS2R20*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 770 bp. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.
(JPG)

S15 Fig. Gene expression of *TAS2R30*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was

used as a no-template control. The expected band size is 603 bp. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.

(JPG)

S16 Fig. Gene expression of *TAS2R31*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 661 bp. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.

(JPG)

S17 Fig. Gene expression of *TAS2R38*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 766 bp. Multiple bands are likely because of non-specific binding. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.

(JPG)

S18 Fig. Gene expression of *TAS2R39*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 841 bp. The experiment was replicated, but results were omitted because of non-specific binding.

(JPG)

S19 Fig. Gene expression of *TAS2R40*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 685 bp. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.

(JPG)

S20 Fig. Gene expression of *TAS2R41*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 738 bp. Multiple bands are likely because of non-specific binding. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.

(JPG)

S21 Fig. Gene expression of *TAS2R42*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 871 bp. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.

(JPG)

S22 Fig. Gene expression of *TAS2R43*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 698 bp. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.

(JPG)

S23 Fig. Gene expression of *TAS2R45*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 709 bp. Multiple bands are likely because of non-specific binding. The experiment was replicated (bottom panel) because taste

receptors are not abundant and can have variable results.
(JPG)

S24 Fig. Gene expression of *TAS2R46*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 606 bp. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.
(JPG)

S25 Fig. Gene expression of *TAS2R50*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 710 bp. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.
(JPG)

S26 Fig. Gene expression of *TAS2R60*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The expected band size is 748 bp. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.
(JPG)

S27 Fig. Gene expression of *GNAT3*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The primer set is intron-spanning, so there is no expected band size for genomic DNA, while there is an expected band size of 386 bp for cDNA. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.
(JPG)

S28 Fig. Gene expression of *KRT10*. PCR was performed with genomic DNA from skin (gDNA), cDNA from fungiform papillae (FP), and cDNA from nine skin samples. Water was used as a no-template control. The primer set is intron-spanning, so there is no expected band size for genomic DNA and an expected band size of 290 bp for cDNA. The experiment was replicated (bottom panel) because taste receptors are not abundant and can have variable results.
(JPG)

S1 Table. Kruskal-Wallis test statistics for GTEx data comparing effects of presumed sun exposure for each gene of interest (N = 598).
(PDF)

S2 Table. Kruskal-Wallis test statistics for GTEx data comparing effects of sex for each gene of interest in not sun-exposed tissue (N = 406).
(PDF)

S3 Table. Kruskal-Wallis test statistics for GTEx data comparing effects of sex for each gene of interest in sun-exposed tissue (N = 508).
(PDF)

Acknowledgments

Dr. Aimee Payne and the staff at the University of Pennsylvania Department of Dermatology are acknowledged for providing samples and technical assistance. This work was supported in

part by the Penn Skin Biology and Diseases Resource-based Center with grant funding from the National Institute of Arthritis and Musculoskeletal and Skin Diseases (P30-AR069589). The contents are solely the responsibility of the authors and do not necessarily represent the official views of the NIH. We are thankful to Dr. Casey Trimmer for sharing her expertise and providing advice for the study. Dr. Mary Matsui, and Dr. Ed Pelle from the Estée Lauder Companies are also acknowledged for their contribution to our preliminary work and for their insight on the study. We are grateful to Nora Ruth from The Estée Lauder Companies for revising the manuscript.

Author Contributions

Conceptualization: Lauren Shaw, Corrine Mansfield, Jaime Ferreira, Jaime Emmetsberger, Danielle R. Reed.

Data curation: Lauren Shaw, Corrine Mansfield, Cailu Lin, Danielle R. Reed.

Formal analysis: Lauren Shaw, Corrine Mansfield, Cailu Lin, Danielle R. Reed.

Funding acquisition: Lauren Shaw, Jaime Ferreira, Jaime Emmetsberger, Danielle R. Reed.

Investigation: Lauren Shaw, Danielle R. Reed.

Methodology: Lauren Shaw, Corrine Mansfield, Cailu Lin, Danielle R. Reed.

Project administration: Danielle R. Reed.

Resources: Lauren Colquitt, Cailu Lin, Danielle R. Reed.

Software: Cailu Lin.

Supervision: Jaime Ferreira, Jaime Emmetsberger, Danielle R. Reed.

Validation: Lauren Shaw, Danielle R. Reed.

Visualization: Lauren Shaw.

Writing – original draft: Lauren Shaw, Danielle R. Reed.

Writing – review & editing: Lauren Shaw, Corrine Mansfield, Lauren Colquitt, Cailu Lin, Jaime Ferreira, Jaime Emmetsberger, Danielle R. Reed.

References

1. Reed DR, Knaapila A. Genetics of taste and smell: poisons and pleasures. *Prog Mol Biol Transl Sci.* 2010; 94:213–40. Epub 2010/11/03. B978-0-12-375003-7.00008-X [pii] [https://doi.org/10.1016/S1877-1173\(10\)94008-8](https://doi.org/10.1016/S1877-1173(10)94008-8) PMID: 21036327; PubMed Central PMCID: PMC3342754.
2. Lee RJ, Cohen NA. Taste receptors in innate immunity. *Cell Mol Life Sci.* 2015; 72(2):217–36. <https://doi.org/10.1007/s00018-014-1736-7> PMID: 25323130; PubMed Central PMCID: PMC34286424.
3. Zheng K, Lu P, Delpapa E, Bellve K, Deng R, Condon JC, et al. Bitter taste receptors as targets for tocolytics in preterm labor therapy. *FASEB J.* 2017; 31(9):4037–52. <https://doi.org/10.1096/fj.201601323RR> PMID: 28559440; PubMed Central PMCID: PMC5572693.
4. Shah AS, Ben-Shahar Y, Moninger TO, Kline JN, Welsh MJ. Motile cilia of human airway epithelia are chemosensory. *Science.* 2009; 325:1131–4. <https://doi.org/10.1126/science.1173869> PMID: 19628819.
5. Janssen S, Laermans J, Verhulst PJ, Thijs T, Tack J, Depoortere I. Bitter taste receptors and -gustducin regulate the secretion of ghrelin with functional effects on food intake and gastric emptying. *Proc Natl Acad Sci USA.* 2011. <https://doi.org/10.1073/pnas.1011508108> PMID: 21245306
6. Foster SR, Porrello ER, Purdue B, Chan HW, Voigt A, Frenzel S, et al. Expression, regulation and putative nutrient-sensing function of taste GPCRs in the heart. *PLoS One.* 2013; 8(5):e64579. Epub 2013/05/23. <https://doi.org/10.1371/journal.pone.0064579> PMID: 23696900; PubMed Central PMCID: PMC3655793.

7. Jang HJ, Kokrashvili Z, Theodorakis MJ, Carlson OD, Kim BJ, Zhou J, et al. Gut-expressed gustducin and taste receptors regulate secretion of glucagon-like peptide-1. *Proc Natl Acad Sci USA*. 2007; 104(38):15069–74. <https://doi.org/10.1073/pnas.0706890104> PMID: 17724330.
8. Lee RJ, Xiong G, Kofonow JM, Chen B, Lysenko A, Jiang P, et al. T2R38 taste receptor polymorphisms underlie susceptibility to upper respiratory infection. *J Clin Invest*. 2012; 122(11):4145–59. Epub 2012/10/09. <https://doi.org/10.1172/JCI64240> PMID: 23041624; PubMed Central PMCID: PMC3484455.
9. Denda M. Newly discovered olfactory receptors in epidermal keratinocytes are associated with proliferation, migration, and re-epithelialization of keratinocytes. *J Invest Dermatol*. 2014; 134(11):2677–9. <https://doi.org/10.1038/jid.2014.229> PMID: 25318430.
10. Wolfle U, Elsholz FA, Kersten A, Haarhaus B, Muller WE, Schempp CM. Expression and functional activity of the bitter taste receptors TAS2R1 and TAS2R38 in human keratinocytes. *Skin Pharmacol Physiol*. 2015; 28(3):137–46. <https://doi.org/10.1159/000367631> PMID: 25573083.
11. Reszka E, Nowakowska-Swirta E, Kupczyk M, Dudek W, Swierczynska-Machura D, Wittczak T, et al. Expression of bitter taste receptors in the human skin in vitro. *Journal of Clinical Research and Bioethics*. 2015; 6(2).
12. Lion T. Appropriate controls for RT-PCR. *Leukemia*. 1996; 10(11):1843. PMID: 8892692.
13. Cottle WI, Proper S. Mohs' surgery, fresh-tissue technique. Our technique with a review. *J Dermatol Surg Oncol*. 1982; 8(7):576–87. PMID: 6181110.
14. Edqvist PH, Fagerberg L, Hallstrom BM, Danielsson A, Edlund K, Uhlen M, et al. Expression of human skin-specific genes defined by transcriptomics and antibody-based profiling. *J Histochem Cytochem*. 2015; 63(2):129–41. <https://doi.org/10.1369/0022155414562646> PMID: 25411189; PubMed Central PMCID: PMC34305515.
15. Tracy LE, Minasian RA, Cateson EJ. Extracellular matrix and dermal fibroblast function in the healing wound. *Adv Wound Care (New Rochelle)*. 2016; 5(3):119–36. <https://doi.org/10.1089/wound.2014.0561> PMID: 26989578; PubMed Central PMCID: PMC34779293.
16. Behrens M, Foerster S, Staehler F, Raguse JD, Meyerhof W. Gustatory expression pattern of the human TAS2R bitter receptor gene family reveals a heterogenous population of bitter responsive taste receptor cells. *J Neurosci*. 2007; 27(46):12630–40. <https://doi.org/10.1523/JNEUROSCI.1168-07.2007> PMID: 18003842.
17. Lossow K, Hubner S, Roudnitzky N, Slack JP, Pollastro F, Behrens M, et al. Comprehensive analysis of mouse bitter taste receptors reveals different molecular receptive ranges for orthologous receptors in mice and humans. *J Biol Chem*. 2016. <https://doi.org/10.1074/jbc.M116.718544> PMID: 27226572.
18. Meyerhof W, Batram C, Kuhn C, Brockhoff A, Chudoba E, Bufe B, et al. The molecular receptive ranges of human TAS2R bitter taste receptors. *Chem Senses*. 2010; 35(2):157–70. <https://doi.org/10.1093/chemse/bjp092> PMID: 20022913.
19. Tizzano M, Gulbransen BD, Vandenbeuch A, Clapp TR, Herman JP, Sibhatu HM, et al. Nasal chemosensory cells use bitter taste signaling to detect irritants and bacterial signals. *Proc Natl Acad Sci USA*. 2010; 107:3210–5. <https://doi.org/10.1073/pnas.0911934107> PMID: 20133764.
20. Spielman AI, Pepino MY, Feldman R, Brand JG. Technique to collect fungiform (taste) papillae from human tongue. *J Vis Exp*. 2010; 18(42):2201. Epub 2010/10/26. <https://doi.org/10.3791/2201> PMID: 20972388; PubMed Central PMCID: PMC3168239.
21. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods*. 2001; 25(4):402–8. <https://doi.org/10.1006/meth.2001.1262> PMID: 11846609.