

Review

¹³C-NMR Data of Three Important Diterpenes Isolated from *Euphorbia* Species

Qi-Cheng Wu ¹, Yu-Ping Tang ^{1,*}, An-Wei Ding ^{1,*}, Fen-Qiang You ², Li Zhang ¹
and Jin-Ao Duan ²

¹ Jiangsu Key Laboratory for TCM Formulae Research, Nanjing University of Chinese Medicine, Nanjing 210046, China

² Affiliated Hospital, Nanjing University of Chinese Medicine, Kunshan, China

* Authors to whom correspondence should be addressed; E-Mail: yupingtang@njutcm.edu.cn (Y.-P.T.); ltcmf@njutcm.edu.cn (A.-W.D.).

Received: 21 October 2009; in revised form: 2 November 2009 / Accepted: 4 November 2009 /

Published: 6 November 2009

Abstract: *Euphorbia* species are widely distributed plants, many of which are used in folk medicine. Over the past twenty years, they have received considerable phytochemical and biological attention. Their diterpenoid constituents, especially those with abietane, tiglane, ingenane skeletons, are thought to be the main toxicant and bioactive factors. In this work, the utility of ¹³C-NMR spectroscopy for the structural elucidation of these compounds is briefly discussed.

Keywords: *Euphorbia*; diterpene; abietane; tiglane; ingenane; ¹³C-NMR data

1. Introduction

The *Euphorbia* is the largest genus in the plant family Euphorbiaceae, comprising about 2,000 known species [1]. *Euphorbia* are widely distributed throughout both hemispheres and range in morphology from large desert succulents to trees and even some small herbaceous plant types. Researched parts in various *Euphorbia* species include the roots, seeds, latex, lactiferous tubes, stem wood, stem barks, leaves, and whole plants.

Many studies have suggested that these plants have not only therapeutic relevance but that they also display toxicity [2]. Some constituents of *Euphorbia* species may be promising lead compounds for

drug development. Certain *Euphorbia* species have been reported to possess antitumor activity and have been recommended for use as anticancer remedies [3,4]. Their antitumor activity was mainly attributed to the presence of abietane diterpene derivatives, most of which contain lactone structures reported to possess potent antineoplastic activity towards various cancer cell lines [5–9]. Moreover, some *Euphorbia* species have been also used as medicinal plants for the treatment of skin diseases, gonorrhea, migraines, intestinal parasites, warts and for mediating pain perception [10–12]. Many researchers have shown that *Euphorbia* species also possess antiproliferative activity [13], cytotoxicity [14], antimicrobial activity [15], antipyretic-analgesic activity [16], inhibition of HIV-1 viral infection [17], inhibitory activity on the mammalian mitochondrial respiratory chain [18], etc.

As mentioned, there are also some reports of toxicity in *Euphorbia* species. Their toxic substances originate from the milky sap, which is a deterrent to insects and herbivores [19]. Besides, they may possess extreme proinflammatory and tumor promoting toxicities [20,21]. Severe pain and inflammation can result from contact with the eyes, nose, mouth and even skin, which may be due to the activation of protein kinase C enzyme [22]. The toxic constituents of *Euphorbia* species were considered to be a kind of specific diterpenes, globally called phorboids, which comprise tiglane, ingenane and daphnane diterpene derivatives [23,24].

Terpenes, including diterpenes and triterpenes, have been frequently found in *Euphorbia* species. Steroids, cerebrosides, glycerols, phenolics and flavonoids were also isolated from plants of the genus [10], but the compounds most relevant to the toxicity and considerable biological activities in *Euphorbia* are diterpenes, especially those with abietane, tiglane, and ingenane skeletons [10].

Many researchers have suggested that there was a close relationship between the structures and the biological activity, so the structure elucidation is very important for these diterpenes. In this review article, we summarize the ^{13}C -NMR data of these three important diterpene skeleton types of *Euphorbia* species, covering 42 abietanes, 51 ingenanes and 30 tiglianes. The structure-activity relationship and the features on the chemical shifts were also briefly discussed.

2. Abietane Derivates Isolated from *Euphorbia* Species (Table 1)

Most plants of the genus *Euphorbia* contain abietane diterpenoids, which usually have an extra α,β -unsaturated γ -lactone ring located between C-12 and C-13, and some of which have an epoxy ring at C-8 and C-14, or C-11 and C-12, as is the case of **7–14**. Some carbons of these diterpenes, especially C-8, C-14, C-11 and C-12 are frequently substituted by hydroxyl groups or form double bonds. Compounds **32–36** indicate that the 18-Me and C-3 could form a three-membered ring. In addition, some abietanes (**39–42**) without lactone rings were also isolated from the genus *Euphorbia*. Many abietane diterpenoids exhibit inhibitive activity on various types of tumor cells, such as ANA-1, B16, Jurkat cells [25], K562 cells [7] and LNCaP cells [6]. By comparing the active compound **8** with the inactive one **12**, it could be concluded that the C-11/C-12 epoxy ring system was necessary in mediating cytotoxicity. Compounds **2** and **3** are diastereomers, differing only in the stereochemistry at the chiral centers C-8 and C-14, but only compound **2** showed activity, which suggested that the ring C configuration is also crucial for the activity [25]. The α,β -unsaturated lactone is not the only necessary group for the cytotoxic effects, since compounds **3** and **12** do not show cytotoxicity [25]. In addition, the similar compounds **18**, **19**, **26** and **28** were tested in the inhibition of P-glycoprotein transport

activity. The higher inhibitory effect of **26** might be derived from the carbonyl position at C-2, most probably due to the conformational and functional changes in the P-gp induced by the particular structures of helioscopinolides [26].

Table 1. Abietane diterpenoids isolated from *Euphorbia* species.

No	Name	Species	Ref
1	7 β -Hydroxy-8 α ,14-dihydro jolkinolide E	<i>E. terracina</i>	[27]
2	Yuexiandajisu D	<i>E. ebracteolata</i>	[25]
3	Yuexiandajisu E	<i>E. ebracteolata</i>	[25]
4	ent-8 β ,14 α -Dihydroxy-13(15)-ene-16(12 β)-abietanolide	<i>E. wallichii</i>	[28]
5	ent-8 β ,14 β -Dihydroxy-13(15)-ene-16(12 β)-abietanolide	<i>E. wallichii</i>	[28]
6	Ebracteolatanolide B	<i>E. ebracteolata</i>	[25]
7	Ebracteolatanolide A	<i>E. ebracteolata</i>	[25]
8	Jolkinolide B	<i>E. fischeriana</i> , <i>E. sessiliflora</i>	[29] [30]
9	17-Hydroxyjolkinolide B	<i>E. fischeriana</i>	[31]
10	17-Acetoxyjolkinolide B	<i>E. fischeriana</i>	[31]
11	17-Acetoxyjolkinolide A	<i>E. fischeriana</i>	[32]
12	Jolkinolide A	<i>E. wallichii</i> <i>E. fischeriana</i> <i>E. fidjiana</i> <i>E. guyoniana</i>	[28] [29] [33] [34]
13	17-Hydroxyjolkinolide A	<i>E. fischeriana</i> <i>E. fidjiana</i>	[32] [33]
14	3 α -Hydroxyjolkinolide A	<i>E. wallichii</i>	[28]
15	7 β -Hydroxy-ent-abieta-8(14),13(15)-dien-12 α ,16-olide	<i>E. seguieriana</i>	[35]
16	7 β ,9 β -Dihydroxy-ent-abieta-8(14),13(15)-dien-12 α ,16-olide	<i>E. seguieriana</i>	[35]
17	ent-Abieta-8(14),13(15)-dien-16,12-olide [Jolkinolide E]	<i>E. fidjiana</i> <i>E. characias</i> <i>E. guyoniana</i>	[33] [34] [36]
18	Helioscopinolide A	<i>E. pubescens</i> <i>E. semiperfoliata</i> <i>E. helioscopia</i>	[37] [38] [39]
19	Helioscopinolide B	<i>E. pubescens</i> <i>E. semiperfoliata</i> <i>E. helioscopia</i> <i>E. calypttrata</i>	[37] [38] [39] [40]
20	Helioscopinolides H	<i>E. calypttrata</i>	[40]
21	ent-11 α -Hydroxyabieta-8(14),13(15)-dien-16,12 α -olide	<i>E. ebracteolata</i> <i>E. sessiliflora</i> <i>E. fidjiana</i>	[25] [30] [33]
22	ent-12-Hydroxy-12[R]-abieta-8(14),13(15)-dien-16,12-olide	<i>E. sessiliflora</i>	[30]
23	7 β ,11 β ,12 β -Trihydroxy-ent-abieta-8(14),13(15)-dien-16,12-olide	<i>E. fischeriana</i>	[31]
24	Langduin B	<i>E. fischeriana</i>	[32]
25	Helioscopinolide C	<i>E. helioscopia</i>	[39,41]
26	Helioscopinolides F	<i>E. calypttrata</i>	[40]

Table 1. Cont.

No	Name	Species	Ref
27	Helioscopinolide D	<i>E. calyprate</i>	[42]
28	Helioscopinolide E	<i>E. calyprate</i>	[42]
29	Helioscopinolides I	<i>E. calyprata</i>	[40]
30	8 α ,14-Dihydro-7-oxo-jolkinolide E	<i>E. characias</i>	[36]
31	8 α ,14-Dihydro-7-oxohelioscopinolide A [caudicifolin]	<i>E. sessiliflora</i>	[30]
		<i>E. characias</i>	[36]
		<i>E. semiperfoliata</i>	[38]
32	3,4,18 β -Cyclopropa-8 β -hydroxy-14-oxo- <i>ent</i> -abiet-13,15-en-16,12-olide	<i>E. retusa</i>	[43]
33	3,4,18 β -Cyclopropa-14-oxo- <i>ent</i> -abieta-8,9,13,15-dien-16,12-olide	<i>E. retusa</i>	[43]
34	3,4,18 β -Cyclopropa-14-oxo- <i>ent</i> -abieta-7,13,15-dien-16,12-olide	<i>E. retusa</i>	[43]
35	3,4,18 β -Cyclopropa-7-hydroxy-14-oxo- <i>ent</i> -abieta-8,9,13,15-dien-16,12-olide	<i>E. retusa</i>	[43]
36	3,4,18 β -Cyclopropa-14-oxo- <i>ent</i> -abiet-7-en-16,12-olide	<i>E. retusa</i>	[43]
37	<i>ent</i> -16-Hydroxy-13[R]-pimar-8(14)-ene-3,15-dione	<i>E. fidjiana</i>	[33]
38	<i>ent</i> -12 α ,16-Dihydroxy-13[R]-pimar-8(14)-ene-3,15-dione	<i>E. fidjiana</i>	[33]
39	13 β -Hydroxy- <i>ent</i> -abiet-8(14)-en-7-one	<i>E. fischeriana</i>	[31]
40	Methyl 8 β ,11 β -dihydroxy-12-oxo- <i>ent</i> -abieta-13, 15(17)-dien-16-oate	<i>E. portulacoides</i>	[44]
41	11,16-Epoxy- <i>ent</i> -abieta-8,11,15-triene-13,14-dione	<i>E. guyoniana</i>	[34]
42	11-Hydroxy- <i>ent</i> -abieta-8,11,13-trien-15-one	<i>E. guyoniana</i>	[34]






Figure 1. Abietane diterpenoids isolated from *Euphorbia* species.

Figure 1. Cont.



No.	R
8	H
9	OH
10	OAc
11	OAc
12	H
13	OH



No.	R1	R2	R3	R4
15	H	H	OH	H
16	H	H	OH	OH
17	H	H	H	H
18	OH	H	H	H
19	H	OH	H	H
20	OH	H	H	OH
21	H	H	H	α -OH










Figure 1. Cont.



3. Ingenane Derivates Isolated from *Euphorbia* Species (Table 2)

Ingenane diterpenoids have a very unique structural feature: they all have a same 5/7/7/3-tetracyclic ring system and a ketone bridge between C-8 and C-10. There is a double bond between C-1 and C-2 in ring A, and another double bond between C-6 and C-7 in ring B. A β -hydroxyl group is linked to C-4, so ring A/B must be *trans*-joined. Besides, ring D is a cyclopropane ring. Some positions at C-3, C-5, C-13, C-17 and C-20 may be linked to oxygen-substituted residues, such as hydroxyl, acetyl ester, long-chain alkyl ester, benzoyl ester groups, and so forth. This type of diterpenoids have been widely reported in many *Euphorbia* species. Some researchers have shown that these diterpenoids have antinematodal and termiticidal activity [45,46]. There were also reports about toxicity such as tumor promoting and proinflammatory activity [20,47,48]. Studies on the relationships between structure and irritant activity indicate that presence of a hydroxyl on C-20 is crucial for stimulatory properties. Introduction of an acetyl group in the 20-position results in a lower

irritancy [49]. Some 20-deoxyingenol diterpenes induced cell cleavage arrest, but this activity became weak when C-16 had an acyl residue [50]. Acetylation in the 5-position resulted into a considerable depression of irritancy [49]. The skin tumor promoting and irritant activities of the ingenol-3-esters depend on the length of the aliphatic chain in their ester moiety [51]. In addition, the presence of one free hydroxy group at C-3 or C-5 may play an important role in the antinematodal activity [45].

Table 2. Ingenane diterpenoids isolated from *Euphorbia* species.

No	Name	Species	Ref
43	Ingenol	<i>E. kansui</i>	[45]
		<i>E. paralias</i>	[48]
44	13-O-Dodecanoylingenol	<i>E. kansui</i>	[45]
45	17-[(2Z,4E,6Z)-Deca-2,4,6-trienoyloxy] [ingenol]	<i>E. cauducifolia.</i>	[21]
46	20-Eicosanoate	<i>E. iberica</i>	[52]
47	3,5,20-O-Triacetylingenol	<i>E. kansui</i>	[45]
48	17-Hydroxyingenol tetraacetate	<i>E. kamerunica</i>	[53]
49	5,20-O-Diacetyl-3-O-(2'',3''-dimethylbutanoyl)-13-O-dodecanoylingenol	<i>E. kansui</i>	[45]
50	20-Tetradecanoate-ingenol-3,5-diacetate	<i>E. broteri</i>	[54]
51	17-O-Acetyl-3-O-[(Z)-2-methyl-2-butenoyl]-20-deoxy-17-hydroxy-ingenol	<i>E. trigona</i>	[55]
52	20-O-Acetyl-3-O-[(Z)-2-methyl-2-butenoyl]ingenol	<i>E. trigona</i>	[55]
53	5,17,20-O-Triacetyl-3-O-[(Z)-2-methyl-2-butenoyl]-17-hydroxyingenol	<i>E. trigona</i>	[55]
54	3-O-(2,3-Dimethylbutanoyl)-13-O-dodecanoylingenol	<i>E. kansui</i>	[45]
		<i>E. cyparissias</i>	[46]
55	3-O-(2,3-Dimethylbutanoyl)-13-O-decanoylingenol	<i>E. kansui</i>	[45]
		<i>E. cyparissias</i>	[46]
56	3,20-O-Diacetylingenol 5-O-(2'E,4'Z)-tetradecadienoate	<i>E. petiolata</i>	[56]
57	5,20-O-Diacetylingenol 3-O-(2'E,4'Z)-tetradecadienoa	<i>E. petiolata</i>	[56]
58	Ingenol-3-O-(2'E,4'Z)-tetradecadieno	<i>E. petiolata</i>	[56]
59	5,20-O-Isopropylidenyl ingenol 3-O-(2'Z,4'Z)-tetradecadienoate	<i>E. petiolata</i>	[56]
60	20-O-Acetylingenol-3-O-(2"E,4"Z)-decadienoate	<i>E. petiolata</i>	[57]
61	20-Acetyl-ingenol-3-decadienoate	<i>E. broteri</i>	[54]
62	3-Tetradecanoate-ingenol-5,20-diacetate	<i>E. broteri</i>	[54]
63	5-Tetradecanoate-ingenol-3,20-diacetate	<i>E. broteri</i>	[54]
64	17-Benzoyloxy-3-O-(2,3-dimethylbutanoyl)-20-deoxyingenol	<i>E. esula</i>	[58]
65	17-Benzoyloxy-3-O-(2,3-dimethylbutanoyl)-13-(2,3-dimethylbutanoyloxy)-20-deoxyingenol	<i>E. esula</i>	[58]
66	17-Benzoyloxy-3-O-(2,3-dimethylbutanoyl)-13-(2,3-dimethylbutanoyloxy) ingenol	<i>E. esula</i>	[58]
67	13,17-Dibenzoyloxy-3-O-(2,3-dimethylbutanoyl)ingenol	<i>E. esula</i>	[58]
68	13,17-Dibenzoyloxy-3-O-(2,3-dimethylbutanoyl)-20-deoxyingenol	<i>E. esula</i>	[58]
69	3-O-(2,3-dimethylbutanoyl)-13-octanoyloxyingenol	<i>E. esula</i>	[58]
70	17-Benzoyloxy-3-O-(2,3-dimethylbutanoyl)-13-octanoyloxyingenol	<i>E. esula</i>	[58]
71	17-Benzoyloxy-20-O-(2,3-dimethylbutanoyl)-13-(2,3-dimethylbutanoyloxy)ingenol	<i>E. esula</i>	[58]
72	17-Benzoyloxy-13-octanoyloxyingenol	<i>E. esula</i>	[58]
73	20-O-Benzoyl-17-benzoyloxy-13-octanoyloxyingenol	<i>E. esula</i>	[58]
74	17-Benzoyloxy-20-O-(2,3-dimethylbutanoyl)-13-octanoyloxyingenol	<i>E. esula</i>	[58]
75	3-O-Benzoyl-17-benzoyloxy-13-(2,3-dimethylbutanoyloxy)ingenol	<i>E. esula</i>	[58]
76	3-O-Benzoyl-13,17-dibenzoyloxyingenol	<i>E. esula</i>	[58]
77	3-O-Benzoyl-13-octanoyloxyingenol	<i>E. esula</i>	[58]

Table 2. Cont.

No	Name	Species	Ref
78	3-O-Benzoyl-17-benzoyloxy-13-octanoyloxyingenol	<i>E. esula</i>	[58]
79	3-O-Benzoyl-17-benzoyloxy-13-octanoyloxy-20-deoxyingenol	<i>E. esula</i>	[58]
80	Ingenol-3-angelate-5,20-diacetate	<i>E. canariensis</i>	[59]
		<i>E. acurensis</i>	[60]
81	5-Deoxyingenol-3-angelate-20-acetate	<i>E. canariensis</i>	[59]
82	17-Acetoxyingenol-5,20-diacetate-3-angelate	<i>E. kamerunica</i>	[53]
83	Ingenol-3-angelate	<i>E. canariensis</i>	[59]
84	17-Hydroxyingenol-3-angelate-17-benzoate	<i>E. canariensis</i>	[59]
85	17-Hydroxyingenol-3-angelate-20-acetate-17-benzoate	<i>E. canariensis</i>	[59]
86	17-Acetoxyingenol-20-acetate-3-angelate	<i>E. canariensis</i>	[59]
87	17-Hydroxyingenol 17-benzoate 20-angelate	<i>E. canariensis</i>	[59]
88	3-O-Angeloyl-17-[(2Z,4E,6Z)-deca-2,4,6-trienoyloxy]ingenol	<i>E. cauducifolia</i>	[21]
89	17-Acetyloxy-3-O-angeloyl-ingenol	<i>E. cauducifolia</i>	[21]
90	3-O-Angeloyl-17-(benzoyloxy)ingenol	<i>E. cauducifolia</i>	[21]
91	20-O-Acetyl-3-O-angeloyl-17-hydroxyingenol	<i>E. cauducifolia</i>	[21]
92	20-O-Acetyl-3-O-angeloyl-17-(benzoyloxy)ingenol	<i>E. cauducifolia</i>	[21]
93	3-O-Acetyl-20-O-angeloyl-17-hydroxyingenol	<i>E. cauducifolia</i>	[21]



Figure 2. Ingenane Diterpenoids Isolated from *Euphorbia* Species.


Figure 2. Cont.



51 $R_1 = H$, $R_2 = H$, $R_3 = OAc$


52 $R_1 = OAc$, $R_2 = OAc$, $R_3 = H$

53 $R_1 = OAc$, $R_2 = OAc$, $R_3 = OAc$



54 $R = OCO(CH_2)_{10}CH_3$

55 $R = OCO(CH_2)_8CH_3$



56 $R_1 = Ac$, $R_2 = (2'E,4'Z)$ -tetradecadienoyl, $R_3 = Ac$

57 $R_1 = (2'E,4'Z)$ -tetradecadienoyl, $R_2 = Ac$, $R_3 = Ac$

58 $R_1 = (2'E,4'Z)$ -tetradecadienoyl, $R_2 = H$, $R_3 = H$





59 $R_1 = (2'E,4'Z)$ -tetradecadienoyl, $R_2 = isopropylidenyl$, $R_3 = isopropylidenyl$

60 $R_1 = (2'E,4'Z)$ -tetradecadienoyl, $R_2 = H$, $R_3 = Ac$


61 $R_1 = CO(CH=CH)_2(CH_2)_4Me$, $R_2 = H$, $R_3 = Ac$

62 $R_1 = CO(CH_2)_{12}CH_3$, $R_2 = Ac$, $R_3 = Ac$

63 $R_1 = Ac$, $R_2 = CO(CH_2)_{12}CH_3$, $R_3 = Ac$




No.	R_1	R_2	R_3	R_4	No.	R_1	R_2	R_3	R_4
64	A	H	OB	H	72	H	OC	OB	OH
65	A	OA	OB	H	73	H	OC	OB	OB
66	A	OA	OB	OH	74	H	OC	OB	OA
67	A	OB	OB	OH	75	B	OA	OB	OH
68	A	OB	OB	H	76	B	OB	OB	OH
69	A	OC	H	OH	77	B	OC	H	OH
70	A	OC	OB	OH	78	B	OC	OB	OH
71	H	OA	OB	OA	79	B	OC	OB	H



80 $R_1 = OAc$, $R_2 = H$


81 $R_1 = H$, $R_2 = H$

82 $R_1 = OAc$, $R_2 = OAc$



83 $R = H$


84 $R = OBz$



85 $R = OBz$

86 $R = OAc$

Figure 2. Cont.



4. Tiglane Derivates Isolated from *Euphorbia* Species (Table 3)

The tiglane diterpenoids in *Euphorbia* have a 5/7/6/3-tetracyclic ring system. Rings A/B are usually in *trans*-integrated configuration, as in compounds **94–98** and **100–120**. Only a few tiglane diterpenoids, such as **99** and **121**, are in *cis*-configuration. Rings B/C are joined in *trans*-configuration and Rings B/C in *cis*-configuration. Most tiglane diterpenoids have polyhydroxy groups located on C-4, C-9, C-13 and C-20. C-3 forms a carbonyl group. C1,2 and C6,7 form double bonds, respectively. Like the abietane and the ingenane diterpenoids, the hydroxyl groups of tiglane diterpenoids are easily esterified, as in compounds **98–103**. This type of macrocyclic diterpene, which is widespread in the seeds, roots, latex and stem of *Euphorbia* genus, is the main toxic constituent causing irritant, proinflammatory and tumor promoting activity [18,61,62]. When the C12-OH and C13-OH were esterified as a bis-ester, the tumor promoting activity was reinforced at the same time. For example, 12-*O*-tetradecanoylphorbol 12-acetate (TPA) is well-known as a tumor promotor. The diterpene ester with a saturated aliphatic long chain acyl group exhibited high irritant activity and high tumor promoting activity, and the highly unsaturated analogue exhibits high irritant activity, but very weak tumor promoting activity, suggesting that the irritant activity but not the tumour promoting activity of these diterpenoids is related to the degree of unsaturation of the aliphatic long chain [63]. The absence of a C20-OH is known to be important for the irritant and tumor promoting activities of phorbol esters [64]. Introduction of an acetyl group in the 20-position gives rise to a lower irritancy [65]. Compounds **122** and **123** belong to the daphnane diterpene group, which may be derived from the tiglane diterpenoids by cleavage of ring D and isopropenyl linked on C-13.

Table 3. Tigliane diterpenoids isolated from *Euphorbia* species.

No.	Name	Species	Ref
94	13-Acetoxy-12-deoxyphorbol [Prostratin]	<i>E. fischeriana</i>	[66]
95	20-Hydroxy-12-deoxyphorbol 13-(<i>cis</i> -9,10-methylene)-undecanoate	<i>E. poisonii</i>	[14]
96	20-Hydroxy-12-deoxyphorbol angelate	<i>E. poisonii</i>	[14]
97	12-Deoxyphorbolaldehyde-13-acetate	<i>E. fischeriana</i>	[31]
98	12-Deoxyphorbolaldehyde-13-hexadecacetate	<i>E. fischeriana</i>	[31]
99	4,12-Dideoxy(4 α)phorbol-13-hexadecanoate	<i>E. guyoniana</i>	[67]
100	12-O-(2 <i>Z</i> ,4 <i>E</i> -Octadienyl)-4-deoxyphorbol-13,20-diacetate	<i>E. broteri</i>	[54]
101	4,12,20-Trideoxyphorbol-13-(2,3-dimethyl)butyrate	<i>E. pithyusa subsp</i>	[68]
102	12-O-(2 <i>Z</i> ,4 <i>E</i> -octadienyl)-phorbol-13,20-diacetate	<i>E. broteri</i>	[54]
103	12-Deoxyphorbol-13-(9 <i>Z</i>)-octadecanoate-20-acetate	<i>E. fischeriana</i>	[31]
104	13-O-Acetyl-20-O-benzoyl-12-deoxyphorbol	<i>E. cornigera</i>	[69]
105	13-O-Acetyl-20-O-p-methoxybenzoyl-12-deoxyphorbol	<i>E. cornigera</i>	[69]
106	13-O-Acetyl-20-O-decanoyl-12-deoxyphorbol	<i>E. cornigera</i>	[69]
107	13-O-Butanoyl-20-O-decanoyl-12-deoxyphorbol	<i>E. cornigera</i>	[69]
108	13-O-Hexanoyl-20-O-decanoyl-12-deoxyphorbol	<i>E. cornigera</i>	[69]
109	13-O-Octanoyl-20-O-decanoyl-12-deoxyphorbol	<i>E. cornigera</i>	[69]
110	13,20-Didecanoylphorbol	<i>E. cornigera</i>	[69]
111	13-O-Dodecanoyl-20-O-decanoyl-12-deoxyphorbol	<i>E. cornigera</i>	[69]
112	13-O-Decanoyl-20-O-angelyl-12-deoxyphorbol	<i>E. cornigera</i>	[69]
113	13-O-Decanoyl-20-O-tiglyl-12-deoxyphorbol	<i>E. cornigera</i>	[69]
114	12-Deoxyphorbol 20-acetate 13-angelate	<i>E. poisonii</i>	[14]
115	12-Deoxyphorbol 20-acetate 13-phenylacetate	<i>E. poisonii</i>	[14]
116	4,20-Dideoxyphorbol 12,13-bis(isobutyrate)	<i>E. obtusifolia</i>	[70]
117	4-Deoxyphorbol 12,13-bis(isobutyrate)	<i>E. obtusifolia</i>	[70]
118	17-Acetoxy-4-deoxyphorbol 12,13-bis(isobutyrate)	<i>E. obtusifolia</i>	[70]
119	17-Acetoxy-4,20-dideoxyphorbol 12,13-bis(isobutyrate)	<i>E. obtusifolia</i>	[70]
120	4-Deoxyphorbol 12,13-bis(isobutyrate) 20-acetate	<i>E. obtusifolia</i>	[70]
121	4-Epi-4-Deoxyphorbol 12,13-bis(isobutyrate)	<i>E. obtusifolia</i>	[70]
122	20-(4-Hydroxy-3-methoxyphenylacetate)9,13,14-orthophenylacetate	<i>E. poisonii</i>	[14]
123	20-Hydroxyresiniferol 9,13,14-orthophenylacetate	<i>E. poisonii</i>	[14]




Figure 3. Tigliane diterpenoids isolated from *Euphorbia* species.

Figure 3. Cont.




101




102 $R_1 = \text{OCO}(\text{CH}=\text{CH})_2(\text{CH}_2)_3\text{CH}_3$, $R_2 = \text{Ac}$,

103 $R_1 = \text{H}$, $R_2 = \text{CO}(\text{CH}_2)_7\text{CH}=\text{CH}(\text{CH}_2)_7\text{CH}_3$




104–115




116–121

No.	R ₁	R ₂	No.	R ₁	R ₂
104	Acetyl	Benzoyl	113	Decanoyl	Tigloyl
105	Acetyl	p-Methoxybenzoyl	114	Tigloyl	Ac
106	Acetyl	Decanoyl	115	PhCH ₂ CO	Ac
107	Butanoyl	Decanoyl	116	H	H
108	Hexanoyl	Decanoyl	117	OH	H
109	Octanoyl	Decanoyl	118	OH	OAc
110	Decanoyl	Decanoyl	119	H	OAc
111	Dodecanoyl	Decanoyl	120	OAc	H
112	Decanoyl	Angeloyl	121	4-Epi-117	



122



123

5. ¹³C-NMR Data of Diterpenes

Table 4 shows the ¹³C-NMR data of the diterpenoids **1–123**. All the ¹³C-NMR data were recorded in CDCl₃. The structures and the carbon chemical shifts of the abietane diterpenoids are quite different from each other. Here we only discuss the most frequent abietane lactones **1–35**. Four carbons (C-12, C-13, C-15 and C-16) of the lactone ring are the main feature, and their chemical shifts are around δ_{C} 78.5–80.0, 148.4–165.0, 117.0–132.8 and 167.0–178.0, respectively.

Table 4. ^{13}C -NMR data (in CDCl_3) of diterpenes from *Euphorbia* species.

Carbon	Compound / δ_{C} (in ppm)													
	1	2	3	4	5	6	7	8	9	10	11	12	13	14
1	38.0	40.9	38.3	43.2	40.5	41.9	41.6	41.4	41.4	41.3	40.0	41.4	39.3	37.6
2	18.4	18.0	17.3	20.1	18.7	17.4	17.1	18.5	18.5	18.5	18.4	18.4	18.4	27.0
3	42.0	40.9	41.3	43.2	43.1	39.9	38.6	40.0	39.2	39.0	41.5	39.8	41.5	78.2
4	32.7	32.3	32.4	34.2	34.1	37.8	37.6	33.3	33.5	33.5	33.5	33.4	33.5	39.2
5	46.8	54.7	55.3	56.6	57.0	55.6	55.4	53.2	53.6	53.5	53.5	53.4	53.5	52.9
6	30.2	19.8	16.7	22.0	19.2	18.0	17.7	21.2	21.0	20.9	20.8	20.8	20.9	20.5
7	68.9	39.9	34.9	43.1	36.3	35.5	35.4	36.5	36.6	35.7	33.8	34.0	34.0	33.9
8	35.1	75	74.4	75.7	77.6	75.0	74.8	60.8	66.9	67.4	61.3	61.1	61.3	61.0
9	42.5	62.6	56.3	58.0	47.4	72.0	72.1	66.6	47.0	47.8	51.9	51.7	51.8	51.6
10	38.0	36.9	37.2	40.3	39.0	40.5	40.5	48.2	39.1	39.3	41.6	41.3	41.4	41.1
11	27.5	67.3	65.0	29.9	29.0	57.2	65.4	61.3	61.6	61.9	107.6	104.1	106.4	103.4
12	78.5	79.0	79.7	79.0	79.9	79.7	79.8	85.4	85.5	85.3	149.5	147.4	147.3	147.6
13	163.4	157.5	160.3	165.3	166.5	161.2	160.3	148.4	150.8	154.5	147.2	144.9	146.5	144.8
14	26.7	71.9	71.8	73.5	74.3	65.2	57.4	55.6	53.6	55.3	54.3	54.4	54.4	54.3
15	120.5	124.2	125.9	123.2	125.5	125.0	126.0	130.1	150.8	128.3	122.3	125.1	127.4	125.4
16	175.4	175.4	176.2	178.0	177.8	175.4	175.3	169.8	168.2	167.4	170.5	170.6	169.2	170.4
17	8.4	6.7	7.9	8.2	9.4	9.2	9.0	8.6	56.5	54.9	55.4	8.6	56.3	8.6
18	33.1	33.0	32.4	34.6	34.1	16.7	16.4	33.2	33.5	33.5	33.4	33.4	33.5	28.3
19	21.6	20.8	20.6	22.4	22.4	21.8	21.3	22.1	21.9	21.9	21.9	21.9	21.9	15.5
20	12.6	16.8	15.7	18.1	15.4	33.6	33.7	15.4	15.6	15.1	15.0	14.9	15.1	15.0

Carbon	Compound / δ_{C} (in ppm)													
	15	16	17	18	19	20	21	22	23	24	25	26	27	28
1	41.9	31.7	39.7	37.4	32.1	29.9	39.4	39.0	39.6	40.2	51.2	55.9*	30.5	37.4
2	19.0	18.7	19.1	27.5	25.7	27.3	19.0	18.6	18.8	18.9	209.4	209.4	34.2	34.4
3	39.5	41.6	41.9	78.5	75.6	78.3	41.7	41.7	41.7	41.8	82.4	54.0*	216.4	215.6
4	33.1	33.2	33.6	39.0	37.8	39.0	33.6	33.4	32.9	41.0	45.0	38.7	47.1	47.5
5	47.1	39.9	55.3	54.3	48.4	45.4	55.4	54.0	46.8	46.7	53.4	54.5	46.0	54.8
6	31.0	31.0	23.9	23.4	23.4	23.0	23.8	22.3	29.9	30.7	23.0	23.6	24.1	24.6
7	72.4	74.4	37.2	36.8	37.1	32.7	37.1	36.0	71.5	71.1	36.3	36.4	32.2	36.6
8	151.2	148.4	156.3	151.4	152.0	152.6	152.6	154.4	153.4	155.2	149.4	149.5	152.2	150.2
9	46.7	79.1	51.9	51.5	51.6	77.2	60.8	51.4	54.8	55.3	51.3	51.3	76.9	50.7
10	41.9	44.7	41.6	41.2	41.3	44.2	40.3	38.9	40.9	32.7	46.9	46.2	43.8	40.9
11	27.2	38.4	27.5	27.5	27.5	39.7	64.6	31.2	69.6	70.2	27.6	27.6	40.0	27.8
12	76.1	77.2	76.1	75.9	76.0	77.1	79.4	102.4	102.7	104.3	75.3	75.3	76.9	75.6
13	155.1	153.9	152.3	156.0	156.0	154.7	150.1	154.2	152.8	156.0	155.0	155.0	154.6	155.5
14	115.9	118.8	113.9	114.2	114.1	115.7	113.5	113.4	114.7	115.4	115.2	114.9	115.9	114.8
15	118.9	130.0	116.2	116.4	116.4	117.9	118.2	116.3	121.0	124.0	117.5	117.3	118.1	117.1
16	174.9	174.3	175.4	175.3	175.2	174.7	175.4	173.1	173.6	172.1	174.9	174.7	174.6	175.1
17	8.5	8.6	8.3	8.2	28.7	28.9	8.5	8.1	8.4	55.5	8.3	33.5	27.1	26.5
18	33.6	33.8	33.9	28.6	22.2	16.0	33.9	33.5	32.9	32.9	29.5	23.0	21.7	21.8
19	21.7	22.0	21.8	15.6	16.7	17.5	21.8	22.0	21.6	21.4	16.4	17.3	17.8	16.2
20	16.1	17.4	16.8	16.7	8.2	8.4	17.3	14.6	14.3	14.4	17.3	8.2	8.3	8.3

Table 4. Cont.

Carbon	Compound / δ_c (in ppm)													
	29	30	31	32	33	34	35	36	37	38	39	40	41	42
1	23.4*	37.9	28.2	33.6	30.4	31.0	29.8	31.4	37.5	37.4	38.9	39.3	36.6	36.1
2	145.2	18.2	26.8	18.8	19.3	19.1	19.2	19.2	34.7	34.6	18.4	18.31	19.0	19.2
3	200.2	41.4	78.1	19.1	18.5	19.9	18.4	19.9	216.5	216	41.8	41.5	41.3	41.3
4	44.0*	32.7	37.2	15.7	16.4	14.7	16.6	14.8	47.8	47.8	33.2	33.1	33.6	33.7
5	52.6	53.9	43.8	51.1	47.6	44.8	41.6	44.1	55.0	54.7	49.8	54.2	52.0	52.8
6	23.2	38.9	35.7	22.7	20.6	27.5	28.9	27.6	23.1	22.7	37.5	19.21	17.3	18.9
7	36.7	209.8	209.4	33.5	24.4	140.4	62.5	139.9	35.5	34.6	200.7	41.01	26.2	33.0
8	148.9	44.2	49.4	76.1	134.6	136.8	135.7	134.8	139.9	139.9	138.9	69.5	143.4	139.1
9	48.3	50.1	53.0	52.6	160.5	41.6	164.9	40.6	49.0	50.0	51.8	61.3	150.3	141.5
10	41.7*	37.4	39.1	36.6	38.9	34.3	39.8	35.2	38.1	37.7	35.9	37.8	39.3	39.8
11	27.6	28.1	23.8	27	34.2	27.2	33.9	27.7	18.6	26.3	18.6	71.7	152.0	154.6
12	75.4	77.5	77.0	79.5	78.8	77.9	78.4	76.7	31.0	69.8	29.7	197.5	125.4	112.8
13	154.9	160.7	160.4	153.3	150.6	151.1	149.9	52.9	46.7	52.1	71.8	136.9	176.4	135.0
14	116.0	24.0	38.5	196	185.7	187.5	187.0	196.2	123.7	121.9	139.5	151.4	184.8	123.0
15	117.7	121.9	122.0	131.8	131.1	132.5	132.8	40.0	214.1	215.1	37.8	137.3	120.2	198.0
16	174.9	174.8	174.8	172.9	172.8	173.5	173.3	178.2	64.8	65.2	17.6	–	144.2	26.4
17	27.0	8.4	8.4	9.4	9.8	10.0	9.9	16.2	23.8	17.5	16.2	128.8	8.5	33.7
18	2.0	13.1	27.6	21.5	22.3	20.5	22.2	20.4	25.8	25.6	32.6	33.8	33.5	22.2
19	19.4	33.5	14.9	23.9	23.2	24.5	23.2	24.7	22.3	22.3	21.2	21.9	21.9	19.6
20	8.3	21.0	13.1	16.9	16.8	11.5	15.7	12.4	14.7	14.3	14.1	17.8	20.4	–

Carbon	Compound / δ_c (in ppm)													
	43	44	45	46	47	48	49	50	51	52	53	54	55	56
1	130.0	129.0	131.7	130.0	132.2	131.8	130.5	132.2	132.1	132.2	131.6	131.4	131.4	132.1
2	138.8	139.3	136.3	138.8	133.2	133.6	136.0	133.4	135.8	135.8	136.0	136.2	136.2	133.3
3	80.5	80.3	80.2	80.6	82.2	82.0	81.7	82.1	82.9	82.7	81.6	82.5	82.5	82.1
4	84.3	84.0	74.7	84.4	85.8	85.7	85.7	86.0	85.0	84.9	85.8	84.5	84.4	86.0
5	75.3	75.2	75.0	73.8	74.8	74.6	74.8	75.0	77.3	74.8	74.8	76.7	76.7	75.0
7	127.4	126.2	128.4	128.3	131.9	131.0	129.7	131.8	123.1	129.5	130.9	127.2	127.3	128.2
8	44.0	43.2	42.9	44.1	43.6	43.1	42.9	43.7	43.0	43.6	43.2	42.6	42.6	43.7
9	207.8	206.5	205.2	208.0	205.4	204.6	204.7	205.4	206.0	206.2	204.7	205.9	205.9	205.6
10	72.4	72.6	72.0	72.6	71.9	71.8	71.8	72.1	72.0	72.1	72.0	71.8	71.8	72.1
11	39.8	38.6	37.7	39.6	38.6	38.6	37.9	38.7	38.7	38.5	38.6	37.4	37.4	38.6
12	30.8	35.0	35.2	31.0	31.1	30.8	35.0	31.1	30.8	31.2	30.7	35.0	35.0	31.2
13	23.1	68.8	29.1	23.9	23.1	23.2	69.1	23.1	24.0	24.0	24.1	69.0	69.0	23.2
14	22.9	28.2	29.4	23.2	22.9	24.0	28.0	23.3	23.7	23.3	23.2	28.2	28.2	23.0
15	24.0	30.2	30.1	22.7	24.4	27.7	30.7	24.4	27.4	28.5	27.6	30.3	30.3	24.3
16	28.5	22.5	24.7	28.5	28.4	24.2	22.2	28.5	24.3	23.1	24.2	22.4	22.5	28.4
17	15.4	16.7	66.8	15.5	15.5	65.7	16.8	15.4	65.6	15.5	65.7	16.7	16.7	15.5
18	17.3	18.4	18.2	17.4	17.0	16.6	18.0	17.1	16.7	17.3	16.6	18.2	18.2	17.1
19	15.5	15.4	15.5	15.4	15.4	15.4	15.3	15.6	15.6	15.6	15.4	15.4	15.5	15.4
20	67.2	66.8	62.3	66.4	65.8	65.7	65.7	65.9	21.9	66.8	65.7	67.1	67.2	65.9

Table 4. Cont.

Carbon	Compound / δ_C (in ppm)													
	57	58	59	60	61	62	63	64	65	66	67	68	69	70
1	132.2	132.0	132.2	132.2	132.2	132.0	132.3	132.1	131.5	131.0	131.2	131.8	131.5	131.1
2	133.4	135.9	135.9	136.1	136.1	—	133.5	136.2	136.3	136.8	136.8	136.4	136.2	136.7
3	82.5	82.7	81.8	82.8	82.5	82.4	82.4	82.0	82.7	82.1	82.2	82.9	82.6	82.0
4	86.0	84.8	84.1	85.0	85.1	—	85.9	85.2	84.9	84.8	84.9	85.0	84.5	84.8
5	78.5	76.7	74.0	74.9	75.0	75.0	75.1	76.7	77.0	76.0	76.5	77.4	76.9	76.0
6	135.8	139.4	136.6	136.1	136.2	—	135.5	138.4	138.2	140.2	140.1	138.3	139.6	140.1
7	128.4	128.2	128.3	129.3	129.4	132.0	131.7	122.8	122.0	125.6	126.1	122.2	127.3	125.7
8	43.7	43.5	43.6	43.9	43.8	43.7	43.7	42.8	42.6	42.7	43.0	42.9	42.7	42.6
9	205.7	207.0	207.6	206.2	206.1	—	205.3	206.7	205.1	205.2	204.9	205	205.8	205.1
10	72.2	72.0	72.3	72.1	72.3	—	72.1	71.5	71.9	71.9	71.9	72.0	71.9	71.7
11	38.5	38.4	37.6	38.8	38.6	38.6	38.7	38.5	38.1	37.8	38.0	38.4	37.5	37.8
12	31.1	31.2	31.2	31.2	31.3	31.2	31.2	30.5	35.6	35.6	35.3	35.3	34.4	35.0
13	23.2	23.3	23.5	23.3	23.4	23.1	23.1	23.8	68.3	68.3	69.3	69.3	69.0	68.5
14	23.0	23.0	23.0	23.1	23.3	23.3	23.3	23.4	28.7	28.5	28.6	29.0	28.3	28.4
15	24.3	24.0	24.0	24.0	24.0	—	24.4	27.5	33.9	34.0	34.5	34.4	30.3	34.0
16	28.4	28.5	28.5	28.5	28.5	28.5	28.5	24.2	18.7	18.5	18.7	18.7	22.5	18.5
17	15.6	15.5	15.6	15.5	15.5	15.4	15.4	66.2	65.5	65.5	65.6	65.7	16.7	65.5
18	17.1	17.3	17.5	17.3	17.3	17.1	17.1	16.4	18.0	18.1	18.0	17.9	18.2	17.9
19	15.4	15.5	15.5	15.6	15.5	15.6	15.6	15.3	15.5	15.4	15.6	15.6	15.4	15.5
20	66.1	67.1	64.3	66.8	66.7	65.9	65.6	21.6	21.7	66.6	67.0	21.8	67.2	66.7

Carbon	Compound / δ_C (in ppm)													
	71	72	73	74	75	76	77	78	79	80	81	82	83	84
1	128.6	128.6	128.4	128.8	131.4	131.5	132.0	131.5	132.0	132.0	132.0	131.5	132.1	131.6
2	139.7	139.8	139.7	139.6	136.8	136.8	136.0	136.8	136.0	136.0	136.0	133.7	135.7	136.2
3	80.4	80.2	80.1	80.4	83.0	83.0	83.3	83.0	83.6	81.8	85.7	81.5	82.5	82.2
4	84.3	84.3	84.3	84.3	84.9	85.1	84.6	84.9	85.1	85.8	80.5	85.7	84.7	84.9
5	73.8	75.2	74.0	73.8	77.3	76.3	76.9	77.2	77.2	74.9	43.7	74.6	77.1	76.6
6	137.7	141.2	137.5	137.8	139.8	140.2	139.0	139.8	138.0	133.0	132.0	135.9	139.0	139.7
7	126.8	125.2	125.5	126.7	126.4	126.0	128.0	126.4	122.0	132.0	129.0	130.8	128.6	127.2
8	43.2	43.3	43.3	43.3	42.9	43.0	42.7	42.9	42.8	43.6	44.2	43.1	43.5	43.2
9	204.9	205.4	205.3	204.8	205.0	205.1	206.0	204.9	205.0	206.0	208.0	204.6	206.6	205.9
10	72.9	72.7	72.8	72.7	72.2	72.0	72.0	72.0	72.1	72.0	75.0	71.9	72.0	72.0
11	38.8	38.8	38.8	38.9	38.0	38.2	37.6	38.1	38.5	38.6	37.0	38.5	38.3	38.4
12	35.9	35.4	35.6	35.4	35.9	35.3	35.2	35.4	35.4	31.1	31.5	30.6	31.1	30.9
13	68.1	68.4	68.3	68.3	69.4	69.3	69.0	68.6	68.6	23.1	23.7	23.1	23.3	24.3
14	28.7	28.7	28.6	28.7	28.6	28.6	28.3	28.6	28.8	22.9	23.2	23.9	23.0	23.5
15	33.8	33.9	33.9	33.9	34.0	34.4	30.3	34.1	34.1	24.3	23.6	27.5	24.0	27.7
16	18.8	18.7	18.6	18.7	18.9	18.6	22.5	18.7	18.7	28.4	28.6	24.1	28.5	24.6
17	65.4	65.7	65.5	65.5	65.5	65.6	16.7	65.5	65.6	15.4	15.5	65.6	15.5	66.2
18	18.5	18.2	18.2	18.2	18.4	18.1	18.4	18.1	18.1	17.0	18.2	16.5	17.3	16.9
19	15.2	15.4	15.3	15.3	15.6	15.6	15.6	15.6	15.6	15.5	15.6	15.3	15.5	15.6
20	65.7	66.6	66.4	65.7	67.1	66.9	67.4	67.1	21.8	65.9	68.5	65.6	67.5	67.2

Table 4. Cont.

Carbon	Compound / δ_c (in ppm)													
	85	86	87	88	89	90	91	92	93	94	95	96	97	98
1	131.6	131.7	131.3	131.3	131.7	131.7	131.7	131.7	131.7	160.6	161.4	161.3	160.4	160.5
2	136.2	136.4	136.8	136.8	136.3	136.3	136.7	136.3	136.3	132.9	132.7	132.7	133.5	133.5
3	82.3	82.5	84.7	84.7	84.2	84.4	83.6	83.9	84.1	209.2	209.3	209.4	208.3	208.4
4	84.9	84.9	74.2	74.2	74.2	74.5	74.3	74.4	74.1	73.8	73.8	73.8	72.8	72.8
5	74.7	74.9	74.0	74.0	74.5	74.2	74.3	74.3	75.0	38.7	38.6	38.5	34.4	34.6
6	136.4	136.3	136.8	136.8	136.7	136.5	136.7	136.3	136.6	140.4	139.8	140.0	142.9	142.9
7	128.2	128.7	128.5	128.5	128.3	128.4	128.0	128.1	128.4	130.4	130.6	130.4	158.1	158.2
8	43.2	43.3	42.1	42.1	42.9	42.9	42.4	42.8	42.9	39.1	39.2	39.1	41.4	41.5
9	205.5	205.5	207.2	207.2	206.0	205.7	205.0	205.4	205.2	76.0	76.0	76.2	77.1	77.1
10	72.0	72.1	71.7	71.7	71.9	71.9	71.7	71.8	71.9	56.2	55.8	55.7	55.8	55.8
11	38.5	38.5	37.2	37.2	37.7	37.6	37.6	37.6	37.7	36.6	36.3	36.3	36.5	36.5
12	30.9	30.9	34.9	34.9	35.1	35.1	35.2	35.3	35.2	32.3	32.0	31.9	31.7	31.8
13	24.3	24.3	29.4	29.4	29.4	29.3	29.5	29.8	29.2	63.8	63.2	63.2	63.0	63.0
14	23.5	23.5	29.1	29.1	29.6	29.4	29.8	29.6	29.7	32.8	32.6	32.8	32.0	32.1
15	27.7	27.5	29.7	29.7	30.2	29.8	30.0	30.0	30.0	22.5	26.6	22.9	22.9	22.7
16	24.5	24.4	24.8	24.8	24.8	24.9	24.7	24.9	24.7	23.2	23.1	23.6	23.1	23.2
17	66.1	65.6	66.2	66.2	66.3	66.3	62.5	66.3	62.2	15.3	15.4	15.4	15.3	15.3
18	16.9	17.0	18.5	18.5	18.6	18.7	18.8	18.2	18.4	18.8	18.6	18.6	18.5	18.6
19	15.6	15.6	16.1	16.1	15.3	16.0	16.0	15.9	15.5	9.9	10.1	10.1	10.1	10.1
20	66.5	66.7	62.2	62.2	62.4	62.3	65.5	66.3	66.4	67.9	68.3	68.2	193.8	193.8

Carbon	Compound / δ_{C} (in ppm)												
	99	100	101	102	103	104	105	106	107	108	109	110	111
1	156.9	159.7	161.0	160.8	161.3	160.3	160.3	160.3	160.3	160.3	160.3	160.3	160.3
2	143.0	137.3	138.3	135.7	132.9	136.3	136.3	136.2	136.2	136.3	136.3	136.2	136.3
3	213.8	208.9	203.0	208.6	208.9	210.2	210.3	210.3	210.3	210.3	210.3	210.3	210.3
4	50.1	42.6	44.4	73.6	73.7	44.5	44.5	44.5	44.5	44.6	44.5	44.6	44.6
5	25.1	35.1	34.0	38.8	38.9	34.0	34.0	34.1	34.0	34.0	34.0	34.1	34.1
6	136.3	136.5	136.2	132.9	135.2	139.6	139.6	139.7	139.7	139.6	139.6	139.6	139.7
7	127.7	130.2	126.8	132.7	133.7	125.0	125.7	125.7	125.7	125.7	125.6	125.6	125.6
8	41.0	42.3	41.9	39.4	39.5	42.2	42.2	42.2	42.2	42.2	42.2	42.2	42.2
9	75.5	77.8	75.2	78.2	75.9	77.9	77.9	77.9	77.9	77.9	77.9	77.9	77.9
10	47.1	54.1	53.9	56.2	55.8	54.3	54.3	54.3	54.4	54.3	54.3	54.3	54.3
11	37.1	44.1	46.2	43.2	36.3	42.3	42.3	42.3	42.3	42.3	42.3	42.4	42.4
12	30.5	76.1	31.8	76.1	31.9	56.7	56.7	54.8	56.8	56.7	56.7	56.7	52.7
13	62.7	65.4	62.8	65.7	63.6	65.0	65.0	65.0	65.1	65.0	65.0	65.0	65.0
14	33.1	35.4	32.0	36.1	32.4	35.9	35.8	35.9	36.0	35.9	35.9	36.0	36.0
15	22.5	25.7	22.5	25.7	22.6	25.8	25.8	25.8	25.8	25.8	25.8	25.8	25.8
16	23.7	23.8	15.2	23.8	23.2	23.9	23.9	23.9	24.0	23.9	24.0	23.9	23.9
17	15.2	16.7	22.9	16.7	15.3	17.0	17.0	16.9	16.9	17.0	16.9	16.9	16.8
18	15.9	15.1	19.0	14.4	18.6	15.1	15.1	15.1	15.2	15.1	15.2	15.2	15.0
19	10.4	10.2	10.0	10.1	10.1	10.2	10.2	10.3	10.3	10.2	10.3	10.3	10.3
20	69.5	68.9	25.2	69.4	69.4	62.2	66.4	67.4	68.2	66.9	66.7	66.5	65.9

Table 4. Cont.

Carbon	Compound / δ_C (in ppm)											
	112	113	114	115	116	117	118	119	120	121	122	123
1	160.3	160.2	161.5	161.4	160.2	159.8	160.0	159.4	159.8	156.2	158.2	158.3
2	136.2	136.2	132.8	132.8	136.3	136.4	136.6	136.5	136.5	143.3	136.6	136.5
3	210.3	210.3	209.1	209.0	210.2	209.7	209.7	210.1	209.6	213.3	208.4	209.0
4	44.5	44.5	73.6	73.6	44.5	44.2	44.0	44.3	44.1	49.6	73.3	73.5
5	34.1	34.1	39.0	39.0	34.0	29.6	29.0	33.6	30.0	25.1	40.0	39.8
6	139.6	139.7	134.8	134.8	139.0	142.0	142.4	139.3	137.2	137.0	135.0	138.9
7	125.7	125.6	134.1	133.2	125.8	126.5	125.3	125.0	130.3	126.5	130.8	130.8
8	42.2	42.2	39.5	39.4	42.2	42.1	42.4	42.6	42.2	40.7	39.1	38.9
9	77.9	77.9	76.0	75.9	77.9	77.8	—	77.5	77.8	78.1	81.1	81.2
10	54.3	54.3	55.7	56.7	54.3	54.2	53.8	53.9	54.0	47.4	55.4	55.5
11	42.4	42.4	36.4	36.3	42.3	42.4	42.4	42.5	42.3	43.2	33.0	33.1
12	53.8	56.8	31.9	31.7	76.7	76.7	76.1	76.2	76.4	75.3	35.7	35.7
13	65.0	65.1	63.1	63.9	65.0	65.0	65.2	65.3	64.8	64.8	84.4	84.5
14	36.0	35.9	32.7	32.3	35.9	35.8	36.4	36.5	35.5	37.1	80.6	80.8
15	25.8	25.8	22.9	23.0	25.8	25.9	30.0	29.8	25.8	25.3	146.5	146.4
16	23.9	24.0	23.6	23.0	23.9	23.8	—	19.5	23.8	24.2	110.7	110.7
17	17.0	16.9	15.4	15.3	16.9	16.9	63.3	63.5	16.8	16.5	18.8	18.8
18	15.2	15.2	18.6	18.5	15.1	15.1	15.2	15.2	15.0	11.9	19.8	19.9
19	10.3	10.3	10.1	10.1	10.2	10.2	10.3	10.3	10.3	10.5	10.2	10.2
20	67.2	65.9	69.8	69.7	25.4	67.5	67.1	25.4	68.9	69.3	70.4	69.3

(—) Data not observed; *interchangeable.

Affected by the α,β -unsaturated γ -lactone, the carbon chemical shifts of Me-17 is very low at δ_C 6.7–10.0, as shown in **1–7**, **12**, **14–18**, **21–23**, **25** and **30–34**. Thus, the assignment of the four methyl groups (C-17, C-18, C-19 and C-20) in compounds **19–20** and **26–29** [40,42] were doubtful. In their ^{13}C -NMR spectra, chemical shifts around δ_C 8.0 should be assigned to C-17 instead of C-20, δ_C 27.0–34.0 should be assigned to C-18 instead of C-17, and δ_C 16.0–23.0 should be assigned to C-19 instead of C-18. Other positions, such as C-2, C-3, C-7, C-8, C-11 and C-14 are usually substituted by oxygen groups, whose carbon chemical shifts are around δ_C 65.0–78.0. Values close to δ_C 115.0 can be assigned to tertiary carbon (C-14) on double bond in **15–29**. Some carbons of the abietane diterpenoids (**27**, **28** and **30–36**) may be carbonylated when their chemical shifts are above δ_C 185.0.

The carbon chemical shifts do not show very characteristic features for ingenane skeleton type. Their characteristic carbon chemical shifts are observed around δ_C 127.0–140.0 and 204.0–208.0, which are assigned to four carbons (C-1, C-2, C-6 and C-7) on two double bonds and the bridged carbonyl (C-9). The carbon chemical shift around δ_C 71.8–72.8 is assigned to the quaternary carbon (C-10) near the bridged carbonyl. The carbon chemical shifts around δ_C 74.0–86.0 (assigned to C-3, C-4 and C-5 with oxygen substituted) are registered in the ^{13}C -NMR spectra of these compounds. Besides, there is another carbon (C-20) usually oxygen substituted, but it show little lower values (δ_C 62.3–67.2), while its value is around δ_C 21.6–21.8 without oxygen substitution, as the compounds **64**, **65** and **68**. The carbon chemical shifts of **65–79** show C-13 and C-17 are registered δ_C 65.5–69.1 after acylation.

The structures of tiglane diterpenes can be confirmed by carbon chemical shifts around δ_C 203.0–210.3, 156.2–160.6, 132.7–138.3, 132.9–142.9 and 125.0–158.2 assigned to C-3, C-1, C-2, C-6 and C-7, respectively (except **99** and **121**). Values close to δ_C 73.0 and 77.0 are assigned to the carbons C-4 and C-9 substituted by hydroxyl groups. The carbon chemical shift of C-20 is at δ_C 193.8 when it is substituted by hydroxyl group (**97** and **98**), as well as the chemical shift of C-7 is obvious higher than other tiglane diterpenes because of conjugated effect. Compounds **99** and **121** are rare A/B *cis*-integrated compounds, and the structure of these isomers can be confirmed from the data of ^{13}C -NMR, for the chemical shifts of C-2 and C-3 are 4–7 ppm higher than that of the A/B *trans*-integrated ones.

Acknowledgements

The authors thank National Natural Science Foundation of China (No.30672678, 30973940) and “Qinglan Project” Scientific and Technological Innovation Team Training Program of Jiangsu College and University for their financial support.

References and Notes

1. Jassbi, A.R. Chemistry and biological activity of secondary metabolites in *Euphorbia* from Iran. *Phytochemistry* **2006**, *67*, 1977–1984.
2. Hohmann, J.; Molnár, J. Euphorbiaceae diterpenes: Plant toxins or promising molecules for the therapy? *Acta Pharm. Hung.* **2004**, *74*, 149–157.
3. Ahmad, W.; Nazir, M.; Khan, S.A. The chemical composition of various *Euphorbia* species for industrial applications. Part II. Neutral lipids of *Euphorbia cauducifolia*. *Pak. J. Sci. Res.* **1988**, *31*, 85–89.
4. Duarte, N.; Gyémánt, N.; Abreu, P.M.; Molnár, J.; Ferreira, M.J.U. New macrocyclic lathyrane diterpenes, from *Euphorbia lagascae*, as inhibitors of multidrug resistance of tumour cells. *Planta Med.* **2006**, *72*, 162–168.
5. Yan, S.S.; Li, Y.; Wang, Y.; Shen, S.S.; Gu, Y.; Wang, H.B.; Qin, G.W.; Yu, Q. 17-Acetoxyjolkinolide B irreversibly inhibits I-kappaB kinase and induces apoptosis of tumor cells. *Mol. Cancer Ther.* **2008**, *7*, 1523–1532.
6. Liu, W.K.; Ho, J.C.K.; Qin, G.W.; Che, C.T. Jolkinolide B induces neuroendocrine differentiation of human prostate LNCaP cancer cell line. *Biochem. Pharmacol.* **2002**, *63*, 951–957.
7. Luo, H.Y.; Wang, A.Q. Induction of apoptosis in K562 cells by jolkinolide B. *Can. J. Physiol. Pharmacol.* **2006**, *84*, 959–965.
8. Duarte, N.; Loge, H.; Ferreira, M.J.U. Antiproliferative activity of *ent*-abietane lactones against resistant human cancer cell lines. *Planta Med.* **2008**, *74*, 984–985.
9. Kigoshi, H.; Ichino, T.; Yamada, K.; Ijuin, Y.; Makita, S.F.; Uemura, D. Stereostructure and bioactivities of jolkinolide D. *Chem. Lett.* **2001**, *6*, 518–519.
10. Shi, Q.W.; Su, X.H.; Kiyota, H. Chemical and pharmacological research of the plants in genus *Euphorbia*. *Chem. Rev.* **2008**, *108*, 4295–4327.
11. Singla, A.K.; Pathak, K. Phytoconstituents of *Euphorbia* species. *Fitoterapia* **1990**, *61*, 483–516.

12. Appendino, G.; Szallasi, A. Euphorbium: Modern research on its active principle, resiniferatoxin, revives an ancient medicine. *Life Sci.* **1997**, *60*, 681–696.
13. Xu, Z.H.; Sun, J.; Xu, R.S.; Qin, G.W. Casbane diterpenoids from *Euphorbia ebracteolata*. *Phytochemistry* **1998**, *49*, 149–151.
14. Fatope, M.O.; Zeng, L.; Ohayaga, J.E.; Shi, G.; McLaughlin, J.L. Selectively cytotoxic diterpenes from *Euphorbia poisonii*. *J. Med. Chem.* **1996**, *39*, 1005–1008.
15. Murugan, S.; Anand, R.; Uma-Devi, P.; Vidhya, N.; Rajesh, K.A. Efficacy of *Euphorbia milli* and *Euphorbia pulcherrima* on aflatoxin producing fungi (*Aspergillus flavus* and *Aspergillus parasiticus*). *Afr. J. Biotechnol.* **2007**, *6*, 718–719.
16. Ma, Q.G.; Liu, W.Z.; Wu, X.Y.; Zhou, T.X.; Qin, G.W. Chemical studies of Lang-Du, a traditional Chinese medicine. 1. Diterpenoids from *Euphorbia fischeriana*. *Phytochemistry* **1997**, *44*, 663–666.
17. Hezareh, M. Prostratin as a new therapeutic agent targeting HIV viral reservoirs. *Drug News Perspect.* **2005**, *18*, 496–500.
18. Betancur-Galvis, L.; Palomares, E.; Marco, J.A.; Estornell, E. Tiglane diterpenes from the latex of *Euphorbia obtusifolia* with inhibitory activity on the mammalian mitochondrial respiratory chain. *J. Ethnopharmacol.* **2003**, *85*, 279–282.
19. Singh, A.; Agarwal, R.A. Possibility of using latex of euphorbiales for snail control. *Sci. Total Environ.* **1988**, *77*, 231–236.
20. Vogg, G.; Mattes, E.; Rothenburger, J.; Hertkorn, N.; Stefan, A.; Sandermann, J.H. Tumor promoting diterpenes from *Euphorbia leuconeura* L. *Phytochemistry* **1999**, *51*, 289–295.
21. Baloch, I.B.; Baloch, M.K.; Saqib, Q.N. Tumor-Promoting Diterpene Esters from Latex of *Euphorbia cauducifolia* L. *Helv. Chim. Acta.* **2005**, *88*, 3145–3150.
22. Kedei, N.; Lundberg, D.J.; Toth, A.; Welburn, P.; Garfield, S.H.; Blumberg, P.M. Characterization of the Interaction of Ingenol 3-Angelate with Protein Kinase C. *Cancer Res.* **2004**, *64*, 3243–3255.
23. Cateni, F.; Falsone, G.; Zilic, J. Terpenoids and glycolipids from Euphorbiaceae. *Mini-Rev. Med. Chem.* **2003**, *3*, 425–437.
24. Zayed, S.; Farghaly, M.; Soliman, S.M.; Gotta, H.; Sorg, B.; Hecker, E. Dietary cancer risk from conditional cancerogens (tumor promoters) in produce of livestock fed on species of spurge (Euphorbiaceae) V. Skin irritant and tumor-promoting diterpene ester toxins of the tiglane and ingenane type in the herbs *Euphorbia nubica* and *Euphorbia helioscopia* contaminating fodder of livestock. *J. Cancer Res. Clin. Oncol.* **2001**, *127*, 40–47.
25. Shi, H.M.; Williams, I.D.; Sung, H.H.Y.; Zhu, H.X.; Ip, N.Y.; Min, Z.D. Cytotoxic diterpenoids from the roots of *Euphorbia ebracteolata*. *Planta Med.* **2005**, *71*, 349–354.
26. Ferreira, M.U.; Gyémánt, N.; Madureira, A.M.; Molnár, J. Inhibition of P-glycoprotein transport activity in a resistant mouse lymphoma cell line by diterpenic lactones. *Anticancer Res.* **2005**, *25*, 3259–3262.
27. Marco, J.A.; Sanz-Cervera, J.F.; Yuste, A.; Jakupovic, J. Isoterracinolides A and B, novel bishomoditerpene lactones from *Euphorbia terracina*. *J. Nat. Prod.* **1999**, *62*, 110–113.
28. Wang, H.; Zhang, X.F.; Cai, X.H.; Ma, Y.B.; Luo, X.D. Three new diterpenoids from *Euphorbia wallichii*. *Chin. J. Chem.* **2004**, *22*, 199–202.

29. Pan, Q.; Shi, M.F.; Min, Z.D. Studies on the 2D NMR spectra of Jolkinolide diterpenoids from *Euphorbia fischriana*. *J. Chin. Pharm. Univ.* **2004**, *35*, 16–19.
30. Sutthivaiyakit, S.; Thapsut, M.; Prachayasittikul, V. Constituents and bioactivity of the tubers of *Euphorbia sessiliflora*. *Phytochemistry* **2000**, *53*, 947–950.
31. Wang, Y.B.; Huang, R.; Wang, H.B.; Jin, H.Z.; Lou, L.G.; Qin, G.W. Diterpenoids from the roots of *Euphorbia fischeriana*. *J. Nat. Prod.* **2006**, *69*, 967–970.
32. Che, C.T.; Zhou, T.X.; Ma, Q.G.; Qin, G.W.; Williams, I.D.; Wu, H.M.; Shi, Z.S. Diterpenes and aromatic compounds from *Euphorbia fischeriana*. *Phytochemistry* **1999**, *52*, 117–21.
33. Lal, A.R.; Cambie, R.C.; Rutledge, P.S.; Woodgate, P.D. Chemistry of fijian plants. 6. Ent-pimarane and ent-abietane diterpenes from *Euphorbia fidjiana*. *Phytochemistry* **1990**, *29*, 2239–2246.
34. Haba, H.; Lavaud, C.; Marcourt, L.; Long, C.; Harkat, H.; Benkhaled, M. Ent-abietane diterpenoids from *Euphorbia guyoniana* Boiss. & Reut. *Biochem. Syst. Ecol.* in press.
35. Jeske, F.; Jakupovic, J.; Berendsohn, W. Diterpenes from *Euphorbia seguieriana*. *Phytochemistry* **1995**, *40*, 1743–1750.
36. Appendino, G.; Belloro, E.; Cesare-Tron, G.; Jakupovic, J.; Ballero, M. Polycyclic diterpenoids from *Euphorbia characias*. *Fitoterapia* **2000**, *71*, 134–142.
37. Valente, C.; Pedro, M.; Duarte, A.; Nascimento, M.S.J.; Abreu, P.M.; Ferreira, M.J.U. Bioactive diterpenoids, a new jatrophane and two ent-abietanes, and other constituents from *Euphorbia pubescens*. *J. Nat. Prod.* **2004**, *67*, 902–904.
38. Appendino, G.; Jakupovic, S.; Tron, G.C.; Jakupovic, J.; Milon, V.; Ballero, M. Macrocyclic diterpenoids from *Euphorbia semiperfoliata*. *J. Nat. Prod.* **1998**, *61*, 749–756.
39. Shizuri, Y.; Kosemura, S.; Yamamura, S.; Ohba, S.; Ito, M.; Saito, Y. Isolation and structures of helioscopinolides, new diterpenes from *Euphorbia helioscopia* L. *Chem. Lett.* **1983**, *12*, 65–68.
40. Crespi-Perellino, N.; Garofano, L.; Arlandini, E.; Pinciroli, V.; Minghetti, A.; Vincieri, F.F.; Danieli, B. Identification of new diterpenoids from *Euphorbia calypttrata* cell cultures. *J. Nat. Prod.* **1996**, *59*, 773–776.
41. Zhang, W.; Guo, Y.W. Chemical Studies on the constituents of the Chinese medicinal herb *Euphorbia helioscopia* L. *Chem. Pharm. Bull.* **2006**, *54*, 1037–1039.
42. Borghi, D.; Baumer, L.; Ballabio, M.; Arlandini, E.; Perellino, N.C.; Minghetti, A.; Vincieri, F.F. Structure elucidation of helioscopinolides D and E from *Euphorbia calypttrata* cell cultures. *J. Nat. Prod.* **2004**, *54*, 1503–1508.
43. Haba, H.; Lavaud, C.; Magid, A.A.; Benkhaled, M. Diterpenoids and triterpenoids from *Euphorbia retusa*. *J. Nat. Prod.* **2009**, *72*, 1258–1264.
44. Morgenstern, T.; Bittner, M.; Silva, M.; Aqueveque, P.; Jakupovic, J. Diterpenes and phloracetophenones from *Euphorbia portulacoides*. *Phytochemistry* **1996**, *41*, 1149–1153.
45. Shi, J.S.; Li, Z.X.; Nitoda, T.; Minoru, I.; Hiroshi, K.; Naomichi, B.; Kazuyoshi, K.; Shuhei, N. Antinematodal activities of ingenane diterpenes from *Euphorbia kansui* and their derivatives against the pine wood nematode. *Z. Naturforsch., C: Biosci.* **2008**, *63*, 59–65.
46. Shi, J.X.; Li, Z.X.; Izumi, M.; Baba, N.; Nakajima, S. Termiticidal activity of diterpenes from the roots of *Euphorbia kansui*. *Z. Naturforsch., C: Biosci.* **2008**, *63*, 51–58.

47. Matsumoto, T.; Cyong, J.C.; Yamada, H. Stimulatory effects of ingenols from *Euphorbia kansui* on the expression of macrophage Fc receptor. *Planta Med.* **1992**, *58*, 255–258.
48. Sayed, M.D.; Ritzk, A.; Hammouda, F.M.; Elmissiry, M.M.; Williamson, E.M.; Evans, F.J. Constituents of Egyptian Euphorbiaceae. 9. Irritant and cyto-toxic ingenane esters from *Euphorbia paralias* L. *Experientia* **1980**, *36*, 1206–1207.
49. Marston, A.; Hecker, E. On the active principles of the Euphorbiaceae. 6. Isolation and biological activities of seven milliamines from *Euphorbia milii*. *Planta Med.* **1983**, *47*, 141–147.
50. Wang, L.Y.; Wang, N.L.; Yao, X.S.; Miyata, S.; Kitanaka, S. Studies on the bioactive constituents in Euphorbiaceae, part 3. Diterpenes from the roots of *Euphorbia kansui* and their in vitro effects on the cell division of *Xenopus*. *Chem. Pharm. Bull.* **2003**, *51*, 935–941.
51. Sorg, B.; Schmidt, R.; Hecker, E. Structure activity relationships of polyfunctional diterpenes of the ingenane type. I. Tumor-promoting activity of homologous, aliphatic 3-esters of ingenol and of $\Delta^7,8$ -isoinngenol-3-tetradecanoate. *Carcinogenesis* **1987**, *8*, 1–4.
52. Oksuz, S.; Ulubelen, A.; Barla, A.; Kohlbau, H.J.; Voelter, W. Triterpenoids and a diterpene from *Euphorbia iberica*. *Planta Med.* **1999**, *65*, 475–477.
53. Connolly, J.D.; Facunle, C.O.; Rycroft, D.S. Five ingol esters and a 17-hydroxyingenol ester from the latex of *Euphorbia kamerunica*. Assignment of esters using ^{13}C -NMR. methods. *Tetrahedron Lett.* **1984**, *25*, 3773–3776.
54. Urones, J.G.; Barcala, P.B.; Cuadrado, M.J.S.; Marcos, I.S. Diterpenes from the Latex of *Euphorbia broteri*. *Phytochemistry* **1988**, *27*, 207–212.
55. Tada, M.; Seki, H. Toxic diterpernes from *Euphorbia trigona* (Saiunkaku: an indoor foliage plant in Japan). *Agric. Biol. Chem.* **1989**, *53*, 425–430.
56. Shi, Y.P.; Jia, Z.J.; Ma, B.; Saleh, S.D.; Lahham, J. Ingenane diterpenes from *Euphorbia petiolata*. *Planta Med.* **1996**, *62*, 260–262.
57. Shi, J.G.; Jia, Z.J.; Yang, L. Lathyrane and ingenane diterpenoids from *Euphorbia micractina*. *Planta Med.* **1994**, *60*, 588–589.
58. Lu, Z.Q.; Yang, M.; Zhang, J.Q.; Chen, G.T.; Huang, H.L.; Guan, S.H.; Ma, C.; Liu, X.; Guo, D.A. Ingenane diterpenoids from *Euphorbia esula*. *Phytochemistry* **2008**, *69*, 812–819.
59. Marco, J.A.; Sanz-Cervera, J.F.; Yuste, A. Ingenane and lathyrane diterpenes from the latex of *Euphorbia canariensis*. *Phytochemistry* **1997**, *45*, 563–570.
60. Marco, J.A.; Sanz-Cervera, J.F.; Roperio, F.J.; Checa, J.; Fraga, B.M. Ingenane and lathyrane diterpenes from the latex of *Euphorbia acurensis*. *Phytochemistry* **1998**, *49*, 1095–1099.
61. Ott, H.H.; Hecker, E. Highly irritant ingenane type diterpene esters from *Euphorbia cyparissias* L. *Experientia* **1981**, *37*, 88–91.
62. Wu, D.G.; Sorg, B.; Hecker, E. Oligocyclic and macrocyclic diterpenes in thymelaeaceae and Euphorbiaceae occurring and utilized in Yunnan (Southwest China). 6. Tiglane type diterpene esters from latex of *Euphorbia prolifera*. *Phytother. Res.* **1994**, *8*, 95–99.
63. Marston, A.; Hecker, E. Active principles of the Euphorbiaceae. 7. Milliamine H and milliamine I, peptide esters of 20-deoxy-5 ξ -hydroxyphorbol from *Euphorbia milii*. *Planta Med.* **1984**, *50*, 319–322.
64. Wu, D.G.; Sorg, B.; Adolf W.; Seip, E.H.; Hecker, E. Oligo- and macrocyclic diterpenes in thymelaeaceae and Euphorbiaceae occurring and utilized in Yunnan (Southwest China). 2.

- Ingenane type diterpene esters from *Euphorbia nematocypha* Hand.-Mazz. *Phytother. Res.* **1992**, *6*, 237–240.
65. Furstenberger, G.; Hecker, E. On the active principles of the Euphorbiaceae, XII. Highly unsaturated irritant diterpene esters from *Euphorbia tirucalli* originating from Madagascar. *J. Nat. Prod.* **1986**, *49*, 386–397.
66. Liu, W.Z.; Wu, X.Y.; Yang, G.J.; Ma, Q.G.; Zhou, T.X.; Tang, X.C.; Qin, G.W. 12-deoxyphorbol esters from *Euphorbia fischeriana*. *Chin. Chem. Lett.* **1996**, *7*, 917–918.
67. Haba, H.; Lavaud, C.; Harkat, H.; Alabdul-Magid, A.; Marcourt, L.; Benkhalel, M. Diterpenoids and triterpenoids from *Euphorbia guyoniana*. *Phytochemistry* **2007**, *68*, 1255–1260.
68. Appendino, G.; Belloro, E.; Tron, G.C.; Jakupovic J, Ballero M. Diterpenoids from *Euphorbia pithyusa subsp. cupanii*. *J. Nat. Prod.* **1999**, *62*, 1399–1404.
69. Baloch, I.B.; Baloch, M.K.; Saqib, Q. Anti-tumor 12-deoxyphorbol esters from *Euphorbia cornigera*. *Eur. J. Med. Chem.* **2008**, *43*, 274–281.
70. Marco, J.A.; Sanz-Cervera, J.F.; Checa, J.; Palomares, E.; Fraga, B.M. Jatrophone and tiglane diterpenes from the latex of *Euphorbia obtusifolia*. *Phytochemistry* **1999**, *52*, 479–485.

Sample Availability: Samples of the compounds **8**, **12**, **61** and **94** are available from the authors.

© 2009 by the authors; licensee Molecular Diversity Preservation International, Basel, Switzerland. This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).