



Cyberchondria During the Coronavirus Pandemic: The Effects of Neuroticism and Optimism

Alexandra Maftei* and Andrei Corneliu Holman

Department of Psychology, Alexandru Ioan Cuza University, Iaşi, Romania

Health anxiety during the current coronavirus pandemic can be a serious psychological issue, amplified by the medical uncertainty around this disease and social isolation. As older people are especially at risk of becoming severely ill, it is important to examine the personal factors that make members of this age group more prone to health anxiety. Previous studies indicated that cyberchondria, i.e., the repeated online search for medical information, exacerbates health anxiety. The present research investigated the effect of two opposing traits, optimism and neuroticism, on cyberchondria during the COVID-19 pandemic. The associations of cyberchondria with demographic factors (age, gender, and education) were also examined. A sample of 880 participants, aged 15-67, 65% of whom were female, participated in an online survey. Results show that neuroticism, age, and being female are positively associated with cyberchondria. Optimism was found to be related to cyberchondria, but this effect was qualified by a significant interaction with age. Further analysis revealed that the effect of optimism was significant only in the highest age group. Moreover, among these elderly participants, the psychologically protective influence of optimism against cyberchondria emerged as larger than the opposite effect of neuroticism. This demonstrates the mental benefits of encouraging a positive outlook on the current health crisis and on one's personal resilience in facing it, especially among the elderly. Conversely, among people who use the Internet as a major source of medical information, those high in neuroticism may be more prone to cyberchondria.

Keywords: cyberchondria, coronavirus, neuroticism, optimism, age

INTRODUCTION

The coronavirus disease-2019 (COVID-19) emerged in December 2019, spread rapidly across the globe, and brought major changes to our lives, with countries all over the world imposing confinement measures (e.g., lockdowns and the closure of non-essential businesses) in order to avoid a rapid, uncontrolled spread of the virus and an overwhelming of medical systems. However, at the time of writing (August 2020), almost 24 million people have been infected across the globe, and more than 800,000 have already died. The COVID-19 outbreak brought an unprecedented disruption to people's personal and social lives, with complex psychological implications (i.e., distress, anxiety, depression, financial worry, loneliness, confusion, and anger) (Brooks et al., 2020; Lin et al., 2020; Tull et al., 2020; Zhang et al., 2020).

OPEN ACCESS

Edited by:

Daniel Bressington, Hong Kong Polytechnic University, Hong Kong

Reviewed by:

Paula Maria Di Nota, Justice Institute of British Columbia (JIBC), Canada Serena Giunta, University of Palermo, Italy

> *Correspondence: Alexandra Maftei psihologamaftei@gmail.com

Specialty section:

This article was submitted to Health Psychology, a section of the journal Frontiers in Psychology

Received: 29 May 2020 Accepted: 11 September 2020 Published: 30 October 2020

Citation:

Maftei A and Holman AC (2020) Cyberchondria During the Coronavirus Pandemic: The Effects of Neuroticism and Optimism. Front. Psychol. 11:567345. doi: 10.3389/fpsyg.2020.567345

1

The information rate shared via the Internet increased significantly during the pandemic, as in similar previous health crises (Sharma et al., 2017). Using social media sites or news platforms, people accessed news and various articles about the pandemic and COVID-19 related information and shared their experiences and concerns. The medical uncertainty around COVID-19, as well as the social isolation measures, raised severe psychological concerns, and online searches for information related to specific health symptoms (e.g., loss of smell or chest pain) increased during the pandemic (Walker et al., 2020). For example, people's searches for information about COVID-19 increased in the US by 36% 1 day after the announcement of the first COVID-19 case (Bento et al., 2020). Khasawneh et al. (2020) revealed that over 80% of medical students used social media platforms and online search engines as their primary source of information on COVID-19. In Romania, searches about symptoms, health-related issues, and treatment for COVID-19 increased exponentially, with a maximum reached in March 20201 (Google Trends, accessed August 23, 2020). However, research also suggested that online information about COVID-19 also contains a large amount of misleading information (Li et al., 2020), and public health agencies should aim to control the spread of misinformation concerning the virus to be able to more efficiently manage the pandemic.

In most cases, COVID-19 (the disease caused by the novel coronavirus) causes mild symptoms (such as dry cough, fever, or tiredness), though some people do not develop any symptoms. However, according to World Health Organization [WHO] (2020), 1 in 6 people may become seriously ill. The differences concerning the incidence and the severity of COVID-19 are multifaced and based on several complex factors (e.g., sociobiological characteristics or socioeconomic factors) (e.g., Hatef et al., 2020; Kopel et al., 2020); the elderly—especially those with pre-existing medical conditions (such as diabetes, cancer, lung disease, or high blood pressure)—are generally more prone to develop more severe cases of COVID-19 infections, compared to other groups (Fischer et al., 2020; Nanda et al., 2020; Niu et al., 2020).

CYBERCHONDRIA AND ASSOCIATED FACTORS

Though accessing medical information using online sources is a common, useful, and accessible strategy for most people, in some cases, when online searching becomes excessive and repetitive, it can turn into a pathological behavior (i.e., cyberchondria). Vismara et al. (2020) provided a systematic review of cyberchondria (CYB), confirming its significant role in the increase of health anxiety, distress, and obsessivecompulsive related behaviors. Although there is still no consensus on the definition of CYB, most researchers agree that this type of behavior is often driven by distress or anxiety about one's health (Starcevic and Berle, 2013), which subsequently increases both distress and anxiety (e.g., Belling, 2006; Recupero, 2010). More importantly, this search for online medical and health-related information is compulsive and hard to resist (Vismara et al., 2020).

Various international surveys suggested that almost 80% of Internet users used the Internet for medical appointments (Aiken et al., 2012; Fox and Duggan, 2013), while a sample of > 12,000 individuals suggested that almost half of the participants used the Google search engine for self-diagnosis (Mcdaid and Park, 2011). The constant coverage on both online and offline media of the COVID-19 pandemic might have contributed to an increase in health anxiety, particularly for people with CYB. The constant reminders of guidelines for preventions (i.e., wearing protective masks and gloves, washing hands, and avoiding social contact), along with updates related to the novel treatments for COVID-19, and infection and death rates, fueled health anxiety and exacerbated behaviors associated with CYB (Farooq et al., 2020; Hongbo et al., 2020).

Data suggests that younger individuals (aged 30–44) are the most active users when seeking health-related information via the Internet (Andreassen et al., 2007). However, the associations between age, gender, and CYB are scarce and contradictory (Vismara et al., 2020). Some studies have found no direct relationship between age and CYB (Barke et al., 2016); meanwhile, Doherty-Torstrick et al. (2016) suggested that older participants were less likely to experience an increase in anxiety due to the search for medical information, compared to younger participants.

In terms of gender, education, and their relation to CYB, Doherty-Torstrick et al. (2016) reported no significant differences between males and females, similar to Bajcar et al. (2019) and Akhtar and Fatima (2020). Zarcadoolas et al. (2002) suggested that medical information is one of the main topics researched online by the less educated. Meanwhile, Atkinson et al. (2009) reported that higher educated women (those with a Bachelor's degree) look for more health information online, compared to men and to less educated people.

DISPOSITIONAL OPTIMISM, NEUROTICISM, AND HEALTH

Dispositional optimism refers to "the generalized, relatively stable tendency to expect good outcomes across important life domains" (Scheier and Carver, 2018, p. 1082) and is considered a stable trait over time. The effects of dispositional optimism on physical health were widely examined. For example, Carver and Scheier (1981, 1998) were among the first researchers to suggest that optimistic people engage more in efforts to fight difficulties when they experience adversity and are more likely to achieve better outcomes. Previous research suggested that individuals higher in dispositional optimism generally engage in more protective health-related behaviors than people low in this specific trait (Carvajal et al., 1998; Giltay et al., 2007; Krane et al., 2018). Scheier et al. (1989) explored the associations between dispositional optimism and an objective physical health outcome (i.e., a heart attack). They found that individuals higher

¹https://trends.google.com/trends/explore?date=today%203-m&geo=RO&q= coronavirus%20simptome%20tratament

in optimism were significantly less likely to suffer a heart attack during a medical procedure (such as surgery).

Additionally, other researchers confirmed the links between optimism and the progression of certain diseases (e.g., atherosclerosis, Matthews et al., 2004; coronary heart disease, Tindle et al., 2009; stroke, Kim et al., 2011). People high in dispositional optimism also seem to have higher levels of cognitive functioning following traumatic brain injury (Lee et al., 2019). Dispositional optimism seems to be as important for one's mental health as it is for their physical health. For example, Liu et al. (2016) suggested that participants' higher levels of dispositional optimism were associated with lower levels of perceived stress and depression. Similar findings were reported by He et al. (2016), who highlighted the mediating role of optimism on the relationship between perceived social support and depression. Finally, optimism is generally negatively associated with maladaptive coping strategies (Segerstrom, 2006), and positively correlated to self-confidence, a general better adjustment in the face of adverse and traumatic life events, and healthier coping strategies compared to individuals low in dispositional optimism (Nes, 2016; Reed, 2016). During the COVID-19 pandemic, optimism was found to be significantly associated with a higher level of preventive behaviors (Jovančević and Milićević, 2020).

According to McCrae and Costa (1994), neuroticism is a temperamental, stable predisposition toward dysfunctions "that reflects the tendency to experience negative emotions, cognitions, and maladaptive behaviors" (Bajcar and Babiak, 2020, p. 1). Neuroticism was found to be a significant risk factor for CYB by Bajcar and Babiak (2020), in line with previous research that linked neuroticism and other personality traits (i.e., extraversion and conscientiousness) to health-related seeking behaviors (e.g., Lagoe and Atkin, 2015; Jacobs et al., 2017; Fergus and Spada, 2018), which are generally mediated by health anxiety (e.g., Lagoe and Atkin, 2015). Neuroticism might also harm a person's immune system through "a "predisposition model," wherein effects of stressors on the immune system are dependent on personality (Khosravi, 2020, p. 1.).

Concerning the COVID-19 pandemic, neuroticism was suggested as a marker of vulnerability to COVID-19 infection by Khosravi (2020), in line with Zajenkowski et al. (2020), and Abdelrahman (2020), who suggested that neuroticism may be positively associated with adopting social distancing to avoid potential COVID-19 infection. Additionally, higher neuroticism was related to more concerns and longer duration estimates related to the COVID-19 pandemic (Aschwanden et al., 2020), in line with similar findings (e.g., Weiss and Deary, 2020). Finally, Kroencke et al. (2020) suggested that individuals high in neuroticism might experience more negative effects during the COVID-19 pandemic.

THE PRESENT STUDY

Although various studies suggested the significant associations between optimism, neuroticism, and health-related behaviors, and a growing number of COVID-19 studies point to the importance of psychological traits for predicting pandemicrelated behavior, there is still a scarcity of research concerning the relationship between dispositional optimism, neuroticism, and cyberchondria. Therefore, this study aimed to investigate the effects of optimism and neuroticism on CYB during the COVID-19 crisis. Additionally, we examined the associations between CYB and age, gender, and education. We also investigated whether the effects of the two psychological traits assessed on cyberchondria are moderated by age, given the increased COVID-19–related risk faced by the elderly.

METHODS

Procedure and Participants

We ran a web-based, cross-sectional survey at the beginning of April 2020, a few weeks after governments all over the world, including Romania, imposed numerous restrictions (i.e., social confinement) to prevent the spread of COVID-19. The study was approved by the Research Ethics Committee from the institution where the authors are affiliated. The survey was distributed to and by students enrolled at the university where the authors are affiliated, as a course credit requirement. The survey link was available for 10 days and posted in academic and social media groups, or by using other media channels (such as e-mails or other online communication groups). Participation was voluntary. Our final convenience sample consisted of 880 participants (57 were dropped from the study due to missing data). Their age varied from 15 to 67 (M = 34.36, SD = 10.17), most of them being females (64.8%), with a Bachelor's degree (48.6%). Participants answered anonymously after being presented with an informed consent form describing the aim of the study and ensuring the confidentiality of their answers.

Instruments

Participants answered the revised form of **The Life Orientation Test** (LOT; Scheier et al., 1994) which measures dispositional optimism on a 5-point Likert scale (1-strongly disagree, 5strongly agree) using 10 items (out of which four were filler items). Cronbach's alpha indicated satisfying reliability (0.715) of the scale.

We used the **Cyberchondria Severity Scale** (CSS; McElroy and Shevlin, 2014) to assess people's excessive online searching for health-related issues, and the way this conduct affects their daily routine. Participants in our sample answered on a 5-point Likert scale (1-Never, 5-Always) to 33 items further divided into five different dimensions—*Compulsion, Distress, Excessiveness, Reassurance,* and *Mistrust of medical professionals*—assessing their related behavior within the past 2 weeks. High total scores indicated high levels of CYB. Cronbach's alpha indicated high reliability (0.947) of the scale.

The Neuroticism Scale from the international personality item pool—IPIP (1996)—was used to measure psychological distress. The 10-item instrument was translated and adapted for the Romanian population by Iliescu et al. (2015). Cronbach's alpha indicated good reliability (0.865) of the scale. A demographic scale assessed participants' age, gender, and education (i.e., high school, Bachelor's degree, or postgraduate).

RESULTS

We computed the Spearman correlations between study variables, as the results of the Shapiro-Wilk tests of normality indicated that CYB, LOT, NS, and age are not normally distributed. These correlations and normality statistics are presented in **Table 1**, together with descriptive statistics on all variables. CYB was found to have significant but weak negative associations to optimism (i.e., LOT) and age, and to be positively related, albeit to the same weak order of magnitude, to neuroticism (i.e., NS) and gender, with female participants scoring higher than males.

Next, we investigated the effect of neuroticism, optimism, age, gender, and education on CYB through hierarchical multiple regression analysis. The three sociodemographic variables were entered in the first step, while neuroticism and optimism were entered in the second. In order to explore whether age moderates the effect of these two psychological dimensions on CYB, we introduced the interaction term (i.e., product) between neuroticism and age, as well as the interaction between optimism and age in the third step.

Results, summarized in Table 2, showed that in the third model age, gender, neuroticism, and optimism were significant predictors of CYB. However, while neuroticism was positive, as in the previous correlational analysis, the relationships of the other three variables to cyberchondria emerged as opposed to those indicated by simple correlations. Thus, when controlling the other dimensions, the standardized coefficients in the third step of the regression analysis indicate that male and older participants score higher on CYB than their female and younger counterparts. Moreover, optimism was indicated as having a positive association with CYB, but this main effect was qualified by a significant interaction between LOT and age, while the interaction term between neuroticism and age did not significantly account for cyberchondria. In order to explore the interaction between optimism and age, we analyzed the effect of optimism at each age level through multiple regression analyses. To this aim, we used the mean and standard deviation of the age distribution to split the sample into three groups.

TABLE 2 | Hierarchical regression model for cyberchondria (CYB).

	Step 1	Step 2	Step 3	
	β	β	β	
CSS				
Age	-0.08*	-0.02	0.72*	
Gender	0.11**	0.07*	0.07*	
Education	-0.07*	-0.06	-0.06	
NS		0.28**	0.50**	
LOT		-0.07	0.32*	
NS x Age			-0.26	
LOT x Age			-0.73**	
ΔR^2	0.02**	0.10**	0.01*	

*p < 0.01; **p < 0.001.

LOT, Life Orientation Test; NS, Neuroticism Scale.

TABLE 3 Hierarchical regression model for cyberchondria (CN	ŕВ)
at each age level.	

	Lowest age (under 24 years) <i>N</i> = 166	Medium age (25–44 years) <i>N</i> = 572	Highest age (over 45 years) <i>N</i> = 142	
css				
Age	-0.12	-0.06	0.10	
Gender	0.10	0.05	0.08	
Education	-0.14	-0.01	-0.21**	
NS	0.34**	0.29**	0.18*	
LOT	0.10	-0.08	-0.22*	
R^2	0.14**	0.13**	0.19**	

*p < 0.01; **p < 0.001.

LOT, Life Orientation Test; NS, Neuroticism Scale.

The results of the analysis regression of CYB on the three sociodemographic variables, neuroticism, and optimism in each of the three groups defined by their age level are presented in **Table 3**. Results indicate that neuroticism is a significant positive factor of CYB at every age level. Optimism did not emerge as a significant predictor in the first two age groups, but it negatively and significantly predicted CYB in participants with the highest age. Moreover, at this age level, the effect of optimism on CYB, as indicated by standardized regression coefficients, emerged as stronger than that of neuroticism. The association of education to CYB scores was found to be similar to that of

	1	2	3	4	5	6	М	SD	Shapiro-Wilk W statistic
1. CSS	1	-0.26**	0.33**	-0.08*	0.10*	-0.05	2.29	0.73	0.97**
2. LOT		1	-0.59**	0.17**	0.006	0.12**	3.61	0.73	0.98**
3. NS			1	-0.18**	0.13**	-0.03	2.30	0.76	0.98**
4. Age				1	0.06	0.21**	34.36	10.17	0.98**
5. Gender					1	0.17**	35.2% males, 64.8% females		
6. Education						1	23.2% highschool, 48.6% Bachelor's degree		

*p < 0.05: **p < 0.01.

CSS, Cyberchondria Severity Scale; LOT, Life Orientation Test; NS, Neuroticism Scale.

optimism across the three age groups, as education emerged as a significant negative factor of cyberchondria only in participants in the highest age group.

DISCUSSION

In a survey-based, cross-sectional study, we examined the effect of two opposite dispositional traits, i.e., optimism and neuroticism, and a series of demographic factors (gender, education, and age) on cyberchondria during the COVID-19 pandemic. We were also interested in the way age moderates the effects of the two psychological factors examined, given the fact that COVID-19 is especially dangerous for older people. Results suggested that optimism decreases the likelihood that people in the most vulnerable COVID-19 group (i.e., older individuals) will experience high levels of cyberchondria, emphasizing the importance of this psychological trait in overcoming the health-related anxiety that may be associated with CYB and amplified by the current crisis, in this specific age group. These findings extend the previously documented positive influence on optimism on protective health-related behaviors (e.g., Krane et al., 2018), suggesting that this active stance in the area of health management also protects highly optimistic individuals from excessive online searching for medical information during the current COVID-19 crisis, and from the psychological consequences of this behavior.

Given this benefit of optimism during the current pandemic, it is essential to consider ways in which it might be enhanced and/or pessimism reduced, especially among the elderly. For example, previous studies suggested that engaging in social activities, religious involvement, social support, physical activities, or practicing gratitude might enhance optimism and coping with adverse life situations (e.g., Greenglass et al., 2006; Giltay et al., 2007; Prati and Pietrantoni, 2009; Greene and McGovern, 2017; Progovac et al., 2017; Oberle et al., 2018; Wong et al., 2018). Health professionals and national authorities fighting the pandemic may benefit from the findings by focusing on the significant role of optimism for older people who use the Internet as a major source of medical information during the current health crisis.

The pattern of findings indicates that people high in neuroticism are especially prone to develop CYB manifestations irrespective of their age, in line with past studies on the relationship of this personality trait with CYB (Bajcar and Babiak, 2020), as well as with health anxiety and healthrelated behaviors (Lagoe and Atkin, 2015; Fergus and Spada, 2018). Results also suggest that when controlling for the effects of neuroticism, optimism, gender, and education, CYB is positively related to age, a result that contradicts previous findings (Doherty-Torstrick et al., 2016), while also highlighting the psychologically vulnerable status of the elderly during the COVID-19 crisis.

Besides other limits, such as its reliance on a convenience sample, which undermines the generalizability of the current findings, one of the limits of this study is that we did not measure health anxiety. This precludes us from concluding the degree to which the relationships that emerged in our results extend more generally in people's health anxiety. Further research should investigate the relationships between age, CYB, and health anxiety in the context of the COVID-19 pandemic and its psychological effects. For instance, McMullan et al. (2019) suggested that age might moderate the relationship between health anxiety and CYB. Future studies could examine whether this pattern of findings is altered by increased health risks faced by the older population during the current pandemic. Furthermore, to increase the generalizability of the current findings, future studies should rely on non-convenient, larger samples. Longitudinal approaches might also express in more detail whether and how the evolution of the current pandemic affects the nature and direction of relationships in the main variables explored in the current research.

To conclude, this study found that optimism is a psychologically protective factor against CYB during the current health crisis in the most vulnerable age group, i.e., the elderly. This effect runs against the general influence of neuroticism, which amplifies the risk of CYB. The practical implications of the current findings are various and mainly important for policy makers, clinicians, and healthcare systems in general. For example, public policies should find effective ways to promote rapid and strategic ways to increase optimism as a protective measure against CYB during and post the COVID-19 pandemic. Additionally, as Khosravi (2020) suggested, given the important association between neuroticism and CYB, self-report scales might be useful for the initial screening of individuals high in neuroticism, to launch personality-tailored prevention campaigns that might reduce CYB and its negative consequences during the pandemic. (such as disseminating information about ways to fight the current crisis, especially among individuals of high neuroticism, using optimism-enhancing strategies, and advertising the negative impact of excessive and repetitive online searching for COVID-19 related information).

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation, to any qualified researcher.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by the Institutional Review Board of Alexandru Ioan Cuza University in Iasi. Written informed consent to participate in this study was provided by the participants, and where necessary, the participants' legal guardian/next of kin.

AUTHOR CONTRIBUTIONS

Both authors contributed equally to conceive and design the main goal of the study, analyze the data, and write the manuscript.

REFERENCES

- Abdelrahman, M. (2020). Personality Traits, Risk Perception, and Protective Behaviors of Arab Residents of Qatar During the COVID-19 Pandemic. Int. J. Ment. Health Addict. 2020, 1–12.
- Aiken, M., Kirwan, G., Berry, M., and O'Boyle, C. O. (2012). The age of cyberchondria. Royal Coll. Surg. Ireland 5, 71–74.
- Akhtar, M., and Fatima, T. (2020). Exploring cyberchondria and worry about health among individuals with no diagnosed medical condition. J. Pakistan Med. Assoc. 70, 90–95. doi: 10.5455/JPMA.8682
- Andreassen, H. K., Bujnowska-Fedak, M. M., Chronaki, C. E., Dumitriu, R. C., Pudule, I., Santana, S., et al. (2007). European citizens' use of E-health services: A study of seven countries. *BMC Publ. Health* 7:53. doi: 10.1186/1471-24 58-7-53
- Aschwanden, D., Strickhouser, J. E., Sesker, A. A., Lee, J., Luchetti, M., Stephan, Y., et al. (2020). Psychological and Behavioural Responses to Coronavirus Disease 2019: The Role of Personality. *Eur. J. Personal.* 2020:2281 doi: 10.1002/per.2281
- Atkinson, N., Saperstein, S., and Pleis, J. (2009). Using the Internet for healthrelated activities: findings from a national probability sample. *J. Med. Internet Res.* 11:å4. doi: 10.2196/jmir.1035
- Bajcar, B., and Babiak, J. (2020). Neuroticism and cyberchondria: The mediating role of intolerance of uncertainty and defensive pessimism. *Personal. Individ. Diff.* 162:110006 doi: 10.1016/j.paid.2020.110006
- Bajcar, B., Babiak, J., and Olchowska-Kotala, A. (2019). Cyberchondria and its measurement. The polish adaptation and psychometric properties of the cyberchondria severity scale CSS-PL. *Pol.Psych.* 53, 49–60. doi: 10.12740/PP/ 81799
- Barke, A., Bleichhardt, G., Rief, W., and Doering, B. K. (2016). The Cyberchondria Severity Scale (CSS): German validation and development of a short form. *Int. J. Behav. Med.*23, 595–605. doi: 10.1007/s12529-016-9549-8
- Belling, C. (2006). Hypochondriac hermeneutics: medicine and the anxiety of interpretation. *Literat. Med.* 25, 376–401. doi: 10.1353/lm.2007.0000
- Bento, A. I., Nguyen, T., Wing, C., Lozano-Rojas, F., Ahn, Y.-Y., and Simon, K. (2020). Evidence from internet search data shows informationseeking responses to news of local COVID-19 cases. *Proc. Natl. Acad. Sci.* 117:202005335. doi: 10.1073/pnas.2005335117
- Brooks, S. K., Webster, R. K., Smith, L. E., Woodland, L., Wessely, S., Greenberg, N., et al. (2020). The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet* 395, 912–920. doi: 10.1016/S0140-6736(20) 30460-8
- Carvajal, S. C., Garner, R. L., and Evans, R. I. (1998). Dispositional optimism as a protective factor in resisting HIV exposure in sexually active inner-city minority adolescents. *J. Appl. Soc. Psychol.* 28, 2196–2211. doi: 10.1111/j.1559-1816.1998. tb01367.x
- Carver, C. S., and Scheier, M. F. (1981). Attention and self-regulation: A controltheory approach to human behavior. New York, NY: Springer Verlag.
- Carver, C. S., and Scheier, M. F. (1998). On the self-regulation of behavior. New York,NY: Cambridge University Press.
- Doherty-Torstrick, E. R., Walton, K. E., and Fallon, B. A. (2016). Cyberchondria: parsing health anxiety from online behavior. *Psychosomatics* 57, 390–400. doi: 10.1016/j.psym.2016.02.002
- Farooq, A., Laato, S., and Islam, A. K. M. N. (2020). Impact of Online Information on Self-Isolation Intention During the COVID-19 Pandemic: Cross-Sectional Study. J. Med. Internet Res. 22:e19128. doi: 10.2196/19128
- Fergus, T. A., and Spada, M. M. (2018). Moving toward a metacognitive conceptualization of cyberchondria: Examining the contribution of metacognitive beliefs, beliefs about rituals, and stop signals. J. Anxiety Disord. 60, 11–19. doi: 10.1016/j.janxdis.2018.09.003
- Fischer, F., Raiber, L., Boscher, C., and Winter, M. H.-J. (2020). COVID-19 and the Elderly: Who Cares? *Front.Public Health* 8:151. doi: 10.3389/fpubh.2020. 00151
- Fox, S., and Duggan, M. (2013). Health Online 2013. Pew Research Center's Internet & American Life Project, Vol. 15. Available online at: http://bibliobase.sermais. pt:8008/BiblioNET/Upload/PDF5/003820.pdf (accessed January 15, 2013).
- Giltay, E. J., Geleijnse, J. M., Zitman, F. G., Buijsse, B., and Kromhout, D. (2007). Lifestyle and dietary correlates of dispositional optimism in men: The Zutphen Elderly Study. *J. Psychosom. Res.* 63, 483–490. doi: 10.1016/j.jpsychores.2007. 07.014

- Greene, N., and McGovern, K. (2017). Gratitude, psychological well-being, and perceptions of posttraumatic growth in adults who lost a parent in childhood. *Death stud.* 41, 436–446. doi: 10.1080/07481187.2017.1296505
- Greenglass, E., Fiksenbaum, L., and Eaton, J. (2006). The relationship between coping, social support, functional disability and depression in the elderly. *Anxiety Stress Coping Int. J.* 19, 15–31. doi: 10.1080/14659890500436430
- Hatef, E., Chang, H.-Y., Kitchen, C., Weiner, J. P., and Kharrazi, H. (2020). Assessing the Impact of Neighborhood Socioeconomic Characteristics on COVID-19 Prevalence Across Seven States in the United States. *Front. Public Health* 2020:571808.doi: 10.3389/fpubh.2020.571808
- Hongbo, L., Iqbal, A., and Waqas, M. A. (2020). Psychological predictors of anxiety in response to the covid-19 pandemic: evidence from Pakistan. Available online at: https://psyarxiv.com/gk6mt/ (accessed May 21, 2020).
- Iliescu, D., Popa, M., and Dimache, R. (2015). Adaptarea românească a Setului International de Itemi de Personalitate: IPIP-Ro [The Romanian adaptation of the International Personality Item Pool: IPIP-Ro]. *Psihol. Res. Umane* 13, 83–112.
- IPIP (1996). International Personality Item Pool. A Scientific Collaboratory for the Development of Advanced Measures of Personality and Other Individual Differences. Available online at: https://ipip.ori.org/ (accessed August 23, 2020).
- Jacobs, W., Amuta, A. O., and Jeon, K. C. (2017). Health information seeking in the digital age: An analysis of health information seeking behavior among US adults. *Cogent Soc. Sci* 3, 1–11. doi: 10.1080/23311886.2017.1302785
- Jovančević, A., and Milićević, N. (2020). Optimism-pessimism, conspiracy theories and general trust as factors contributing to COVID-19 related behavior – A cross-cultural study. *Personal. Individ. Differ*. 167:110216. doi: 10.1016/j.paid. 2020.110216
- Khasawneh, A. I., Humeidan, A. A., Alsulaiman, J. W., Bloukh, S., Ramadan, M., Al-Shatanawi, T. N., et al. (2020). Medical Students and COVID-19: Knowledge, Attitudes and Precautionary Measures. A Descriptive Study from Jordan. Front. Publ. Health 8:253. doi: 10.3389/fpubh.2020.00253
- Khosravi, M. (2020). Neuroticism as a Marker of Vulnerability to COVID-19 Infection. Psych. Investig. 17, 710–711. doi: 10.30773/pi.2020.0199
- Kim, E. S., Park, N., and Peterson, C. (2011). Dispositional optimism protects older adults from stroke: The Health and Retirement Study. *Stroke* 42, 2855–2859. doi: 10.1161/strokeaha.111.613448
- Kopel, J., Perisetti, A., Roghani, A., Aziz, M., Gajendran, M., and Goyal, H. (2020). Racial and Gender-Based Differences in COVID-19. *Front. Publ. Health* 8:418. doi: 10.3389/fpubh.2020.00418
- Krane, A., Terhorst, L., Bovbjerg, D. H., Scheier, M. F., Kucinski, B., Geller, D. A., et al. (2018). Putting the life in lifestyle: Lifestyle choices after a diagnosis of cancer predicts overall survival. *Cancer* 124, 3417–3426. doi: 10.1002/cncr. 31572
- Kroencke, L., Geukes, K., Utesch, T., Kuper, N., and Back, M. (2020). Neuroticism and emotional risk during the COVID-19 pandemic. PsyArXiv [Preprint]. doi: 10.31234/osf.io/8c6nh
- Lagoe, C., and Atkin, D. (2015). Health anxiety in the digital age: An exploration of psychological determinants of online health information seeking. *Comput. Hum. Behav.* 52, 484–491.. doi: 10.1016/j.chb.2015.06.003
- Lee, E., Jayasinghe, N., Swenson, C., and Dams-O'Connor, K. (2019). Dispositional optimism and cognitive functioning following traumatic brain injury. *Brain Injury* 33, 985–990. doi: 10.1080/02699052.2019.1606448
- Li, H. O., Bailey, A., Huynh, D., and Chan, J. (2020). YouTube as a source of information on COVID-19: a pandemic of misinformation? *BMJ Glob.Health* 5: e002604. doi: 10.1136/bmjgh-2020-002604
- Lin, Y., Hu, Z., Alias, H., and Wong, L. P. (2020). Knowledge, Attitudes, Impact, and Anxiety Regarding COVID-19 Infection Among the Public in China. *Front. Publ. Health* 8:236. doi: 10.3389/fpubh.2020.00236
- Liu, B., Pu, J., and Hou, H. (2016). Effect of perceived stress on depression of Chinese "Ant Tribe" and the moderating role of dispositional optimism. *J. Health Psychol.* 21, 2725–2731. doi: 10.1177/1359105315583373
- Matthews, K. A., Raikkonen, K., Sutton-Tyrrell, K., and Kuller, L. H. (2004). Optimistic attitudes protect against progression of carotid atherosclerosis in healthy middle-aged women. *Psychosomatic Med.* 66, 640–644. doi: 10.1097/ 01.psy.0000139999.99756.a5
- Mcdaid, D., and Park, A. L. (2011). *Online Health: Untangling the Web. BUPA Health Pulse 2010**. Available online at: www.bupa.com/healthpulse (accessed August 23, 2020).

- McElroy, E., and Shevlin, M. (2014). The development and initial validation of the cyberchondria severity scale (CSS). J. Anxiety Disord.28, 259–265. doi: 10.1016/ j.janxdis.2013.12.007
- McMullan, R. D., Berle, D., Arnaez, S., and Startevic, V. (2019). The relationships between health anxiety, online health information seeking, and cyberchondria: Systematic review and meta-analysis. J. Affect. Disord. 245, 270–278. doi: 10. 1016/j.jad.2018.11.037
- Nanda, A., Vura, N., and Gravenstein, S. (2020). COVID-19 in older adults. Aging Clin. Exp. Res.32, 1199–1202. doi: 10.1007/s40520-020-01581-5
- Nes, L. S. (2016). "Optimism, pessimism, and stress," in Stress: Concepts, cognition, emotion, and behavior, ed. G. Fink (Cambridge,MA: Elsevier Academic Press), 405–411. doi: 10.1016/b978-0-12-800951-2.00052-2
- Niu, S., Tian, S., Lou, J., Kang, X., Zhang, L., Lian, H., et al. (2020). Clinical characteristics of older patients infected with COVID-19: A descriptive study. *Archiv. Gerontol. Geriat.* 89:104058. doi: 10.1016/j.archger.2020.104058
- Oberle, E., Guhn, M., Gadermann, A. M., Thomson, K., and Schonert-Reichl, K. A. (2018). Positive mental health and supportive school environments: A population-level longitudinal study of dispositional optimism and school relationships in early adolescence. *Soc. Sci. Med.* 214, 154–161. doi: 10.1016/j. socscimed.2018.06.041
- Prati, G., and Pietrantoni, L. (2009). Optimism, social support, and coping strategies as factors contributing to posttraumatic growth: A meta-analysis. J. Loss Trauma 14, 364–388. doi: 10.1080/15325020902724271
- Progovac, A. M., Donohue, J. M., Matthews, K. A., Chang, C. H., Habermann, E. B., Kuller, L. H., et al. (2017). Optimism predicts sustained vigorous physical activity in postmenopausal women. *Prevent. Med. Rep.*8, 286–293. doi: 10.1016/ j.pmedr.2017.10.008
- Recupero, P. R. (2010). The mental status examination in the age of the Internet. J. Am. Acad. Psych. Law 38, 15–26.
- Reed, D. J. (2016). Coping with occupational stress: the role of optimism and coping flexibility. *Psychol. Res. Behav. Manage.* 9, 71–79. doi: 10.2147/prbm. s97595
- Scheier, M. F., and Carver, C. S. (2018). Dispositional Optimism and Physical Health: A Long Look Back, a Quick Look Forward. Am. Psychol. 73, 1082–1094. doi: 10.1037/amp0000384
- Scheier, M. F., Carver, C. S., and Bridges, M. W. (1994). Distinguishing optimism from neuroticism (and trait anxiety, self-mastery, and self-esteem): A reevaluation of the Life Orientation Test. J. Personal. Soc. Psychol. 67, 1063–1078. doi: 10.1037/0022-3514.67.6.1063
- Scheier, M. F., Matthews, K. A., Owens, J., Magovern, G. J. Sr., Lefebvre, R. C., Abbott, R. A., et al. (1989). Dispositional optimism and recovery from coronary artery bypass surgery: The beneficial effects on physical and psychological wellbeing. *J. Personal. Soc. Psychol.* 57, 1024–1040. doi: 10.1037/0022-3514.57.6. 1024
- Segerstrom, S. C. (2006). How does optimism suppress immunity? Evaluation of three affective pathways. *Health Psychol.* 25, 653–657. doi: 10.1037/0278-6133. 25.5.653
- Sharma, M., Yadav, K., Yadav, N., and Ferdinand, K. C. (2017). Zika virus pandemic-analysis of Facebook as a social media health information platform. Am. J. Infect. Control 45, 301–302. doi: 10.1016/j.ajic.2016. 08.022

- Starcevic, V., and Berle, D. (2013). Cyberchondria: towards a better understanding of excessive health-related internet use. *Exp. Rev. Neurother.* 13, 2015–2213. doi: 10.1586/ern.12.162
- Tindle, H. A., Chang, Y. F., Kuller, L. H., Manson, J. E., Robinson, J. G., Rosal, M. C., et al. (2009). Optimism, cynical hostility, and incident coronary heart disease and mortality in the Women's Health Initiative. *Circulation* 120, 656–662. doi: 10.1161/circulationaha.108.827642
- Tull, M. T., Edmonds, K. A., Scamaldo, K. M., Richmond, J. R., Rose, J. P., and Gratz, K. L. (2020). Psychological Outcomes Associated with Stay-at-Home Orders and the Perceived Impact of COVID-19 on Daily Life. *Psychiatry Res.* 289:113098. doi: 10.1016/j.psychres.2020.113098
- Vismara, M., Caricasole, V., Starcevic, V., Cinosi, E., Dell'Osso, B., Martinotti, G., et al. (2020). Is cyberchondria a new transdiagnostic digital compulsive syndrome? A systematic review of the evidence. *Compr. Psych.* 99:152167. doi: 10.1016/j.comppsych.2020.152167
- Walker, A., Hopkins, C., and Surda, P. (2020). Use of Google Trends to investigate loss-of-smell-related searches during the COVID-19 outbreak. *Int. Allergy Rhinol.* 10, 839–847. doi: 10.1002/alr.22580
- Weiss, A., and Deary, I. J. (2020). A New Look at Neuroticism: Should We Worry So Much About Worrying? *Curr. Direct. Psychol. Sci.* 29, 92–101. doi: 10.1177/ 0963721419887184
- Wong, Y. J., Owen, J., Gabana, N. T., Brown, J. W., McInnis, S., Toth, P., et al. (2018). Does gratitude writing improve the mental health of psychotherapy clients? Evidence from a randomized controlled trial. *Psychother. Res. J. Soc.Psychother. Res.* 28, 192–202. doi: 10.1080/10503307.2016.1169332
- World Health Organization [WHO] (2020). Q&A: Older people and COVID-19. Available online at: https://www.who.int/news-room/q-a-detail/q-a-onon-covid-19-for-older-people (accessed 8 May 2020).
- Zajenkowski, M., Jonason, P. K., Leniarska, M., and Kozakiewicz, Z. (2020). Who complies with the restrictions to reduce the spread of COVID-19? Personality and perceptions of the COVID-19 situation. *Personal. Individ. Differ.* 166:110199.8.
- Zarcadoolas, C., Blanco, M., Boyer, J. F., and Pleasant, A. (2002). Unweaving the Web: An Exploratory Study of Low-Literate Adults'. Navigation Skills on the World Wide Web. J. Health Commun.7, 309–324. doi: 10.1080/ 10810730290088157
- Zhang, S. X., Wang, Y., Rauch, A., and Wei, F. (2020). Unprecedented disruption of lives and work: Health, distress and life satisfaction of working adults in China one month into the COVID-19 outbreak. *Psychiatry Res*.288:112958. doi: 10.1016/j.psychres.2020.112958

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Maftei and Holman. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.