



Review Article

Saccular Kommerell aneurysm, a potential pitfall on MDCT imaging – A review of imaging features and potential mimics

Hui Lin Wong*, Charlene Jin Yee Liew, Angeline Choo Choo Poh

Diagnostic Radiology, Changi General Hospital, 2 Simei Street 3, 529889, Singapore

ARTICLE INFO

Keywords:

Kommerell aneurysm
Multi-detector computed tomography
Vascular
Thoracic aorta
Diverticulum

ABSTRACT

Saccular Kommerell aneurysm represents a potential pitfall on Multidetector CT (MDCT) imaging, mimicking conditions such as saccular aneurysm of the thoracic aorta, ductus diverticulum and dilated Kommerell diverticulum. Accurate diagnosis of this condition is critical in the management of this potentially fatal condition. This paper reviews the MDCT imaging features of Kommerell aneurysms and its mimics and demonstrates how to make an accurate diagnosis through a series of four cases. MDCT features of Kommerell aneurysms, either saccular or fusiform types arising from a Kommerell diverticulum with atherosclerotic plaque and mural thrombus are discussed.

1. Introduction

Aberrant right subclavian artery is encountered in approximately 1% of the population and Kommerell diverticulum is present in about 60% of cases of aberrant subclavian artery, hence both conditions are not rare. Kommerell diverticulum was first eponymously used in 1936 to describe a diverticulum at the origin of an aberrant right subclavian artery in a left-sided aortic arch [1], although it may be used to describe a diverticulum in either an aberrant right or left subclavian artery arising from a left or right-sided aortic arch. Aneurysms of this diverticulum are seen in 3–8% of imaging studies, usually representing fusiform aneurysmal dilation of the diverticulum at the origin of the aberrant subclavian artery or transformation of the diverticulum into a saccular aneurysm [2]. An aneurysm of a Kommerell diverticulum is often abbreviated to the term Kommerell aneurysm.

Occasionally, as depicted in a case below, saccular Kommerell aneurysms may grow to such a size as to completely obliterate the normal appearance of its underlying diverticulum – a configuration easily confused with a saccular aneurysm of the descending thoracic aorta. The importance of making this distinction in the radiology reading room is twofold – Kommerell aneurysms and saccular aneurysms of the descending thoracic aorta are managed differently, and Kommerell aneurysms have a predilection for rupture [3,4]. With this in mind, the objective of this paper is to review the MDCT imaging features of Kommerell aneurysms and its mimics and to demonstrate how to make an accurate diagnosis through a series of cases.

2. Clinical presentation

Most patients with Kommerell aneurysms present with symptoms of dysphagia, followed by dyspnea and chest pain [3,5–7]. A significant proportion of patients are completely asymptomatic. Symptoms are determined by the location of the aneurysm and the course of the aberrant subclavian artery, which has been extensively described in the literature as being retro-esophageal (80%), between the esophagus and trachea (15%), or pre-tracheal (5%). [8–13].

2.1. Management

Kommerell aneurysms and diverticula are normally repaired in a two-staged thoracotomy procedure in which the descending aorta and aberrant subclavian artery are approached separately [14–20] and re-constructed with grafts. On the other hand, saccular aneurysms of the descending thoracic aorta are commonly managed with a one-stage endovascular graft repair [21–23] or open surgery [24–27].

2.2. Prognosis

Kommerell aneurysms are prone to rupture and dissection, an observation not lost on Dr Benjamin Felson, who astutely remarked that a surprisingly high incidence of rupture was noted in aneurysms of an anomalous right subclavian artery in 1989 [4]. Pathological specimens have been observed to contain mural abnormalities such as medial necrosis in up to 50% of cases [28,29], which may account for this

Abbreviations: MDCT, multidetector CT; 3D, three-dimensional

* Corresponding author.

E-mail address: huilin.wong@mohh.com.sg (H.L. Wong).

<http://dx.doi.org/10.1016/j.ejro.2017.06.003>

Received 24 May 2017; Received in revised form 25 June 2017; Accepted 26 June 2017

Available online 11 July 2017

2352-0477/ © 2017 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Table 1
Features of Kommerell aneurysm and mimics on MDCT imaging.

	Origin of the aneurysm/outpouching	Outpouching contents	Morphology
Kommerell aneurysm	Arising from a Kommerell diverticulum	Atherosclerotic plaque, mural thrombus common in saccular form	Saccular or fusiform
Kommerell diverticulum	From the aortic arch	Mural thrombus uncommon	Fusiform
Saccular aneurysm of the aortic arch	May be close to but not involving the origin of the aberrant right subclavian artery	Atherosclerotic plaque always present	Saccular
Ductus diverticulum	Inferior anteromedial aspect of the aorta at site of the ligamentum arteriosum	Mural thrombus usually not present, although atherosclerotic plaque may be seen	Smooth focal bulge with obtuse angles with the aortic wall

prédisposition. Rupture of a Kommerell aneurysm occurs in about 19% of affected cases, and is almost universally fatal [3,30]. Hence, it has been suggested that Kommerell diverticula should be removed in children before these undergo aneurysmal transformation. Due to the higher rate of complications in large aneurysms, some authors also advocate operating on those aneurysms larger than 3 cm in size [10].

2.3. Mimics of Kommerell aneurysm

The main task of the radiologist when presented with a case of Kommerell aneurysm is to identify the condition and its potential complications accurately, as well as to exclude other similar aortic conditions which may mimic it. These mimics include the uncomplicated diverticulum of Kommerell, saccular aneurysm of the thoracic aorta, and ductus diverticulum.

Differentiating between these diagnoses is not always an easy process. It requires careful examination of the anatomy of the anomalous vessel and morphology of the outpouching, as well as the relationship of the neck of the aneurysm to the aberrant subclavian artery as summarized in Table 1. Multiplanar reformation and three-dimensional (3D) volume-rendered projections may prove invaluable in these circumstances [31–33].

A useful distinguishing feature between Kommerell aneurysms, Kommerell diverticula and ductus diverticula, is that Kommerell aneurysms have the typical imaging features of atherosclerotic aneurysms, including calcified plaque and mural thrombus, whereas plaque and thrombus is not commonly seen in uncomplicated Kommerell diverticula and ductus diverticula [4,34]. We present four cases below which illustrate the imaging features of Kommerell aneurysm and its imaging mimics.




Fig. 1. 78-year-old male with Kommerell aneurysm. Invasive catheter angiogram revealed mediastinal widening during cardiac catheterization. Note the tracheal deviation to the right (arrowheads) due to mass effect from the aneurysm.

3. Cases

3.1. Case 1: Kommerell aneurysm

A 78-year-old male ex-smoker presented with chest pain and ST-elevation on electrocardiogram in the anterior and lateral leads to the emergency department. He was reviewed by the cardiology team and diagnosed as having an acute myocardial infarction. Mediastinal widening was noted during cardiac catheterization (Fig. 1) and a gated CT aortogram of the chest was performed using a 320-slice single-source scanner (Aquilion ONE; Toshiba Medical Systems Corporation, Tochigi, Japan) with an injection of 100 ml of a low-osmolar contrast agent, iohexol (Omnipaque 350, GE Healthcare Inc.) at 4 ml/s followed by 50 ml saline bolus.

The study showed a small, hyperdense (38 Hounsfield units) pericardial effusion suggestive of hemopericardium, with a saccular aneurysm measuring 4.1 cm in diameter and 1.7 cm in craniocaudal dimension arising from a Kommerell diverticulum of an anomalous right subclavian artery in a left-sided aortic arch (Fig. 2). The ascending aorta and descending thoracic aorta were normal in caliber. A subsequent chest radiograph was obtained which demonstrated mediastinal widening with bilateral rounded and smooth paratracheal soft-tissue densities (Fig. 2a). The patient was deemed unfit for surgery and died three days later from an unrelated cause. Post-mortem examination confirmed an unruptured aneurysm arising from a diverticulum of Kommerell and left ventricular rupture which was secondary to myocardial infarction.

Radiologic appearance – On frontal chest radiography, the anomalous right subclavian artery has been classically described as an oblique density contiguous with the medial margin of the aortic knuckle, ending as a rounded right paratracheal density [4,35]. On the lateral radiograph, it may appear as a mass in the anteroinferior aspect of Raider's triangle [4,13]. A diverticulum or aneurysm of Kommerell produces a similar appearance, except that the density will appear wider and may cause deviation of the trachea [35,36]. Catheter angiogram may show widening of the mediastinum, and if the aneurysm is large enough, tracheal deviation due to mass effect. On contrast-enhanced MDCT of the chest, Kommerell aneurysms appear as saccular outpouchings from a Kommerell diverticulum if they are atherosclerotic in origin, or they may appear as grossly expanded fusiform Kommerell diverticula [4]. Fusiform aneurysms of Kommerell diverticula are associated with poststenotic dilatation of the subclavian artery arising from a punctate opening in the diverticulum [4].

Teaching points – It may be difficult to distinguish a saccular Kommerell aneurysm from a dilated fusiform Kommerell diverticulum; by convention, a dilatation greater than 1.5 times the expected normal diameter of the artery would qualify as an aneurysm [37]. Atherosclerotic plaque and mural thrombi are common in the saccular forms of Kommerell aneurysm [34,38], as these are essentially atherosclerotic in origin [34].

3.2. Case 2: Kommerell diverticulum

A 68-year-old male presented with an incidental lung nodule




Fig. 2. 78-year-old male with Kommerell aneurysm. (a) Frontal chest radiograph in the same patient demonstrating mediastinal widening with a smooth right paratracheal density (arrow) representing the aberrant right subclavian artery and tracheal deviation (arrowheads) caused by the Kommerell aneurysm. (b) axial, (c) coronal and (d) sagittal oblique post-IV contrast-enhanced MDCT demonstrates a saccular aneurysm (arrow) containing mural thrombus (dashed arrow), arising from the Kommerell diverticulum of an anomalous right subclavian artery (arrowheads) in a left-sided aortic arch. Hemopericardium was present, which was due to left ventricular rupture.

detected on chest radiograph. CT thorax was performed using a 320-slice single-source scanner (Aquilion ONE; Toshiba Medical Systems Corporation, Tochigi, Japan) with an injection of 50 ml of a low-osmolar contrast agent, iohexol (Omnipaque 350, GE Healthcare Inc.) at 1.5 ml/s. The study showed a right-sided aortic arch with an aberrant left subclavian artery. There was fusiform dilatation of the origin of the aberrant left subclavian artery from the aorta, measuring 2.3 cm, in keeping with a large Kommerell diverticulum (Fig. 3).

Radiologic appearance – on MDCT images Kommerell diverticulum is a smooth, fusiform diverticular outpouching from the aortic origin of an aberrant subclavian artery. The diameter of a Kommerell diverticulum should not be more than 50% larger than the diameter of the normal subclavian artery, otherwise by naming convention it would be defined as an aneurysm [37].

Teaching points – Mural thrombus is not usually seen even in large diverticula, presumably due to the hemodynamics of blood flow through an uncomplicated diverticulum which retains a laminar pattern [39]. However, if the subclavian artery arises from a punctate origin in the diverticulum, there is often post-stenotic dilatation with concurrent

mural thrombus formation [4]. Surgical management is usually not considered until the diverticulum becomes aneurysmal [40].

3.3. Case 3: saccular aneurysm of the proximal descending thoracic aorta

A 69-year-old male presented with a left-sided thunderclap bursting headache. A CT carotid angiogram was performed to exclude carotid dissection using a 320-slice single-source scanner (Aquilion ONE; Toshiba Medical Systems Corporation, Tochigi, Japan) with an injection of 70 ml of a low-osmolar contrast agent, iohexol (Omnipaque 350, GE Healthcare Inc.) at 4.0 ml/s. The study showed a 2.9 × 0.9 cm saccular outpouching pointing left antero-laterally from the proximal descending thoracic aorta just after the takeoff of the left subclavian artery, in keeping with a saccular aneurysm (Fig. 4). This patient did not have an aberrant right subclavian artery.

Radiologic appearance – On MDCT, saccular aneurysms of the thoracic aorta appear as outpouchings from the aortic wall. It is important to note that saccular aneurysms in the thoracic aorta may be related to underlying aortitis either due to vasculitis or infection (e.g.




Fig. 3. 68-year-old male with a large Kommerell diverticulum. (a) Axial and (b) coronal post-IV contrast-enhanced MDCT showed a right-sided aortic arch with an aberrant left subclavian artery. There was fusiform dilatation of the origin of the aberrant left subclavian artery (arrows), in keeping with a Kommerell diverticulum. Note the absence of mural thrombus within the outpouching.

mycotic or syphilitic aneurysms), as these are more prone to rupture and carry a high mortality [16,41–44]. In these cases, the imaging features may include additional findings of periaortic soft tissue and an enhancing rim [45].

Teaching points – Thoracic saccular aneurysms tend to demonstrate the typical imaging features of atherosclerotic aneurysms, containing mural thrombus and calcified plaque [46,47]. Those that arise in the proximal descending thoracic aorta just after the subclavian origin may mimic Kommerell aneurysms. Meticulous attention should be paid to the exact location of the neck of the aneurysm, which may be close to, but does not involve the origin of the aberrant right subclavian artery.

3.4. Case 4: ductus diverticulum in a right-sided aortic arch

A 74-year-old female presented with dyspepsia. A CT of the thorax was performed using a 320-slice single-source scanner (Aquilion ONE; Toshiba Medical Systems Corporation, Tochigi, Japan) with an injection of 50 ml of a low-osmolar contrast agent, iohexol (Omnipaque 350,




Fig. 4. 69-year-old male with a saccular thoracic aneurysm. (a) Axial-oblique, (b) coronal-oblique reformatted post-IV contrast enhanced MDCT and (c) three-dimensional (3D) volume-rendered projection shows a saccular outpouching (arrows) pointing left anterolaterally from the proximal descending thoracic aorta just after the origin of the left subclavian artery, in keeping with a saccular aneurysm.




Fig. 5. 74-year-old female with right-sided aortic arch and ductus diverticulum. (a) Axial and (b) coronal post-IV contrast-enhanced MDCT demonstrates a smooth focal bulge (arrows) in the aortic wall at the level of the aortic isthmus, just distal to the right subclavian artery, in keeping with a ductus diverticulum.

GE Healthcare Inc.) at 1.5 ml/s. The study showed a right-sided aortic arch with mirror-image branching. Incidentally there was also a smooth focal bulge of the aortic wall at the level of the isthmus of the aortic arch just distal to the right subclavian artery, forming obtuse angles with the aortic wall, in keeping with a ductus diverticulum (Fig. 5).

Radiologic appearance— On MDCT, ductus diverticulum is a smooth focal bulge which forms obtuse angles with the wall of the aorta [48]. Typically it is found at the inferior anteromedial aspect of the aortic isthmus at the ligamentum arteriosum attachment site [49].

Teaching points—A ductus diverticulum is a developmental out-pouching of the thoracic aorta and represents the vestigial infundibular section of the ductus arteriosus [50]. It is vital to distinguish this from an aneurysm of the aortic arch or traumatic aortic injury to avoid unnecessary surgery [51–53].

4. Summary of imaging features for Kommerell aneurysm and mimics

The discriminating features between Kommerell aneurysm, uncomplicated diverticulum of Kommerell, saccular aneurysm of the thoracic aorta, and ductus diverticulum are summarized in Table 1 and Fig. 6.

5. Conclusion

Saccular Kommerell aneurysm may present as an out-pouching from a diverticulum of Kommerell and is a potential pitfall on MDCT imaging, mimicking other conditions such as saccular aneurysm of the thoracic aorta and dilated Kommerell Diverticula. Accurate diagnosis plays a vital role in successful management of this potentially fatal condition. This review examines the pertinent MDCT findings for Kommerell aneurysm, Kommerell diverticulum, ductus diverticulum and saccular aneurysm of the thoracic aorta.

Conflicts of interest

None.

Funding

The authors state that this work has not received any funding.

IRB statement

IRB approval is not required.




Fig. 6. Schematic diagram of Kommerell aneurysm and mimics. The outpouching (arrows) may involve the aberrant right subclavian artery (ARSCA) or arise separately from the descending thoracic aorta. The shape of the outpouching is important; it may be either fusiform or saccular. Multiplanar reconstructions and three-dimensional (3D) volume-rendered reformations are crucial in making an accurate diagnosis (Diagram by Charlene Liew).

References

- [1] B. Kommerell, Verlagerung des Ösophagus durch eine abnorm verlaufende Arteria subclavia dextra (Arteria lusoria), *Fortschr Geb Roentgenstrahlen* 54 (1936) 590–595.
- [2] N.T. Kouchoukos, P. Masetti, Aberrant subclavian artery and Kommerell aneurysm: Surgical treatment with a standard approach, *J. Thorac. Cardiovasc. Surg.* 133 (2007) 888–892.
- [3] E.H. Austin, W.G. Wolfe, Aneurysm of aberrant subclavian artery with a review of the literature, *J. Vasc. Surg.* 2 (1985) 571–577.
- [4] B. Felson, Ruptured anomalous right subclavian artery: aneurysm or diverticulum, *Semin. Roentgenol.* 24 (1989) 121–126.
- [5] J.M. Miller, K.S. Miller, A note on the historical aspects of dysphagia lusoria, *Am. Surg.* 58 (1992) 502–503.
- [6] D.L. Brown, W.C. Chapman, W.H. Edwards, W.H. Coltharp, W.S. Stoney, Dysphagia lusoria: aberrant right subclavian artery with a Kommerell's diverticulum, *Am. Surg.* 59 (1993) 582–586.
- [7] M.H. Yang, Z.C. Weng, Y.G. Weng, H.H. Chang, A right-sided aortic arch with Kommerell's diverticulum of the aberrant left subclavian artery presenting with syncope, *J. Chinese Med Assoc.* 72 (2009) 275–277.
- [8] W.E. Berdon, Rings, slings, and other things: vascular compression of the infant trachea updated from the midcentury to the millennium—the legacy of Robert E. Gross, MD, and Edward B.D. Neuhauser, MD, *Radiology* 216 (2000) 624–632.
- [9] L.F. Donnelly, R.J. Fleck, P. Pacharn, M.A. Ziegler, B.L. Fricke, R.T. Cotton, Aberrant subclavian arteries: cross-sectional imaging findings in infants and children referred for evaluation of extrinsic airway compression, *Am. J. Roentgenol.* 178 (2002) 1269–1274.
- [10] C.S. Cinà, H. Althani, J. Pasenau, L. Abouzahr, Kommerell's diverticulum and right-sided aortic arch: a cohort study and review of the literature, *J. Vasc. Surg.* 39 (2004) 131–139.
- [11] M. Hara, M. Satake, M. Itoh, et al., Radiographic findings of aberrant right subclavian artery initially depicted on CT, *Radiat. Med.* 21 (2003) 161–165.
- [12] T. Franquet, J.J. Erasmus, A. Giménez, S. Rossi, R. Prats, The retrotracheal space: normal anatomic and pathologic appearances, *RadioGraphics* 22 (2002) S231–S246.
- [13] L. Raider, The retrotracheal triangle, *Chest* 63 (1973) 835–838.
- [14] T. Ota, K. Okada, S. Takanashi, S. Yamamoto, Y. Okita, Surgical treatment for Kommerell's diverticulum, *J. Thorac. Cardiovasc. Surg.* 131 (2006) 574–578.
- [15] R. Kopp, I. Wizgall, E. Kreuzer, et al., Surgical and endovascular treatment of symptomatic aberrant right subclavian artery (arteria lusoria), *Vascular* 15 (2007) 84–91.
- [16] T. Attmann, M. Brandt, S. Müller-Hülsbeck, J. Cremer, Two-stage surgical and endovascular treatment of an aneurysmal aberrant right subclavian (Lusoria) artery, *Eur. J. Cardiothorac. Surg.* 27 (2005) 1125–1127.
- [17] V. Lacroix, P. Astarci, D. Philippe, et al., Endovascular treatment of an aneurysmal aberrant right subclavian artery, *J. Endovasc. Ther.* 10 (2003) 190–194.
- [18] J.S. Corral, C.G. Zúñiga, J.B. Sánchez, J.O. Guaita, A.M. Basail, C.C. Gimeno, Treatment of aberrant right subclavian artery aneurysm with endovascular exclusion and adjunctive surgical bypass, *J. Vasc. Interv. Radiol.* 14 (2003) 789–792.
- [19] M. Davidian, S.T. Kee, N. Kato, et al., Aneurysm of an aberrant right subclavian artery: treatment with PTFE covered stentgraft, *J. Vasc. Surg.* 28 (1998) 335–339.
- [20] H. Shennib, E.B. Diethrich, Novel approaches for the treatment of the aberrant right subclavian artery and its aneurysms, *J. Vasc. Surg.* 47 (2008) 1066–1070.
- [21] R. Greenberg, T. Resch, U. Nyman, et al., Endovascular repair of descending thoracic aortic aneurysms: an early experience with intermediate-term follow-up, *J. Vasc. Surg.* 31 (2000) 147–156.
- [22] C.T. Shen, M.K. Razavi, C.P. Semba, M.D. Dake, Use of endoluminal stent-grafts in acute aortic rupture, *Emerg. Radiol.* 6 (1999) 197–203.
- [23] M.S. Makaroun, E.D. Dillavou, S.T. Kee, et al., Endovascular treatment of thoracic aortic aneurysms: results of the phase II multicenter trial of the GORE TAG thoracic endoprosthesis, *J. Vasc. Surg.* 41 (2005) 1–9.
- [24] L.G. Svensson, N.T. Kouchoukos, D.C. Miller, et al., Expert consensus document on the treatment of descending thoracic aortic disease using endovascular stent-grafts, *Ann. Thorac. Surg.* 85 (2008) S1–S41.
- [25] R. Greenberg, W. Risher, Clinical decision making and operative approaches to thoracic aortic aneurysms, *Surg. Clin. North Am.* 78 (1998) 805–826.
- [26] J.A. Elefteriades, Natural history of thoracic aortic aneurysms: indications for surgery, and surgical versus nonsurgical risks, *Ann. Thorac. Surg.* 74 (2002) S1877–S1880.
- [27] R.K. Greenberg, Q. Lu, E.E. Roselli, et al., Contemporary Analysis of Descending Thoracic and Thoracoabdominal Aneurysm Repair: a comparison of endovascular and open technique, *Circulation* 118 (2008) 808–817.
- [28] M. Panagiotou, V. Filias, C. Prokakis, E. Koletsis, Asymptomatic pseudo-aneurysm of the aortic arch in a patient with aberrant right subclavian artery. A complication of Kommerell's diverticulum? *Interact Cardiovasc. Thorac. Surg.* 7 (2008) 730–731.
- [29] D. Luciano, J. Mitchell, A. Fraise, H. Lepidi, B. Kreitmann, C. Ovaert, Kommerell diverticulum should be removed in children with vascular ring and aberrant left subclavian artery, *Ann. Thorac. Surg.* 100 (2015) 2293–2297.
- [30] G.C. Knight, J.E. Codd, Anomalous right subclavian artery aneurysms report of 3 cases with a review of the literature, *Texas Hear Inst. J.* 18 (1991) 209–218.
- [31] T. Schertler, S. Wildermuth, N. Teodorovic, D. Mayer, B. Marincek, T. Boehm, Visualization of congenital thoracic vascular anomalies using multi-detector row computed tomography and two- and three-dimensional post-processing, *Eur. J. Radiol.* 61 (2007) 97–119.
- [32] G.C. Jakanani, W. Adair, Frequency of variations in aortic arch anatomy depicted on multidetector CT, *Clin. Radiol.* 65 (2010) 481–487.
- [33] F. Alper, M. Akgun, M. Kantarci, et al., Demonstration of vascular abnormalities compressing esophagus by MDCT: Special focus on dysphagia lusoria, *Eur. J. Radiol.* 59 (2006) 82–87.
- [34] T.J. Welch, M.A. McKusick, Cardiovascular case of the day. aneurysm of aberrant right subclavian artery arising from diverticulum of Kommerell, *AJR Am. J. Roentgenol.* 160 (1993) 1314–1316.
- [35] A. Proto, N. Cuthbert, Raidert, Aberrant right subclavian artery: further observations, *Am. J. Roentgenol.* 148 (1987) 253–257.
- [36] P.D. Kiernan, J. Dearani, W.D. Byrne, et al., Aneurysm of an aberrant right subclavian artery: case report and review of the literature, *Mayo Clin. Proc.* 68 (1993) 468–474.
- [37] D. Litmanovich, A.A. Bankier, L. Cantin, V. Raptopoulos, P.M. Boiselle, CT and MRI in diseases of the aorta, *Am. J. Roentgenol.* 193 (2009) 928–940.
- [38] C.L. Siegel, R.H. Cohan, M. Korobkin, M.B. Alpern, D.L. Courneya, R.A. Leder, Abdominal aortic aneurysm morphology: CT features in patients with ruptured and nonruptured aneurysms, *AJR Am. J. Roentgenol.* 163 (1994) 1123–1129.
- [39] D. Bluestein, L. Niu, R.T. Schoepfoerster, M.K. Dewanjee, Fluid mechanics of arterial stenosis: relationship to the development of mural thrombus, *Ann. Biomed. Eng.* 25 (1997) 344–356.
- [40] C.L. Backer, H.M. Russell, K.C. Wurlitzer, J.C. Rastatter, C.K. Rigsby, Primary resection of Kommerell diverticulum and left subclavian artery transfer, *Ann. Thorac. Surg.* 94 (2012) 1612–1617.
- [41] S.L. Brown, R.W. Busuttill, J.D. Baker, H.I. Machleder, W.S. Moore, W.F. Barker, Bacteriologic and surgical determinants of survival in patients with mycotic aneurysms, *J. Vasc. Surg.* 1 (1984) 541–547.
- [42] J.J. McNamara, V.M. Pressler, Natural history of arteriosclerotic thoracic aortic aneurysms, *Ann. Thorac. Surg.* 26 (1978) 468–473.
- [43] B.T. Müller, O.R. Wegener, K. Grabitz, M. Pillny, L. Thomas, W. Sandmann, Mycotic aneurysms of the thoracic and abdominal aorta and iliac arteries: experience with anatomic and extra-anatomic repair in 33 cases, *J. Vasc. Surg.* 33 (2001) 106–113.
- [44] S.R. Sheeran, S.J.A. Sclafani, Syphilitic aneurysm of descending thoracic aorta causing vertebral body erosion and spastic paraparesis, *Emerg. Radiol.* 7 (2000) 245–247.
- [45] C.L. Siegel, R.H. Cohan, CT of abdominal aortic aneurysms, *AJR Am. J. Roentgenol.* 163 (1994) 17–29.
- [46] K. Machida, A. Tasaka, CT patterns of mural thrombus in aortic aneurysms, *J. Comput. Assist. Tomogr.* 4 (1980) 840–842.
- [47] K. Takahashi, W. Stanford, Multidetector CT of the thoracic aorta, *Int. J. Cardiovasc. Imaging* 21 (2005) 141–153.
- [48] P.C. Goodman, R.B. Jeffrey, H. Minagi, M.P. Federle, A.N. Thomas, Angiographic evaluation of the ductus diverticulum, *Cardiovasc. Intervent. Radiol.* 5 (1982) 1–4.
- [49] R.G. Fisher, M. Sanchez-Torres, C.J. Whigham, J.W. Thomas, Lumps and bumps; that mimic acute aortic and brachiocephalic vessel injury, *RadioGraphics* 17 (1997) 825–834.
- [50] E. Salomonowitz, J.E. Edwards, D.W. Hunter, et al., The three types of aortic diverticula, *AJR Am. J. Roentgenol.* 142 (1984) 673–679.
- [51] S.E. Mirvis, S.O. Pais, K. Shanmuganathan, Atypical results of thoracic aortography performed to exclude aortic injury, *Emerg. Radiol.* 1 (1994) 24–31.
- [52] S.S. Morse, M.G. Glickman, L.H. Greenwood, et al., Traumatic aortic rupture: false-positive aortographic diagnosis due to atypical ductus diverticulum, *AJR Am. J. Roentgenol.* 150 (1988) 793–796.
- [53] S.E. Mirvis, K. Shanmuganathan, Diagnosis of blunt traumatic aortic injury 2007: Still a nemesis, *Eur. J. Radiol.* 64 (2007) 27–40.