them at increased risk for poor health-related outcomes. Walking is widely supported as a cost-effective, accessible exercise and way to maintain physical fitness and mitigate stress levels. There is a critical gap in addressing health promotion strategies in caregivers. The specific aims of this health-coaching (HC) walking study using wireless pedometers in family caregivers of persons with dementia were to: 1) establish the feasibility of HC and wireless pedometer use, 2) examine outcomes of well-being, stress and activity level and 3) understand the experience of participation. This 2-group comparative study used a repeated measure design and mixed methods approach. We enrolled 27 females and 5 male caregivers (n=32), μ age 57 years: with both 16 in the control in the intervention arm. Prepost measures of general health (body mass index [BMI], blood pressure, heart rate, cognition, well-being, stress and perceived activity level) were obtained from caregivers at baseline and again at 12 weeks. Results indicate that those who received HC had a statistically significant improvement in BMI (p = .01). There were no other statistically significant improvements in outcomes in either group. Qualitative findings suggest that participants reported many stresses that made self-care challenging. In summary, this was a feasible intervention that resulted in improved BMI in the HC group. Work is needed to understand the long-term impact of this outcome. Further exploration of other health-promoting interventions that may be beneficial for this population is essential.

QUALITY OF LIFE OF GRANDPARENTS RAISING GRANDCHILDREN IN NIGERIA

Joshua O. Aransiola,¹ Funmi Togonu-Bickersteth,¹ Kolawole Aliyu,² Mojirayo Afolabi,² and Akanni Akinyemi², 1. Obafemi Awolowo University, Ile-Ife, Ile-Ife, Osun State, Nigeria, 2. Obafemi Awolowo University, Ile-Ife, Osun State, Nigeria

This article examined the personal and household characteristics influencing the quality of life (QoL) of grandparents caring for grandchildren in Skipped Generation Households in Nigeria with a sample of 2, 144 grandparents in Imo, Lagos and Kano. Chi square and multinomial logistic regression were employed to understand the relationship between the dependent variable (QoL) and independent variables (personal and household characteristics). The level of the QoL of the grandparents almost spread evenly among low (34.3%), average (34.3%) and high (31.4%). Five domains of QoL were examined including level of independence (LI), psychological well-being (PW), social relation (SR), physical health (PH), environment (ENV) and engagement in income generating activities (IGA). Personal characteristics including; state of residence was significantly associated with all the domains except LI and PW, age was associated with IGA, LI and ENV, sex was associated with SR and ENV and level of education was associated with all the domains except IGA and ENV while religious affiliation was associated with IGA. Household characteristics including; sex of household head was significantly associated with PH, SR and ENV, age of household head was associated with IGA and LI and wealth index was associated with all the domains while the number of household members was significantly associated with ENV. The regression analysis shows that only state of residence and wealth index significantly influence the QoL of the grandparents ($P \le 0.05$). The state of residence and wealth index are therefore important in any policy intervention for this category of elderly persons in Nigeria.

SUPPORTING FAMILY CAREGIVERS: HOW DOES RELATIONSHIP STRAIN OCCUR IN CAREGIVING DYADS? A QUALITATIVE STUDY

Elizabeth S. Avent,¹ Laura Rath,² Kylie Meyer,³ Donna Benton,¹ and Paul Nash⁴, 1. University of Southern California, Los Angeles, California, United States, 2. Archstone Foundation, Long Beach, California, United States, 3. University of Texas Health Science Center San Antonio, San Antonio, Texas, United States, 4. University of Southern California, LA, California, United States

Family members and spouses are usually the primary caregivers for older adults. Providing direct care can be stressful, strenuous, and time-consuming for caregivers, potentially leading to frustration and anger towards care recipients. This can be detrimental to the relationship quality of the caregiving dyad. Though caregiver strain and burden have been extensively studied, there is limited information on the development of relationship strain. To explore how relationship strain occurs between caregivers and care recipients, 8 focus groups (N=62) and 8 semi-structured telephone interviews were conducted with caregivers in Los Angeles, inquiring about relationship quality with their care recipients and when frustration and anger occurs. Inductive coding was used to create coding schemas. Findings showed that most caregivers reported relationship strain occurring after taking on the caregiving role, and frustration and anger arose when providing ADLs, especially during bathing and toileting. Although these caregivers had initially experienced strain in their relationships, a recurring theme that emerged was that they developed strategies to decrease frustration and anger and improve the quality of their relationships with their care recipients. Direct communication with caregivers is important in designing a structured and effective intervention. These findings help inform an intervention for new caregivers to help them identify what can lead to relationship strain, as well as teach them reliable strategies to manage frustration and anger towards their care recipients.

SESSION 1360 (POSTER)

FRAILTY, FALLS AND FALL PREVENTION

THE EFFECT OF YOGA ON FALL PREVENTION OUTCOMES IN ADULTS 50 YEARS AND OLDER: A SYSTEMATIC REVIEW OF THE LITERATURE

Chinelo K. Nsobundu,¹ Margaret J. Foster,² and Yan Hong¹, 1. Texas A&M University - School of Public Health, College Station, Texas, United States, 2. Texas A&M University, College Station, Texas, United States

Falls constitute a multitude of injuries irrespective of age. To combat these challenges, older adults are