

RESEARCH ARTICLE

Open Access

# Evolution at increased error rate leads to the coexistence of multiple adaptive pathways in an RNA virus

Laura Cabanillas<sup>†</sup>, María Arribas<sup>†</sup> and Ester Lázaro<sup>\*</sup>

## Abstract

**Background:** When beneficial mutations present in different genomes spread simultaneously in an asexual population, their fixation can be delayed due to competition among them. This interference among mutations is mainly determined by the rate of beneficial mutations, which in turn depends on the population size, the total error rate, and the degree of adaptation of the population. RNA viruses, with their large population sizes and high error rates, are good candidates to present a great extent of interference. To test this hypothesis, in the current study we have investigated whether competition among beneficial mutations was responsible for the prolonged presence of polymorphisms in the mutant spectrum of an RNA virus, the bacteriophage Q $\beta$ , evolved during a large number of generations in the presence of the mutagenic nucleoside analogue 5-azacytidine.

**Results:** The analysis of the mutant spectra of bacteriophage Q $\beta$  populations evolved at artificially increased error rate shows a large number of polymorphic mutations, some of them with demonstrated selective value. Polymorphisms distributed into several evolutionary lines that can compete among them, making it difficult the emergence of a defined consensus sequence. The presence of accompanying deleterious mutations, the high degree of recurrence of the polymorphic mutations, and the occurrence of epistatic interactions generate a highly complex interference dynamics.

**Conclusions:** Interference among beneficial mutations in bacteriophage Q $\beta$  evolved at increased error rate permits the coexistence of multiple adaptive pathways that can provide selective advantages by different molecular mechanisms. In this way, interference can be seen as a positive factor that allows the exploration of the different local maxima that exist in rugged fitness landscapes.

**Keywords:** RNA viruses, Interference, Beneficial mutations, Mutagenesis, 5-azacytidine, Epistasis, Polymorphisms

## Background

Beneficial mutations are the substrate upon which natural selection acts to drive adaptive evolution. For a beneficial mutation to fix in a population it is necessary that it survives genetic drift and that under the influence of selection displaces the rest of genomes. This means that any beneficial mutation remains as a polymorphism for a certain period of time, which in asexual populations lacking recombination is determined not only by its selective coefficient but also by the probability that other beneficial mutations are present at the same time

in the population [1,2]. When beneficial mutations are scarce, they can be fixed before a secondary mutation arises, and adaptation takes place through the sequential fixation of mutations [3]. In contrast to this, when beneficial mutations are frequent, their fixation usually does not take place before one or more secondary beneficial mutations are generated. Genomes carrying the secondary mutations spread simultaneously with the genomes containing the primary one, delaying its fixation, and occasionally causing its elimination [4-10]. This competition among beneficial mutations is known as dynamics of interference. Whether a secondary mutation will displace a previously spreading one depends, among other factors, on the frequency reached by the first mutation when the second one arises, and on their relative

\* Correspondence: lazaro@cab.inta-csic.es

<sup>†</sup>Equal contributors

Centro de Astrobiología (CSIC-INTA) Ctra de Ajalvir Km 4, Torrejón de Ardoz, Madrid 28850, Spain

selective values, which in the last term are determined by the distribution of beneficial mutation effects.

The frequency of beneficial mutations, and thus the intensity of interference, depends on several factors, among which some of the most relevant are the population size, the error rate, and the degree of adaptation of the population [11,12]. In large populations a high number of genomes replicate simultaneously, increasing the probability of generating beneficial mutations that can compete for fixation. In addition, the time necessary to fix any substitution is longer in large populations than in smaller ones, which also increases the opportunities for interference [13,14]. The increase of the error rate is associated with higher numbers of both deleterious and beneficial mutations per genome. In this situation, the fixation of beneficial mutations can be delayed not only by the coexistence of several lines carrying different mutations but also because at high error rate beneficial mutations arise in deleterious backgrounds, thus hampering the action of selection [14-16].

The dynamics of interference has been the focus of a number of theoretical and experimental studies. Theoretical studies establish different assumptions concerning the fitness effects of beneficial mutations and the genetic backgrounds where they can arise. The clonal interference model [13,14] assumes that most beneficial mutations have different selective values and always appear in the previously fixed background. The assumption that secondary mutations never appear in the genomes bearing a mutation that is in the process to fixation means that in this model competition takes place among genomes differing in a single beneficial mutation. By contrast, the multiple mutations model [17-19] assumes that beneficial mutations have a single characteristic selective value and can appear in any genetic background, including those containing beneficial mutations generated previously and still not fixed. In this way, competition can also occur among genomes differing in the number of beneficial mutations. Finally, the full interference model [20,21] tries to eliminate the limitations above by considering that beneficial mutations of different selective values can arise in any genetic background.

Each of the theoretical approaches predicts a particular dynamics of adaptation as a function of the population size and the error rate, and experimental studies have been designed to verify whether the behaviour of real populations of bacteria [11,22,23], yeast [17], or viruses [24] fit the theoretical predictions. Interference has also been studied at the genetic level by tracing specific mutations with particular labels [20,25], and by sequencing ensembles of individual genomes isolated from adapting populations [23,26-30]. This latter scenario allows analyzing the variation in the frequency and distribution of particular beneficial mutations along the

adaptation process, which can shed much light about how interference affects the fate of beneficial mutations.

In this work we analyze the results of a previous evolution experiment carried out with a population of an RNA virus, the bacteriophage Q $\beta$ , evolved for a large number of generations in the presence of the mutagenic nucleoside analogue 5-azacytidine (AZC) [31] which increases the virus error rate [32]. The initial objective of the experiment was to characterize the possible mechanisms providing resistance to AZC in bacteriophage Q $\beta$ . It is important to note that during adaptation to mutagens, the same agent that acts as selective pressure also can interfere with the fixation of adaptive mutations, due to the increased number of errors generated in its presence. The analysis of the populations obtained at different points of the evolutionary process showed the fixation of two substitutions, A2187C that has a general beneficial effect, and A1746U with a selective advantage in the presence of AZC and a fitness cost in its absence [31]. In addition to these, six other mutations were detected as polymorphisms that, despite the demonstrated selective value of at least two of them, did not reach fixation after a large number of transfers in the presence of AZC. Since the conditions under which bacteriophage Q $\beta$  was propagated in this experiment (high mutation rate and large population sizes) favour the coexistence of multiple beneficial mutations, we thought that interference among beneficial mutations could be one of the reasons underlying the prolonged presence of polymorphisms. To further study whether this process operates during the evolution of bacteriophage Q $\beta$  at increased error rate, we have analyzed how polymorphic mutations are distributed in individual virus genomes isolated at different points of the evolutionary series. This approach has allowed us to identify several competing lines carrying different combinations of polymorphic mutations which differ in their fitness values and in their ability to fix when present in a simpler mutant spectrum. The presence of additional mutations accompanying the polymorphic ones, the high frequency of recurrent mutations, and the occurrence of epistatic interactions contribute to generate a highly complex dynamics that would likely require improved theoretical models to be successfully described.

## Methods

### Viruses and bacteria. Standard procedures for infection

Bacteriophage Q $\beta$  was routinely propagated by infecting log-phase cultures of *Escherichia coli*, strain Hfr (Hayes) in NB medium (8 g/l Nutrient Broth from Merck and 5 g/l NaCl). The virus was adapted to replication in liquid culture medium in our laboratory as described [32].

Infections in liquid medium were always carried out using fresh exponential phase *E. coli* cultures (with an

optical density at 550 nm between 0.6 and 0.8) that were infected with the virus at the multiplicity of infection (moi) indicated in each experiment. After 2 h of incubation at 37°C with good aeration, cultures were treated with 1/20 vol of chloroform for 15 min at 37°C with shaking (300 rpm). Virus supernatants were harvested upon centrifugation at 13000 × g for 10 min and maintained at 4°C for short-term use (less than 15 days) or at -80°C for long-term storage. Virus titres were determined by plaque assay and expressed as the number of plaque forming units (pfu) per ml of the phage suspension.

Virus populations were used to obtain biological clones that correspond to lytic plaques obtained in semisolid agar. Virus clones were isolated by punching and removing the top and the bottom agar around well-separated lytic plaques. The agar containing the lytic plaque was transferred into an eppendorf tube with 1 ml of phage buffer (1 g/l gelatine, 0.05 M Tris-HCl, pH 7.5, and 0.01 M MgCl<sub>2</sub>) and 50 µl of chloroform, and incubated for 1 h at 28°C with shaking (300 rpm). After centrifugation at 13000 × g for 15 min to clarify the supernatant, the latter was stored over 25 µl of chloroform.

#### Serial transfers of bacteriophage Qβ

**Prior transfers:** A population of bacteriophage Qβ, previously adapted to replicate in our laboratory (population Qβ<sub>0</sub>), was used to infect two parallel cultures of *E. coli* in exponential phase at an initial moi = 1 pfu/cell in a volume of 10 ml either in the absence of AZC (population Qβ-control) or in the presence of a gradually increased AZC concentration (population Qβ-AZC) (Figure 1a). After 2 h of incubation at 37°C with good aeration, the virus supernatants were collected as described above, and 1 ml of each phage suspension was used to infect a fresh *E. coli* culture. Virus titres were determined each 10 transfers, which allowed us to estimate the number of viruses used to initiate each subsequent transfer. This procedure was repeated for a total of 70 transfers in both the control population and in the AZC-exposed population. Virus populations were isolated throughout the transfer series and the number of transfers experienced by each of them was indicated in brackets beside the name of the population (Figure 1a) [31].

**New transfers:** The population Qβ-AZC(t70) was subjected to 20 additional transfers, the first 10 in the presence of 80 µg/ml of AZC and the last 10 in the presence of 100 µg/ml of AZC (Figure 1b). Transfers were carried out as described above. The number of viruses used to initiate each subsequent transfer was always above 10<sup>7</sup> pfu.

#### RNA extraction, cDNA synthesis, PCR amplification and nucleotide sequencing


Virus RNA was prepared following standard procedures [31,32] from both complex populations, to determine

the consensus sequence (from nucleotide 180 to nucleotide 4180), and biological clones, to determine individual virus sequences (from nucleotide 1485 to nucleotide 4028). Sequences were deposited in NCBI GenBank with accession numbers KC137648- KC137682.

RNAs were amplified by RT-PCR using Avian Myeloblastosis Virus RT (Promega) and Expand High Fidelity DNA polymerase (Roche). The cDNAs were purified with a Qiagen purification kit and subjected to cycle sequencing with Big Dye Chemistry (Applied Biosystems; Perkin Elmer). The following pairs of oligonucleotide primers were used for RT-PCR: P1 forward (5'CGAATCTTCCGACACGCATCC3') with P1 reverse (5'AAACG GTAACACGCTTCTCCAG3') to amplify from nucleotide position 150 to 1497; P2 forward (5'CTCAATCCGCGTGGGGTAAATCC3') with P2 reverse (5'CA GAAAATCGGCAGTGACGCAACA3') to amplify from nucleotide position 1407 to 2817; P3 forward (5'GTGC CATACCGTTTGACT3') with P3 reverse (5'TCGTG CCCTGGAAGACC3') to amplify from nucleotide position 2254 to 4095; and P4 forward (5'GCGGCAAG CACTACTATTCT3') with P4 reverse (5'GATCCCC CTCTCACTCGT3') to amplify from nucleotide position 3541 to 4195. Sequences were aligned with the consensus sequence of the wild type phage with Clustal W. Mutations relative to the consensus sequence were identified using the program BioEdit. Nucleotides were numbered according to the sequence of a cDNA of bacteriophage Qβ cloned in the plasmid pBR322 [33].

#### Preparation of virus clones through directed mutagenesis of an expression vector of bacteriophage Qβ

The plasmid pBRT7Qβ, which contains the cDNA of the bacteriophage Qβ cloned in the plasmid pBR322 [33] was used to express the wild type virus (Qβ<sub>wt</sub>) and also mutant viruses containing specific substitutions. Mutagenesis was carried out using a QuickChange II Site-Directed Mutagenesis Kit from Stratagene. Primers 5'CTTAGAC TCGTCTGAGGTGACTGTTTACGGAGACGA3', 5'CC TCTTAGGGGTCCATCGAGTTGCGATTCTGCGG3', 5'CCATCGATCAGCTTATCTGCAGGAGTAATCCTA CGAAG3', and their complementary were used to introduce the substitutions C3413G, G3945A, and U3989C respectively. The primers used to build the mutants containing A1746U, A2982G, and U3582C had been described previously [31]. The procedures to isolate the site-directed mutant viruses were as detailed in [31]. Each of the experiments carried out with viruses obtained upon expression of the mutagenized infectious clone was initiated with a single lytic plaque whose sequence had been previously analyzed to check the presence of the desired mutation.



### Determination of relative fitness values of bacteriophage Q $\beta$ mutant viruses

Growth rate values were used as a surrogate of fitness. To determine them, liquid cultures containing  $10^8$  bacteria growing in exponential phase were inoculated with  $10^4$  pfu of the site-directed mutant indicated in a final volume of 1 ml. After two hours at  $37^\circ\text{C}$  with good aeration, the virus supernatants were collected as described above and titrated to estimate the virus yield. Preliminary assays showed that the virus used as reference (the wild type virus, Q $\beta_{\text{wt}}$ , obtained upon expression of the infectious clone of bacteriophage Q $\beta$  described in the previous section) grew exponentially during this time interval. Growth rate determinations for each virus were carried out in triplicate in an assay which always

included the reference virus. Absolute fitness for a given mutant was calculated as the change in  $\log_2$  of the virus titre, and relative fitness was defined as the ratio between the absolute fitness of the mutant assayed and the absolute fitness of the reference virus.

### Competition between bacteriophage Q $\beta$ virus clones

Competitions between bacteriophage Q $\beta$  virus clones (Q $\beta_{\text{wt}}$  and the site-directed mutants indicated in each experiment) were carried out by mixing equal amounts of each competitor virus ( $10^6$  or  $10^7$  pfu from each of them) which were used to infect  $10^9$  bacteria in a final volume of 10 ml. Each new transfer was initiated with 1 ml of a dilution of the previous virus supernatant to maintain a moi about 0.01. All the competitions were



carried out in duplicate. Evolution took place either in the presence (60 µg/ml) or in the absence of AZC. The population obtained after the number of transfers indicated was sequenced to determine whether one of the viruses had become dominant.

## Results

### Previous results

In our previous work [31] a population of bacteriophage Q $\beta$  was evolved during 70 transfers under two parallel transmission regimes that differed in the presence of the mutagenic nucleoside analogue AZC. In this way we obtained the population Q $\beta$ -control(t70) evolved at the standard error rate of the virus, and the population Q $\beta$ -AZC(t70) evolved at artificially increased error rate (Figure 1a). These populations had in common the fixation of substitution A2187C, and differed in the presence of substitutions A1746U, which was only fixed in population Q $\beta$ -AZC(t70), and substitution G1773A, which was exclusive of population Q $\beta$ -control(t70) (Figure 1a). Population Q $\beta$ -AZC(t70) showed a statistically significant higher mutation frequency in the mutant spectrum than population Q $\beta$ -control(t70) ( $3 \times 10^{-3}$  versus  $2.5 \times 10^{-4}$  substitutions per nucleotide). In addition, 6 polymorphic mutations (U1520C in the coat protein gene, and A2982G, C3413G+U, U3582C, G3945A, and U3989C in the replicase gene) were detected in population Q $\beta$ -AZC(t70). In contrast to this result, population Q $\beta$ -control(t70) showed no polymorphisms in its consensus sequence, and only one substitution was represented at a frequency of 0.2 when the mutant spectrum was analyzed through sequencing of 10 biological clones.

Competition experiments between the wild type virus and two single-directed mutants containing the replicase polymorphic substitutions A2982G and U3582C showed that both could fix in only five transfers in the presence of AZC, which clearly demonstrates that they have a selective advantage under this condition, and can fix when the complexity of the mutant spectrum is reduced. These results motivated us to investigate how the complexity of the mutant spectra generated at increased error rate interferes with the fixation of beneficial mutations.

### Extended evolution of population Q $\beta$ -AZC(t70) in the presence of AZC leads to a further increase in the number of polymorphisms

Population Q $\beta$ -AZC(t70) described in the previous section was subjected to 20 additional transfers, the first 10 in the presence of 80 µg/ml of AZC and the last 10 in the presence of 100 µg/ml of AZC (Figure 1b). In this way we obtained the population Q $\beta$ -AZC(t90), evolved during 90 transfers in the presence of AZC. The analysis of the consensus sequence of this population revealed

that none of the polymorphic mutations previously identified was fixed at transfer number 90. Moreover, the analysis of 15 biological clones isolated from population Q $\beta$ -AZC(t90) revealed the presence of 9 additional substitutions that were represented at high frequency ( $\geq 0.2$ ) in the mutant spectrum (Table 1).

To study how polymorphic mutations distribute in individual virus genomes along the evolutionary series carried out in the presence of AZC, we also analyzed the mutant spectrum of two previous populations [Q $\beta$ -AZC(t60) and Q $\beta$ -AZC(t70), see Figure 1]. We isolated 10 biological clones from each population and sequenced the genomic region where polymorphic substitutions had been identified in our prior work [31]. We sequenced biological clones instead of molecular clones to ensure the analysis of viable viruses and to identify possible associations among mutations in the same genome. The whole list of polymorphisms, as well as their distribution in the virus genomes analyzed is shown in Table 1. All the genomes listed in Table 1 also carry substitutions A2187C, which was fixed a transfer number 30, and A1746U, which was fixed at transfer number 40, together with a number of additional mutations that were exclusive of each of them (see Additional file 1). All sequences were submitted to NCBI GenBank. Their accession numbers are KC137673-KC137682 (virus clones from population Q $\beta$ -AZC(t60), KC137663-KC137672 (virus clones from population Q $\beta$ -AZC(t70), and KC137648-KC137662 (virus clones from population Q $\beta$ -AZC(t90)).

Another criterion to identify polymorphic mutations is their presence as double bands, consisting of a mixture of the mutated and the wild nucleotides, in the chromatograms of the consensus sequences of the corresponding virus populations. In good agreement with this expectation, all substitutions represented at a frequency  $\geq 0.2$  in the mutant spectrum of a given population appeared also as double peaks in the chromatogram of the consensus sequence of the same population. This approach was also used to identify the transfer number at which polymorphisms could be first detected. To this end we examined the chromatograms corresponding to the consensus sequences of populations Q $\beta$ -AZC(t20), Q $\beta$ -AZC(t30), Q $\beta$ -AZC(t40), and Q $\beta$ -AZC(t50). We observed that some polymorphisms could be detected as double bands in the chromatograms at transfers as early as 30, whereas others could not be detected until transfer number 90 (Table 2). Substitution U3989C which in our previous work [31] was identified as a polymorphism at transfer number 40 together with A3945G, was actually already present as a double band at transfer number 30.

The location of polymorphic substitutions in the genome of bacteriophage Q $\beta$  shows a clear preference for the replicase gene (Table 2). However, we must take into

**Table 1 Distribution of polymorphic mutations in individual viruses isolated from bacteriophage Q $\beta$  populations evolved in the presence of AZC**

Nt <sup>1</sup>	1520	1604	2059	2277	2378	2384	2471	2982	3413	3545	3582	3879	3945	3989	4006
Wt <sup>2</sup>	U	C	C	A	C	C	C	A	C	C	U	C	G	U	A
Population Q $\beta$ -AZC(t60) <sup>3</sup>															
C <sub>1</sub>	C	G	.	.	.	.	.	G	U	.	.	.	A	C	.
C <sub>2</sub>	C	.	.	.	.	.	.	G	.	.	.	.	A	C	.
C <sub>3</sub>	C	.	.	.	.	.	.	G	.	.	.	.	A	C	.
C <sub>4</sub>	C	G	.	.	.	.	G	G	.	.	.	.	A	C	.
C <sub>5</sub>	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
C <sub>6</sub>	.	.	.	.	.	.	.	.	G	.	.	.	.	.	G
C <sub>7</sub>	.	.	.	G	G	.	.	.	.	.	.	.	.	.	.
C <sub>8</sub>	.	.	.	.	.	G	.	.	.	.	.	.	.	.	.
C <sub>9</sub>	.	.	.	.	.	G	G	.	.	.	.	.	.	.	.
C <sub>10</sub>	.	.	.	.	.	.	.	.	.	.	.	.	.	C	.
Population Q $\beta$ -AZC(t70) <sup>3</sup>															
C <sub>1</sub>	C	.	.	.	.	.	.	G	.	.	.	.	A	C	.
C <sub>2</sub>	C	.	G	.	.	.	.	G	.	.	.	.	A	C	.
C <sub>3</sub>	C	.	.	.	.	G	.	G	G	.	.	.	A	C	.
C <sub>4</sub>	.	.	.	.	.	.	.	.	.	.	C	.	.	.	.
C <sub>5</sub>	.	.	.	.	.	.	G	.	.	.	C	.	.	.	.
C <sub>6</sub>	.	.	.	.	.	.	.	.	G	.	C	.	A	C	.
C <sub>7</sub>	.	.	G	.	.	G	.	.	G	.	.	.	.	.	.
C <sub>8</sub>	.	.	.	.	.	.	.	.	G	.	.	.	.	.	.
C <sub>9</sub>	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
C <sub>10</sub>	.	.	.	.	.	.	G	.	.	.	.	G	.	.	.
Population Q $\beta$ -AZC(t90) <sup>3</sup>															
C <sub>1</sub>	C	.	.	.	.	.	.	G	.	.	.	.	A	C	.
C <sub>2</sub>	C	G	.	.	.	.	G	G	.	A	.	.	A	C	G
C <sub>3</sub>	C	.	G	G	G	.	.	G	.	.	.	.	A	C	.
C <sub>4</sub>	C	G	.	.	.	.	.	G	.	A	.	.	A	C	G
C <sub>5</sub>	C	.	G	.	.	U	.	G	.	.	.	.	A	C	.
C <sub>6</sub>	C	.	G	G	.	.	G	G	.	.	.	.	A	C	.
C <sub>7</sub>	C	.	.	G	.	.	.	G	.	A	.	G	A	C	.
C <sub>8</sub>	C	G	G	.	.	.	G	G	G	A	.	.	A	C	G
C <sub>9</sub>	.	.	.	.	G	.	.	.	.	.	C	G	.	.	.
C <sub>10</sub>	.	.	.	.	G	.	.	.	.	.	C	.	.	.	G
C <sub>11</sub>	.	.	G	G	.	U	.	.	G	.	C	.	A	C	.
C <sub>12</sub>	.	.	.	G	.	G	.	.	.	.	.	.	.	.	.
C <sub>13</sub>	.	G	.	.	G	U	.	.	.	.	.	.	.	.	.
C <sub>14</sub>	C	.	.	G	.	G	.	.	.	.	.	.	A	.	.
C <sub>15</sub>	.	.	.	.	.	.	.	.	G	.	.	G	A	C	.

<sup>1</sup>Nucleotide positions where polymorphisms were detected.

<sup>2</sup>Nucleotides present at the positions indicated in the wild type virus.

<sup>3</sup>Nucleotides present at the positions indicated in the virus clones (indicated as C<sub>x</sub>, where x is a number arbitrarily assigned) isolated from that populations. The region sequenced comprises from nucleotide 1485 to nucleotide 4028.

account that the lysis gene and the starting of the coat protein gene were only analyzed at the level of consensus sequences. Although we did not observe any double band in the chromatograms corresponding to these regions, we cannot exclude that some polymorphism could have been detected upon the analysis of the mutant spectrum.

### Polymorphic mutations differ in their effects on fitness and in their ability to fix when present in a simple mutant spectrum

An explanation for the abundance of polymorphic mutations is that they have a selective advantage that permits them to increase their frequency thanks to the action of selection. To check this assumption we determined separately the ratio  $ds/dn$  for the polymorphic and non polymorphic mutations present in each population analyzed (Table 3). When mutations repeated in several genomes of the same population were counted only once, we found that the average ratio  $ds/dn$  was larger than 1 in both cases, rejecting the action of positive selection. However, the average  $ds/dn$  value obtained for the polymorphic mutations was significantly lower ( $p < 0.05$ , Student's  $t$  test for the difference of means) than that estimated for the non polymorphic substitutions. A similar analysis carried out counting the repeated mutations the

number of times that they appear in each population led to different results. In this case the average  $ds/dn$  obtained for the polymorphic substitutions was reduced to a value close to 1, whereas the average for the non polymorphic was kept almost unaltered ( $p > 0.05$ , Student's  $t$  test for the difference of means) (Table 3). These findings suggest that positive selection acts at a higher extent in the substitutions represented as polymorphisms than in the non polymorphic.

There are, however, two particular situations in which deleterious or neutral mutations can also reach high frequencies. The first one is the occurrence of population bottlenecks, a circumstance that reduces the genetic diversity, leading to the fixation of mutations independently of their selective value [2,36-38]. The second one is hitchhiking with beneficial mutations. Since the bacteriophage Q $\beta$  populations analyzed in this work were propagated using large population sizes (above  $10^7$  pfu, see Methods), we can discard the first possibility. We neither found clear evidences of hitchhiking for most of the substitutions analyzed. If a mutation had reached high frequency because of its presence in the same genome where a beneficial mutation is generated, both mutations should appear always linked and should have been first detected at the same transfer number. The only two substitutions that meet these two requirements were U1520C and A2982G (see Tables 1 and 2). Since A2982G is beneficial in the presence of AZC [31], the only mutation that seems to have reached high frequency in our experiment because of hitchhiking is U1520C. However, as we will explain in the discussion, we cannot discard that some neutral or deleterious mutations could have achieved high frequency due to hitchhiking, even in the absence of a perfect association with a beneficial mutation.

To ascertain whether some of the polymorphic substitutions shown in Table 1 have selective advantages we prepared single site-directed mutants containing A1746U and the polymorphic substitutions that had been identified in population Q $\beta$ -AZC(t70) (Q $\beta$ <sub>A1746U</sub>, Q $\beta$ <sub>A2982G</sub>, Q $\beta$ <sub>U3582C</sub>, Q $\beta$ <sub>C3413G</sub>, Q $\beta$ <sub>G3945A</sub>, and Q $\beta$ <sub>U3989C</sub>), and calculated their relative fitness values using the wild type virus, Q $\beta$ <sub>wt</sub>, as reference (see Methods). We observed that, with the only exception of Q $\beta$ <sub>G3945A</sub>, all the mutants assayed had lower fitness than the virus Q $\beta$ <sub>wt</sub> in the absence of AZC (Table 4). In contrast to this, mutant viruses Q $\beta$ <sub>A2982G</sub>, Q $\beta$ <sub>U3582C</sub>, and Q $\beta$ <sub>G3945A</sub> had higher fitness than the virus Q $\beta$ <sub>wt</sub> in the presence of AZC (Table 4). To check whether the mutations contained in the site-directed mutants assayed can fix when present in a simple mutant spectrum, we carried out competition experiments of these viruses with the virus Q $\beta$ <sub>wt</sub>. After a number of transfers either in the presence or in the absence of AZC we determined the consensus sequences of the new populations generated to check the status of the substitutions analyzed (Table 4). In good

**Table 2 Substitutions that remain as polymorphisms at transfer number 90 in the bacteriophage Q $\beta$  evolved in the presence of AZC**

Substitution	Gene <sup>1</sup>	change <sup>2</sup>	First detected <sup>3</sup>
U1520C	Coat	Syn	50
C1604G	Coat	Syn	60
C2059G	Read-through	T/S	60
A2277G	Read-through	K/E	90
C2378G	Replicase	N/K	90
C2384(G+U)	Replicase	Syn	60
C2471G	Replicase	Syn	30
A2982G	Replicase	T/A	50
C3413(U+G)	Replicase	Syn	50
C3545A	Replicase	Syn	70
U3582C	Replicase	Y/H	50
C3879G	Replicase	L/V	70
G3945A	Replicase	G/S	40
U3989C	Replicase	Syn	30
A4006G	Replicase	K/R	60

<sup>1</sup>Location of each substitution in the genome of the bacteriophage Q $\beta$ .

<sup>2</sup>Type of change produced by each nucleotide substitution. In the case of non synonymous changes, the amino acid substitution is indicated.

<sup>3</sup>Transfer number at which each substitution was first detected as a double band in the chromatogram corresponding to the consensus sequence of that population. The genomic region analyzed comprised from nucleotide 180 to 4180.

**Table 3 Ratio ds/dn for the nucleotide substitutions found in the bacteriophage Qβ populations evolved in the presence of AZC**

Population	ds/dn <sup>1</sup>			
	Repeated mutations counted once		Repeated mutations counted all the times they appear	
	Non polymorphic substitutions <sup>2</sup>	Polymorphic substitutions <sup>2</sup>	Non polymorphic Substitutions <sup>2</sup>	Polymorphic Substitutions <sup>2</sup>
Qβ-AZC(t60)	5.7	4.0	5.3	1.7
Qβ-AZC(t70)	5.7	2.7	6.9	1.5
Qβ-AZC(t90)	7.2	3.2	7.1	1.6
Average <sup>3</sup>	6.2 ± 0.9	3.3 ± 0.7	6.4 ± 1.0	1.6 ± 0.1

<sup>1</sup>The value ds/dn indicates the ratio of synonymous to non synonymous substitutions corrected by the potential number of synonymous and non synonymous positions. The genomic region analyzed comprised from nucleotide 1485 to nucleotide 4028, excluding the non coding region (from nt 2331 to nt 2351). The number of synonymous and non synonymous positions was evaluated using the program SNAP [34,35].

<sup>2</sup>Polymorphic substitutions (those shown in Table 1 together with the fixed substitutions A1746U and A2187C) and non polymorphic substitutions (those shown in Additional file 1) were analyzed separately. All these mutations were placed in coding regions.

<sup>3</sup>Average value of the ratio ds/dn for the 3 populations analyzed. The difference between the average values obtained for non polymorphic and polymorphic substitutions was significant ( $p < 0.01$  in both analysis, Student's *t* test for the difference of means).

agreement with their relative fitness values, mutants Qβ<sub>A2982G</sub>, Qβ<sub>U3582C</sub>, and Qβ<sub>G3945A</sub> fixed in the presence of AZC, whereas only Qβ<sub>G3945A</sub> fixed in the absence of AZC. These results indicate that substitution G3945A has a general beneficial fitness effect that is independent of the presence of AZC. Taking into account that A2982G and U3582C were fixed in only five transfers in the presence of AZC, whereas G3945A needed 15 transfers (Table 4), we can state that the last substitution has lower selective advantage under this condition than the first ones. Substitution C3413G remained as a polymorphism after 15 transfers in both the presence and the absence of AZC, which agrees with their fitness values close to neutrality. There is a disagreement between the relative fitness value of Qβ<sub>A1746U</sub> in the presence of AZC (lower than 1) and the fact that this virus was able to displace the virus Qβ<sub>wt</sub> when both competed under this condition. This discrepancy suggests that the growth rate of a virus cannot always be a good predictor of its behaviour when it propagates in the presence of competitor genomes.

The results showing that substitution U3989C was deleterious (Table 4) were puzzling since this substitution reached high frequency in the absence of apparent hitchhiking with any other substitution (Table 1). A possibility is that substitution U3989C is a compensatory mutation that raised high frequency because it reduces the fitness cost of another mutation previously selected. U3989C was generated in the genomic context of A2187C and A1746U. Since A1746U had a fitness cost in the absence of AZC (Table 4) [31], this substitution was the most probable candidate to be compensated by U3989C. To investigate this point we built a site-directed double mutant containing both substitutions A1746U and U3989C (Qβ<sub>A1746U+U3989C</sub>), and determined its relative fitness in the presence and in the absence of AZC. The value obtained in the presence of AZC (0.89 ± 0.2) was significantly higher than those obtained for the single mutants Qβ<sub>A1746U</sub> and Qβ<sub>U3989C</sub> ( $p < 0.05$ , Student's *t* test for the difference of means). However, the relative fitness value obtained in the

**Table 4 Relative fitness of bacteriophage Qβ site-directed mutants**

Site-directed mutant	Relative fitness <sup>1</sup>		Dominant virus in competition with Qβ <sub>wt</sub> <sup>2</sup>	
	+AZC	-AZC	+AZC	-AZC
Qβ <sub>A1746U</sub>	0.71 ± 0.02*	0.58 ± 0.03*	Qβ <sub>A1746U</sub>	Qβ <sub>wt</sub>
Qβ <sub>A2982G</sub>	1.24 ± 0.08*	0.89 ± 0.05*	Qβ <sub>A2982G</sub>	Qβ <sub>wt</sub>
Qβ <sub>U3582C</sub>	1.31 ± 0.03*	0.89 ± 0.06*	Qβ <sub>U3582C</sub>	Qβ <sub>wt</sub>
Qβ <sub>C3413G</sub> <sup>3</sup>	0.95 ± 0.04*	0.92 ± 0.07*	Qβ <sub>C3413G</sub> + Qβ <sub>wt</sub> (transfer 15)	Qβ <sub>C3413G</sub> + Qβ <sub>wt</sub> (transfer15)
Qβ <sub>G3945A</sub> <sup>3</sup>	1.1 ± 0.06*	1.02 ± 0.05	Qβ <sub>G3945A</sub> (transfer15)	Qβ <sub>G3945A</sub> (transfer15)
Qβ <sub>U3989C</sub>	0.79 ± 0.04*	0.88 ± 0.05*	Qβ <sub>wt</sub>	Qβ <sub>wt</sub>

<sup>1</sup>Relative fitness was evaluated with respect to the virus Qβ<sub>wt</sub> either in the absence or the presence of AZC (60 µg/ml). The absolute fitness value estimated for Qβ<sub>wt</sub> was 15.42 ± 0.50 in the absence of AZC and 9.68 ± 0.22 in its presence. The asterisk indicates that the difference between the fitness values of the site-directed mutant and the virus Qβ<sub>wt</sub> was significant ( $p$  value < 0.05, Student's *t* test for the difference of means).

<sup>2</sup>Competition between the site-directed mutants indicated and the virus Qβ<sub>wt</sub> was carried out as described in Methods during 5 transfers either in the absence or the presence of AZC (60 µg/ml).

<sup>3</sup>Since no virus became dominant at transfer number 5, competitions carried out with viruses Qβ<sub>C3413G</sub> and Qβ<sub>G3945A</sub> were extended until transfer number 15, yielding the results shown in the table.




absence of AZC ( $0.80 \pm 0.1$ ) was only significantly higher than that obtained for the single mutant  $Q\beta_{A1746U}$ . Competition experiments of the double mutant with the virus  $Q\beta_{wt}$  showed that after five transfers, the wild virus again dominated in the absence of AZC, indicating that under this condition substitution U3989C was not able to compensate the fitness cost of A1746U (Figure 2a). In contrast to this, the double mutant was selected in the presence of AZC, showing that in this case the combined effect of both substitutions was beneficial. Only with this result we cannot distinguish whether the advantage provided by A1746U in the presence of AZC is strong enough to cause the hitchhiking of U3989C or the double mutant  $Q\beta_{A1746U+U3989C}$  is more advantageous than the single mutant  $Q\beta_{A1746U}$ . To test the last possibility we performed another competition experiment between the double mutant  $Q\beta_{A1746U+U3989C}$  and the single mutant  $Q\beta_{A1746U}$  (Figure 2b). After 10 transfers in both the presence and the absence of AZC, both viruses remained in the population. The result indicates that the effect of substitution U3989C was less deleterious in populations where substitution A1746U was previously fixed than in the mutational context of the wild type virus.

#### Some polymorphic substitutions appear in multiple mutational contexts whereas other ones cannot coexist in the same genome

The analysis of the virus sequences displayed in Table 1 shows that most polymorphic substitutions are found in

multiple mutational contexts, which can be indicative of a high frequency of recurrent mutations in bacteriophage  $Q\beta$ . In contrast to this result, substitutions A2982G and U3582C were never located in the same genome. This finding is even more surprising if we take into account that A2982G and U3582C have selective advantages in the presence of AZC and, hence, their combination in the same genome could lead to a better adapted virus. Therefore, we were interested in analyzing how the behaviour of the double mutant would be. To this end, we introduced both mutations in the infectious clone of bacteriophage  $Q\beta$  (see Methods), and tried to recover the viruses expressed. Sequencing of the plasmid DNA extracted from five *E. coli* transformed colonies showed that the site-directed mutagenesis had been successful. However, using the same conditions that let us obtain viruses when transformation took place with the infectious clone containing any of the two single mutations, we were not able to recover virus from any of the bacteria transformed with the infectious clone containing both substitutions. This result strongly suggests that the presence of both A2982G and U3582C in the same genome either is lethal for the virus or highly deleterious, a conclusion that agrees with the absence of any mutant containing both substitutions in the mutant spectrum of the populations analyzed in this work (Table 1), and also with the fitness costs that both substitutions have in the absence of AZC (Table 4) [31].



### **Bacteriophage Q $\beta$ populations evolved in the presence of AZC are composed by genomes that can be grouped into multiple evolutionary lines**

The observation that the combination in the same genome of substitutions A2982G and U3582C, both with clear selective advantages in the presence of AZC, is lethal or highly deleterious, and, hence, they are never associated in the same genome suggests that during the evolution of bacteriophage Q $\beta$  under mutagenic conditions, at least two evolutionary lines that evolve independently are generated. The first line would comprise virus clones C<sub>1</sub> to C<sub>4</sub> in population Q $\beta$ -AZC(t60), C<sub>1</sub> to C<sub>3</sub> in population Q $\beta$ -AZC(t70), and C<sub>1</sub> to C<sub>8</sub> in population Q $\beta$ -AZC(t90) (Table 1). Genomes from these viruses have in common the presence of substitution A2982G, accompanied by U1520C, G3945A, and U3989C. The second line would contain the virus clones C<sub>4</sub> to C<sub>6</sub> in population Q $\beta$ -AZC(t70), and C<sub>9</sub> to C<sub>11</sub> in population Q $\beta$ -AZC(t90), all carrying substitution U3582C (Table 1). Finally, a third line could be established containing the remaining genomes, which lack a clear set of polymorphic mutations in common.

A phylogenetic analysis carried out with the 35 genomes analyzed in this work (Figure 3) shows that all the genomes that we had previously included in line 1 (those carrying substitution A2982G, together with U1520C, G3945A, and U3989C) group into a cluster with a high bootstrap value (cluster a). This cluster is a part of another one (cluster b) that contains some additional genomes carrying substitution U3989C. The rest of genomes groups into 4 independent clusters differing in their polymorphic substitutions (Figure 3). It is remarkable that whereas all the genomes containing substitution A2982G are grouped into the same cluster, genomes containing substitution U3582C are placed in different ones, suggesting that this substitution has appeared repeatedly in different evolutionary lines.

Taking into account the transfer number at which each polymorphism was first detected (Table 2), a feasible sequence of events for the generation of the different lines can be established. Genomes containing A2187C (fixed at transfer number 30) and A1746U (fixed at transfer number 40) evolved through the acquisition of additional substitutions that can become represented at high frequency. One of the mutations appearing first was U3989C, which spreaded giving rise to the genomes included in clusters a and b (Figure 3). The remaining genomes continued replicating, and in this process they acquired some of the other polymorphic mutations listed in Table 1. Genomes belonging to different lines can compete among them, delaying the fixation of their mutations and allowing for the prolonged presence of a high number of polymorphisms. Moreover, genomes within each line can differ in additional polymorphic substitutions (see Table 1), and also in mutations that

are exclusive of each of them (Additional file 1), which can modify their selective advantages adding further complexity to the competition process.

### **The presence in different genomes of mutations having a selective advantage in the presence of AZC interferes with their fixation**

To demonstrate whether the coexistence of multiple evolutionary lines carrying different mutations with selective advantages in the presence of AZC interferes with their fixation, we compared the results of competition experiments carried out between different site-directed mutants (Q $\beta$ <sub>A1746U</sub>, Q $\beta$ <sub>A2982G</sub>, and Q $\beta$ <sub>U3582C</sub>) and between each one of the mutants and the virus Q $\beta$ <sub>wt</sub> (Table 5).

Our results show that the virus Q $\beta$ <sub>A1746U</sub> became dominant after 5 transfers when it competed with either Q $\beta$ <sub>A2982G</sub> or Q $\beta$ <sub>U3582C</sub>, as it happened when the competitor was the wild type virus (Table 5). In contrast to this result, competition between the site-directed mutants Q $\beta$ <sub>A2982G</sub> and Q $\beta$ <sub>U3582C</sub> rendered polymorphisms at both positions after 20 transfers in the presence of AZC (Table 5), indicating that no virus was dominant in the population at this point. This result represents a clear delay relative to the fixation of both mutations in only 5 transfers when the site-directed mutants competed with the wild type virus.

We also analyzed the behaviour of the site-directed double mutants, Q $\beta$ <sub>A1746U+A2982G</sub> and Q $\beta$ <sub>A1746U+U3582C</sub> in competition experiments with the single mutant Q $\beta$ <sub>A1746U</sub>. In both cases the double mutant completely displaced the single mutant after 5 transfers in the presence of AZC (Table 5). However, similar competition experiments carried out by mixing the two double mutants or the two double mutants together with the single mutant Q $\beta$ <sub>A1746U</sub>, showed that no virus became dominant after 20 transfers in the presence of AZC (Table 5). This result reinforces the view that the spreading of the genomes containing substitution A2982G interferes with the spreading of the genomes carrying U3582C.


### **Discussion**

In this work we analyze how beneficial mutations spread in a virus population when replication takes place at increased error rate. To carry out this study we have chosen an RNA virus, the bacteriophage Q $\beta$ , which as a consequence of its high error rate [40,41] constitutes extremely heterogeneous populations composed by a complex mutant spectrum [42]. The presence of AZC in the growing medium increased the mutation frequency in viable viruses about an order of magnitude relative to that observed in virus populations evolved in the absence of AZC [31]. The mutation excess (Table 1 and Additional file 1) probably distributes among deleterious, beneficial and neutral mutations, although the much

higher frequency of errors having a negative effect on fitness suggests that many of them must be deleterious. Thus, it is expected that beneficial mutations generated in the presence of AZC arise in unfavourable genomic contexts, which can reduce their selective advantages [14-16]. In addition, when the error rate is high enough several beneficial mutations can expand simultaneously, which constitutes another important factor delaying their spreading in the population [12,13]. Thus, although replication at high error rate provides important adaptive advantages to RNA viruses [43,44], it is also true that when subjected to increased mutagenesis, they are

frequently pushed to the edge of extinction [45-47], as a result of both the increase in the number of deleterious mutations and the difficulties for the expansion of the beneficial ones.

The analysis of the consensus sequences and the mutant spectra of the AZC-evolved populations showed the fixation of two mutations and the presence of a number of polymorphisms (Table 1) higher than that found during the evolution of the same virus in the absence of AZC (see the two first subsections of Results). It should also be noted that, given the relatively low number of genomes analyzed in each population, the frequency of



**Figure 3 Phylogenetic analysis of the virus genomes isolated from bacteriophage Q $\beta$  populations evolved in the presence of AZC.** Different colours are used to distinguish the genomes from each population included in the analysis: green for Q $\beta$ -AZC(t60), pink for Q $\beta$ -AZC(t70), and orange for Q $\beta$ -AZC(t90). Virus genomes corresponded to those shown in Table 1 and Additional file 1, and are identified using the same notation. The tree was derived by maximum likelihood methods (PhyML, program seaview 4) [39] using the sequence of the wild type virus to root the tree. Numbers at each node represent the bootstrap value (carried out with 100 replicates). Clusters described in the main text are highlighted, and the mutations common to all the genomes included in each one are also indicated.

**Table 5 Competition among different site-directed mutants of bacteriophage Q $\beta$  in the presence of AZC**

Competitor viruses	Viruses present in the final population <sup>1</sup>	
	Transfer number 5	Transfer number 20
Q $\beta$ <sub>wt</sub> + Q $\beta$ <sub>A1746U</sub>	Q $\beta$ <sub>A1746U</sub>	ND <sup>2</sup>
Q $\beta$ <sub>wt</sub> + Q $\beta$ <sub>A2982G</sub>	Q $\beta$ <sub>A2982G</sub>	ND
Q $\beta$ <sub>wt</sub> + Q $\beta$ <sub>U3582C</sub>	Q $\beta$ <sub>U3582C</sub>	ND
Q $\beta$ <sub>A1746U</sub> + Q $\beta$ <sub>A2982G</sub>	Q $\beta$ <sub>A1746U</sub>	ND
Q $\beta$ <sub>A1746U</sub> + Q $\beta$ <sub>U3582C</sub>	Q $\beta$ <sub>A1746U</sub>	ND
Q $\beta$ <sub>A2982G</sub> + Q $\beta$ <sub>U3582C</sub>	Q $\beta$ <sub>A2982G</sub> + Q $\beta$ <sub>U3582C</sub>	Q $\beta$ <sub>A2982G</sub> + Q $\beta$ <sub>U3582C</sub>
Q $\beta$ <sub>A1746U</sub> + Q $\beta$ <sub>A1746U</sub> + A2982G	Q $\beta$ <sub>A1746U</sub> + A2982G	ND
Q $\beta$ <sub>A1746U</sub> + Q $\beta$ <sub>A1746U</sub> + U3582C	Q $\beta$ <sub>A1746U</sub> + U3582C	ND
Q $\beta$ <sub>A1746U</sub> + A2982G + Q $\beta$ <sub>A1746U</sub> + U3582C	Q $\beta$ <sub>A1746U</sub> + A2982G + Q $\beta$ <sub>A1746U</sub> + U3582C	Q $\beta$ <sub>A1746U</sub> + A2982G + Q $\beta$ <sub>A1746U</sub> + U3582C
Q $\beta$ <sub>A1746U</sub> + Q $\beta$ <sub>A1746U</sub> + A2982G + Q $\beta$ <sub>A1746U</sub> + U3582C	(Q $\beta$ <sub>A1746U</sub> ) <sup>3</sup> + Q $\beta$ <sub>A1746U</sub> + A2982G + Q $\beta$ <sub>A1746U</sub> + U3582C	(Q $\beta$ <sub>A1746U</sub> ) <sup>3</sup> + Q $\beta$ <sub>A1746U</sub> + A2982G + Q $\beta$ <sub>A1746U</sub> + U3582C

<sup>1</sup>Competitions were carried out as described in Methods. The viruses present in the populations obtained at the transfer indicated were identified through the determination of the consensus sequences at the nucleotide positions that distinguish the competitor viruses.

<sup>2</sup>ND means not determined.

<sup>3</sup>In this case, the determination of the consensus sequence do not allow distinguishing whether Q $\beta$ <sub>A1746U</sub> is present in the populations analyzed or not. We show this virus in brackets to indicate this uncertainty.

polymorphic mutations could be even higher than we have reported. Adaptation of bacteriophage MS2 to different selective pressures in the absence of mutagenic conditions also led to the presence of polymorphisms, although to a lower extent to that reported in the current study [26,29]. The findings obtained in studies performed with DNA viruses, which replicate with lower error rate than RNA viruses, are more difficult to compare to our results [30]. Nevertheless, evolution of bacteriophage T7 in the presence of a mutagen also showed a dramatic increase in the number of polymorphic substitutions [48]. Therefore, the high frequency of polymorphisms observed during the evolution of bacteriophage Q $\beta$  in the presence of AZC seems to be due, at least in part, to the increase of the error rate.

The high population sizes used for the transmission of the virus (above 10<sup>7</sup> pfu) together with the absence of clear evidences of hitchhiking suggests that most polymorphic mutations provide a selective advantage, at least in the selective environment provided by the mutagen. The lower values of the ratios ds/dn obtained for the polymorphic mutations than for the non polymorphic ones (Table 3) also supports that selection can be responsible, at least in part, of the high frequency reached by some substitutions. In addition, synonymous changes can also have a fitness effect mediated through *cis* interactions that may be significant in the case of highly compacted genomes, as it is the one of bacteriophage Q $\beta$  [49-51]. In contrast to our assumption that most polymorphic mutations are beneficial, there are some

studies showing that in natural populations of RNA viruses many high frequency mutations are deleterious, and will be later purged by natural selection [52]. The easiest way for these deleterious mutations to reach high frequency is their linkage with a beneficial mutation. In the absence of clear associations among mutations, this explanation could still be possible if mutations recur frequently as it seems to happen in bacteriophage Q $\beta$  (see below).

We have demonstrated that one of the mutations fixed (A2187C) and at least one of the polymorphic substitutions (G3945A) are beneficial in both the presence and the absence of AZC (Table 4) [31], whereas others (A2982G, and U3582C) only provide an advantage in the selective medium, having a fitness cost under standard replication conditions [31]. The results obtained with mutant Q $\beta$ <sub>A1746U</sub> deserve particular attention. This mutant has lower fitness than the virus Q $\beta$ <sub>wt</sub> when they grow independently. However, Q $\beta$ <sub>A1746U</sub> fixes in only 5 transfers when both viruses propagate together in the presence of AZC (Table 4). Fitness is a complex parameter that involves many traits, among which some of the most relevant are the lysis time, the adsorption rate, the burst size, and the stability of the particles outside of the host [53]. It is possible that some of these features influence differently the performance of a virus when it grows isolated or in competition. In good agreement with these considerations Springman et al. [48] reported fitness gains in a population of bacteriophage T7 that, however, showed a clear decline in the burst size.

Our results also showed that the fitness advantages provided by the substitutions assayed were of different magnitude (Tables 4 and 5), which is in disagreement with the proposal of the multiple mutations model which posits a single characteristic value for the coefficient of selection of all beneficial mutations [17,18]. The situation becomes more complex due to mutations whose effects are context dependent. This is the case of substitutions A2982G and U3582C which were beneficial when present in separate genomes, and, however, were lethal or highly deleterious when present in the same genome, providing a clear example of antagonistic epistasis. Another substitution whose effect varied depending on the mutational context and on the competitor genomes was U3989C. This substitution behaved as deleterious in both the presence and the absence of AZC when it was present in the mutational context of the wild type virus (Table 4), and, however, it could be propagated when it was present in the mutational context of substitution A1746U (Figure 2b). These findings support the notion that epistatic interactions among mutations are very frequent in RNA viruses [6,54-58] and can influence the adaptive trajectories followed and the intensity of interference. The increase of the error rate may exacerbate the frequency of this type of interactions, which are not included in any of the current models developed to explain the fixation of beneficial mutations in asexual populations.

The fixation of substitution A1746U was not delayed by the presence in different genomes of either A2982G or U3582C (Table 5). As a consequence, this substitution was probably little affected by interference and could get fixed in population Q $\beta$ -AZC(t40). By contrast, substitutions A2982G and U3582C that fixed in 5 transfers when they competed with the wild type virus remained as polymorphisms during at least 20 transfers when they competed with each other (Table 5). A similar competition could be established among the different mutants present in the populations analyzed in this work, supporting that interference among mutations is one of the reasons underlying the sustained presence of polymorphisms.

A phylogenetic analysis carried out with the complete set of genomes obtained at different stages of the evolutionary series allowed us to group them into several clusters that represent independent evolutionary lines that may compete among them (Figure 3). In addition to the mutations representatives of each line, genomes can also contain different mutations among those represented at high frequency, and others that are exclusive of each virus. These mutations can modify the fitness of the genomes where they appear, adding further complexity to the process of interference, and making it difficult to fix any beneficial mutation and

the emergence of a defined consensus sequence. The generation of new beneficial mutations in genomes carrying others still not fixed, together with the different magnitude of the effects of beneficial mutations, is more in line with the model of complete interference [21] than with the models of clonal interference [13,14] or multiple mutations [17,18].

Another important finding is that most polymorphic mutations were present in different genomic contexts, suggesting that they were generated repeated times. Results obtained in two adaptation experiments carried out with the bacteriophage MS2 also showed a high number of beneficial mutations in different mutational contexts [26,29]. A theoretical analysis demonstrated that this repeated presence of mutations was more probably due to their generation multiple times than to recombination [26]. The low capability of Q $\beta$  replicase to switch between templates does not allow for homologous recombination to be observed in most of the systems assayed [59], as it would be masked by the much higher frequency of punctual mutations [60]. Therefore, we think that the presence of polymorphic mutations in different mutational contexts is more likely due to a high recurrence of punctual mutations than to recombination. This fact would allow combining several beneficial mutations in the same genome, alleviating in this way the costs of interference [61]. However, our results showing that two of the mutations having high selective value in the presence of AZC (A2982G and U3582C) cannot coexist in the same genome suggest that the fitness landscape for bacteriophage Q $\beta$  is highly rugged, with the existence of several local maxima where the virus could be trapped without reaching the best adaptive solution.

The high incidence of recurrent mutations could also account for the few hitchhiking mutations found in this study. Given the high mutation frequencies of the bacteriophage Q $\beta$  populations evolved in the presence of AZC, it would seem reasonable that almost each genome with a beneficial mutation in the process to fixation will also carry a set of hitchhiking mutations. However, the repeated occurrence of the beneficial mutations on different backgrounds also means that no single set of accompanying mutations gets fixed. A similar result was reported for bacteriophage MS2 adapting to cold temperatures [29], and was also supported by theoretical calculations [62].

Although at a first glance the interference among mutations can be seen as a negative feature that delays adaptation, there are also positive consequences that deserve to be pointed out. Maybe one of the most relevant is the coexistence in the population of multiple adaptive possibilities that can provide selective advantages by different molecular mechanisms. Population bottlenecks



occurring during the propagation of the virus can lead to the fixation of different adaptive solutions. In this way, interference could be seen as a positive factor contributing to the diversification of populations, and permitting the exploration of the different local maxima that exist in rugged fitness landscapes, such as those described for RNA viruses [63,64].

Interference among mutations can also play a relevant role in the extinction of viruses through lethal mutagenesis, a new antiviral strategy that derives from theoretical considerations [65-67] and that consists in the treatment of virus infections through the artificial increase of the virus error rate [46,47,68,69]. Our results show that virus replication under mutagenic conditions can lead to the simultaneous presence in the mutant spectrum of multiple mutations conferring different advantages in the presence of the mutagen. The fixation of these mutations in particular individuals upon transmission of the virus through population bottlenecks, as indicated above, can lead to the co-circulation of viruses differing in their adaptive properties, jeopardizing in this way the efficacy of further treatments. Given the high error rates of RNA viruses, similar situations could also occur during the treatment with some replication inhibitors. Therefore, it is expected that future research on the evolutionary consequences of the interference among mutations also provides significant benefits to clinic and epidemiologic virology.

## Conclusions

Evolution of bacteriophage Q $\beta$  at artificially increased error rate by means of the use of a mutagenic nucleoside analogue leads to the prolonged permanence of multiple polymorphisms which could be detected in both the consensus sequences and the mutant spectra.

Polymorphic mutations have different effects on fitness. Some of them provide selective advantages in both the presence and the absence of AZC, whereas others only are beneficial under selective conditions, having a fitness cost under standard replication conditions. Epistatic interactions also play a role in deciding whether a particular mutation will reach high frequency or not.

Polymorphic mutations distribute into multiple evolutionary lines that compete among them making it difficult the emergence of a defined consensus sequence. Each evolutionary line can provide a selective advantage by a different molecular mechanism leading to the coexistence of multiple adaptive pathways in the same population.

Antagonistic epistasis determines that two of the replicase mutations providing selective advantages in the presence of AZC cannot associate in the same genome. As a consequence, genomes carrying each of these

mutations spread simultaneously delaying the fixation of any of them.

The variety of mutational contexts in which most of the polymorphic mutations have been detected indicates that beneficial mutation recur frequently in bacteriophage Q $\beta$ . This circumstance could alleviate the disadvantages caused by interference. However, the fact that some of the adaptive substitutions cannot be combined in the same genome limits the potential benefits of the repeated generation of beneficial mutations.

The fixation of mutations in bacteriophage Q $\beta$  evolved in the presence of AZC is better approached by the model of complete interference than by the models of clonal interference or multiple mutations. However, the presence of epistatic interactions and the high frequency of recurrent mutations contribute to generate a highly complex interference dynamics, which would require further improvements of the theoretical models to be successfully described.

## Additional file

**Additional file 1: Additional substitutions to those represented at high frequency in the virus genomes analyzed.** Description: It contains all the substitutions that are not represented at high frequency in the mutant spectra of the virus populations analyzed. Genomes correspond to those indicated in Table 1.

## Competing interests

The authors declare that they do not have competing interests.

## Authors' contributions

MA and LC contributed equally to this paper. EL designed the study. MA and LC carried out the experiments. EL, MA, and LC interpreted the results. EL wrote the paper. All the authors read and approved the final manuscript.

## Acknowledgments

We are indebted to SC Manrubia and E Domingo for useful discussions throughout the development of this study and for the critical reading of the manuscript. Manuel J Gómez is also acknowledged for helping with the phylogenetic analysis. Work was financially supported by the Spanish MICINN through projects FIS2011-27569 and BFU2011-25271. We acknowledge support of the publication fee by the CSIC Open Access Publication Support Initiative through its Unit of Information Resources for Research (URICI).

Received: 1 August 2012 Accepted: 11 January 2013

Published: 16 January 2013

## References

1. Muller HJ: Some genetic aspects of sex. *Am Nat* 1932, **68**:118-138.
2. Muller HJ: The relation of recombination to mutational advance. *Mutat Res* 1964, **106**:2-9.
3. Atwood KC, Schneider LK, Ryan FJ: Periodic selection in *Escherichia coli*. *Proc Natl Acad Sci USA* 1951, **37**:146-155.
4. Papadopoulos D, Schneider D, Meier-Eiss J, Arber W, Lenski RE, Blot M: Genomic evolution during a 10,000-generation experiment with bacteria. *Proc Natl Acad Sci USA* 1999, **96**:3807-3812.
5. Wichman HA, Badgett MR, Scott LA, Boulianne CM, Bull JJ: Different trajectories of parallel evolution during viral adaptation. *Science* 1999, **285**:422-424.
6. Holder KK, Bull JJ: Profiles of adaptation in two similar viruses. *Genetics* 2001, **159**:1393-1404.

7. Rozen DE, de Visser JA, Gerrish PJ: **Fitness effects of fixed beneficial mutations in microbial populations.** *Curr Biol* 2002, **12**:1040–1045.
8. Shaver AC, Dombrowski PG, Sweeney JY, Treis T, Zappala RM, Sniegowski PD: **Fitness evolution and the rise of mutator alleles in experimental *Escherichia coli* populations.** *Genetics* 2002, **162**:557–566.
9. de Visser JA, Rozen DE: **Clonal interference and the periodic selection of new beneficial mutations in *Escherichia coli*.** *Genetics* 2006, **172**:2093–2100.
10. Lang GI, Botstein D, Desai MM: **Genetic variation and the fate of beneficial mutations in asexual populations.** *Genetics* 2011, **188**:647–661.
11. de Visser JA, Rozen DE: **Limits to adaptation in asexual populations.** *J Evol Biol* 2005, **18**:779–788.
12. Sniegowski PD, Gerrish PJ: **Beneficial mutations and the dynamics of adaptation in asexual populations.** *Philos Trans R Soc Lond B Biol Sci* 2010, **365**:1255–1263.
13. Gerrish PJ, Lenski RE: **The fate of competing beneficial mutations in an asexual population.** *Genetica* 1998, **102–103**:127–144.
14. Wilke CO: **The speed of adaptation in large asexual populations.** *Genetics* 2004, **167**:2045–2053.
15. Orr HA: **The rate of adaptation in asexuals.** *Genetics* 2000, **155**:961–968.
16. Orr HA: **The distribution of fitness effects among beneficial mutations.** *Genetics* 2003, **163**:1519–1526.
17. Desai M, Fisher DS: **Beneficial mutation-selection balance and the effect of linkage on positive selection.** *Genetics* 2007, **176**:1759–1798.
18. Desai M, Fisher DS, Murray AW: **The speed of evolution and maintenance of variation in asexual populations.** *Curr Biol* 2007, **17**:385–394.
19. Brunet E, Rouzine IM, Wilke CO: **The stochastic edge in adaptive evolution.** *Genetics* 2008, **179**:603–620.
20. Hegreness M, Shores N, Hartl D, Kishony R: **An equivalence principle for the incorporation of favorable mutations in asexual populations.** *Science* 2006, **17(311)**:1615–1617.
21. Park SC, Krug J: **Clonal interference in large populations.** *Proc Natl Acad Sci USA* 2007, **104**:18135–18140.
22. de Visser JA, Zeyl CW, Gerrish PJ, Blanchard JL, Lenski RE: **Diminishing returns from mutation supply rate in asexual populations.** *Science* 1999, **283**:404–406.
23. Barrick JE, Lenski RE: **Genome-wide mutational diversity in an evolving population of *Escherichia coli*.** *Cold Spring Harb Symp Quant Biol* 2009, **74**:119–29.
24. Miralles R, Gerrish PJ, Moya A, Elena SF: **Clonal interference and the evolution of RNA viruses.** *Science* 1999, **285**:1745–1747.
25. Kao KC, Sherlock G: **Molecular characterization of clonal interference during adaptive evolution in asexual populations of *Saccharomyces cerevisiae*.** *Nat Genet* 2008, **40**:1499–1504.
26. Bollback JP, Huelsenbeck JP: **Clonal interference is alleviated by high mutation rates in large populations.** *Mol Biol Evol* 2007, **24**:1397–1406.
27. Gresham D, Desai MM, Tucker CM, Jenq HT, Pai DA, Ward A, DeSevo CG, Botstein D, Dunham MJ: **The repertoire and dynamics of evolutionary adaptations to controlled nutrient-limited environments in yeast.** *PLoS Genet* 2008, **4(12)**:e1000303.
28. Barrick JE, Yu DS, Yoon SH, Jeong H, Oh TK, Schneider D, Lenski RE, Kim JF: **Genome evolution and adaptation in a long-term experiment with *Escherichia coli*.** *Nature* 2009, **461**:1243–1247.
29. Betancourt AJ: **Genomewide patterns of substitution in adaptively evolving populations of the RNA bacteriophage MS2.** *Genetics* 2009, **181**:1535–1544.
30. Miller CR, Joyce P, Wichman HA: **Mutational effects and population dynamics during viral adaptation challenge current models.** *Genetics* 2011, **187**:185–202.
31. Arribas M, Cabanillas L, Lázaro E: **Identification of mutations conferring 5-azacytidine resistance in bacteriophage Q $\beta$ .** *Virology* 2011, **417**:343–352.
32. Cases-González C, Arribas M, Domingo E, Lázaro E: **Beneficial effects of population bottlenecks in an RNA virus evolving at increased error rate.** *J Mol Biol* 2008, **384**:1120–1129.
33. Barrera I, Schuppli D, Sogo JM, Weber H: **Different mechanisms of recognition of bacteriophage Q $\beta$  plus and minus strand RNAs by Q $\beta$  replicase.** *J Mol Biol* 1993, **232**:512–521.
34. *The HIV databases.* www.hiv.lanl.gov.
35. Korber B: **HIV Signature and Sequence Variation Analysis.** In *Computational Analysis of HIV Molecular Sequences.* Edited by Rodrigo AG, Learn GH. Dordrecht, Netherlands: Kluwer Academic Publishers; 2000:55–72.
36. Lynch M, Gabriel W: **Mutational load and the survival of small populations.** *Evolution* 1990, **44**:1725–1737.
37. Lázaro E, Escarmis C, Pérez-Mercader J, Manrubia SC, Domingo E: **Resistance of virus to extinction on bottleneck passages: study of a decaying and fluctuating pattern of fitness loss.** *Proc Natl Acad Sci USA* 2003, **100**:10830–10835.
38. Escarmis C, Lázaro E, Manrubia SC: **Population bottlenecks in quasispecies dynamics.** *Curr Top Microbiol Immunol* 2006, **299**:141–70.
39. Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O: **New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0.** *Syst Biol* 2010, **59**:307–321.
40. Drake JW: **Rates of spontaneous mutation among RNA viruses.** *Proc Natl Acad Sci USA* 1993, **90**:4171–4175.
41. Drake JW, Charlesworth B, Charlesworth D, Crow JF: **Rates of spontaneous mutation.** *Genetics* 1998, **148**:1667–1686.
42. Domingo E, Sabo D, Taniguchi T, Weissmann C: **Nucleotide sequence heterogeneity of an RNA phage population.** *Cell* 1978, **13**:735–744.
43. Pfeiffer JK, Kirkegaard K: **Increased fidelity reduces poliovirus fitness and virulence under selective pressure in mice.** *PLoS Pathog* 2005, **1(2)**:e11.
44. Vignuzzi M, Stone JK, Arnold JJ, Cameron CE, Andino R: **Quasispecies diversity determines pathogenesis through cooperative interactions in a viral population.** *Nature* 2006, **439**:344–348.
45. Holland JJ, Domingo E, de la Torre JC, Steinhauer DA: **Mutation frequencies at defined single codon sites in vesicular stomatitis virus and poliovirus can be increased only slightly by chemical mutagenesis.** *J Virol* 1990, **64**:3960–3962.
46. Loeb LA, Essigmann JM, Kazazi F, Zhang J, Rose KD, Mullins JL: **Lethal mutagenesis of HIV with mutagenic nucleoside analogs.** *Proc Natl Acad Sci USA* 1999, **96**:1492–1497.
47. Domingo E: **Virus entry into error catastrophe as a new antiviral strategy.** *Virus Res* 2005, **107**:115–228.
48. Springman R, Keller T, Molineux IJ, Bull JJ: **Evolution at a high imposed mutation rate: adaptation obscures the load in phage T7.** *Genetics* 2010, **184**:221–232.
49. Mills DR, Priano C, Merz P, Binderow BD: **Q $\beta$  RNA bacteriophage: mapping cis-acting elements within an RNA genome.** *J Virol* 1990, **64**:3872–3881.
50. Klovinis J, Berzins V, van Duin J: **A long-range interaction in Qbeta RNA that bridges the thousand nucleotides between the M-site and the 3' end is required for replication.** *RNA* 1998, **4**:948–957.
51. Klovinis J, van Duin J: **A long-range pseudoknot in Qbeta RNA is essential for replication.** *J Mol Biol* 1999, **294**:875–884.
52. Pybus OG, Rambaut A, Belshaw R, Freckleton RP, Drummond AJ, Holmes EC: **Phylogenetic evidence for deleterious mutation load in RNA viruses and its contribution to viral evolution.** *Mol Biol Evol* 2007, **24**:845–852.
53. Shao Y, Wang IN: **Bacteriophage adsorption rate and optimal lysis time.** *Genetics* 2008, **180**:471–482.
54. Sanjuán R, Moya A, Elena SF: **The contribution of epistasis to the architecture of fitness in an RNA virus.** *Proc Natl Acad Sci USA* 2004, **101**:15376–15379.
55. Sanjuán R, Cuevas JM, Moya A, Elena SF: **Epistasis and the adaptability of an RNA virus.** *Genetics* 2005, **170**:1001–1008.
56. Sanjuán R, Elena SF: **Epistasis correlates to genomic complexity.** *Proc Natl Acad Sci USA* 2006, **103**:14402–14405.
57. Weinreich DM, Watson RA, Chao L: **Perspective: Sign epistasis and genetic constraint on evolutionary trajectories.** *Evolution* 2005, **59**:1165–1174.
58. Betancourt AJ: **Lack of evidence for sign epistasis between beneficial mutations in an RNA bacteriophage.** *J Mol Evol* 2010, **71**:437–443.
59. Chetverin AB, Chetverina HV, Demidenko AA, Ugarov VI: **Nonhomologous RNA recombination in a cell-free system: evidence for a transesterification mechanism guided by secondary structure.** *Cell* 1997, **88**:503–513.
60. Palasingam K, Shaklee PN: **Reversion of Q beta RNA phage mutants by homologous RNA recombination.** *J Virol* 1992, **66**:2435–2442.
61. Kim Y, Orr HA: **Adaptation in sexuals vs. asexuals: clonal interference and the Fisher-Muller model.** *Genetics* 2005, **171**:1377–1386.
62. Charlesworth B, Morgan MT, Charlesworth D: **The effect of deleterious mutations on neutral molecular variation.** *Genetics* 1993, **134**:1289–1303.
63. Elena SF, Sanjuán R: **RNA viruses as complex adaptive systems.** *Biosystems* 2005, **81**:31–41.
64. Lalić J, Elena SF: **Magnitude and sign epistasis among deleterious mutations in a positive-sense plant RNA virus.** *Heredity* 2012, **109**:71–7.

65. Eigen M, Schuster P: *The hypercycle. A principle of natural self-organization*. Berlin: Springer; 1979.
66. Nowak M, Schuster P: Error thresholds of replication in finite populations mutation frequencies and the onset of Muller's ratchet. *J Theor Biol* 1989, **137**:375–395.
67. Biebricher CK, Eigen M: The error threshold. *Virus Res* 2005, **107**:117–127.
68. Eigen M: Error catastrophe and antiviral strategy. *Proc Natl Acad Sci USA* 2002, **99**:13374–13376.
69. Lázaro E: *RNA Viruses: Control, Mutagenesis and Extinction*, eLS. Ltd: Chichester: John Wiley & Sons; doi:10.1002/9780470015902.a0023276.

doi:10.1186/1471-2148-13-11

**Cite this article as:** Cabanillas et al.: Evolution at increased error rate leads to the coexistence of multiple adaptive pathways in an RNA virus. *BMC Evolutionary Biology* 2013 **13**:11.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

