



Short Communication

The principles of ultra-low contrast percutaneous coronary intervention

Babu Ezhumalai*

Department of Cardiology, Fortis Malar Hospital, Adyar, Chennai, India



ARTICLE INFO

Article history:

Received 10 February 2022

Received in revised form

21 March 2022

Accepted 9 April 2022

Available online 12 April 2022

Keywords:

Ultra-low contrast PCI

Zero contrast PCI

Contrast-induced acute kidney injury

IVUS-Guided PCI

PCI in Chronic kidney disease

ABSTRACT

Ultra-low contrast percutaneous coronary intervention (ULCPCI) can be performed electively in advanced chronic kidney disease. Engage guide catheter and advance guidewire into the coronary artery without using contrast. IVUS-guided PCI can reduce the contrast load. Perform co-registration of distal and proximal radio-opaque marker bands of intravascular ultrasound (IVUS) catheter. Deploy the stent at the target lesion under fluoroscopic guidance of these co-registered position of the IVUS-marking images. Complete the ULCPCI procedure with a final angiography using minimal contrast. Newer contrast sparing techniques and intravascular imaging technologies provide opportunities to perform ULCPCI efficiently with good results and the least complications.

© 2022 Cardiological Society of India. Published by Elsevier, a division of RELX India, Pvt. Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Contrast-induced acute kidney injury (CI-AKI) is a serious issue following coronary angiography (CAG) and percutaneous coronary interventions (PCI) in patients with chronic kidney disease (CKD). The development of CI-AKI prolongs the duration of hospitalization, increases the financial burden and adversely affects both short-term and long-term mortality rates especially in those with advanced CKD (estimated glomerular filtration rate <30 ml/min).¹ Ultra-low contrast PCI (ULCPCI) can be a solution for this situation; this technique is also known as Zero Contrast PCI.^{2–5}

The following are the general principles for performing ULCPCI in advanced CKD patients:

1. Pre-procedural hydration of the patient is very essential.¹ Perform baseline coronary angiography (CAG) with non-ionic iso-osmolar contrast (iodixanol) prior to the elective ULCPCI on a different day with the least possible contrast volume (CV) preferably less than the estimated glomerular filtration rate (eGFR) value. Maintain a CV/eGFR ratio <1.⁴ In order to reduce the volume of contrast, preferably use small diameter catheter (5 Fr or less) without side holes, small syringes (3–5 cc) and biplane imaging system.⁶ Dilute the

contrast with saline and optimally engage the vessel to make the most use of contrast. Use increased acquisition rates (30 frames/sec) to improve vessel visualization with minimal contrast load during CAG and to evaluate the final result after ULCPCI.⁴

2. Whenever possible, perform ULCPCI as a staged procedure 7 days after CAG to allow complete washout of contrast from the body.⁷ Transfemoral approach may preserve radial artery for fistula and dialysis in the future in CKD patients.⁴ 7F guide catheter allows simultaneous passage of multiple guidewires and intravascular ultrasound (IVUS) catheter.^{4,5} Smaller French guide catheters including slender PCI technique may reduce the contrast volume.⁶ Confirm successful engagement of the guide catheter by noting transient ST segment depression in the ECG monitor by injecting 10 ml of saline or by-passing coronary guidewire down into the coronary artery.⁸ Take additional guidance from fluoroscopic markers like coronary calcification, post-surgical clips etc.⁷ Avoid frequent contrast puffs and eliminate the contrast in the diagnostic/guiding catheter by aspiration or back-bleeding prior to exchanging the hardwares or administering drugs.^{4,9}
3. With previous baseline CAG as reference, advance guidewire distally into the coronary artery and cross the target lesion without using contrast.⁹ Park another guidewire in any adjacent side branch as a landmark. If possible, park two coronary guidewires in the side branches proximal and distal

* Department of Cardiology, Fortis Malar Hospital, 52, First Main Road, Gandhi Nagar, Adyar, Chennai, 600020, India

E-mail address: drebabu@gmail.com.

to lesion and these will serve as landmarks for the stenotic coronary segment; this is called 'metallic silhouette technique'.^{4,5,7} Another technique to get orientation of coronary artery is using the 'marking wire technique' wherein two wires (operating wire and marking wire) are inserted through a double Y connector. The operating wire is advanced through the main port of this connector while the marking wire advanced via the side port serves as a guide for positioning the stent. Using IVUS, the distal landing zone is determined and marked with the tip of this marking wire. Then, secure this marking wire by screwing the connector's side port and attach a torquer on this wire close to the side port. The proximal landing zone is determined by IVUS and the stent length is estimated with the help of the marking wire.⁴

4. Contrast-sparing strategies such as dual axis rotational angiography, dynamic coronary roadmap, stent enhancement techniques (StentBoost or ClearStent), intracoronary imaging and co-registration techniques can help in reducing the contrast load.⁹
5. IVUS-guided PCI can reduce the contrast load to a great extent. Perform automated IVUS pullback to assess the length of lesion, reference lumen size and the landing zones (percent plaque area less than 50%). Based on these measurements choose a suitable stent preferably a little longer stent.^{2,3,5}
6. Reintroduce IVUS catheter manually and perform a 'dry cine' angiogram for at least two heartbeats without panning the cath table. This allows co-registration of distal and proximal radio-opaque marker bands of IVUS catheter as reference segments for stent placement.^{4,11} After acquiring these IVUS-marking images do not move the flat panel detector arm or the cath table until the procedure is completed.⁵ Avoid breath-holding during ULCPCI as it may alter the reference position.⁴
7. Deploy the stent at the target lesion under fluoroscopic imaging with reference to the position of the IVUS-marking images displayed on the auxiliary monitor. This stent deployment is performed without injecting any contrast or occasionally with only a tiny amount of contrast.²
8. After stenting, perform a final IVUS imaging and post-dilatation as required.^{2,5}
9. Perform a final angiography with less than 3 ml of contrast particularly to recognize any guidewire/dilatation induced coronary artery perforation, slow-flow/no-reflow phenomena, and side branch occlusion.⁴
10. Optical Coherence Tomography (OCT) with dextran has been used instead of IVUS by some operators.^{2,10} Coronary flow physiology assessments like Fractional Flow Reserve (FFR), Coronary flow Reserve (CFR) or instantaneous wave-free ratio (iFR) can also be performed to determine functional outcome post-stenting.^{4,9}
11. Flush the catheter with heparinized saline at regular intervals to reduce the risk of clot formation. Look for pericardial effusion in post-procedure echocardiography to rule out any significant perforation.^{5,9}
12. Although not optimal for beginners, a few experienced operators have been successful in performing ULCPCI in situations like chronic total occlusion lesions that require contralateral injection, calcified lesions that require debulking with Rotational atherectomy or severe stenotic lesions that will not allow IVUS to pass through.^{11,12}

CKD patients are definitely high-risk patients but cannot be ignored. ULCPCI is better performed electively as a planned procedure in advanced CKD patients. Newer contrast sparing techniques and intravascular imaging technologies provide opportunities to perform ULCPCI efficiently with good results and the least complications.

Funding/grants

No funding/grant was obtained for this study.

Declaration of competing interest

None to be declared for all the authors of this study.

References

1. Gupta R, Gurm HS, Bhatt DL, et al. Renal failure after percutaneous coronary intervention is associated with high mortality. *Cathet Cardiovasc Interv.* 2005;64(4):442–448.
2. Mariani J, et al. Intravascular ultrasound guidance to minimize the use of iodine contrast in percutaneous coronary intervention the MOZART (minimizing contrast utilization with US guidance in coronary angioplasty) randomized controlled trial. *JACC Cardiovasc Interv.* 2014;7:1287–1293.
3. Sakai K, et al. Impact of intravascular ultrasound-guided minimum-contrast coronary intervention on 1-year clinical outcomes in patients with stage 4 or 5 advanced chronic kidney disease. *Cardiovasc Inter Ther.* 2018;34(3):234–241.
4. Sacha J, Gierlotka M, Feusette P, et al. Ultra-low contrast coronary angiography and zero-contrast percutaneous coronary intervention for prevention of contrast-induced nephropathy: step-by-step approach and review. *Postepy Kardiol Interwencyjne.* 2019;15(2):127–136.
5. Kumar P, Jino B, Shafeeq A, et al. IVUS-guided zero-contrast PCI in CKD patients: safety and short-term outcome in patients with: complex demographics and/or lesion characteristics. *Hindawi Intervention Cardiol.* 2021;4(1–7).
6. Nozue T, Michishita I, Mizuguchi I. Impact of catheter down-sizing and power injector use on the amount of contrast medium delivered. *Cardiovasc Interv Ther.* 2010;25:24–28.
7. Ali ZA, Galougahi KK, Nazif T, et al. Imaging- and physiology guided percutaneous coronary intervention without contrast administration in advanced renal failure: a feasibility, safety and outcome study. *Eur Heart J.* 2016;37(40):3090–3095.
8. Kim JK, Kim NH, Shin IS, et al. Alteration of ventricular repolarization by intracoronary infusion of normal saline in patients with variant angina. *Korean Circ J.* 2009;39:223–227.
9. Almdarez M, Gurm HS, Mariani Jr J, et al. Procedural strategies to reduce the incidence of contrast-induced acute kidney injury during percutaneous coronary intervention. *JACC Cardiovasc Interv.* 2019 Oct 14;12(19):1877–1888.
10. Azzalini L, Laricchia A, Regazzoli D, et al. Ultra-low contrast percutaneous coronary intervention to minimize the risk for contrast-induced acute kidney injury in patients with severe chronic kidney disease. *J Invasive Cardiol.* 2019 Jun;31(6):176–182.
11. Karimi Galougahi K, Mintz GS, Karpaliotis D, Ali ZA. Zero-contrast percutaneous coronary intervention on calcified lesions facilitated by rotational atherectomy. *Cathet Cardiovasc Interv.* 2017;90:E85–E89.
12. Koide M, et al. Successful complete revascularization with PCI using super-low volume of contrast medium in a patient with three vessel disease including 2 chronic total occlusions with severe renal dysfunction. *Int Heart J.* 2017;58:624–628.