



**POSTER PRESENTATION**

**Open Access**

# Usefulness of semi-quantification of ischemic myocardium after adenosine stress magnetic resonance

Lorenzo Monti<sup>1,2\*</sup>, Marco Tramarin<sup>2</sup>, Margherita Calcagnino<sup>1</sup>, Veronica Lisignoli<sup>1</sup>, Barbara Nardi<sup>1</sup>, Luca Balzarini<sup>2</sup>

From 16th Annual SCMR Scientific Sessions  
San Francisco, CA, USA. 31 January - 3 February 2013

## Background

Adenosine stress MR shows high sensitivity and intermediate-to-high specificity; therefore, there is a number of false positive patients that are referred to useless coronary angiography studies after a positive adenosine stress MR. We sought to verify whether the (semi) quantification of the amount of ischemic myocardium can improve the management of patients after a positive adenosine stress MR study. According to previous nuclear medicine evidence, we defined a 10% LV mass cut-off for the diagnosis of clinically relevant ischemia.

## Methods

80 MR studies were classified as positive or negative according to the presence or absence of a reversible perfusion defect ("old approach"). We re-analyzed the perfusion studies dividing each one of the 16 AHA segment of the 3 perfusion slices in 2 similar sub-segments (subepicardial and subendocardial); each one of the resulting 32 segments represent about 3% of the myocardial mass. We re-defined ("new approach") as ischemic the patients with > 3 subsegments (>10% of LV mass) with reversible lesions, and as non-ischemic the patients with 3 or less positive sub-segments. Patients were classified as affected by clinically relevant ischemia if the post-adenosine MR coronary angiography confirmed coronary lesions > 50%, and non-affected if the angio was negative and/or no MACE were observed after a > 6 months follow-up.

## Results

Mean age 60.7 years old. Mean follow-up = 18 months (5-37). Pre-test prevalence of CHD: 64%. Prevalence of

clinically relevant ischemia = 17.5%. With the old approach we obtained a sensitivity and specificity of 92.9% and 72.7 respectively. With the new approach we obtained a sensitivity and specificity of 85.7% and 90.9% respectively. Global accuracy of the stress MR exam increased from 76.2 to 90%.

## Conclusions

A visual semi-quantification of the ischemic myocardium improves patient management after a positive adenosine stress MR study.

## Funding

None.

## Author details

<sup>1</sup>Cardiology, I.R.C.C.S. Istituto Clinico Humanitas, Rozzano(MI), Italy.

<sup>2</sup>Radiology, I.R.C.C.S. Istituto Clinico Humanitas, Rozzano (MI), Italy.

Published: 30 January 2013

doi:10.1186/1532-429X-15-S1-E58

Cite this article as: Monti et al.: Usefulness of semi-quantification of ischemic myocardium after adenosine stress magnetic resonance. *Journal of Cardiovascular Magnetic Resonance* 2013 15(Suppl 1):E58.

<sup>1</sup>Cardiology, I.R.C.C.S. Istituto Clinico Humanitas, Rozzano(MI), Italy  
Full list of author information is available at the end of the article