

The complete chloroplast genome of *Epimedium enshiense* B. L. Guo et Hsiao (Berberidaceae)

Xiaoming Li^{a,b,*}, Tianrong Xu^{b,c,*}, Yu Yao^b, Qianru Yang^b, Chaoqun Xu^b, Fengmei Suo^b, Cheng Zhang^b, Guoan Shen^b, Baolin Guo^b, Xiang Liu^d and Shenghua Wei^a

^aSchool of Pharmacy, Guizhou University of Traditional Chinese Medicine, Guiyang, China; ^bInstitute of Medicinal Plant Development, Chinese Academy of Medical Science, Peking Union Medical College, Beijing, China; ^cCollege of Pharmacy, Tianjin University of Traditional Chinese Medicine, Tianjin, China; ^dChongqing Key Laboratory of Traditional Chinese Medicine Resource, Chongqing Academy of Chinese Materia Medica, Chongqing, China

ABSTRACT

The genus of *Epimedium* belongs to Berberidaceae family, which is famous for their medicinal and ornamental value. In recent years, *Epimedium* has attracted increasing attention due to their medicinal and nutritive value. In this research, we reported the complete chloroplast (cp) genome of *Epimedium enshiense*. The complete chloroplast of this species is 157,076 bp in length, including a pair of inverted repeat regions (IRS) (25,833 bp) that is divided by a large single copy area (LSC) (88,340 bp) and a small single copy area (SSC) (17,070 bp). The circular chloroplast genome of *E. enshiense* contains 112 unique genes, composing of 78 protein-coding genes, 30 tRNA, and four rRNA genes. Phylogenetic analysis indicates that *E. enshiense* has a closer relationship with *E. dolichostmon*.

ARTICLE HISTORY

Received 28 January 2020
Accepted 11 February 2020

KEYWORDS

Chloroplast genome;
Epimedium enshiense;
Berberidaceae




Epimedium is a genus of herbaceous plants in the Berberidaceae family, which was used as traditional medicinal plants for more than 2000 years in China (Zhao et al. 2012). *Epimedium enshiense* B. L. Guo et Hsiao is a rare species with yellow spider-like flowers, which only inhabits in the Enshi County, Hubei Province, China (Guo and Xiao 1993). In this study, we reported the complete chloroplast sequence of *E. enshiense*, and analyzed the relationship between *E. enshiense* and other *Epimedium* species by phylogenetic analysis.

The sample of *E. enshiense* was collected from the Enshi County, Hubei province, China (E108°42', N28°26'). The voucher specimen (Guo0464) was deposited at the Herbarium of the Institute of Medicinal Plant (IMPLAD), Beijing, China. Total genomic DNA was extracted from the fresh leaves of *E. enshiense* with the modified CTAB method (Doyle and Doyle 1987). DNA library was sequenced, and 150 bp paired-end reads were generated on an Illumina Novaseq PE150 platform. The clean reads were assembled by using the program GetOrganelle v1.5 (Jin et al. 2018) with the reference chloroplast genome of *E. acuminatum* (GenBank accession number: KU522469). The chloroplast genome annotation was conducted through the online program CPGAVAS2 (Shi et al. 2019) and GeSeq (Tillich et al. 2017). The annotated chloroplast genomic sequence has been registered in GenBank with an accession number (MN937557).

The complete chloroplast genome of *E. enshiense* is 157,076 bp in length, and has a typical quadripartite

construction, which contains two inverted repeat regions (IRa and IRb) of 25,833 bp that is insulated by a large single-copy (LSC, 88,340 bp) and a small single-copy (SSC, 17,070 bp). The total GC content of complete chloroplast genome, LSC, SSC, IR regions is 38.81%, 37.43%, 32.74% and 43.19%, respectively. The complete chloroplast genome of *E. enshiense* contains 112 unique genes, including 78 protein-coding genes, 30 tRNA genes and four rRNA genes. Most of these genes are single-copy genes. However, four protein-coding genes (*rps7*, *ndhB*, *ycf2*, and *rp123*), seven tRNAs (*trnI-CAU*, *trnL-CAA*, *trnV-GAC*, *trnI-GAU*, *trnA-UGC*, *trnR-ACG*, and *trnN-GUU*), and four rRNAs (*rrn16*, *rrn23*, *rrn4.5*, and *rrn5*) are duplicated in the IR regions. One tRNA gene (*trnQ-UUG*) repeated in the LSC regions. In these genes, 15 genes (six tRNA genes and nine protein-coding genes) contain one intron, whereas three genes (*ycf3*, *clpP*, and *rps12*) contain double introns. The *rps12* gene is trans-spliced, with the 5' end located in the LSC region, and the 3' end duplicated in the IR region.

To confirm the phylogenetic position of *E. enshiense*, the complete chloroplast genomes of 19 other plant species were downloaded from the NCBI GenBank database. The sequences were aligned using MAFFT v7 (Katoh et al. 2017), and then the maximum likelihood tree (Figure 1) was constructed using raxmlGUI1.5b (v8.2.10) (Silvestro and Michalak 2012). Phylogenetic analysis shows that *E. enshiense* is closely related to *E. dolichostemon*. The published *E. enshiense*

CONTACT Xiang Liu  zysliux@163.com  Chongqing Academy of Chinese Materia Medica, No. 34 Nanshan Road, Nanan District, Chongqing, 400065, China; Shenghua Wei  weishenghua6512@126.com  School of Pharmacy, Guizhou University of Traditional Chinese Medicine, Dongqing South Road, Anxin District, Guiyang, 550025, China

*These authors contributed equally to the work.

© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. The Maximum likelihood (ML) phylogenetic tree based on complete chloroplast genomes of 20 species, with *Papaver rhoeas* and *Papaver orientale* as out-group. The numbers above the lines represent ML bootstrap values.

chloroplast genome provides useful information for phylogenetic and evolutionary studies in Berberidaceae.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

This work was supported by the CAMS Innovation Fund for Medical Sciences (CIFMS) [2017-12M-3-013], the National Science Foundation of China [81473302], the Chongqing Science and Technology Bureau [cstc2018jcyjAX0316, cc-cstc-CA-19-2], and the Traditional Chinese Medicine Science and the Technology Projects of Chongqing Health and Family Planning Commission [ZY201802117].

References

Doyle JJ, Doyle JL. 1987. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem Bull.* 19:11–15.

- Guo B, Xiao P. 1993. A new taxa of the genus *Epimedium*. *J Syst Evol.* 31: 194–196.
- Jin JJ, Yu WB, Yang JB, Song Y, Yi TS, Li DZ. 2018. GetOrganelle: a simple and fast pipeline for de novo assembly of a complete circular chloroplast genome using genome skimming data. *bioRxiv.* 4: 256479.
- Katoh K, Rozewicki J, Yamada KD. 2017. MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. *Brief Bioinform.* 4:1–7.
- Shi L, Chen H, Jiang M, Wang L, Wu X, Huang L, Liu C. 2019. CPGAVAS2, an integrated plastome sequence annotator and analyzer. *Nucleic Acids Res.* 47(W1):W65–W73.
- Silvestro D, Michalak I. 2012. raxmlGUI: a graphical front-end for RAxML. *Org Divers Evol.* 12:335–337.
- Tillich M, Lehwark P, Pellizzer T, Ulbricht-Jones ES, Fischer A, Bock R, Greiner S. 2017. GeSeq versatile and accurate annotation of organelle genomes. *Nucleic Acids Res.* 45(W1):W6–W11.
- Zhao Y, Song H, Fei J, Liang Y, Zhang B, Liu Q, Wang J, Hu P. 2012. The effects of Chinese yam-epimedium mixture on respiratory function and quality of life in patients with chronic obstructive pulmonary disease. *J Tradit Chin Med.* 32(2):203–207.