

Differential Roles of MAPK Kinases MKK3 and MKK6 in Osteoclastogenesis and Bone Loss

David L. Boyle¹, Deepa Hammaker^{1*}, Meghan Edgar¹, Mario M. Zaiss², Stefan Teufel³, Jean Pierre David³, Georg Schett⁴, Gary S. Firestein¹

1 Medicine, University of California San Diego, La Jolla, California, United States of America, **2** Global Health Institute, École Polytechnique Fédérale de Lausanne, Switzerland, **3** Institute of Osteology and Biomechanics, University Medical Center Hamburg-Eppendorf, Hamburg, Germany, **4** Department of Internal Medicine 3, University of Erlangen-Nuremberg, Friedrich Alexander University, Erlangen, Germany

Abstract

Bone mass is maintained by osteoclasts that resorb bone and osteoblasts that promote matrix deposition and mineralization. Bone homeostasis is altered in chronic inflammation as well as in post-menopausal loss of estrogen, which favors osteoclast activity that leads to osteoporosis. The MAPK p38 α is a key regulator of bone loss and p38 inhibitors preserve bone mass by inhibiting osteoclastogenesis. p38 function is regulated by two upstream MAPK kinases, namely MKK3 and MKK6. The goal of this study was to assess the effect of MKK3- or MKK6-deficiency on osteoclastogenesis in vitro and on bone loss in ovariectomy-induced osteoporosis in mice. We demonstrated that MKK3 but not MKK6, regulates osteoclast differentiation from bone marrow cells in vitro. Expression of NFATc1, a master transcription factor in osteoclastogenesis, is decreased in cells lacking MKK3 but not MKK6. Expression of osteoclast-specific genes Cathepsin K, osteoclast-associated receptor and MMP9, was inhibited in MKK3 $-/-$ cells. The effect of MKK-deficiency on ovariectomy-induced bone loss was then evaluated in female WT, MKK3 $-/-$ and MKK6 $-/-$ mice by micro-CT analysis. Bone loss was partially inhibited in MKK3 $-/-$ as well as MKK6 $-/-$ mice, despite normal osteoclastogenesis in MKK6 $-/-$ cells. This correlated with the lower osteoclast numbers in the MKK-deficient ovariectomized mice. These studies suggest that MKK3 and MKK6 differentially regulate bone loss due to estrogen withdrawal. MKK3 directly mediates osteoclastogenesis while MKK6 likely contributes to pro-inflammatory cytokine production that promotes osteoclast formation.

Citation: Boyle DL, Hammaker D, Edgar M, Zaiss MM, Teufel S, et al. (2014) Differential Roles of MAPK Kinases MKK3 and MKK6 in Osteoclastogenesis and Bone Loss. PLoS ONE 9(1): e84818. doi:10.1371/journal.pone.0084818

Editor: Dominique Heymann, Faculté de médecine de Nantes, France

Received: September 10, 2013; **Accepted:** November 21, 2013; **Published:** January 6, 2014

Copyright: © 2014 Boyle et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Supported by the NIH grants AI-070555 and AI-067752. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: dhammaker@ucsd.edu

Introduction

Homeostatic maintenance of bone mass and strength requires the concerted actions of bone-forming osteoblasts and bone-resorbing osteoclasts [1]. Imbalances in bone remodeling that favor osteoclast differentiation, survival and activation occur in chronic inflammatory diseases such as rheumatoid arthritis (RA) as well as in post-menopausal osteoporosis [2,3]. Pro-inflammatory cytokines such as IL-1, TNF, and IL-6 in chronic inflammation or the loss of estrogen in menopause regulate bone resorption by increasing the expression of Receptor Activator of NF- κ B ligand (RANKL) on osteoblasts and stromal cells or by amplifying RANKL-mediated differentiation of monocytes to osteoclasts [4–7]. The interaction of RANKL with its cognate receptor RANK, expressed on osteoclasts, initiates signaling that enhances osteoclast activity, leading to accelerated bone loss and fractures [8].

RANK and RANKL belong to the TNF receptor superfamily [9,10]. The binding of RANKL to RANK recruits the adapter protein TNF receptor associated factor-6 (TRAF6) to the plasma membrane [11]. RANK, RANKL, and TRAF6 are essential for osteoclastogenesis as mice lacking these molecules have profound defects in bone resorption [11]. The RANK/TRAF6 complex activates several pathways including NF- κ B and the MAPKs, JNK

and p38, which induce the expression of NFATc1, considered one of the master transcription factors of osteoclastogenesis [12,13]. NFATc1, in conjunction with the transcription factors AP-1, PU.1 and microphthalmia transcription factor (MITF), is required for the expression of osteoclast specific genes such as tartrate-resistant acid phosphatase (TRAP) and Cathepsin K that promote bone resorption and demineralization [14,15].

The p38 family of MAPKs, most notably p38 α , is an essential regulator of RANKL-mediated osteoclastogenesis, which makes it a potential therapeutic target for osteoporosis [16]. Several studies have shown that p38 inhibitors prevented inflammatory bone destruction in animal models and in some cases, reversed the existing disease [17–19]. MAPK kinases, MKK3 and MKK6, are two upstream enzymes that differentially regulate p38 function [20]. Our previous studies have shown that MKK3 is required for optimal p38 α activation while MKK6 regulates pro-inflammatory cytokine production in response to LPS and TNF [21]. In addition, either MKK3- or MKK6-deficiency reduces arthritis severity and joint destruction in passive serum transfer K/BxN arthritis or collagen-induced arthritis [21–23]. The goal of this study was to assess the role of MKK3 and MKK6 in p38-regulated bone resorption. Therefore, we evaluated the effect of MKK3- and MKK6-deficiency on osteoclastogenesis in vitro and on bone




Figure 1. Effect of MKK-deficiency in osteoclastogenesis in vitro. (A) WT, MKK3^{-/-} and MKK6^{-/-} bone marrow cells were cultured to induce osteoclastogenesis and analyzed by TRAP staining (n=3/group). The number of osteoclasts were counted in 5 pre-defined locations per well and the data are represented as mean ± SEM. Numbers of osteoclasts were reduced in SB-treated WT and MKK3^{-/-} cells compared with WT. Osteoclastogenesis in MKK6^{-/-} group were comparable to WT cells. *p<0.05 (B) Phosphorylation of p38 in MKK-deficient osteoclasts. Western blot analysis showed that P-p38 levels were significantly lower in MKK3^{-/-} but not in MKK6^{-/-} compared with WT group. *p<0.05. doi:10.1371/journal.pone.0084818.g001

preservation in vivo using an ovariectomy model of osteoporosis. These studies show that blocking either MKK3 or MKK6 has beneficial effects on bone preservation by differentially regulating osteoclastogenesis or function.

Materials and Methods

Mice

Female wildtype (WT) C57Bl/6 mice (6–8 weeks old) were purchased from Charles River Laboratories (Wilmington, MA).

Female MKK3^{-/-} and MKK6^{-/-} mice (C57Bl/6 background) were obtained from Dr. Richard Flavell (Yale University, CT).

Ethics Statement

All experimental animal protocols were reviewed and approved by the University of California San Diego Institutional Animal Care and Use Committee. This study was carried out in strict accordance with the recommendations in the Guide for the Care




Figure 2. Bone resorption by MKK3- and MKK6-deficient osteoclasts. WT, MKK3^{-/-} and MKK6^{-/-} bone marrow cells were differentiated into osteoclasts on a calcium matrix and subjected to Von Kossa staining to determine resorption activity. The resorption pit area was calculated in eight areas/well and data are shown as percent resorbed area/total area. Resorption activity was significantly reduced in MKK3^{-/-} but not MKK6^{-/-} osteoclasts compared with WT. n = 3/group, *p < 0.05. doi:10.1371/journal.pone.0084818.g002

and Use of Laboratory Animals of the National Institutes of Health. All surgeries were performed under isoflurane vaporizer, and all efforts were made to minimize suffering. Mice were housed using standard housing conditions and allowed access to food and water ad libitum.

Generation of Osteoclasts

Bone marrow was isolated from WT, MKK3^{-/-}, and MKK6^{-/-} femora and tibiae. The red blood cells were lysed and the remaining cells were counted and seeded in T75 flasks at 1×10^6 cells/ml. The cells were cultured for 24 h in α MEM (1X) containing GlutaMAX (Invitrogen, Carlsbad, CA) supplemented with 10% FBS, 5% penicillin-streptomycin, and 30 ng/ml of recombinant murine M-CSF (Peprotech, Rocky Hill, NJ). After 24 hours, the non-adherent cells were seeded in 24-well plates (1×10^6 cells/ml) with 30 ng/ml of M-CSF and 50 ng/ml of receptor activator of NF- κ B ligand (RANKL) (R&D, Minneapolis, MN). The culture medium was replaced every 72 h and supplemented with M-CSF and RANKL for 5–8 days until multinucleated osteoclasts were observed. In some experiments, cells were cultured for 5 days with 3 μ M SB203580 (p38 inhibitor) or DMSO (EMD Chemicals, Gibbstown, NJ).

Osteoclast Differentiation and Activity

Tartrate-resistant acid phosphatase (TRAP) staining: Differentiated cells were stained using the Acid Phosphatase, Leukocyte (TRAP) kit (Sigma-Aldrich, St. Louis, MO). Osteoclasts were defined as multinucleated (≥ 3 nuclei), TRAP positive cells and

were counted in 5 pre-defined locations in each well. The data are represented as average count in each well.

Bone resorption assay: Bone marrow monocytes (BMM) were seeded at 3×10^5 cells/well for 8 days on an OsteologicTM Bone Cell Culture System (BD BiocoatTM, Bedford, MA) (4). The cell culture system was then subjected to von Kossa staining. Eight defined areas of each well were imaged and the resorption pits were quantified using Image Pro Plus software (Media Cybernetics, Bethesda, MD). The average resorption was calculated per area for each well and the data are expressed as a percentage of resorbed area/total area.

Ovariectomy

At 16 weeks of age, female WT, MKK3^{-/-}, and MKK6^{-/-} mice were subjected to ovariectomy or sham-surgery and allowed to recover for 6 weeks. To confirm ovariectomy, leutinizing hormone (LH) was measured in serum by sandwich ELISA (Ligand assay and analysis core lab at the University of Virginia).

Micro-Computed Tomography (micro-CT) and Bone Histomorphometry

Micro-CT images of the tibiae were acquired as described previously [13]. Briefly, the tibiae were harvested, fixed in 10% neutral buffered formalin for 24 h and stored in 70% ethanol. All quantifications were performed with digital image analysis (OsteoMeasure; Osteometrics, Atlanta, GA). The following parameters were measured: fraction of bone volume of the total




Figure 3. Gene expression in MKK3^{-/-} and MKK6^{-/-} osteoclasts. WT, MKK3^{-/-} and MKK6^{-/-} osteoclast RNA was isolated and subjected to qPCR using gene-specific primer probe sets (n=3/group). Expression of NFATc1, cFos, Cathepsin K, OSCAR and MMP9 was significantly lower in MKK3^{-/-} but not MKK6^{-/-} cells, compared with WT. RANK expression was comparable in WT, MKK3^{-/-} and MKK6^{-/-} groups. *p<0.05. doi:10.1371/journal.pone.0084818.g003

sample volume (bone volume/tissue volume (BV/TV)), trabecular number (Tb.N), trabecular thickness (Tb.Th), and trabecular separation (Tb.Sp).

Histomorphometric analysis was performed on methacrylate-embedded undecalcified plastic sections of bone and were stained using von Kossa’s stain and TRAP stain. To measure bone formation rate (BFR), all mice were injected with calcein green solution (30 mg/kg, Sigma, MO) at 10 days and 2 days prior to harvesting. The tibias were prepared as described above and the measurements were performed using an image analysis system (OsteoMeasure) and the bone formation rate was calculated and the data are shown as $\mu\text{m}^3/\mu\text{m}^2/\text{year}$.

Gene Expression Analysis

Total RNA was isolated from cells and 500 ng RNA was used for first-strand cDNA synthesis (Life Technologies, NY). Quantitative real-time PCR was performed using Taqman gene expression assay primer/probe sets and GeneAmp 7300 system (Applied Biosystems, Foster City, CA) [24]. Ct values were normalized to hypoxanthine-guanine phosphoribosyl transferase 1 (Hprt1) using the $\Delta\Delta\text{Ct}$ method.

Western Blot Analysis

Osteoclasts were lysed using a lysis buffer containing 50 mM HEPES (pH 7.4), 150 mM NaCl, 25 mM MgCl₂, 1 mM EDTA




Figure 4. Ovariectomy-induced bone loss in MKK3^{-/-} and MKK6^{-/-} mice. (A) WT, MKK3^{-/-} and MKK6^{-/-} mice were subjected to ovariectomy and bone loss was calculated using micro-CT analysis (n=5/group). Representative images are shown. Significant bone loss was observed in all three groups. (B) Bone volume and trabecular number were elevated in MKK3^{-/-} and MKK6^{-/-} sham-treated or OVX mice compared with respective WT controls. Trabecular thickness was similar in sham-treated and OVX mice. Trabecular separation was significantly higher in WT and MKK3^{-/-} mice after OVX. MKK3^{-/-} and MKK6^{-/-} mice had high bone mass phenotype compared with WT in sham-treated or OVX groups.

doi:10.1371/journal.pone.0084818.g004

(pH8.0), 10% glycerol, 1% Triton X-100, 20 mM β-glycerophosphate, 10 mM NaF, 1 mM Na₂VO₄, 0.4% cocktail protease inhibitor (Roche, Indianapolis, IN). Protein was quantified using a Micro BCA protein assay kit (Thermo Scientific, Waltham, MA) and separated using a Nupage[®] Bis-Tris 4–12% polyacrylamide gel (Invitrogen) and transferred onto a PVDF membrane. The membrane was probed with antibodies specific for mouse phospho-p38 (Sigma), total p38 (Cell Signaling Technology, Danvers, MA), and Actin (Sigma, MO). Western blots were developed using the Immun-star[™] Western C Kit and images

were analyzed using the Versadoc Imaging System and Quantity One software (Bio-Rad Laboratories, Hercules, CA).

Statistical Analysis

Comparisons between WT, MKK3^{-/-}, and MKK6^{-/-} cells were analyzed by one-way ANOVA and Tukey’s or Kruskal-Wallis multiple comparison tests, unless otherwise stated. Ovariectomy data were analyzed by two-way ANOVA and Tukey’s multiple comparison test. Data were analyzed using GraphPad




Figure 5. Histomorphometric analysis of bone in ovariectomized MKK3^{-/-} and MKK6^{-/-} mice. (A) Tibial bone sections from sham and OVX-treated mice were stained and representative figures are shown (n = 5/group) (B) Histomorphometric analysis showed that bone volume and trabecular number were elevated in MKK6^{-/-} mice compared with WT after sham surgery or OVX. Trabecular thickness was unaltered after OVX in the three groups. TRAP-staining showed lower osteoclast numbers (NOc/B.Pm) after OVX in MKK3^{-/-} and MKK6^{-/-} compared with WT mice. Osteoblast numbers were unaltered after OVX in the three groups. (C) Bone formation rate was evaluated by injecting mice with calcein green solution at 10 days and 2 days prior to harvesting. The tibias were evaluated for bone formation rate (n = 3/group). Bone formation rate was unaltered by MKK-deficiency in sham- or OVX-treated mice compared with WT controls. *p < 0.05. doi:10.1371/journal.pone.0084818.g005

Prism 6.0 and the comparisons were considered statistically significant if *p* < 0.05.

Results

Effect of MKK3- and MKK6-deficiency on Osteoclastogenesis in vitro

To evaluate the role of MKK3 and MKK6 in osteoclast differentiation, we cultured WT and MKK3- and MKK6-

deficient bone marrow cells in the presence of M-CSF and RANKL. WT cells were cultured in the presence or absence of the p38 inhibitor SB203580 (SB). Osteoclastogenesis was significantly decreased in MKK3^{-/-} ($p=0.0012$) and SB-treated WT group ($p<0.0001$) compared with WT control (Figure 1A, $n=3$ mice/group). MKK3^{-/-} osteoclast numbers were also significantly lower than MKK6^{-/-} cells ($p=0.0012$). Surprisingly, osteoclast numbers in MKK6^{-/-} group were comparable to the WT control. We next determined the level of phosphorylation of p38 (P-p38) in cultured WT, MKK3^{-/-} and MKK6^{-/-} osteoclasts by Western blot analysis. MKK3^{-/-} cells had significantly reduced P-p38 levels compared with WT or MKK6^{-/-} (Figure 1B, $p=0.0009$ and $p=0.005$, respectively). These data show that MKK3, but not MKK6, is an important regulator of osteoclast differentiation *in vitro*.

Bone Resorption Activity in MKK3- and MKK6-deficient Osteoclasts

Bone resorption of MKK-deficient osteoclasts was then determined by their ability to demineralize matrix and form resorption pits. Resorption was visualized using Von Kossa staining (Figure 2). MKK3^{-/-} osteoclasts had a significantly lower number of resorptive pits than WT ($p=0.0061$) or MKK6^{-/-} cells ($p=0.027$, Kruskal-Wallis test), suggesting that MKK3 is a key regulator of osteoclast differentiation *in vitro*.

Gene Expression in MKK3^{-/-} and MKK6^{-/-} Osteoclasts

Osteoclast differentiation requires the expression of a number of well-characterized markers such as NFATc1 and Cathepsin K, which regulate osteoclast function. To determine the effect of MKK-deficiency on these markers, we compared gene expression in WT, MKK3^{-/-}, and MKK6^{-/-} osteoclasts by qPCR (Figure 3). Expression of Cathepsin K and osteoclast-associated receptor (OSCAR) was significantly reduced in MKK3^{-/-} osteoclasts compared with WT ($p<0.001$, respectively) or MKK6^{-/-} cells ($p<0.001$, respectively). Similarly, NFATc1 expression was significantly reduced in MKK3^{-/-} cells compared with WT ($p=0.004$) or MKK6^{-/-} osteoclasts ($p=0.009$). Expression of c-fos and MMP9 (gelatinase B) was inhibited only in MKK3^{-/-} cells ($p=0.037$ and $p=0.031$, respectively, t-test) compared with WT. In contrast, RANK expression was not decreased in MKK-deficient cells. These data show that *in vitro* expression of osteoclast-specific markers is inhibited in the absence of MKK3.

Ovariectomy-induced Bone Loss in MKK3- and MKK6-deficient Mice

Loss of estrogen in ovariectomized (OVX) mice is associated with increased bone turnover with bone resorption exceeding formation, leading to a net bone loss. Thus far, our data have shown that MKK3-deficiency decreases osteoclastogenesis and bone resorption *in vitro*, suggesting that a lack of MKK3 might preserve bone *in vivo*. We therefore examined the effect of MKK-deficiency on bone preservation after ovariectomy. The efficiency of the surgery was confirmed by showing increased levels of luteinizing hormone (LH) in post-surgical sera (Figure S1). Trabecular bone density at the proximal tibia was evaluated in OVX or sham control mice using micro-CT (Figure 4A, $n=5$ /group).

Micro-CT analysis showed significantly higher bone volume (BV/TV) and trabecular number (Tb.N.) in sham MKK3^{-/-} mice compared with WT sham control (Figure 4B, BV: $p=0.003$,

Tb.N: $p=0.006$). Surprisingly, sham MKK6^{-/-} mice had a significantly elevated bone volume and Tb.N. compared with sham WT control (BV: $p<0.0001$, Tb.N: $p<0.0001$). Together, these data suggest that both MKK3 and MKK6 play a role in physiological bone homeostasis. While a significant bone loss was observed after OVX in all three genotypes, bone volume and Tb.N were modestly elevated in OVX MKK3^{-/-} and significantly increased in MKK6^{-/-} mice compared to the OVX WT control. These results suggest that MKK6^{-/-} mice were partially protected from OVX-mediated bone loss, maintaining a bone volume similar to sham-treated WT controls. Trabecular thickness was virtually unchanged in OVX or sham control WT and MKK^{-/-} mice. In contrast, trabecular separation (Tb. Sp) was significantly higher after OVX in WT ($p=0.002$) and MKK3^{-/-} ($p=0.005$) but not in MKK6^{-/-}, compared with their respective sham controls. Trabecular separation was significantly lower in OVX MKK3^{-/-} ($p=0.003$) and MKK6^{-/-} ($p<0.0001$) compared with OVX WT mice, indicating a higher bone mass in MKK-deficient mice. Together, the *in vivo* data demonstrate that MKK3- and MKK6-deficiency are associated with a high bone mass phenotype, which is partially maintained after OVX.

Histomorphometric Analysis of Bone in Ovariectomized MKK3- and MKK6-deficient Mice

To further evaluate the effect of MKK3- and MKK6-deficiency in estrogen-deficient mice, undecalcified tibial bone sections from OVX and sham control mice were stained and subjected to histomorphometric analysis (Figure 5A, B). Consistent with the micro-CT data, bone volume (%BV/TV) was significantly higher in sham- and OVX-treated MKK3^{-/-} ($p<0.05$ for each) and MKK6^{-/-} mice compared to respective WT controls ($p<0.001$, for each). OVX led to a reduction of bone volume in the WT, MKK3^{-/-} and MKK6^{-/-} groups ($n=5$ /treatment, $p<0.001$, respectively). Nonetheless, bone volume and trabecular number were at least partially maintained in MKK-mutants, indicating a similar bone volume after OVX as sham-treated WT mice. Trabecular thickness (Tb.Th) was similar in WT, MKK3^{-/-}, and MKK6^{-/-} mice, both after sham-treatment and OVX.

TRAP staining showed higher osteoclast numbers (Noc/bone perimeter) in OVX WT mice compared with sham-treated WT mice ($p<0.001$). However, MKK3- and MKK6-deficient mice had significantly lower osteoclast numbers compared with sham-treated or OVX WT mice (MKK3^{-/-}: $p<0.01$, MKK6^{-/-}: $p<0.001$). Interestingly, the number of osteoblasts (NOb) and bone formation rate was not altered in MKK3 and MKK6 mutants. These results suggest that MKK-deficiency inhibits osteoclast formation *in vivo* but does not alter numbers of osteoblasts or bone formation (Figure 5C). Together, these data show that MKK-deficiency induced high bone mass by lowering osteoclast numbers *in vivo*.

Discussion

Osteoclast differentiation and activation is regulated by p38 MAPK, at least in part by mediating the effects of pro-inflammatory cytokines [25,26]. Several studies have shown that p38 inhibitors protect against bone resorption in inflammation, possibly through the direct effects on osteoclasts as well as through inhibition of p38-induced cytokine expression. For instance, p38 α activation is essential for TNF-mediated inflammatory bone loss, and a selective p38 α inhibitor prevented trabecular bone loss and increased cortical bone area in ovariectomized rats [13,27]. Our previous studies demonstrated clinical benefits of MKK3- and

MKK6-deficiency in acute and chronic models of arthritis [21–23]. Inflammation-induced bone loss in the joints was markedly reduced in the absence of MKK3 or MKK6, with a corresponding decrease in serum IL-6 and synovial MMP [22]. In the present study, we extend our observations on bone protection by evaluating the effect of MKK3- and MKK6-deficiency on osteoclastogenesis and bone loss due to estrogen withdrawal in ovariectomized mice.

Our data show that MKK3 and MKK6 differentially regulate osteoclastogenesis. MKK3 is required for optimal osteoclastogenesis and function. MKK6 was unable to effectively compensate for the absence of MKK3 in the osteoclasts. MKK6-deficiency, on the other hand, did not affect osteoclastogenesis *in vitro*. Consistent with our previous studies, P-p38 levels were markedly reduced in MKK3^{-/-} but normal in MKK6^{-/-} osteoclasts [21]. Osteoclast numbers and resorption activity correlated with p38 activation in MKK3^{-/-} and MKK6^{-/-} cells and are in agreement with previous studies [16]. Expression of NFATc1 and c-fos were reduced in MKK3^{-/-} osteoclasts, indicating that p38 function is necessary for the induction of these transcription factors that are involved in osteoclast differentiation [28]. Cathepsin K, OSCAR, and matrix metalloproteinase 9 (MMP9) expression was reduced in MKK3-deficient osteoclasts. This is consistent with the requirement of NFATc1 in the transcription of these osteoclast-specific markers [15,29,30]. Together, these results indicate that MKK3, rather than MKK6, is a principal regulator of RANKL-induced signaling in osteoclastogenesis *in vitro*. These data are in accordance with other studies showing that p38 regulates the early phases of osteoclast differentiation [31,32]. However, those studies suggested that MKK6, but not MKK3, is necessary for osteoclastogenesis *in vitro*, where it enhances cell survival but not bone resorption. The differences might be partly due to differences in the methodology, such as the efficiency of inhibiting MKK3 and MKK6 using viral transduction of dominant-negative MKK3 and MKK6 constructs compared with using bone marrow-derived osteoclasts of MKK3^{-/-} and MKK6^{-/-} mice.

Osteoclastogenesis *in vivo* is regulated by a number of factors such as vitamin D, parathyroid hormone, prostaglandins and most notably, estrogen [33–36]. Estrogen is a key modulator of both the immune system and bone homeostasis [37]. It regulates bone remodeling by decreasing osteoclast life span by promoting apoptosis [38,39]. Estrogen inhibits expression of RANKL on osteoblasts, stromal cells, and lymphocytes, and increases production of osteoprotegerin (OPG), an inhibitor of RANK signaling [1]. In menopause, the loss of estrogen is associated with elevated levels of IL-1, IL-6, TNF, and M-CSF, which support bone resorption [40,41]. Estrogen deficiency also triggers the differentiation of Th17 cells that produce the pro-inflammatory cytokine IL-17 [42]. IL-17 induces osteoblast and stromal cell production of IL-6, and TNF, which induce RANKL expression, accelerate osteoclastogenesis and lead to osteoporosis [43].

References

- Khosla S (2001) Minireview: the OPG/RANKL/RANK system. *Endocrinology* 142: 5050–5055.
- Kimble RB, Srivastava S, Ross FP, Matayoshi A, Pacifici R (1996) Estrogen deficiency increases the ability of stromal cells to support murine osteoclastogenesis via an interleukin-1 and tumor necrosis factor-mediated stimulation of macrophage colony-stimulating factor production. *J Biol Chem* 271: 28890–28897.
- Manolagas SC (1995) Role of cytokines in bone resorption. *Bone* 17: 63S–67S.
- Wei S, Kitaura H, Zhou P, Ross FP, Teitelbaum SL (2005) IL-1 mediates TNF-induced osteoclastogenesis. *J Clin Invest* 115: 282–290.
- Kim JH, Jin HM, Kim K, Song I, Youn BU, et al. (2009) The mechanism of osteoclast differentiation induced by IL-1. *J Immunol* 183: 1862–1870.
- Faienza MF, Ventura A, Marzano F, Cavallo L (2013) Postmenopausal osteoporosis: the role of immune system cells. *Clin Dev Immunol* 2013: 575936.
- David JP, Schett G (2010) TNF and bone. *Curr Dir Autoimmun* 11: 135–144.
- Feng X (2005) RANKing intracellular signaling in osteoclasts. *IUBMB Life* 57: 389–395.
- Raggatt LJ, Partridge NC (2010) Cellular and molecular mechanisms of bone remodeling. *J Biol Chem* 285: 25103–25108.
- Boyle WJ, Simonet WS, Lacey DL (2003) Osteoclast differentiation and activation. *Nature* 423: 337–342.
- Asagiri M, Takayanagi H (2007) The molecular understanding of osteoclast differentiation. *Bone* 40: 251–264.

In this study, we showed that both MKK3- and MKK6-deficiency is associated with high bone mass and at least partially inhibited bone loss after ovariectomy, even though our *in vitro* studies did not show major effects of MKK6 on *in vitro* osteoclastogenesis. Histomorphometric data demonstrated that high bone mass in MKK3 and MKK6-deficiency correlates with reduced osteoclast numbers. The decrease in osteoclast numbers in MKK6-deficient mice could be due to decreased p38-regulated cytokine production, such as IL-6, since a direct effect on osteoclasts was not observed. The bone protection is consistent with our previous studies showing reduced inflammation and joint destruction in MKK3^{-/-} and MKK6^{-/-} mice, where a direct effect on osteoclasts for MKK3 and an indirect effect by blocking cytokines for MKK6 were found [22].

The study also found no major effect on osteoblast number and bone formation rate in MKK3- and MKK6-deficient mice. The regulation of bone homeostasis and skeletal formation was recently evaluated in MKK3- and MKK6-deficient mice [44]. That study showed that both MKK3^{-/-} and MKK6^{-/-} mice had reduced bone mass resulting from defective osteoblast differentiation. The reason for the discrepancy is unclear but might be partly due to strain variations in mice and the different ages of the mice used, 5 weeks compared with 20 weeks for our study.

In conclusion, MKK3 and MKK6 affect bone mass through regulation of osteoclast numbers *in vivo*. MKK6 and to a lesser extent MKK3 mitigate bone loss in conjunction with estrogen deficiency. While MKK3 likely mediates direct effects on osteoclasts through RANKL signaling, MKK6 likely contributes to the production of pro-inflammatory cytokines such as IL-6 and IL-17 that increase RANKL expression. Targeting upstream p38 kinases, such as MKK3 and MKK6 might modulate bone loss during inflammation and estrogen deficiency.

Supporting Information

Figure S1 Serum leutinizing hormone (LH) in ovariectomized mice. The efficiency of the ovariectomy was evaluated by measuring LH in sera. LH levels were significantly elevated in ovariectomized WT, MKK3^{-/-}, and MKK6^{-/-} sera compared with sham-treated sera (n = 8/group, *p<0.05). (TIF)

Acknowledgments

The authors thank Dr. Pamela Mellon and Dr. Christine Glidewell-Kenney for advice on ovariectomy surgeries and on the LH confirmations.

Author Contributions

Conceived and designed the experiments: DLB GS GSF. Performed the experiments: ME MMZ ST. Analyzed the data: DLB DH ME JD GS GSF. Contributed reagents/materials/analysis tools: MMZ ST JD GS. Wrote the paper: DH DLB JD GS GSF.

12. Strait K, Li Y, Dillehay DL, Weitzmann MN (2008) Suppression of NF-kappaB activation blocks osteoclastic bone resorption during estrogen deficiency. *Int J Mol Med* 21: 521–525.
13. Zverina J, Hayer S, Redlich K, Bobacz K, Kollias G, et al. (2006) Activation of p38 MAPK is a key step in tumor necrosis factor-mediated inflammatory bone destruction. *Arthritis Rheum* 54: 463–472.
14. Sharma SM, Bronisz A, Hu R, Patel K, Mansky KC, et al. (2007) MITF and PU.1 recruit p38 MAPK and NFATc1 to target genes during osteoclast differentiation. *J Biol Chem* 282: 15921–15929.
15. Troen BR (2006) The regulation of cathepsin K gene expression. *Ann N Y Acad Sci* 1068: 165–172.
16. Bohm C, Hayer S, Kilian A, Zaiss MM, Finger S, et al. (2009) The alpha-isoform of p38 MAPK specifically regulates arthritic bone loss. *J Immunol* 183: 5938–5947.
17. Hill RJ, Dabbagh K, Phippard D, Li C, Suttman RT, et al. (2008) Pamipomod, a novel p38 mitogen-activated protein kinase inhibitor: preclinical analysis of efficacy and selectivity. *J Pharmacol Exp Ther* 327: 610–619.
18. Caverzasio J, Higgins L, Ammann P (2008) Prevention of trabecular bone loss induced by estrogen deficiency by a selective $\alpha\alpha$ inhibitor. *J Bone Miner Res* 23: 1389–1397.
19. Medicherla S, Ma JY, Mangadu R, Jiang Y, Zhao JJ, et al. (2006) A selective p38 alpha mitogen-activated protein kinase inhibitor reverses cartilage and bone destruction in mice with collagen-induced arthritis. *J Pharmacol Exp Ther* 318: 132–141.
20. Inoue T, Hammaker D, Boyle DL, Firestein GS (2005) Regulation of p38 MAPK by MAPK kinases 3 and 6 in fibroblast-like synoviocytes. *J Immunol* 174: 4301–4306.
21. Yoshizawa T, Hammaker D, Boyle DL, Corr M, Flavell R, et al. (2009) Role of MAPK kinase 6 in arthritis: distinct mechanism of action in inflammation and cytokine expression. *J Immunol* 183: 1360–1367.
22. Hammaker D, Topolewski K, Edgar M, Yoshizawa T, Fukushima A, et al. (2012) Decreased collagen-induced arthritis severity and adaptive immunity in MKK-6-deficient mice. *Arthritis Rheum* 64: 678–687.
23. Inoue T, Boyle DL, Corr M, Hammaker D, Davis RJ, et al. (2006) Mitogen-activated protein kinase kinase 3 is a pivotal pathway regulating p38 activation in inflammatory arthritis. *Proc Natl Acad Sci U S A* 103: 5484–5489.
24. Boyle DL, Rosengren S, Bugbee W, Kavanaugh A, Firestein GS (2003) Quantitative biomarker analysis of synovial gene expression by real-time PCR. *Arthritis Res Ther* 5: R352–360.
25. Matsumoto M, Sudo T, Saito T, Osada H, Tsujimoto M (2000) Involvement of p38 mitogen-activated protein kinase signaling pathway in osteoclastogenesis mediated by receptor activator of NF-kappa B ligand (RANKL). *J Biol Chem* 275: 31155–31161.
26. Mbalaviele G, Anderson G, Jones A, De Cicchi P, Settle S, et al. (2006) Inhibition of p38 mitogen-activated protein kinase prevents inflammatory bone destruction. *J Pharmacol Exp Ther* 317: 1044–1053.
27. Matsumoto M, Sudo T, Maruyama M, Osada H, Tsujimoto M (2000) Activation of p38 mitogen-activated protein kinase is crucial in osteoclastogenesis induced by tumor necrosis factor. *Febs Lett* 486: 23–28.
28. Huang H, Chang EJ, Ryu J, Lee ZH, Lee Y, et al. (2006) Induction of c-Fos and NFATc1 during RANKL-stimulated osteoclast differentiation is mediated by the p38 signaling pathway. *Biochem Biophys Res Commun* 351: 99–105.
29. So H, Rho J, Jeong D, Park R, Fisher DE, et al. (2003) Microphthalmia transcription factor and PU.1 synergistically induce the leukocyte receptor osteoclast-associated receptor gene expression. *J Biol Chem* 278: 24209–24216.
30. Sundaram K, Nishimura R, Sem J, Youssef RF, London SD, et al. (2007) RANK ligand signaling modulates the matrix metalloproteinase-9 gene expression during osteoclast differentiation. *Exp Cell Res* 313: 168–178.
31. Yamashita T, Kobayashi Y, Mizoguchi T, Yamaki M, Miura T, et al. (2008) MKK6-p38 MAPK signaling pathway enhances survival but not bone-resorbing activity of osteoclasts. *Biochem Biophys Res Commun* 365: 252–257.
32. Huang H, Ryu J, Ha J, Chang EJ, Kim HJ, et al. (2006) Osteoclast differentiation requires TAK1 and MKK6 for NFATc1 induction and NF-kappaB transactivation by RANKL. *Cell Death Differ* 13: 1879–1891.
33. Khosla S, Oursler MJ, Monroe DG (2012) Estrogen and the skeleton. *Trends Endocrinol Metab* 23: 576–581.
34. Armas LA, Recker RR (2012) Pathophysiology of osteoporosis: new mechanistic insights. *Endocrinol Metab Clin North Am* 41: 475–486.
35. Shibata T, Shira-Ishi A, Sato T, Masaki T, Masuda A, et al. (2002) Vitamin D hormone inhibits osteoclastogenesis in vivo by decreasing the pool of osteoclast precursors in bone marrow. *J Bone Miner Res* 17: 622–629.
36. Bolon B, Carter C, Daris M, Morony S, Capparelli C, et al. (2001) Adenoviral delivery of osteoprotegerin ameliorates bone resorption in a mouse ovariectomy model of osteoporosis. *Mol Ther* 3: 197–205.
37. Krum SA, Brown M (2008) Unraveling estrogen action in osteoporosis. *Cell Cycle* 7: 1348–1352.
38. Krum SA, Miranda-Carboni GA, Hauschka PV, Carroll JS, Lane TF, et al. (2008) Estrogen protects bone by inducing Fas ligand in osteoblasts to regulate osteoclast survival. *EMBO J* 27: 535–545.
39. Nakamura T, Imai Y, Matsumoto T, Sato S, Takeuchi K, et al. (2007) Estrogen prevents bone loss via estrogen receptor alpha and induction of Fas ligand in osteoclasts. *Cell* 130: 811–823.
40. Passeri G, Girasole G, Jilka RL, Manolagas SC (1993) Increased interleukin-6 production by murine bone marrow and bone cells after estrogen withdrawal. *Endocrinology* 133: 822–828.
41. Srivastava S, Weitzmann MN, Kimble RB, Rizzo M, Zahner M, et al. (1998) Estrogen blocks M-CSF gene expression and osteoclast formation by regulating phosphorylation of Egr-1 and its interaction with Sp-1. *J Clin Invest* 102: 1850–1859.
42. Tyagi AM, Srivastava K, Mansoori MN, Trivedi R, Chattopadhyay N, et al. (2012) Estrogen deficiency induces the differentiation of IL-17 secreting Th17 cells: a new candidate in the pathogenesis of osteoporosis. *PLoS One* 7: e44552.
43. Tokuda H, Kanno Y, Ishisaki A, Takenaka M, Harada A, et al. (2004) Interleukin (IL)-17 enhances tumor necrosis factor-alpha-stimulated IL-6 synthesis via p38 mitogen-activated protein kinase in osteoblasts. *J Cell Biochem* 91: 1053–1061.
44. Greenblatt MB, Shim JH, Zou W, Sitara D, Schweitzer M, et al. (2010) The p38 MAPK pathway is essential for skeletogenesis and bone homeostasis in mice. *J Clin Invest* 120: 2457–2473.