

Trends of Low Back Pain Research in Older and Working-Age Adults from 1993 to 2023: A Bibliometric Analysis

Daniel KY Zheng¹, Gregory N Kawchuk², André E Bussi eres^{3,4}, Fadi M Al Zoubi¹, Jan Hartvigsen⁵, Siu Ngor Fu¹, Katie de Luca⁶, Debra K Weiner^{7,8}, Jaro Karppinen^{9,10}, Dino Samartzis¹¹, Manuela L Ferreira¹², Jinlong Wu¹³, Liz Dennett¹², Arnold YL Wong¹

¹Hong Kong Polytechnic University, Hung Hom, Hong Kong SAR, People's Republic of China; ²University of Alberta, Edmonton, Canada; ³Universit e McGill, Montreal, Canada; ⁴Universit e du Qu ebec   Trois-Rivi eres, Trois-Rivi eres, Canada; ⁵University of Southern Denmark, Odense, Denmark; ⁶Central Queensland University, Brisbane, Australia; ⁷University of Pittsburgh, Pittsburgh, PA, USA; ⁸VA Pittsburgh Healthcare System, Pittsburgh, PA, USA; ⁹Research Unit of Health Sciences and Technology, University of Oulu, Oulu, Finland; ¹⁰Rehabilitation Services of South Karelia Social and Health Care District, Lappeenranta, Finland; ¹¹Rush University Medical Center, Chicago, IL, USA; ¹²The University of Sydney, Sydney, Australia; ¹³Southwest University, Chongqing, People's Republic of China

Correspondence: Arnold YL Wong, Department of Rehabilitation Sciences, The Hong Kong Polytechnic University, Hung Hom, Hong Kong SAR, People's Republic of China, Tel +852 2766 6741, Email arnold.wong@polyu.edu.hk

Abstract: Although the number of publications focusing on low back pain in older adults (LBP-O) and working-age adults (LBP-W) has been growing for decades, comparative research trends in these two populations, which may help to guide future investigation, have not been rigorously explored. This analysis aimed to describe publication patterns and trends of research targeting LBP-O and LBP-W over the last three decades. Peer-reviewed LBP-O and LBP-W articles published between 1993 and 2023 were retrieved from the Web of Science, which provided the details of annual publication volume, and prominent journals/countries/institutions. The relationship between the annual publication volumes and years was analyzed by Spearman correlation analysis. The hot topics and emerging trends were analyzed by VOSviewer and CiteSpace, respectively. A total of 4217 LBP-O-related and 50,559 LBP-W-related documents were included. The annual publication volumes of LBP-O and LBP-W articles increased over the years ($r=0.995$ to 0.998 , $p<0.001$). The United States had the highest number of prominent institutions publishing relevant articles. The most prolific journal for LBP-O (5.4%) and LBP-W-related (6.1%) papers is the journal "Spine". Cognitive behavioral therapy, intervertebral disc (IVD) degeneration, physiotherapy, physical activity, and walking were the recent hot topics and physical activity was an emerging trend in LBP-O, while surgery and IVD degeneration (also a hot topic) were emerging trends in LBP-W. This study highlights the paucity of LBP-O-related research in the past. The United States and the journal *Spine* stand out in LBP research. The research trend of physical activity in LBP-O is consistent with the recognized importance of physical activity for older adults in general, and for managing LBP-O in particular. Conversely, the emerging trends of surgery and intervertebral disc degeneration in LBP-W research highlight a focus on the biomedical model of LBP despite LBP being a biopsychosocial condition.

Keywords: low back pain, older adults, working-age adults, bibliometrics, intervertebral disc degeneration, physical activity, surgery

Introduction

Low back pain (LBP) is one of the most common biopsychosocial problems in high- and low-income countries.^{1,2} It is the main cause of years lived with disability at all ages.³ The reported prevalence of LBP in working-age adults (LBP-W) aged 18–64 years is between 4.2% and 19.6%, whereas that in older adults (LBP-O) aged ≥ 65 years is up to 36.1%, although this varies across countries, cultures, and races.^{4–7} While LBP-W is the fourth largest productivity burden for the United States, LBP-O is the third leading cause of chronic disability in Canada.^{3,8}

Given the high prevalence and negative impacts of LBP, numerous LBP-related clinical studies have been conducted. However, recent systematic reviews reveal a lack of research specifically on LBP-O. Paeck et al³ and Levy et al⁹ found

that up to 53% of randomized clinical trials excluded adults aged 65 years and older. Many clinicians assume that findings from LBP-W studies can be generalized to LBP-O.³ Because older adults have a greater burden of physical and cognitive comorbidities than working-age adults, they may respond differently to medications, surgery, exercise, and other forms of interventions, thus an effective treatment for LBP-W may not be equally effective for LBP-O.¹⁰ Therefore, clinical trials should include an adequate number of LBP-O and LBP-W to afford meaningful subgroup analyses, although this rarely has been the case.

Several notable research trends and characteristics have been noted in the LBP-W literature. For example, Modic changes, intervertebral disc (IVD) degeneration, and paraspinal muscles have become common research focuses in LBP-W.^{4,11–13} Further, given the relatively small or no effects of some pain medications (eg, nonsteroidal anti-inflammatory drugs¹⁴ or paracetamol¹⁵), recent research has been conducted to quantify the effects of using exercises to treat LBP-W.^{16,17} However, whether similar research trends occur in LBP-O remains unclear. Some potential old age-related topics, such as physical inactivity, and cognitive decline have seldom been highlighted.^{18–20}

Bibliometric analysis is a quantitative and pragmatic approach to visualize the publication patterns and trends in a given field. It can identify the leading journals, countries, institutions, and research foci.^{21–23} CiteSpace and VOSviewer are freely available tools for bibliometric analysis. Web of Science (WoS), a multidisciplinary platform encompassing different categories of scientific knowledge,^{24,25} is considered as the preferred bibliographic source for bibliometric analysis.^{22,26} Huang et al²⁵ and Guo et al²⁷ also conducted bibliometric analyses using records from WoS to analyze the research trends of LBP from 2000 to 2022, and the evolution of LBP-W-related publications between 2000 and 2020, respectively. However, they were limited by no direct comparisons of publication trends between LBP-O and LBP-W research, which may inform knowledge gaps in both fields.

Against this background, comprehensive investigations of research trends in LBP-O and LBP-W fields are warranted to summarize previous research foci, as well as to inform future research priorities and allocations of research resources to key LBP-related research areas. Importantly, the current bibliometric analysis can foster collaborations among researchers and provide information regarding relevant LBP-related journals. This study aimed to: (1) reveal the publication characteristics of LBP-O and LBP-W-related articles since 1993; (2) compare the research topics; and (3) identify potential emerging trends.

Materials and Methods

Literature Search and Retrieval

Two researchers (KZ and AW) and two experienced research librarians (LD and JB) developed the search strategy ([Supplementary Material 1](#)). The search was conducted in the Web of Science (WoS) Core Collection ([Figure 1](#)). The keywords and medical subject headings included LBP, working-age adults, and older adults. As CiteSpace only allows bibliometric analysis over a 31-year period, the search was conducted from January 1, 1993, to May 14, 2023. Using the “Article and Review” filter,²² relevant papers were identified for inclusion because they were peer-reviewed and could be considered “certified knowledge”, enhancing the results credibility. The full record and cited references of the search results were downloaded in a plain text format. LBP-O and LBP-W papers were searched and analyzed separately.

Data Processing

The relevant information of the included articles (eg, journals, countries, institutions, and keywords) and their statistics (eg, number of included papers, total number of citations, average citations per paper, journal impact factor (IF)) were extracted by an independent researcher (KZ) from WoS. This was followed by a verification process involving the corresponding author (AW) and obtaining the agreement from all co-authors. Based on the number of included papers, the most prolific journals, countries, and institutions were identified.²⁸ For the comparisons of countries and institutions, all authors in the included publications were taken into account. The annual publication volumes were plotted against time using an Excel program. SPSS version 23.0 software (SPSS Inc., Chicago, IL, USA) was used to measure the strength and direction of the association between the annual publication volumes and years by Spearman correlation analysis with correlation coefficient (r) (where 0.8–1.0, 0.6–0.8, 0.4–0.6, 0.2–0.4, and 0.0–0.2 mean very strong, strong,




Figure 1 Literature retrieval flow chart.

Abbreviations: LBP-O, low back pain in older adults; LBP-W, low back pain in working-age adults.

moderate, weak, and no to very weak correlation, respectively).²⁹ The annual publication trends for each population were evaluated and fitted a variation curve using local polynomial regression (locally estimated scatterplot smoothing, LOESS) with R (version 4.3.0).²⁸ The number of citations may indicate the impact of a given paper, while the average citations per paper measures the average impact of its authors.³⁰ Higher citation counts implied that an author's scientific works were more acknowledged by his/her peers, although outcome measure-related papers or reviews are more likely to be cited. Additionally, the journal's IF was calculated based on the latest edition of the Journal Citation Reports. IF was regarded as an indicator of the academic influence of a scientific journal. The analysis of journals, countries and institutions can help researchers to select journals for future submissions and promote international collaboration.

The search results were further analyzed by VOSviewer (version 1.6.19, Universiteit Leiden) and CiteSpace (version 6.2.2, Drexel University). Specifically, VOSviewer was used to count the frequency of keywords and to determine the average publication year in which a keyword or a term occurs or the average publication year of the documents published by a source (eg, a country). CiteSpace was applied to identify keywords with co-occurrence burst, which implies an explosive increase of co-occurrence frequency of keywords in a certain period of time.³¹ Co-occurrence burst has two characteristics, namely strength and duration, which indicate that researchers pay more attention to a specific research work or topic over a particular time period. Compared to keywords with the latest average publication year, latest burst keywords have stronger timeliness. The keywords with the strongest co-occurrence burst lasting until 2023 represent potential emerging trends.²⁸

Results

The search resulted in 4217 LBP-O-related and 50,559 LBP-W-related articles for the bibliometric analysis.

Chronological Changes in Publications

The chronological distributions of papers are plotted in [Figure 2](#), where a geometric growth could be clearly observed for both LBP-O and LBP-W. The cumulative number of LBP-O-related papers was 4217, which was only 8.3% of that of LBP-W-related papers. The annual publication volume of LBP-O-related papers started to grow rapidly in the past decade and reached its highest volume (383 papers) in 2022, averaging 136 papers per year. Likewise, the annual number of LBP-W-related articles was 290 in 1993, and it increased by around 14 times (4105 papers) in 2021, averaging 1631 papers per year. There were very strong correlations between years and annual publication volume in both fields from 1993 to 2022 (r of LBP-O and LBP-W were 0.995 and 0.998, respectively).




Figure 2 Yearly publication trends. **(A)** The yearly publication trend regarding low back pain in older adults; **(B)** The yearly publication trends concerning low back pain in working-age adults.

Notes: The gray area represents the 95% confidence interval. For the lack of full-year data and a better fitted curve, data for 2023 (107 for low back pain in older adults and 980 for low back pain in working-age adults) are not shown in this figure.

Comparisons of Journals

A total of 996 journals published LBP-O-related papers, whereas 3571 journals published LBP-W-related papers. Of the top 10 prolific journals (Table 1), *Spine* published the greatest volume of LBP-O and LBP-W-related papers, accounting for 5.4% (228 publications) and 6.1% (3066 publications), respectively. Ranked Quartile 1 in the Anesthesiology, Clinical Neurology, and Neurosciences categories, *Pain* was the journal with the highest IF (7.4) in the top 10 list of LBP-O research, followed by *The Spine Journal* (IF = 4.5) and *PLOS One* (IF = 3.7). The IFs of the top 10 prolific journals for LBP-W research were very similar, ranging from 1.3 to 7.4.

Comparisons Across Countries

Figure 3 shows that a total of 90 and 157 countries contributed to LBP-O- and LBP-W-related publications indexed in WoS in the last three decades, respectively. Many countries in Asia and Africa were not involved in the LBP-O research. Compared to LBP-O-related research, LBP-W-related research was conducted more globally, but half of African countries were still not involved. Among the top 10 countries in Table 2, the United States published the highest paper volume, while researchers in the Netherlands had the highest average citations per paper in both fields. Although China and the UK had similar numbers of papers in the second and third place in both fields, respectively, the UK had nearly three times more total citations and average citations than China. The average publication years of British LBP-O- and LBP-W-related studies were 2014 and 2012, respectively, while the average publication year of articles from China in both fields was 2018.

Comparisons of Institutions

Our analysis identified 4213 and 25,058 institutions contributing to LBP-O- and LBP-W-related research, respectively. Of the top 10 productive institutions for LBP-O-related research, University of California was the most active institution

Table 1 Top 10 Prolific Journals from 1993 to 2023

Rank	Low Back Pain in Older Adults					Low Back Pain in Working-Age Adults				
	Journal	Papers (%)	IF 2022 ^a	5-Year IF ^b	WoS Categories (Q) ^c	Journal	Papers (%)	IF 2022 ^a	5-Year IF ^b	WoS Categories (Q) ^c
1	<i>Spine</i>	228 (5.4)	3	3.5	Clinical Neurology (3); Orthopedics (2)	<i>Spine</i>	3066 (6.1)	3	3.5	Clinical Neurology (3); Orthopedics (2)
2	<i>Pain Medicine</i>	125 (3.0)	3.1	3.4	Anesthesiology (2); Medicine, General & Internal (2)	<i>European Spine Journal</i>	1435 (2.8)	2.8	3.2	Clinical Neurology (3); Orthopedics (2)
3	<i>European Spine Journal</i>	105 (2.5)	2.8	3.2	Clinical Neurology (3); Orthopedics (2)	<i>BMC Musculoskeletal Disorders</i>	1016 (2.0)	2.3	2.8	Orthopedics (2); Rheumatology (4)
4	<i>BMC Musculoskeletal Disorders</i>	93 (2.2)	2.3	2.8	Orthopedics (2); Rheumatology (4)	<i>Pain</i>	911 (1.8)	7.4	7.7	Anesthesiology (1); Clinical Neurology (1); Neurosciences (1)
5	<i>Pain</i>	81 (1.9)	7.4	7.7	Anesthesiology (1); Clinical Neurology (1); Neurosciences (1)	<i>Spine Journal</i>	760 (1.5)	4.5	4.6	Clinical Neurology (2); Orthopedics (1)
6	<i>Spine Journal</i>	75 (1.8)	4.5	4.6	Clinical Neurology (2); Orthopedics (1)	<i>Pain Medicine</i>	684 (1.4)	3.1	3.4	Anesthesiology (2); Medicine, General & Internal (2)
7	<i>Clinical Journal of Pain</i>	55 (1.3)	2.9	3.8	Anesthesiology (2); Clinical Neurology (3)	<i>Clinical Journal of Pain</i>	644 (1.3)	2.9	3.8	Anesthesiology (2); Clinical Neurology (3)
8	<i>Osteoporosis International</i>	48 (1.1)	4	4.6	Endocrinology & Metabolism (2)	<i>Journal of Manipulative and Physiological Therapeutics</i>	640 (1.3)	1.3	1.8	Health Care Sciences & Services (4) Integrative & Complementary Medicine (4); Rehabilitation (4)
9	<i>PLOS One</i>	47 (1.1)	3.7	3.8	Multidisciplinary Sciences (2)	<i>Journal of Back and Musculoskeletal Rehabilitation</i>	545 (1.1)	1.6	1.7	Orthopedics (3); Rehabilitation (3)
10	<i>Journal of Manipulative and Physiological Therapeutics</i>	45 (1.1)	1.3	1.8	Health Care Sciences & Services (4) Integrative & Complementary Medicine (4); Rehabilitation (4)	<i>Pain Physician</i>	522 (1.0)	3.7	3.9	Anesthesiology (2); Clinical Neurology (2)

Notes: ^aIF and ^cQ in category according to Journal Citation Reports™ 2022.

Abbreviations: ^aIF, impact factor; ^b5-year, 2018–2022; ^cQ, quartile.




Figure 3 Distribution of countries/territories. **(A)** Distribution of countries/territories based on the accumulative number of publications concerning low back pain in older adults from 1993 to 2023; **(B)** Distribution of countries/territories based on the accumulative number of publications concerning low back pain in working-age adults from 1993 to 2023.

Notes: Papers from Hong Kong, Macau and Taiwan were reclassified to China, although Taiwan was not highlighted in the figure because of flaws in the mapping software; papers from England, Scotland, North Ireland, and Wales were reclassified to the UK.

with 157 papers and approximately 62 citations per paper (Table 3). In the LBP-W-related field, Harvard University was ranked number one with 1248 relevant papers, while the University of Washington had the highest average citations per paper (800 papers with approximately 91 citations per paper).

Analyses of Keywords

Table 4 shows the 20 most common keywords. Papers dealing with surgery were more common in LBP-O-related field (7.4%) than in LBP-W-related field (5.2%), as were osteoarthritis (7.1% in LBP-O, 2.6% in LBP-W) and spinal stenosis (7.1% in LBP-O, 3.5% in LBP-W). Figure 4 presents the time-based keyword analysis by average publication year, where the colors of keywords evolve from purple to yellow, indicating changes in hot topics over time. Cognitive-behavioral therapy, intervertebral disc (IVD) degeneration, physiotherapy, physical activity, and walking were the recent

Table 2 Top 10 Prolific Countries from 1993 to 2023

Rank	Low Back Pain in Older Adults					Low Back Pain in Working-Age Adults				
	Country	Papers (%)	Citations (WoS) ^a	Citations Per Paper	Average Publication Year	Country	Papers (%)	Citations (WoS) ^a	Citations Per Paper	Average Publication Year
1	USA	1522 (36.1)	59,006	38.77	2013	USA	15,348 (30.4)	609,476	39.71	2012
2	China	464 (11.0)	6975	15.03	2018	China	4747 (9.4)	71,865	15.14	2018
3	UK	375 (8.9)	18,763	50.03	2014	UK	4701 (9.3)	219,361	46.66	2012
4	Japan	344 (8.2)	7252	21.08	2015	Canada	3940 (7.8)	178,660	45.35	2012
5	Australia	303 (7.2)	10,455	34.5	2016	Australia	3820 (7.6)	156,418	40.95	2014
6	Germany	274 (6.5)	8031	29.31	2013	Germany	3714 (7.3)	110,639	29.79	2013
7	Canada	243 (5.8)	9901	40.74	2014	Netherlands	2882 (5.7)	158,089	54.85	2012
8	South Korea	163 (3.9)	2215	13.59	2016	Japan	2012 (4.0)	39,521	19.64	2014
9	Netherlands	134 (3.2)	6864	51.22	2014	Sweden	1869 (3.7)	82,339	44.06	2011
10	Sweden	131 (3.1)	3698	28.23	2014	South Korea	1785 (3.5)	26,338	14.76	2015

Notes: Papers from Hong Kong, Macau and Taiwan were reclassified to China; papers from England, Scotland, North Ireland, and Wales were reclassified to the UK.

Abbreviation: ^aWoS, Web of Science.

Table 3 Top 10 Prolific Institutions from 1993 to 2023

Rank	Low Back Pain in Older Adults					Low Back Pain in Working-Age Adults				
	Institution	Country	Papers (%)	Citations (WoS) ^a	Citations Per Paper	Institution	Country	Papers (%)	Citations (WoS) ^a	Citations Per Paper
1	University of California	USA	157 (3.7)	9716	61.89	Harvard University	USA	1248 (2.5)	70,583	56.56
2	Harvard University	USA	139 (3.3)	6624	47.65	The University of Sydney	Australia	1124 (2.2)	59,387	52.84
3	University of Pittsburgh	USA	139 (3.3)	7438	53.51	University of California	USA	1088 (2.2)	53,428	49.11
4	The University of Sydney	Australia	120 (2.8)	4637	38.64	Vrije Universiteit Amsterdam	Netherlands	940 (1.9)	60,251	64.1
5	University of Washington	USA	117 (2.8)	6934	59.26	University of Washington	USA	800 (1.6)	73,105	91.38
6	Duke University	USA	75 (1.8)	2477	33.03	University of Toronto	Canada	796 (1.6)	53,896	67.71
7	Keele University	UK	65 (1.5)	2454	37.75	University of London	UK	719 (1.4)	35,287	49.08
8	University of Southern Denmark	Denmark	65 (1.5)	3124	48.06	Karolinska Institutet	Sweden	652 (1.3)	24,141	37.03
9	University of London	UK	64 (1.5)	2427	37.92	The University of Queensland	Australia	645 (1.3)	40,225	62.36
10	University of Toronto	Canada	62 (1.5)	2637	42.53	Maastricht University	Netherlands	622 (1.2)	47,948	77.09

Abbreviation: ^aWoS, Web of Science.

Table 4 The Top 20 Keywords Ranked by Frequency from 1993 to 2023

Rank	Low Back Pain in Older Adults		Low Back Pain in Working-Age Adults	
	Keyword	Co-Occurrences (%)	Keyword	Co-Occurrences (%)
1	Low-back-pain	1578 (37.4)	Low-back-pain	20,196 (40.0)
2	Older-adults	796 (18.9)	Low back pain	8239 (16.3)
3	Prevalence	638 (15.1)	Disability	5215 (10.3)
4	Back-pain	544 (12.9)	Management	4633 (9.2)
5	Disability	544 (12.9)	Back-pain	4509 (8.9)
6	Low back pain	528 (12.5)	Prevalence	4311 (8.5)
7	Management	407 (9.7)	Spine	3522 (7.0)
8	Pain	382 (9.1)	Pain	3472 (6.9)
9	Health	359 (8.5)	Back pain	3188 (6.3)
10	Quality-of-life	347 (8.2)	Reliability	3102 (6.1)
11	Risk-factors	313 (7.4)	Lumbar spine	3063 (6.1)
12	Surgery	312 (7.4)	Chronic pain	3011 (6.0)
13	Osteoarthritis	301 (7.1)	Risk-factors	2912 (5.8)
14	Spinal stenosis	296 (7.0)	Surgery	2609 (5.2)
15	Chronic pain	291 (6.9)	Outcomes	2593 (5.1)
16	Back pain	284 (6.7)	Rehabilitation	2377 (4.7)
17	Outcomes	264 (6.3)	Questionnaire	2146 (4.2)
18	Elderly	263 (6.2)	Health	2075 (4.1)
19	Exercise	262 (6.2)	Follow-up	2068 (4.1)
20	Population	253 (6.0)	Therapy	1967 (3.9)

hot topics with the latest average publication year in the LBP-O-related field (Figure 4A). For LBP-W, IVD degeneration and nucleus pulposus cells were the recent hot topics (Figure 4B). In the LBP-O-related field, some recent burst keywords (ie, lumbar spinal stenosis, and physical activity) were expected to be used more frequently in the coming years, which signified potential emerging trends, while “surgery” was a burst keyword during 1993–2005 and 2007–2010 (Table 5). For the LBP-W-related field, some recent burst keywords, such as surgery and IVD degeneration, signified potential emerging trends (Table 6).

Discussion

This is the first bibliometric analysis to comprehensively map LBP research over the past 3 decades and compare the similarities and differences in the development of LBP-O- and LBP-W-related research. Compared to LBP-W, LBP-O has not been prioritized, which does not reflect the demographic development. The number of journals, countries, and institutions dedicated to the LBP-O-related field was far less than that to the LBP-W-related field. Moreover, the research focuses of LBP-O-related studies differed from those of LBP-W-related research.

The analysis of temporal publication distribution revealed similar increasing annual trends in both fields, with pronounced escalations in the annual publication output after 1993. These surges might be attributed to the publications of many seminal studies after 1990, which laid the conceptual and methodological foundation for clinicians and researchers to develop.^{32,33} For example, Boden et al identified surgical indications (eg, herniated nucleus pulposus, spinal stenosis) in people without symptoms of LBP, sciatica, or neurogenic claudication, and concluded that imaging abnormalities had to be clearly associated with symptoms before surgery;³⁴ Jensen et al linked LBP to disc pathology and concluded that bulges or protrusions in patients with LBP may often be coincidental.³⁵ The disproportionately small number of LBP-O-related studies mirrors the relative lack of clinical interest in managing LBP-O, possibly due to the high prevalence but lack of understanding of its severity and burden, increased comorbidities, and/or lack of funding opportunities.^{3,10,36} Furthermore, it may be partly attributed to the fact that Geriatric Medicine is a relatively young field that arose out of a need rather than a unique body of knowledge. Thus, related research is also relatively new as




Figure 4 Evolution of keywords based on the average publication year. For studies involving: **(A)** low back pain in older adults; and **(B)** low back pain in working-age adults. **Notes:** The year here represents the average publication year, that is, the average year of the articles related to the keywords.

Table 5 Keywords with the Strongest Co-Occurrence Burst on Low Back Pain in Older Adults from 1993 to 2023

Keywords	Year	Strength	Begin	End	1993–2023
Surgical treatment	1993	7.22	1993	2005	
Osteoporosis	1994	12.28	1994	2005	
Symptoms	1995	8.36	1995	2009	
Epidemiology	1995	6.19	1995	2004	
Disorders	1995	4.58	1995	1999	
General population	1996	8.93	1996	2008	
Decompression	1996	8.84	1996	2010	
Bone mineral density	1997	18.65	1997	2008	
Vertebral fractures	1997	10.85	1997	2005	
Diagnosis	1997	7.69	1997	2004	
Laminectomy	1997	6.5	1997	2004	
Deformity	1998	9.25	1998	2005	
Rheumatoid arthritis	1998	8.67	1998	2008	
Nursing home	1998	5.04	1998	2002	
Spondylolisthesis	1998	4.91	1998	2003	
Community	2000	7.75	2000	2007	
Disc degeneration	2000	6.04	2000	2009	
Compression	2000	4.33	2000	2004	
Fusion	2001	4.65	2001	2002	
Postmenopausal women	2002	16.14	2002	2011	
Lumbar spine	1996	22.17	2003	2014	
Validity	2004	6.12	2004	2007	
Spinal stenosis	1993	3.9	2006	2008	
Surgery	1997	5.97	2007	2010	
Arthrodesis	2008	8.65	2008	2010	
Primary care	2000	24.7	2011	2016	
Questionnaire	2000	8.71	2011	2012	
Depression	1995	11.92	2012	2013	
Osteoarthritis	2001	4.7	2013	2014	
Chronic pain	1995	7.78	2015	2016	
Risk	2000	28.06	2019	2021	
Lumbar spinal stenosis	1993	15.05	2019	2023	
Reliability	1998	16.53	2020	2021	
Physical activity	2000	10.97	2020	2023	
Musculoskeletal pain	2002	3.54	2020	2021	
Association	2000	13.16	2021	2023	

Notes: The red bars indicate that the keywords occurred the most frequently; the green bars indicate that the keywords occurred infrequently. Bold keywords represent the potential emerging trends.

compared with the field of pain. Our findings concurred with a previous systematic review that revealed the paucity of clinical trials involving LBP-O.³

LBP-related studies were published in multiple journals. Similar to previous bibliometric analysis studies,^{33,37} our study revealed that *Spine* published the highest publication volume in both fields, accounting for 5.4% and 6.1% of the total number of LBP-O- and LBP-W-related papers, respectively. *Spine* is an international, peer-reviewed, bi-weekly journal focusing on the management of spinal disorders. Among the top 10 journals with the largest number of LBP publications, most have IF < 5.000 and *Pain* has the highest IF in both fields of LBP-O and LBP-W.³¹ *Pain* publishes basic science and clinical research related to the nature, mechanisms, and treatment of pain, which attracts more readers from various disciplines. Journals focusing on LBP treatments (pharmaceutical research in LBP-O and physical rehabilitation in LBP-W) published relatively more LBP-related papers. Although LBP is the number one cause of people living with disabilities and the number of LBP-related publications has been growing exponentially over the last three decades, only a few papers were published in

Table 6 Keywords with the Strongest Co-Occurrence Burst on Low Back Pain in Working-Age Adults from 1993 to 2023

Keywords	Year	Strength	Begin	End	1993–2023
Rheumatoid arthritis	1993	65.45	1993	1997	
Depression	1993	52.61	1993	1997	
Low-back pain	1993	33.88	1993	1995	
Computed tomography	1993	12.98	1993	1994	
Diagnosis	1994	47.96	1994	1999	
Strength	1994	19.48	1994	1996	
Work	1995	81.15	1995	2002	
Sciatica	1996	42.18	1996	1999	
Primary care	1998	152.93	2000	2017	
Population	1995	42.42	2002	2004	
Symptoms	1995	39.83	2002	2003	
Injury	1993	41.13	2003	2004	
Lumbar spine	1993	8.37	2003	2004	
Randomized controlled trial	2004	174.01	2005	2018	
Disorders	1993	40.88	2006	2007	
Double blind	2007	67.29	2007	2008	
Spine	1993	3.98	2007	2008	
Questionnaire	1994	68.67	2012	2013	
Follow up	1994	5.05	2015	2016	
Surgery	1996	146.19	2016	2023	
Outcome	2006	144.08	2016	2023	
Therapy	2000	158.45	2018	2023	
Intervertebral disc degeneration	2017	315.35	2019	2023	
Health	1995	123.81	2019	2020	
Association	2021	183.39	2021	2023	

Notes: The red bars indicate that the keywords occurred frequently; the green bars indicate that the keywords occurred infrequently. Bold keywords represent potential emerging trends.

prominent mainstream medical journals (eg, *Lancet*, *The New England Journal of Medicine*, *Annal Internal Medicine*, and *JAMA Internal Medicine*).^{29,33,38–45} This observation may be ascribed to the perception that LBP is a ubiquitous, but non-life-threatening ailment. Further, LBP research is more related to rehabilitation, which may not be considered for publications in mainstream or specialty medical journals. Despite LBP being a highly costly and debilitating health condition and the fact that primary care providers are nearly always the point of first contact for these patients, insufficient attention is paid to LBP in the general medical literature.^{41–43} In the future, more high-quality LBP studies should be conducted to increase the awareness of medical professionals regarding the prevention and management of LBP.

The highest number of publications in the LBP-O- and LBP-W-related fields over the last three decades originated from the United States because it may have more funding opportunities, and prominent research institutions, which may hire more researchers.^{27,46} Most LBP-O- and LBP-W-related publications were prepared by researchers in developed countries. These papers were cited frequently, which might be attributed to their earlier publication dates. Although the prevalence of LBP is found to be higher in developing countries,⁴⁷ only China was among the top 10 most LBP-prolific countries. The geographical distribution results clearly showed that there were regional differences in the extent of LBP research, and the publication output might be related to the economy and culture.²⁵ Although the incidence of LBP is higher in low-income countries, researchers in these countries may have less funding, interest in, and attention to LBP research. China only started to show dramatic increases in LBP-O- and LBP-W-related publications in the last few years, which might explain the relatively low total number of citations or average citations per paper.

The keywords with recent average publication year help identify the current hot topics. Because IVD degeneration may cause LBP through disc compression, prolapse, and herniation, it has become a common hot topic in both fields. Cognitive-behavioral therapy that incorporates skills of activity rhythm, breathing relaxation, distraction, progressive muscle relaxation, and cognitive restructuring,⁴⁸ has become one of the recent hot topics in LBP-O-related research, while physiotherapy is another hot topic. Moreover, walking and physical activity have recently been recommended as parts of LBP-O management to counteract physical inactivity.⁴⁹ These hot topics highlight the emphasis of conservative LBP management among older adults. Interestingly, nucleus pulposus cells have recently been extensively investigated in LBP-W given their roles in IVD inflammation and degeneration.^{50–52}

The current bibliometric analysis identifies several research priorities. First, as a prevalent and disabling back condition, lumbar spinal stenosis (LSS) was one of the emerging trends in LBP-O research.^{18,53} For example, recent studies have used machine learning to promote individualized LSS treatment and to predict the LSS development.^{54,55} Second, increasing evidence shows that low physical activity is closely related to LBP, although the causation and dose effects remain uncertain.^{56–58} Further studies should determine the optimal dosages of physical activity for both populations. Third, although spine surgery has become an emerging trend in LBP-W-related research, some studies found that certain patients with LBP might not experience significant long-term post-surgery benefits.^{59–64} Future studies should determine the effectiveness of patient-centered precision LBP treatments. Fourth, IVD degeneration is another emerging research trend in LBP-W-related field. The mechanisms of IVD degeneration (eg, miRNAs dysfunction, immunometabolic alterations) and new therapies (eg, stem cell, gene and molecular therapies) may continue to be discovered.^{65–69} Fifth, although our analysis did not reveal any trend in applying artificial intelligence, or big data in LBP research, these approaches have received increasing attention in the last few years,^{70–72} future research should refine and adopt these technologies to improve clinical practice.

The current study had some limitations. First, a small number of studies involving LBP-O or LBP-W might have been missed in the search due to its limitation to WoS alone, without including other databases (eg, Medline, and Embase).²² However, WoS is the most common source for bibliometric analysis and is more advanced in providing detailed information (eg, annual publication volume, journal, country, institution, and citation information).^{73,74} Second, some researchers might have published many papers investigating musculoskeletal or chronic pain conditions, which included LBP. However, because these papers did not use LBP as a keyword, they might have been missed in our search. In addition, although the search strategy was determined by two researchers with content expertise and two experienced librarians, it is plausible that some LBP-W-related papers might include older adults as part of their mixed patient cohorts. That is, given the large number of relevant included literature, our findings should have yielded a good overview of the publication patterns and research trends in the LBP-O and LBP-W fields. Third, this current study did not analyze the Altmetric of the included studies, which is a metric used to measure the social media attention and online engagement of academic articles. Although the citation numbers were commonly used to measure the impact of articles^{75,76} and were correlated with Altmetric,^{77,78} it may not completely represent the impacts of articles. For example, a study about an outcome measure gets cited more frequently, whereas real-life impactful qualitative studies can be cited less. Further studies may consider using Altmetric analysis to evaluate the broader influences of publications. Fourth, there are also some confounding factors, such as the population of the countries and the circulation of the journals, which cannot be excluded.

Implications

The current study provides an overview of the publication patterns and historical research trends, as well as informs research priorities of LBP-O and LBP-W globally. Our findings showed that although there were similar exponential growths in LBP-W- and LBP-O-related publications over the last three decades, the absolute number of LBP-O research was very low as compared to LBP-W research. A recent global burden of LBP study revealed that the prevalence of LBP increases with age, and the highest prevalence of LBP was found in individuals who were approximately 85 years of age.⁷⁹ There is an urgent need to allocate additional research resources towards the prevention and management of LBP-O worldwide (especially in Africa and Asia). Further, our study found that recent research focusing on LBP-W (eg, spinal surgery and IVD degeneration) and LBP-O (eg, physical activity and lumbar spinal stenosis) seemed to be

unrelated to each other. Future efforts should bring researchers in both fields together to discuss the research priorities and synergies so as to optimize the evaluation and management of LBP across the lifespan.

Conclusions

This is the first bibliometric analysis to systematically compare the publication patterns and research trends in LBP research over the past three decades including in LBP-O and LBP-W. Overall, the discrepancy in the publication volume between the two fields has highlighted a relative paucity of LBP-O research in the past. The United States has the highest number of prominent researchers and institutions publishing relevant articles in both populations. The journal *Spine* stands out in LBP research. The shift in the trend of LBP-O-related research from surgery to physical activity suggests the recognition of the importance of physical activity in managing LBP among older adults. Conversely, the growing emphasis on spine surgery and IVD degeneration in LBP-W-related research highlights the strong biomedical focus despite LBP being a biopsychosocial condition.

Abbreviations

IF, impact factor; IVD, intervertebral disc; LBP, low back pain; LBP-O, low back pain in older adults; LBP-W, low back pain in working-age adults; LOESS, locally estimated scatterplot smoothing; LSS, lumbar spinal stenosis; Q, quartile; WoS, Web of Science.

Data Sharing Statement

The data used to support the findings of this study are available from the corresponding author upon request.

Acknowledgments

We would like to thank Jill Boruff, a librarian at McGill University, for giving advice on the search strategy. The abstract of this paper was presented at the “Spineweek 2023” and “The Hong Kong Physiotherapy Association 60th Anniversary Conference” as a conference talk with interim findings.

Funding

This work was supported by GP Batteries Industrial Safety Trust Fund (No.: R-ZDDR).

Disclosure

Professor Jaro Karppinen reports consultation fee from Orion Pharma Ltd outside the submitted work. The authors report no other conflicts of interest in this work.

References

1. Knezevic NN, Candido KD, Vlaeyen JWS, Van Zundert J, Cohen SP. Low back pain. *Lancet*. 2021;398(10294):78–92. doi:10.1016/s0140-6736(21)00733-9
2. Dionne CE, Dunn KM, Croft PR, et al. A consensus approach toward the standardization of back pain definitions for use in prevalence studies. *Spine*. 2008;33(1):95–103. doi:10.1097/BRS.0b013e31815e7f94
3. Paeck T, Ferreira ML, Sun C, Lin CW, Tiedemann A, Maher CG. Are older adults missing from low back pain clinical trials? A systematic review and meta-analysis. *Arthritis Care Res*. 2014;66(8):1220–1226. doi:10.1002/acr.22261
4. Meucci RD, Fassa AG, Faria NM. Prevalence of chronic low back pain: systematic review. *Rev Saude Publica*. 2015;49:1. doi:10.1590/s0034-8910.2015049005874
5. Wong CK, Mak RY, Kwok TS, et al. Prevalence, incidence, and factors associated with non-specific chronic low back pain in community-dwelling older adults aged 60 years and older: a systematic review and meta-analysis. *J Pain*. 2022;23(4):509–534. doi:10.1016/j.jpain.2021.07.012
6. Karjalainen K, Malmivaara A, van Tulder M, et al. Multidisciplinary biopsychosocial rehabilitation for subacute low back pain among working age adults. *Cochrane Database Syst Rev*. 2003;2:Cd002193. doi:10.1002/14651858.Cd002193
7. Kikuchi S, Togo K, Ebata N, et al. A retrospective database study of gastrointestinal events and medical costs associated with nonsteroidal anti-inflammatory drugs in Japanese patients of working age with osteoarthritis and chronic low back pain. *Pain Med*. 2021;22(5):1029–1038. doi:10.1093/pm/pnaa421
8. Vavrek DA, Sharma R, Haas M. Cost analysis related to dose-response of spinal manipulative therapy for chronic low back pain: outcomes from a randomized controlled trial. *J Manipulative Physiol Ther*. 2014;37(5):300–311. doi:10.1016/j.jmpt.2014.03.002

9. Levy B, Kosteas J, Slade M, Myers L. Exclusion of elderly persons from health-risk behavior clinical trials. *Prev Med.* 2006;43(2):80–85. doi:10.1016/j.ypmed.2006.03.019
10. Vadalà G, Russo F, De Salvatore S, et al. Physical activity for the treatment of chronic low back pain in elderly patients: a systematic review. *J Clin Med.* 2020;9(4):1023. doi:10.3390/jcm9041023
11. Bråten LCH, Rolfsen MP, Espeland A, et al. Efficacy of antibiotic treatment in patients with chronic low back pain and Modic changes (the AIM study): double blind, randomised, placebo controlled, multicentre trial. *BMJ.* 2019;367:l5654. doi:10.1136/bmj.l5654
12. Kalichman L, Carmeli E, Been E. The association between imaging parameters of the paraspinal muscles, spinal degeneration, and low back pain. *Biomed Res Int.* 2017;2017:2562957. doi:10.1155/2017/2562957
13. Chiou SY, Koutsos E, Georgiou P, Strutton PH. Association between spectral characteristics of paraspinal muscles and functional disability in patients with low back pain: a cohort study. *BMJ open.* 2018;8(2):e017091. doi:10.1136/bmjopen-2017-017091
14. Qaseem A, Wilt TJ, McLean RM, et al. Noninvasive treatments for acute, subacute, and chronic low back pain: a clinical practice guideline from the American college of physicians. *Ann Intern Med.* 2017;166(7):514–530. doi:10.7326/m16-2367
15. Machado GC, Maher CG, Ferreira PH, et al. Efficacy and safety of paracetamol for spinal pain and osteoarthritis: systematic review and meta-analysis of randomised placebo controlled trials. *BMJ.* 2015;350(mar31 2):h1225. doi:10.1136/bmj.h1225
16. Gobbo S, Bullo V, Bergamo M, et al. Physical exercise is confirmed to reduce low back pain symptoms in office workers: a systematic review of the evidence to improve best practices in the workplace. *J Funct Morphol Kinesiol.* 2019;4(3):43. doi:10.3390/jfkm4030043
17. Zheng K, Chen C, Yang S, Wang X. Aerobic exercise attenuates pain sensitivity: an Event-Related Potential Study. *Front Neurosci.* 2021;15:735470. doi:10.3389/fnins.2021.735470
18. Wong AYL, Karppinen J, Samartzis D. Low back pain in older adults: risk factors, management options and future directions. *Scoliosis Spinal Disord.* 2017;12:14. doi:10.1186/s13013-017-0121-3
19. Chen Q, Hayman LL, Shmerling RH, Bean JF, Leveille SG. Characteristics of chronic pain associated with sleep difficulty in older adults: the Maintenance of Balance, Independent Living, Intellect, and Zest in the Elderly (MOBILIZE) Boston study. *J Am Geriatr Soc.* 2011;59(8):1385–1392. doi:10.1111/j.1532-5415.2011.03544.x
20. Hirase T, Makizako H, Okubo Y, et al. Falls in community-dwelling older adults with lower back or knee pain are associated with cognitive and emotional factors. *Int J Environ Res Public Health.* 2020;17(14):4960. doi:10.3390/ijerph17144960
21. Zheng K, Wang X. Publications on the association between cognitive function and pain from 2000 to 2018: a bibliometric analysis using CiteSpace. *Med Sci Monit.* 2019;25:8940–8951. doi:10.12659/msm.917742
22. AlRyalat SAS, Malkawi LW, Momani SM. Comparing bibliometric analysis using PubMed, Scopus, and Web of Science databases. *J Vis Exp.* 2019;152. doi:10.3791/58494
23. Stout NL, Alfano CM, Belter CW, et al. A bibliometric analysis of the landscape of cancer rehabilitation research (1992–2016). *J Natl Cancer Inst.* 2018;110(8):815–824. doi:10.1093/jnci/djy108
24. Liu X, Zhao S, Tan L, et al. Frontier and hot topics in electrochemiluminescence sensing technology based on CiteSpace bibliometric analysis. *Biosens Bioelectron.* 2022;201:113932. doi:10.1016/j.bios.2021.113932
25. Huang F, Zheng B, Wu C, et al. International publication trends in low back pain research: a bibliometric and visualization analysis. *Front Public Health.* 2022;10:746591. doi:10.3389/fpubh.2022.746591
26. Han X, Zhang J, Chen S, Yu W, Zhou Y, Gu X. Mapping the current trends and hotspots of vascular cognitive impairment from 2000–2021: a bibliometric analysis. *CNS Neurosci Ther.* 2023;29(3):771–782. doi:10.1111/cns.14026
27. Guo F, Liu L, Lv W, Li F, Duffy VG. A bibliometric analysis of occupational low back pain studies from 2000 to 2020. *Arch Environ Occup Health.* 2022;77(5):362–371. doi:10.1080/19338244.2021.1893634
28. Yan W, Zheng K, Weng L, et al. Bibliometric evaluation of 2000–2019 publications on functional near-infrared spectroscopy. *NeuroImage.* 2020;220:117121. doi:10.1016/j.neuroimage.2020.117121
29. Wang R, Weng LM, Peng MS, Wang XQ. Exercise for low back pain: a bibliometric analysis of global research from 1980 to 2018. *J Rehabil Med.* 2020;52(4):jrm00052. doi:10.2340/16501977-2674
30. Du SH, Zheng YL, Zhang YH, Wang MW, Wang XQ. The last decade publications on diabetic peripheral neuropathic pain: a bibliometric analysis. *Front Mol Neurosci.* 2022;15:854000. doi:10.3389/fnmol.2022.854000
31. Zheng KY, Dai GY, Lan Y, Wang XQ. Trends of repetitive transcranial magnetic stimulation from 2009 to 2018: a bibliometric analysis. *Front Neurosci.* 2020;14:106. doi:10.3389/fnins.2020.00106
32. Badhiwala JH, Nassiri F, Witiw CD, et al. Highly cited works in spinal disorders: the top 100 most cited papers published in Spine Journals. *Spine.* 2018;43(24):1746–1755. doi:10.1097/brs.0000000000002735
33. Steinberger J, Skovrlj B, Caridi JM, Cho SK. The top 100 classic papers in lumbar spine surgery. *Spine.* 2015;40(10):740–747. doi:10.1097/brs.0000000000000847
34. Boden SD, Davis DO, Dina TS, Patronas NJ, Wiesel SW. Abnormal magnetic-resonance scans of the lumbar spine in asymptomatic subjects. A prospective investigation. *J Bone Joint Surg Am.* 1990;72(3):403–408. doi:10.2106/00004623-199072030-00013
35. Jensen MC, Brant-Zawadzki MN, Obuchowski N, Modic MT, Malkasian D, Ross JS. Magnetic resonance imaging of the lumbar spine in people without back pain. *N Engl J Med.* 1994;331(2):69–73. doi:10.1056/nejm199407143310201
36. Scheele J, Enthoven WT, Bierma-Zeinstra SM, et al. Course and prognosis of older back pain patients in general practice: a prospective cohort study. *Pain.* 2013;154(6):951–957. doi:10.1016/j.pain.2013.03.007
37. Weng LM, Zheng YL, Peng MS, Chang TT, Wu B, Wang XQ. A bibliometric analysis of nonspecific low back pain research. *Pain Res Manag.* 2020;2020:5396734. doi:10.1155/2020/5396734
38. Cohen KR. Management of chronic low back pain. *JAMA Intern Med.* 2022;182(2):222–223. doi:10.1001/jamainternmed.2021.7359
39. Childs JD, Fritz JM, Flynn TW, et al. A clinical prediction rule to identify patients with low back pain most likely to benefit from spinal manipulation: a validation study. *Ann Intern Med.* 2004;141(12):920–928. doi:10.7326/0003-4819-141-12-200412210-00008
40. Fritz JM, Lane E, McFadden M, et al. Physical therapy referral from primary care for acute back pain with sciatica: a randomized controlled trial. *Ann Intern Med.* 2021;174(1):8–17. doi:10.7326/m20-4187
41. Buchbinder R, van Tulder M, Öberg B, et al. Low back pain: a call for action. *Lancet.* 2018;391(10137):2384–2388. doi:10.1016/s0140-6736(18)30488-4

42. Foster NE, Anema JR, Cherkin D, et al. Prevention and treatment of low back pain: evidence, challenges, and promising directions. *Lancet*. 2018;391(10137):2368–2383. doi:10.1016/s0140-6736(18)30489-6
43. Hartvigsen J, Hancock MJ, Kongsted A, et al. What low back pain is and why we need to pay attention. *Lancet*. 2018;391(10137):2356–2367. doi:10.1016/s0140-6736(18)30480-x
44. Chiarotto A, Koes BW, Solomon CG. Nonspecific Low Back Pain. *N Engl J Med*. 2022;386(18):1732–1740. doi:10.1056/NEJMcp2032396
45. Bagg MK, Wand BM, Cashin AG, et al. Effect of graded sensorimotor retraining on pain intensity in patients with chronic low back pain: a randomized clinical trial. *JAMA*. 2022;328(5):430–439. doi:10.1001/jama.2022.9930
46. Liang YD, Li Y, Zhao J, Wang XY, Zhu HZ, Chen XH. Study of acupuncture for low back pain in recent 20 years: a bibliometric analysis via CiteSpace. *J Pain Res*. 2017;10:951–964. doi:10.2147/jpr.S132808
47. Vos T, Allen C, Arora M. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet*. 2016;388(10053):1545–1602. doi:10.1016/s0140-6736(16)31678-6
48. Goode AP, Taylor SS, Hastings SN, Stanwyck C, Coffman CJ, Allen KD. Effects of a home-based telephone-supported physical activity program for older adult veterans with chronic low back pain. *Phys Ther*. 2018;98(5):369–380. doi:10.1093/ptj/pty026
49. Lewis R, Gómez Álvarez CB, Rayman M, Lanham-New S, Woolf A, Mobasher A. Strategies for optimising musculoskeletal health in the 21(st) century. *BMC Musculoskelet Disord*. 2019;20(1):164. doi:10.1186/s12891-019-2510-7
50. Song XX, Shi S, Guo Z, Li XF, Yu BW. Estrogen receptors involvement in intervertebral discogenic pain of the elderly women: colocalization and correlation with the expression of Substance P in nucleus pulposus. *Oncotarget*. 2017;8(24):38136–38144. doi:10.18632/oncotarget.15421
51. Joyce K, Mohd Isa IL, Krouwels A, Creemers L, Devitt A, Pandit A. The role of altered glycosylation in human nucleus pulposus cells in inflammation and degeneration. *Eur Cell Mater*. 2021;41:401–420. doi:10.22203/eCM.v041a26
52. Liang T, Qiu J, Li S, et al. Inverse agonist of retinoid-related orphan receptor-alpha prevents apoptosis and degeneration in nucleus pulposus cells via upregulation of YAP. *Mediators Inflamm*. 2021;2021:9954909. doi:10.1155/2021/9954909
53. Katz JN, Zimmerman ZE, Mass H, Makhni MC. Diagnosis and management of lumbar spinal stenosis a review. *JAMA*. 2022;327(17):1688–1699. doi:10.1001/jama.2022.5921
54. Liu PY, Wu YT, Xiao ZY, et al. Estimating individualized treatment effects using a risk-modeling approach: an application to epidural steroid injections for lumbar spinal stenosis. *Pain*. 2023;164(4):811–819. doi:10.1097/j.pain.0000000000002768
55. Abbas J, Yousef M, Peled N, Hershkovitz I, Hamoud K. Predictive factors for degenerative lumbar spinal stenosis: a model obtained from a machine learning algorithm technique. *BMC Musculoskelet Disord*. 2023;24(1):218. doi:10.1186/s12891-023-06330-z
56. Vuori IM. Dose-response of physical activity and low back pain, osteoarthritis, and osteoporosis. *Med Sci Sports Exerc*. 2001;33(6 Suppl):S551–S586; discussion 609–10. doi:10.1097/00005768-200106001-00026
57. Tanishima S, Hagino H, Matsumoto H, Tanimura C, Nagashima H. Association between sarcopenia and low back pain in local residents prospective cohort study from the GAINA study. *BMC Musculoskelet Disord*. 2017;18(1):452. doi:10.1186/s12891-017-1807-7
58. Chilibeck PD, Vatanparast H, Cornish SM, Abeysekera S, Charlesworth S. Evidence-based risk assessment and recommendations for physical activity: arthritis, osteoporosis, and low back pain. *Appl Physiol Nutr Metab*. 2011;36(Suppl 1):S49–S79. doi:10.1139/h11-037
59. Kerr D, Zhao W, Lurie JD. What are long-term predictors of outcomes for lumbar disc herniation? A Randomized and Observational Study. *Clin Orthop Relat Res*. 2015;473(6):1920–1930. doi:10.1007/s11999-014-3803-7
60. McGregor AH, Probyn K, Cro S, et al. Rehabilitation following surgery for lumbar spinal stenosis. *Cochrane Database Syst Rev*. 2013;12: Cd009644. doi:10.1002/14651858.CD009644.pub2
61. Fritz JM, Rhon DI, Garland EL, et al. The Effectiveness of a Mindfulness-Based Intervention Integrated with Physical Therapy (MIND-PT) for post-surgical rehabilitation after lumbar surgery: a protocol for a randomized controlled trial as part of the Back Pain Consortium (BACPAC) Research Program. *Pain Med*. 2022. doi:10.1093/pm/pnac138
62. Inose H, Kato T, Sasaki M, et al. Comparison of decompression, decompression plus fusion, and decompression plus stabilization: a long-term follow-up of a prospective, randomized study. *Spine J*. 2022;22(5):747–755. doi:10.1016/j.spinee.2021.12.014
63. Ma XL, Zhao XW, Ma JX, Li F, Wang Y, Lu B. Effectiveness of surgery versus conservative treatment for lumbar spinal stenosis: a system review and meta-analysis of randomized controlled trials. *Int J Surg*. 2017;44:329–338. doi:10.1016/j.ijssu.2017.07.032
64. Gugliotta M, da Costa BR, Dabis E, et al. Surgical versus conservative treatment for lumbar disc herniation: a prospective cohort study. *BMJ open*. 2016;6(12):e012938. doi:10.1136/bmjopen-2016-012938
65. Cazzanelli P, Wuertz-Kozak K. MicroRNAs in intervertebral disc degeneration, apoptosis, inflammation, and mechanobiology. *Int J Mol Sci*. 2020;21(10):3601. doi:10.3390/ijms21103601
66. Francisco V, Pino J, González-Gay M, et al. A new immunometabolic perspective of intervertebral disc degeneration. *Nat Rev Rheumatol*. 2022;18(1):47–60. doi:10.1038/s41584-021-00713-z
67. Wang Y, Che M, Xin J, Zheng Z, Li J, Zhang S. The role of IL-1 β and TNF- α in intervertebral disc degeneration. *Biomed Pharmacother*. 2020;131:110660. doi:10.1016/j.biopha.2020.110660
68. Li B, Yang Y, Wang L, Liu G. Stem cell therapy and exercise for treatment of intervertebral disc degeneration. *Stem Cells Int*. 2021;2021:7982333. doi:10.1155/2021/7982333
69. Kamali A, Ziadlou R, Lang G, et al. Small molecule-based treatment approaches for intervertebral disc degeneration: current options and future directions. *Theranostics*. 2021;11(1):27–47. doi:10.7150/thno.48987
70. Hornung AL, Hornung CM, Mallow GM, et al. Artificial intelligence and spine imaging: limitations, regulatory issues and future direction. *Eur Spine J*. 2022;31(8):2007–2021. doi:10.1007/s00586-021-07108-4
71. Samartzis D, Galbusera F, Wilke HJ. Artificial intelligence, big data and precision spine care: a trend or the holy grail? *Eur Spine J*. 2022;31(8):1941–1942. doi:10.1007/s00586-022-07316-6
72. Wong AYL, Harada G, Lee R, et al. Preoperative paraspinal neck muscle characteristics predict early onset adjacent segment degeneration in anterior cervical fusion patients: a machine-learning modeling analysis. *J Orthop Res*. 2021;39(8):1732–1744. doi:10.1002/jor.24829
73. Miao Y, Xu SY, Chen LS, Liang GY, Pu YP, Yin LH. Trends of long noncoding RNA research from 2007 to 2016: a bibliometric analysis. *Oncotarget*. 2017;8(47):83114–83127. doi:10.18632/oncotarget.20851

74. Chen C. CiteSpace II: detecting and visualizing emerging trends and transient patterns in scientific literature. *J Am Soc Inf Sci Technol.* 2006;57(3):359–377. doi:10.1002/asi.20317
75. Shah SM, Ahmad T, Chen S, Yuting G, Liu X, Yuan Y. A bibliometric analysis of the one hundred most cited studies in psychosomatic research. *Psychother Psychosom.* 2021;90(6):425–430. doi:10.1159/000516185
76. Brandt JS, Hadaya O, Schuster M, Rosen T, Sauer MV, Ananth CV. A bibliometric analysis of top-cited journal articles in obstetrics and gynecology. *JAMA Netw Open.* 2019;2(12):e1918007. doi:10.1001/jamanetworkopen.2019.18007
77. Markusova V, Bogorov V, Libkind A. Usage metrics vs classical metrics: analysis of Russia's research output. *Scientometrics.* 2018;114(2):593–603. doi:10.1007/s11192-017-2597-2
78. Zong Q, Huang Z, Deng Z, Deng W, Xie Y, Yu Y. Identifying combinations of altmetrics and web of science usage that linked to early citations of an article received: a Crisp-Set Qualitative Comparative Analysis (csQCA). *Serials Rev.* 2023;1–10. doi:10.1080/00987913.2023.2234242
79. Ferreira ML, de Luca K, Haile LM; Collaborators GLBP. Global, regional, and national burden of low back pain, 1990–2020, its attributable risk factors, and projections to 2050: a systematic analysis of the Global Burden of Disease Study 2021. *Lancet Rheumatol.* 2023;5(6):e316–e329. doi:10.1016/s2665-9913(23)00098-x

Journal of Pain Research

Dovepress

Publish your work in this journal

The Journal of Pain Research is an international, peer reviewed, open access, online journal that welcomes laboratory and clinical findings in the fields of pain research and the prevention and management of pain. Original research, reviews, symposium reports, hypothesis formation and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-pain-research-journal>