

SCIENTIFIC REPORTS

OPEN

Oxytocin modulates GABA_AR subunits to confer neuroprotection in stroke *in vitro*

Yuji Kaneko¹, Colleen Pappas², Naoki Tajiri¹ & Cesar V. Borlongan¹

Received: 05 August 2016
Accepted: 29 September 2016
Published: 21 October 2016

Oxytocin protects against ischemia-induced inflammation and oxidative stress, and is associated with GABA (γ -aminobutyric acid, an inhibitory neurotransmitter) signaling transduction in neurons. However, the molecular mechanism by which oxytocin affords neuroprotection, especially the interaction between oxytocin receptor and GABA_A receptor (GABA_AR), remains to be elucidated. Primary rat neural cells were exposed to oxytocin before induction of experimental acute stroke model via oxygen-glucose deprivation-reperfusion (OGD/R) injury. Pretreatment with oxytocin increased cell viability, decreased the cell damage against oxidative stress, and prevented the release of high mobility group box1 during OGD/R. However, introduction of oxytocin during OGD/R did not induce neuroprotection. Although oxytocin did not affect the glutathione-related cellular metabolism before OGD, oxytocin modulated the expression levels of GABA_AR subunits, which function to remove excessive neuronal excitability via chloride ion influx. Oxytocin-pretreated cells significantly increased the chloride ion influx in response to GABA and THIP (δ -GABA_AR specific agonist). This study provides evidence that oxytocin regulated GABA_AR subunits in affording neuroprotection against OGD/R injury.

Despite advances in the management and care of stroke, ischemic-reperfusion injury is still a major cause of mortality and morbidity. Male stroke incidence rate and prevalence are significantly higher than females worldwide¹, indicating that this gender difference may be the result of sex hormone, i.e., estrogen. Estrogen regulates oxytocin synthesis in many organs, including the brain. Oxytocin is a typical stress hormone that responds to several acute and chronic stressors, and, together with its receptors, modulates an important array of physiological and biological activities² in central and peripheral nervous systems, such as facilitating birth³. Estrogen, interleukin (IL)-1 β , IL-6, interferon τ , and oxytocin regulate the expression levels of oxytocin receptors⁴.

Plasma oxytocin increases the expression of the peroxisome proliferator-activated receptor gamma gene, a regulator of adipocyte differentiation, and regulates the activity of eukaryotic elongation factor 2⁵, a translation-related protein. Oxytocin controls the differentiation of bone marrow-derived mesenchymal stem cells, regulates the cells' proliferation and carbohydrate metabolism⁶, and promotes lipid metabolism as an energy substrate. Mice deficient in oxytocin receptors have been found to develop obesity⁷.

GABA is the principal inhibitory neurotransmitter in the brain and binds three receptors, GABA_AR, GABA_BR, and GABA_CR. GABA_AR plays a major role in fast synaptic inhibition in the central nervous system (CNS), and is activated by allosteric modulation of interfacial five compositions (2 α /2 β /1 γ , δ , ϵ , θ , π) assembled from 16 subunits (α_{1-6} , β_{1-3} , γ_{1-3} , δ , ϵ , θ , and π) consisting of 20 subtypes⁸. The GABA_AR-mediated hyperpolarization of membrane potential is attributed to the direct activation of an integral anion channel, and the resultant influx of chloride ions along its electrochemical gradient⁹. The equilibrium shift of GABA_AR subtype expression pattern is a key control point for the determination of receptor diversity of the neuronal plasma membrane. Appropriate equilibrium of inhibitory and excitatory neurotransmission regulates the neuronal network in normal brain function. Conversely, an imbalance between inhibitory and excitatory neurotransmission after an ischemic insult creates an excessive secretion of excitatory molecules and suppresses the GABAergic inhibition system, by selectively limiting the trafficking of GABA_ARs on the plasma membrane¹⁰.

Excitotoxicity has been well-documented as a causative factor in ischemia-induced neuronal cell death¹¹. GABA_ARs on neuronal cell membrane are decreased when exposed to oxygen-glucose deprivation (OGD),

¹Center of Excellence for Aging and Brain, Department of Neurosurgery and Brain Repair, University of South Florida College of Medicine, 12901 Bruce B Downs Blvd, Tampa FL 33612, USA. ²School of Aging Studies, University of South Florida, 13301 Bruce B Downs Blvd, Tampa FL 33612, USA. Correspondence and requests for materials should be addressed to Y.K. (email: ykaneko@health.usf.edu) or C.V.B. (email: cborlong@health.usf.edu)




Figure 1. Experimental design. PRNCs; primary rat neural cells. OGD; oxygen and glucose deprivation. OXT; oxytocin. ATS; atosiban.




Figure 2. Oxytocin confers neuroprotective effects and attenuates the oxidative stress against OGD.

(A) Cell viability tested by Calcine-AM/EthD-1 fluorescence and trypan blue dyes. (B) Mitochondrial activity by MTT assay. (C) GSSG activity. (D) Extracellular HMGB1 levels. * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$. Experiments were conducted in triplicate, with $n = 6$ per treatment condition in each run.

suggesting that the number of membrane-bound GABA_ARs could be a pivotal process in the progression of ischemic-induced neuronal cell death¹². Oxytocin regulates GABA_AR-mediated synaptic signaling in the fetal brain during delivery, and reduces brain vulnerability to hypoxic damage¹³. Although oxytocin-induced neuroprotection has been demonstrated in ischemic-reperfusion injury models, the molecular mechanisms underlying such therapeutic benefit, especially how oxytocin interacts with individual GABA_AR subtypes¹⁴, are still unknown.

In this study, we demonstrated that administration of oxytocin in primary rat neural cells (PRNCs) before OGD resulted in robust neuroprotective effects, but not when oxytocin was initiated during OGD/R. We also showed that oxytocin shifted the expression patterns of GABA_AR subunit on the cells, accompanied by increased chloride ion influx. These observations provide evidence that oxytocin modulated GABA_AR in exerting its neuroprotective effects against ischemia-induced neuronal cell death.

Results

Oxytocin exerts neuroprotection against OGD/R. As shown in Fig. 1, PRNCs were exposed to the OGD/R *in vitro* model of stroke^{15,16}. Pretreatment with oxytocin increased cell viability ($F_{(2,34)} = 19.48$; $P < 0.0001$; Fig. 2A), decreased mitochondrial damage ($F_{(2,25)} = 31.81$, $P < 0.0001$; Fig. 2B), reduced oxidative stress ($F_{(2,30)} = 406$, $P < 0.0001$; Fig. 2C), and prevented cell secretion of high mobility group box 1 (HMGB1), a mediator of ischemic progression^{11,17}. Atosiban, the selective oxytocin receptor antagonist¹³, abolished these oxytocin receptor-mediated beneficial effects. Oxytocin administered during OGD/R had no neuroprotective effect (indicated as “With OXT” in Fig. 2).

Biological activity readouts across treatments. Oxytocin acts as an anabolic hormone, and exhibits cell growth^{5,6} and anti-oxidative properties, suggesting its potential therapeutic application in stroke¹⁸. However,




Figure 3. Effects of oxytocin treatment on biological activity before OGD. (A) PRNCs' cell growth. Oxytocin had no significant effects on the cell growth, although oxytocin has been shown to contribute to the differentiation of bone marrow-derived mesenchymal stem cells⁶ (B) G6PD activity. G6PD catalyzes the rate-determining step in the pentose phosphate pathway and produces NADPH to regulate GSH/GSSG levels. G6PD activities across treatments were not significantly different. (C) GAPDH activity. GAPDH regulates the ATP generation phase of glycolysis-derived NAD and functions as a reversible metabolic switch under oxidative stress. GAPDH activities across treatments did not significantly differ. (D) GSSG activity. GSSG is a biomarker of oxidative stress, and is generated by oxidized GSH with reduced NADPH. There were no statistical differences across conditions. Experiments were conducted in triplicate, with $n = 6$ per treatment condition in each run.




Figure 4. Western blot analysis. Following incubation of PRNCs in the absence of reagents (control; CONT), or with $1 \mu\text{M}$ oxytocin + $10 \mu\text{M}$ atosiban (OXT + ATS), or $1 \mu\text{M}$ oxytocin (OXT) for 3 days 37°C . (A) Expression levels of GABA_AR subunits, oxytocin receptor (OXR), bestrophin-1 (BST1), G6PD, and GAPDH. (B) Relative quantification of protein expression levels. Blue bars represent significantly increased protein expression levels, red bar showed significantly decreased levels, and white bars indicate no statistical differences between oxytocin-treated cells and control. * $P < 0.05$ and ** $P < 0.01$. The dotted red line represents combined data from control and OXT + ATS, since these two groups did not significantly differ. Experiments were independently conducted in 3~6 times.

oxytocin administration did not alter cell growth of PRNCs compared with control treatment (Fig. 3A). Next, because peripheral oxytocin participates in glucose metabolism in modulating reactive oxygen species (ROS) production via NADPH (nicotinamide adenine dinucleotide phosphate) pathway, we examined the effects of oxytocin on glutathione (a major antioxidant), and glucose 6-phosphate dehydrogenase (G6PD) and glyceraldehyde 3-phosphate dehydrogenase (GAPDH) (key enzymes that regulate NADPH production)¹⁹. Results revealed that G6PD, GAPDH, and glutathione disulfide (GSSG) activity levels did not significantly differ across treatment conditions (Fig. 3B–D). Pretreating the cells with oxytocin did not change the protein expression levels of G6PD and GAPDH in comparison with control values (Fig. 4).

Oxytocin pretreatment shifts the GABA_AR subunit expression patterns. Oxytocin has been found to alter the subtype expression patterns²⁰ and function of GABA_ARs^{13,21}. We therefore assessed whether oxytocin modulated the expression patterns of GABA_AR subunit on PRNCs (Fig. 4). Treatment of PRNCs with oxytocin significantly increased α_4 , β_3 , δ , and ϵ GABA_AR subunit expression levels, but decreased γ_2 GABA_AR subunit (Fig. 4). Although oxytocin receptor expression has been reported to be increased following oxytocin treatment⁴, we could not detect any differences in oxytocin receptor upregulation between control and oxytocin treatment in this experiment (Fig. 4). Moreover, oxytocin treatment did not affect the expression levels of Bestrophin-1 (BST1), a calcium-activated chloride ion channel normally distributed on synapses adjacent to soma²² and shown to mediate GABA release from astrocytes²³.

Localization of GABA_AR subunits. Electric current and localization patterns of GABA_AR vary depending on the region of interest within the neuron, because glycogen (the main storage form of glucose in the body) is predominantly preserved in the soma where the main production of ATP occurs. In the ischemic brain, the rate of glycogen metabolism is significantly increased²⁴. We observed the localization of α_4 , β_3 , γ_2 , δ , and ϵ GABA_AR subunit expression within subcellular compartments of the neuron. Immunocytochemical analysis showed that (i) δ GABA_AR subunit was mainly located at the axon (Fig. 5A, indicated with box), (ii) ϵ GABA_AR subunit predominantly resided in the soma (Fig. 5B, indicated with box), (iii) γ_2 , α_4 , and β_3 GABA_AR subunits were broadly expressed in the whole neuron (Fig. 5A,B,E,F), (iv) α_4/δ - and β_3/δ -GABA_AR subunits showed co-localization (Fig. 5C,D, indicated with arrow), and (v) α_4/ϵ - and β_3/ϵ -GABA_AR subunits were also co-localized in PRNCs (Fig. 5E,F, indicated with arrow).

Intracellular chloride ion influx kinetics. After binding with GABA, GABA_AR engages a chloride ion selective pore, resulting in chloride ion influx that inhibits the firing of neuron action potentials. The kinetic property of GABA_AR depends on receptor subunit compositions, thereby providing a mechanism for neurons to regulate individual biological activities. We performed a time course study to reveal any differences in GABA-elicited chloride ion influxes between control- and oxytocin-treated PRNCs. Figure 6A revealed that chloride ion influx reached equilibrium at 10 min, but treatment with oxytocin significantly increased the influx at the 20 min period. To assess the differences of both GABA_AR antagonistic conditions, we compared the inhibition dynamics of GABA-induced chloride ion influx in the presence of flumazenil (GABA_AR antagonist, GABA + FLU) or picrotoxin (GABA_AR channel blocker, GABA + PIC). Both reagents inhibited the GABA-induced chloride ion influx (control; Fig. 6B and oxytocin treatment; Fig. 6C). Interestingly, oxytocin-treated cells were more sensitive to picrotoxin inhibition, as evidenced by the Δ value of control cells = 14.0 ± 2.20 AU (Fig. 6B), and that of oxytocin-treated cells = 24.1 ± 1.70 AU (Fig. 6C), $P < 0.001$. Because δ GABA_ARs display increased sensitivity to THIP (δ -GABA_AR specific agonist)²⁵, we tested whether oxytocin-treated cells additively increased THIP-induced chloride ion influx. Results revealed that THIP-evoked chloride ion influx of oxytocin-treated cells was significantly higher than that of control (Fig. 6D).

Discussion

The present study revealed a novel molecular mechanism underlying oxytocin-mediated neuroprotection against ischemic stroke in a cell culture paradigm. We found that oxytocin-induced GABA_AR subunit modification is a predominant factor in conferring neuroprotection against OGD. GABA is the principal inhibitory transmitter in the brain, and its functions are mediated by ubiquitously expressed ligand-opened chloride ion channel GABA_ARs²⁶. Aberrant GABAergic inhibition is a key pathological feature displayed by ischemic neurons in the peri-infarct area (secondary damaged region) after stroke²⁶. Our present results demonstrated that oxytocin reduced ischemic stroke deficits likely by modulating specific GABA_AR subtype signal transduction¹⁴, which parallels studies showing that oxytocin improves stroke outcomes via social interaction pathways¹⁸.

We showed that oxytocin protected PRNCs against OGD (Fig. 2). Ischemic injury is mediated by ROS, generated primarily by damaged mitochondria²⁷, which leads to apoptosis and necrosis. During OGD, cell viability and mitochondrial activity were decreased, and the GSSG activity and extracellular HMGB1 levels were increased. HMGB1, a non-histone DNA-binding protein, is released from necrotic neurons after 2 h OGD¹⁷, and its concentrations in serum are significantly increased in stroke patients due to blood brain barrier (BBB) disruption associated with the disease progression^{11,17}. That oxytocin exerted neuroprotection in OGD, but not in the OGD/R model is consistent with *in vivo* evidence, demonstrating that the subsequent reperfusion after ischemia exacerbates neuronal functions and causes massive brain injuries when oxygen-saturated and nutrient-rich blood suddenly returns to the lesion after a period of ischemia¹¹, suggesting that OGD/R is worse than OGD. Under the OGD condition, pretreatment with oxytocin increased cell viability and mitochondrial activity, decreased the GSSG activity, and prevented HMGB1 secretion from the cells. In the presence of atosiban, this neuroprotection was abolished, indicating that the therapeutic effect was likely mediated by oxytocin receptor signal transduction. HMGB1 is phosphorylated by protein kinase C²⁸ and calcium/calmodulin-dependent protein kinase²⁹. Although oxytocin is capable of activating both kinases³, we could not detect extracellular HMGB1 despite incubating the cells with oxytocin prior OGD. Altogether, these observations suggest that oxytocin could serve as a neuroprotective agent in the acute phase of stroke by acting as an ischemic preconditioning factor in modulating therapeutic protein synthesis.

Oxytocin regulates glucose uptake that is critical for stem cell growth⁶ and antioxidant activity³⁰. However, cell growth of PRNCs was not affected by oxytocin (Fig. 3A). We thus tested whether oxytocin utilized its receptor signal transduction in regulating glutathione-related proteins (G6PD, GAPDH, GSSG). G6PD regulates the antioxidant activity of NADPH³¹, facilitating NADPH to maintain glutathione/GSSG recycling³². GAPDH is not only a key enzyme in glycolysis, but also phosphorylates the α_1 GABA_AR subunit for sustaining the GABA_AR structure and stability, thereby establishing the link of GABAergic inhibition with glucose metabolism. Under




Figure 5. Localization of δ , γ_2 , α_4 , β_3 , and ϵ GABA_AR subunits expression on neurons. Following incubation of PRNCs in the absence of reagent (control; CONT) or with 1 μ M oxytocin (OXT) for 3 days 37 °C. (A) Expression of γ_2 and δ -subunits, (B) γ_2 and ϵ -subunits, (C) δ and α_4 -subunits, (D) δ and β_3 -subunits, (E) ϵ and α_4 -subunits, and (F) ϵ and β_3 -subunits. Scale bars = 10 μ m.




Figure 6. Characterization of chloride ion influx via GABA_AR. Following incubation of PRNCs in the absence of reagents (control; CONT) or with 1 μ M oxytocin (OXT) for 3 days 37 $^{\circ}$ C. **(A)** Time course of chloride ion influx stimulated by 50 μ M GABA. **(B,C)** The cells were pretreated with 1 μ M flumazenil (GABA + FLU), or 1 μ M picrotoxin (GABA + PIC), or PBS (GABA) for 45 min at 37 $^{\circ}$ C, and were stimulated by 50 μ M GABA for 10 min at RT, and then fluorescence intensity was measured. **(B,C)** represent control- and oxytocin treated-cells, respectively. The Δ value was calculated from following equation: fluorescence intensity of (GABA – [GABA + PIC]). **(D)** Time course of chloride ion influx stimulated by 10 nM THIP. * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$. Experiments were conducted in triplicate, with $n = 6$ per treatment condition in each run.

ischemic condition, glycolytic flux increases, while GAPDH activity is reversibly decreased, inhibiting neuronal cells from producing NADPH³¹ as a result of increased G6PD activity. We speculated that oxytocin would normalize G6PD, GAPDH, GSSG activity, but no significant difference were detected across treatment conditions (Fig. 3B–D), suggesting that oxytocin’s neuroprotection is largely independent of G6PD, GAPDH, and GSSG signal transduction pathways.

Maternal oxytocin exerts neuroprotective action on fetal neurons during parturition (a perturbed physiological environment similar to hypoxic-ischemic brain) mediated by GABA_AR signaling pathway¹³. In the present study, oxytocin likely engaged the GABA_AR subunit expression patterns by enhancing α_4 , β_3 , δ , and ϵ GABA_AR subunit expression levels while reducing γ_2 GABA_AR subunit on PRNCs (Fig. 4). Although most GABA_AR subunit expressions were not significantly influenced in the presence of atosiban (OXT + ATS), α_2 GABA_AR subunit expression decreased and γ_1 and π GABA_AR subunits increased (Fig. 4A), implicating that redundant signal impedance of oxytocin receptor could activate alternative signal transduction of these subunit expressions. Conversely, an on/off signaling switch of oxytocin may be tightly regulated by engagement with GABA_AR subunits.

The β_3 and γ_2 GABA_AR subunits on the neuronal membrane are vulnerable to ischemic stroke^{10,33}. In the late stage of rat pregnancy the maternal brain displays increased expression of the ϵ GABA_AR subunit, which is responsible for the respiratory function³⁴. The α_4 GABA_AR subunit, on the other hand, is associated with dendritic development³⁵, and is well co-expressed with δ -subunit in the brain^{8,36}. The α_4 , δ , and ϵ GABA_AR subtypes are extrasynaptic GABA_ARs^{8,36}, which mediate tonic inhibition upon activation by GABA spillover from synaptic sites, as well as by ambient GABA in the extracellular space. The majority of GABA_ARs are $\alpha_1\beta_2\gamma_2$, $\alpha_2\beta_3\gamma_2$, and $\alpha_3\beta_2/\beta_3\gamma_2$, approximately occupying 80% of total GABA_AR expressions in the brain⁸, in contrast, α_4 , δ , and ϵ GABA_ARs represent less than 5%⁸. Oxytocin-treated cells significantly increased α_4 , δ , and ϵ GABA_AR subunit expression levels and decreased γ_2 GABA_AR subunit on PRNCs (Fig. 4). It is conceivable that oxytocin pretreatment led to an upregulation of specific GABA_AR subtype expression levels, which in turn might have modified the neuronal networks towards neuroprotection.

Electrical properties of a neuron vary along the segments of subcellular organization (soma, dendrites, and axon), which is essential for orchestrating cellular function and structure preservation³⁷. GABA_AR-mediated chloride ion fluctuation substantially changes the intracellular chloride ion concentration in the soma and spreads into the dendrites³⁸. Hypoxic ischemia causes retrograde neurodegeneration, which shortens the axonal and dendritic lengths and swells the soma, and produces a rapid and significant loss of axon in the acute phase of injury³⁹. Therefore, it is an important to elucidate the GABA_AR subunit localizations at the subcellular neuron. Interestingly, δ GABA_AR subunit is mainly expressed at axon, and ϵ GABA_AR subunit is primarily distributed on soma which concurred with previous report⁴⁰. In contrast, α_4 , β_3 and γ_2 GABA_AR subunits were broadly expressed throughout the whole neuron⁸. Here, we found that α_4/δ - and β_3/δ -GABA_AR subunits, and α_4/ϵ - and

β_3/ϵ -GABA_AR subunits were well co-localized on neurons (Fig. 5). That appropriate distribution and specific expression of GABA_AR subtypes exist in neurons suggest that oxytocin could elicit neuroprotection by subcellularly targeting specific GABA_AR subunits within ischemic neurons.

Under hypoxic ischemia, the extracellular GABA concentration on/around the synaptic cleft increases and elevates neuronal intracellular chloride ion, which functions as a counter-reaction of depolarization⁴¹. We demonstrated here that oxytocin modulated discrete GABA_AR subunits tasked to monitor chloride ion influx. The kinetic of GABA-stimulated chloride ion influx on oxytocin-treated cells was altered (Fig. 6A), in that while the response to GABA by the control cells was saturated at 10 min, oxytocin-treated cells continuously evoked the ion influx for 20 minutes, suggesting that boosting GABA_AR-mediated neuronal inhibition can afford substantial protection while minimizing the extent of neuronal cell loss during OGD. Following incubation of cells with 50 μ M GABA for 10 min (Fig. 6A), the values of chloride ion influx were similar. We next assessed GABA_AR antagonism and found that Flumazenil inhibited the γ contained GABA_AR subunits in response to GABA, but was not able to antagonize the δ and ϵ GABA_AR subunits⁴². Picrotoxin directly binds the ion pore of GABA_AR, thereby regulating the influx of chloride ion, and inhibiting the whole GABA_AR channel activity⁴². The inhibition ability of flumazenil (GABA + FLU) did not significantly differ between conditions, implicating that oxytocin administration had no effect on the expression levels of γ GABA_AR subtypes (Fig. 6B,C). In contrast, oxytocin-treated cells exhibited significant inhibition of chloride ion influx following picrotoxin treatment (GABA + PIC) compared to control, suggesting that oxytocin significantly increased total GABA_AR expression, especially δ and ϵ GABA_AR subtypes. Of note, to date, there is no specific antagonist for δ and ϵ GABA_AR subunits. The present observation of specialized GABA_AR antagonism is also supported by δ -GABA_AR specific agonist THIP significantly elevating the chloride ion influx of oxytocin-treated cells compared to control (Fig. 6D). In summary, oxytocin induced the shift of GABA_AR subunit expression in cultured PRNCs, which likely changed the kinetics of chloride ion influx in response to GABA.

We demonstrated that oxytocin exerts neuroprotection against ischemic stroke, but requires its treatment initiation prior to injury induction. Oxytocin may serve as a pharmacological ischemic preconditioning factor that can engage GABA_AR towards neuroprotection. The present results provide evidence that oxytocin altered the expression patterns of GABA_AR subunit and the kinetics of GABA-induced chloride ion influx. Our study highlights a close interaction between oxytocin and GABA_AR that should aid in our understanding of stroke pathology and its treatment.

Methods

Cell culture and oxygen-glucose deprivation-reperfusion (OGD/R) progression. Primary rat neural cells (PRNCs; consisted of 40% neurons and 60% astrocytes) were obtained from BrainBit (E18 rat cortex; Springfield, IL, USA). As described elsewhere¹⁵, cells (4×10^4 cells/well) were suspended in 200 μ l Neural Medium (NbActive 4, BrainBit) containing 2 mM *l*-glutamine and 2% B27 in the absence of antibiotics and grown in poly-*l*-lysine-coated 96-well plates at 37 °C in humidified atmosphere containing 95% O₂ and 5% CO₂. After 3 days in culture, PRNCs were exposed to 1 μ M oxytocin (O4375, Sigma-Aldrich, St. Louis, MI, USA), 1 μ M oxytocin + 10 μ M atosiban (A3480, Sigma-Aldrich), 10 μ M atosiban, and the absence of reagents (control) for 3 days at 37 °C. After 6 days in culture (Fig. 1), PRNCs were exposed to OGD as described previously¹⁵. The cells were initially exposed to OGD medium (glucose-free Dulbecco's Modified Eagle Medium), then placed in an anaerobic chamber containing 95% N₂ and 5% CO₂ for 15 min at 37 °C (preincubation), for 90 min at 37 °C (culture medium pH 6.7~6.8; mimicking the acidic environment of ischemic brain *in vivo*). OGD was terminated by adding 5 mM glucose to medium and cell cultures were re-introduced to the regular CO₂ incubator at 37 °C for 2 h. Control cells were incubated in the same buffer containing 5 mM glucose at 37 °C in a regular 95% O₂ and 5% CO₂ incubator.

Measurement of cell viability. Measurement of cell viability was performed using fluorescent live/dead cell assay and trypan blue exclusion method^{15,43}. Following treatment, the cells were incubated with 2 μ M Calcein-AM and 4 μ M EthD-1 (L3224 Invitrogen, Waltham, MA, USA) for 45 min at room temperature (RT) in dark. After washing once with phosphate buffer saline (PBS), the green fluorescence of the live cells was measured by the Gemini EX fluorescence plate reader (Ex/Em = 490/520; Molecular Devices, Sunnyvale, CA). In addition, trypan blue (15250, Gibco, Waltham, MS, USA) exclusion method was conducted and mean viable cell counts were calculated in 16 randomly selected areas (1 mm², n = 10) to reveal the cell viability. Briefly, within 5 min after adding trypan blue, we digitally captured under microscope (200x) 10 pictures (approximately 100 cells/picture) for each condition, then randomly selected 5 pictures, and counted the number of cells for each individual treatment condition. Normalized cell viability was calculated from the following equation: viable cells (%) = [1.00 - (Number of blue cells / Number of total cells)] \times 100. To precisely calibrate the cell viability, the values were standardized from fluorescence intensity and trypan blue data.

Measurement of mitochondrial activity. Following OGD/R, reduction of 3-(4,5-dimethyl-2-thiazoyl)-2,5-diphenyltetrazolium bromide (MTT; 11465007001, Roche, Basel, Switzerland) by mitochondrial dehydrogenases was used as a measure of mitochondrial activity as previously described⁴³. The optical density of solubilized purple formazan was measured at 570 nm on a Synergy HT plate reader (Bio-Tex, Winooski, VT, USA).

Measurement of extracellular high mobility group box1 (HMGB1) levels and glutathione disulfide (GSSH) activity. After OGD/R, culture medium was centrifuged at 3,000 g, 4 °C for 15 min, and the supernatant was processed for detection of HMGB1 using an ELISA kit (amin416082, Antibody, Atlanta, GA, USA) with absorbance measured at 450 nm on a Synergy HT plate reader (Bio-Tex). Cells were treated with oxidized glutathione lysis reagent (V6611, Promega, Fitchburg, WI, USA), and GSSH activity, a biomarker of reactive

oxygen species (ROS) production, was measured by luciferase activity on Spectra Max Gemini EM plate reader (Molecular Devices, Sunnyvale, CA, USA).

Measurement of cell growth, glucose 6-phosphate dehydrogenase (G6PD)-, and glyceraldehyde 3-phosphate dehydrogenase (GAPDH)-activity. Following cell culture, the cleavage of the tetrazolium salt, WST-1 (4-[3(4-Iodophenyl)-2-(4-nitrophenyl)-2H-5-tetrazolio]-1,3-benzene disulfonate; 05015944011, Roche) formazan was used as a measure cell growth. The optical density was measured at 450 nm on a Synergy HT plate reader (Bio-Tex). The levels of G6PDH- and GAPDH-activity were performed according to the manufacturer's protocols for G6PD assay kit (ab102529, Abcam, Cambridge, MA, USA) and GAPDH ELISA kit (ab119627, Abcam), respectively.

Western blot analysis. PRNCs were treated with CellLytic MT mammalian lysis reagent (C3228, Sigma-Aldrich) with protease inhibitor cocktail (I3786, Sigma-Aldrich). The lysate was centrifuged at 3,000 g, 4 °C for 15 min, and the supernatant was stored at -80 °C until analysis. Protein samples (4–35 µg/lane) were processed on 4–14% Tris-Glycine SDS-PAGE gel and then transferred onto a nitrocellulose membrane (162–0112, Bio-Rad, Hercules, CA, USA) at 30 V, 4 °C for 14 h. The nitrocellulose membranes were treated with PBS containing 0.1% Tween-20 and 3% non-fat milk (170–6404, Bio-Rad) for 45 min at RT. Membranes were then incubated with the primary antibodies, anti-oxytocin receptor rabbit antibody (1/10,000, ab181077, Abcam), anti-GABA_AR α₁ subunit rabbit antibody (1/3,000, ab3299, Abcam), anti-GABA_AR α₂ subunit rabbit antibody (1/1,000, ab72445, Abcam), anti-GABA_AR α₃ subunit rabbit antibody (1/1,000, ab72446, Abcam), anti-GABA_AR α₄ subunit rabbit antibody (1/1,000, ab4120, Abcam), anti-GABA_AR α₅ subunit rabbit antibody (1/3,000, ab10098, Abcam), anti-GABA_AR α₆ subunit goat antibody (1/1,000, ab117100, Abcam), anti-GABA_AR β₁ subunit rabbit antibody (1/3,000, ab154822, Abcam), anti-GABA_AR β₂ subunit rabbit antibody (1/30,000, ab16213, Abcam), anti-GABA_AR β₃ subunit rabbit antibody (1/1,000, ab4046, Abcam), anti-GABA_AR γ₁ subunit rabbit antibody (1/5,000, AMIN485542, Antibodies), anti-GABA_AR γ₂ subunit rabbit antibody (1/1,000, ab16213, Abcam), anti-GABA_AR γ₃ subunit rabbit antibody (1/500, ab13861, Abcam), anti-GABA_AR δ subunit rabbit antibody (1/1,000, ab11048, Abcam), anti-GABA_AR ε subunit rabbit antibody (1/500, ab35971, Abcam), anti-GABA_AR π subunit rabbit antibody (1/5,000, ab26055, Abcam), anti-GABA_AR θ subunit rabbit antibody (1/5,000, ARP5283, Antibodies), anti-bestrophin-1 (BST1) mouse antibody (1/3,000, NB300-164, Antibodies), anti-G6PD rabbit antibody (1/10,000, ab993, Abcam), and anti-GAPDH mouse antibody (1/10,000, ab8245, Abcam) at 4 °C for 14 h. After washing with PBS containing 0.1% Tween-20 (PBST), the nitrocellulose membrane was incubated with donkey anti-mouse IRDye800[®]CW secondary antibody (1/5,000, 926-32212, LI-COR, Lincoln, NE, USA), or donkey anti-rabbit IRDye800[®]CW secondary antibody (1/5,000, 926-32213, LI-COR), or donkey anti-goat IRDye800[®]CW secondary antibody (1/5,000, 926-32214, LI-COR) for 90 min at RT in dark. Immunoreactive detection using near-infrared fluorescence was performed according to the protocol of Odyssey[®] Infrared Imaging System (LI-COR[®]).

Immunocytochemistry analysis. PRNCs (8 × 10⁴ cell/well) were cultured in 400 µl Neural medium containing 2 mM *l*- glutamine and 2% B27 in the absence of antibiotics in poly-*l*-lysine 8-chamber (354632, BD Bioscience, Franklin Lakes, NJ, USA) for 3 days, then the cells were exposed to 1 µM oxytocin in the absence of reagents (control) for 3 days and fixed in 4% paraformaldehyde¹⁵. The cells were washed 5 times for 10 min in PBST. Then they were blocked by 5% normal goat serum (50062Z, Invitrogen, Carisbad, CA, USA) in PBST for 1 h at RT. Primary antibodies included anti-GABA_AR α₄ subunit mouse antibody (1/100, SMC-489, StressMarq Bioscience Inc., BC, Canada), anti-GABA_AR β₃ subunit mouse antibody (1/500, ab98968, Abcam), anti-GABA_AR γ₂ subunit mouse antibody (1/250, MABN263, Millipore, Billerica, MA, USA), anti-GABA_AR δ subunit rabbit antibody (1/200, ab111048, Abcam), anti-GABA_AR ε subunit rabbit antibody (1/200, ab35971, Abcam), and anti-microtubule-associated protein 2 (MAP2) chicken antibody (1/10,000, ab5392, Abcam). The cells were incubated overnight at 4 °C in primary antibody with 5% normal goat serum. The cells were washed 5 times for 10 min in PBST and then soaked in 5% normal goat serum in PBST containing corresponding secondary antibodies goat anti-mouse IgG-Alexa 405 (blue; 1/1,000, A31553, Invitrogen), goat anti-rabbit IgG-Alexa 405 (blue; 1/1,000, A31556, Invitrogen), goat anti-mouse IgG-Alexa 488 (green; 1/1,000, A11029, Invitrogen), goat anti-rabbit IgG-Alexa 488 (green; 1/1,000, A11034, Invitrogen), and goat anti-chicken IgG-Alexa 594 (red; 1/1,000, A11042, Invitrogen) for 90 min in the dark. Immunofluorescent images were visualized using confocal microscope (FV1000, Olympus, Tokyo, Japan). Control experiments were performed with the omission of the primary antibodies yielding negative results.

Measurement of intracellular chloride ion influx. The quinolinium salt-based halide-sensitive fluorescence probe *N*-(ethoxycarbonylmethyl)-6-methoxyquinolium bromide (MQAE; ab145418, Abcam) was used as a measure of chloride ion influx activity¹³. Following cell culture, the PRNCs were incubated with 5 mM MQAE for 2 h at RT in the dark, and subsequently washed twice with NbActive 4 (BrainBit). Cells were then treated with 50 µM GABA (A2129, Sigma-Aldrich) or 10 nM GABA analog THIP, δ-GABA_AR specific agonist⁸ (T101, Sigma-Aldrich), then fluorescence intensity was consequently measured at 0, 5, 10, and 20 min at RT. For inhibition assay, the cells were pretreated with 1 µM flumazenil (F6300, Sigma-Aldrich) or 1 µM picrotoxin (P1675, Sigma-Aldrich) or PBS (control) for 45 min at RT, and then stimulated with 50 µM GABA for 10 min at RT. Intercellular MQAE is quenched by 10 µM tributyltin chloride (T50202, Sigma-Aldrich) and 10 µM nigericin sodium salt (N7143, Sigma-Aldrich). The fluorescence intensity was measured by the Gemini EX fluorescence plate reader (Ex/Em = 360/460; Molecular Devices). The kinetic analysis was performed by using GraphPad Prism 6[®] software.

Data analysis. Data were evaluated using one-way analysis of variance (ANOVA) followed by post hoc compromised t-tests (GraphPad Prism 6[®] software). Statistical significance was preset at $P < 0.05$. Data are represented as means \pm SD from quintuplicates of each treatment condition.

References

- Appelros, P., Stegmayr, B. & Terent, A. Sex differences in stroke epidemiology: a systematic review. *Stroke* **40**, 1082–1090 (2009).
- Evans, S. L., Dal Monte, O., Noble, P. & Averbeck, B. B. Intranasal oxytocin effects on social cognition: a critique. *Brain Res* **1580**, 69–77 (2014).
- Shoji, H. & Kaneko, Y. Characterization and expression of oxytocin and the oxytocin receptor. *Mol Genet Metab* **71**, 552–558 (2000).
- Zingg, H. H. & Laporte, S. A. The oxytocin receptor. *Trends Endocrinol Metab* **14**, 222–227 (2003).
- Eckertova, M., Ondrejčáková, M., Krsková, K., Zorad, S. & Jezova, D. Subchronic treatment of rats with oxytocin results in improved adipocyte differentiation and increased gene expression of factors involved in adipogenesis. *Br J Pharmacol* **162**, 452–463 (2011).
- Elabd, C. *et al.* Oxytocin controls differentiation of human mesenchymal stem cells and reverses osteoporosis. *Stem Cells* **26**, 2399–2407 (2008).
- Takayanagi, Y. *et al.* Oxytocin receptor-deficient mice developed late-onset obesity. *Neuroreport* **19**, 951–955 (2008).
- Rudolph, U. & Mohler, H. GABAA receptor subtypes: Therapeutic potential in Down syndrome, affective disorders, schizophrenia, and autism. *Annu Rev Pharmacol Toxicol* **54**, 483–507 (2014).
- Jacob, T. C., Moss, S. J. & Jurd, R. GABA(A) receptor trafficking and its role in the dynamic modulation of neuronal inhibition. *Nat Rev Neurosci* **9**, 331–343 (2008).
- Liu, B. *et al.* Preservation of GABAA receptor function by PTEN inhibition protects against neuronal death in ischemic stroke. *Stroke* **41**, 1018–1026 (2010).
- Jickling, G. C. *et al.* Hemorrhagic transformation after ischemic stroke in animals and humans. *J Cereb Blood Flow Metab* **34**, 185–199 (2014).
- Mielke, J. G. & Wang, Y. T. Insulin exerts neuroprotection by counteracting the decrease in cell-surface GABA receptors following oxygen-glucose deprivation in cultured cortical neurons. *J Neurochem* **92**, 103–113 (2005).
- Tyzio, R. *et al.* Maternal oxytocin triggers a transient inhibitory switch in GABA signaling in the fetal brain during delivery. *Science* **314**, 1788–1792 (2006).
- Bowen, M. T. *et al.* Oxytocin prevents ethanol actions at delta subunit-containing GABAA receptors and attenuates ethanol-induced motor impairment in rats. *Proc Natl Acad Sci USA* **112**, 3104–3109 (2015).
- Kaneko, Y., Tajiri, N., Shoji, H. & Borlongan, C. V. Oxygen-glucose-deprived rat primary neural cells exhibit DJ-1 translocation into healthy mitochondria: a potent stroke therapeutic target. *CNS Neurosci Ther* **20**, 275–281 (2014).
- Tajiri, N., Borlongan, C. V. & Kaneko, Y. Cyclosporine A Treatment Abrogates Ischemia-Induced Neuronal Cell Death by Preserving Mitochondrial Integrity through Upregulation of the Parkinson's Disease-Associated Protein DJ-1. *CNS Neurosci Ther* (2016).
- Zhang, J. *et al.* Anti-high mobility group box-1 monoclonal antibody protects the blood-brain barrier from ischemia-induced disruption in rats. *Stroke* **42**, 1420–1428 (2011).
- Karelina, K. *et al.* Oxytocin mediates social neuroprotection after cerebral ischemia. *Stroke* **42**, 3606–3611 (2011).
- Laschet, J. J. *et al.* Glyceraldehyde-3-phosphate dehydrogenase is a GABAA receptor kinase linking glycolysis to neuronal inhibition. *J Neurosci* **24**, 7614–7622 (2004).
- Koksma, J. J. *et al.* Oxytocin regulates neurosteroid modulation of GABA(A) receptors in supraoptic nucleus around parturition. *J Neurosci* **23**, 788–797 (2003).
- Tyzio, R. *et al.* Oxytocin-mediated GABA inhibition during delivery attenuates autism pathogenesis in rodent offspring. *Science* **343**, 675–679 (2014).
- Jo, S. *et al.* GABA from reactive astrocytes impairs memory in mouse models of Alzheimer's disease. *Nat Med* **20**, 886–896 (2014).
- Lee, S. *et al.* Channel-mediated tonic GABA release from glia. *Science* **330**, 790–796 (2010).
- Dienel, G. A. & Cruz, N. F. Astrocyte activation in working brain: energy supplied by minor substrates. *Neurochem Int* **48**, 586–595 (2006).
- Meera, P., Wallner, M. & Otis, T. S. Molecular basis for the high THIP/gaboxadol sensitivity of extrasynaptic GABA(A) receptors. *J Neurophysiol* **106**, 2057–2064 (2011).
- Clarkson, A. N., Huang, B. S., Macisaac, S. E., Mody, I. & Carmichael, S. T. Reducing excessive GABA-mediated tonic inhibition promotes functional recovery after stroke. *Nature* **468**, 305–309 (2010).
- Fulda, S., Galluzzi, L. & Kroemer, G. Targeting mitochondria for cancer therapy. *Nat Rev Drug Discov* **9**, 447–464 (2010).
- Kang, H. J. *et al.* Non-histone nuclear factor HMGB1 is phosphorylated and secreted in colon cancers. *Lab Invest* **89**, 948–959 (2009).
- Zhang, X. *et al.* Calcium/calmodulin-dependent protein kinase (CaMK) IV mediates nucleocytoplasmic shuttling and release of HMGB1 during lipopolysaccharide stimulation of macrophages. *J Immunol* **181**, 5015–5023 (2008).
- Kim, S. W. *et al.* Glycyrrhizic acid affords robust neuroprotection in the postischemic brain via anti-inflammatory effect by inhibiting HMGB1 phosphorylation and secretion. *Neurobiol Dis* **46**, 147–156 (2012).
- Ralsler, M. *et al.* Dynamic rerouting of the carbohydrate flux is key to counteracting oxidative stress. *J Biol Chem* **282**, 10 (2007).
- Hashida, K., Sakakura, Y. & Makino, N. Kinetic studies on the hydrogen peroxide elimination by cultured PC12 cells: rate limitation by glucose-6-phosphate dehydrogenase. *Biochim Biophys Acta* **1572**, 85–90 (2002).
- Mele, M., Ribeiro, L., Inacio, A. R., Wieloch, T. & Duarte, C. B. GABA(A) receptor dephosphorylation followed by internalization is coupled to neuronal death in *in vitro* ischemia. *Neurobiol Dis* **65**, 220–232 (2014).
- Hengen, K. B. *et al.* Increased GABA(A) receptor epsilon-subunit expression on ventral respiratory column neurons protects breathing during pregnancy. *PLoS One* **7**, e30608 (2012).
- Duveau, V. *et al.* Spatiotemporal specificity of GABAA receptor-mediated regulation of adult hippocampal neurogenesis. *Eur J Neurosci* **34**, 362–373 (2011).
- Brickley, S. G. & Mody, I. Extrasynaptic GABA(A) receptors: their function in the CNS and implications for disease. *Neuron* **73**, 23–34 (2012).
- Dugladze, T., Schmitz, D., Whittington, M. A., Vida, I. & Gloveli, T. Segregation of axonal and somatic activity during fast network oscillations. *Science* **336**, 1458–1461 (2012).
- Kuner, T. & Augustine, G. J. A genetically encoded ratiometric indicator for chloride: capturing chloride transients in cultured hippocampal neurons. *Neuron* **27**, 447–459 (2000).
- Hinman, J. D. The back and forth of axonal injury and repair after stroke. *Curr Opin Neurol* **27**, 615–623 (2014).
- Sergeeva, O. A., Andreeva, N., Garret, M., Scherer, A. & Haas, H. L. Pharmacological properties of GABAA receptors in rat hypothalamic neurons expressing the epsilon-subunit. *J Neurosci* **25**, 88–95 (2005).
- Pond, B. B. *et al.* The chloride transporter Na(+)-K(+)-Cl(-) cotransporter isoform-1 contributes to intracellular chloride increases after *in vitro* ischemia. *J Neurosci* **26**, 1396–1406 (2006).
- Johnston, G. A. GABAA receptor pharmacology. *Pharmacol Ther* **69**, 173–198 (1996).
- Kaneko, Y. *et al.* Kainic Acid-Induced Golgi Complex Fragmentation/Dispersal Shifts the Proteolysis of Reelin in Primary Rat Neuronal Cells: An *In Vitro* Model of Early Stage Epilepsy. *Mol Neurobiol* **53**, 1874–1883 (2016).

Acknowledgements

This work was supported by USF OR&I HSC-18330 & COM HSC-18300 (YK), NIH NINDS RO1 1R01NS071956-01 (CVB), and USF Department of Neurosurgery and Brain Repair Funds (CVB). Part of this manuscript was presented at the American Society for Neural Therapy and Repair 23d Annual Conference, April 28–30, 2016, Clearwater Beach, Florida.

Author Contributions

Y.K. and C.V.B. contributed to study design. Y.K., C.P., N.T. and C.V.B. contributed to data acquisition. Y.K., P.C., N.T. and C.V.B. contributed to data analysis and interpretation. Y.K., C.P., N.T. and C.V.B. contributed to preparation of manuscript.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Kaneko, Y. *et al.* Oxytocin modulates GABA_AR subunits to confer neuroprotection in stroke *in vitro*. *Sci. Rep.* **6**, 35659; doi: 10.1038/srep35659 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016