



# Genome Sequence of Type Strain *Fonsecaea multimorphosa* CBS 980.96<sup>T</sup>, a Causal Agent of Feline Cerebral Phaeohyphomycosis

Aniele C. Ribas Leao,<sup>a,b,c</sup> Vinicius Almir Weiss,<sup>a</sup> Vania Aparecida Vicente,<sup>d</sup> Flavia Costa,<sup>c</sup> Amanda Bombassaro,<sup>d</sup> Roberto Tadeu Raittz,<sup>a</sup> Maria Berenice R. Steffens,<sup>a,b</sup> Fabio Oliveira Pedrosa,<sup>a,b</sup> Renata R. Gomes,<sup>d</sup> Valter Baura,<sup>b</sup> Helisson Faoro,<sup>b,e</sup> Michelle Zibetti Tadra Sfeir,<sup>b</sup> Eduardo Balsanelli,<sup>b</sup> Leandro F. Moreno,<sup>f</sup> M. Javad Najafzadeh,<sup>g</sup> Sybren de Hoog,<sup>f</sup> Emanuel Maltempo Souza<sup>a,b</sup>

Laboratory of Bioinformatics, Professional and Technological Education Sector, Federal University of Paraná, Curitiba, Paraná, Brazil<sup>a</sup>; Department of Biochemistry and Molecular Biology, Federal University of Paraná, Curitiba, Paraná, Brazil<sup>b</sup>; Department of Bioprocess Engineering and Biotechnology, Federal University of Paraná, Curitiba, Paraná, Brazil<sup>c</sup>; Laboratory of Microbiology, Department of Pathology, Federal University of Paraná, Curitiba, Paraná, Brazil<sup>d</sup>; Instituto Carlos Chagas, Fundação Oswaldo Cruz, FIOCRUZ, Curitiba, Paraná, Brazil<sup>e</sup>; CBS-KNAW Fungal Biodiversity Centre, Utrecht, the Netherlands<sup>f</sup>; Department of Parasitology and Mycology, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran<sup>g</sup>

**ABSTRACT** A draft genome sequence of type strain *Fonsecaea multimorphosa* CBS 980.96<sup>T</sup> was obtained. This species was first isolated from a cat with cerebral phaeohyphomycosis in Queensland, Australia.

Pathogenic species of the *Herpotrichiellaceae* are able to cause mutilating skin infections known as chromoblastomycosis as well as disseminated phaeohyphomycosis, and a few species such as *Rhinochrysiella mackenziei* and *Cladophialophora bantiana* can cause cerebral phaeohyphomycosis (1, 2). Clinical symptoms commonly found in patients with cerebral phaeohyphomycosis are severe headache, seizures, paralysis of one or more oculomotor muscles, and weight loss, leading to death (3). Once the fungus reaches the nervous system, the excessive production of melanin leads to inflammation and necrosis of brain tissue. Cerebral phaeohyphomycosis caused by *Fonsecaea* species are rare (4), but *Fonsecaea monophora* has been repeatedly isolated from humans (5) and *F. pugnacious* was isolated only once (6). *F. multimorphosa* CBS 980.96<sup>T</sup> was isolated from an abscess in the left occipital lobe of the cerebrum of a female cat in Queensland, Australia (3). The feline was 18 months of age and had dilated pupils with no reflection, enlarged liver, inflamed bile duct, and congested lungs, with no cutaneous and/or subcutaneous lesions (3).

*F. multimorphosa* CBS 980.96<sup>T</sup> was grown on Sabouraud's glucose agar (SGA) at 28°C for 7 days. DNA was extracted using the CTAB (cetyltrimethylammonium bromide) method and phenol-chloroform/isoamyl alcohol. A microbial DNA UltraClean kit was used for purification of total DNA. A DNA library for sequencing was prepared using the Nextera kit (Illumina) following the manufacturer's recommendations, and sequenced on a MiSeq sequencer using the paired-end (2 × 250 bp) sequencing method. A quality check of the sequenced reads was done using FastQC (7). SPAdes version 3.6.2 (8) was used for *de novo* assembly. Gap closure was performed using FGAP (9) and assembly coverage was determined using Bowtie2 (10). Gene prediction was performed using GeneMarkES version 4.29 (11). Annotation of the protein coding genes was based on best-hit searches against the NR database using RAFTS3 (12) and InterProScan (13). Agoron version 1.2.36 (14) was used for tRNA gene identification.

Received 8 December 2016 Accepted 11 December 2016 Published 16 February 2017

**Citation** Leao ACR, Weiss VA, Vicente VA, Costa F, Bombassaro A, Raittz RT, Steffens MBR, Pedrosa FO, Gomes RR, Baura V, Faoro H, Sfeir MZT, Balsanelli E, Moreno LF, Najafzadeh MJ, de Hoog S, Souza EM. 2017. Genome sequence of type strain *Fonsecaea multimorphosa* CBS 980.96<sup>T</sup>, a causal agent of feline cerebral phaeohyphomycosis. *Genome Announc* 5: e01666-16. <https://doi.org/10.1128/genomeA.01666-16>.

**Copyright** © 2017 Leao et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Sybren de Hoog, s.hoog@cbs.knaw.nl, or Emanuel Maltempo Souza, souzaem@ufpr.br.

A total of 7,445,102 paired-end reads were obtained for *F. multimorphosa* CBS 980.96<sup>T</sup> genome sequencing. The reads were assembled into 163 scaffolds with a G+C content of 52.64% and an  $N_{50}$  of 392,992 bp. The genome size was estimated to be 33.4 Mbp. More than 97% of the reads were used for assembly yielding a genome coverage of 54×. A total of 11,772 protein-coding genes with at least one conserved domain were predicted and 32 tRNAs. The genes were also annotated based on a search of similarity with known proteins of other fungal species.

The genome sequence of *F. multimorphosa* CBS 980.96<sup>T</sup> will be important for a better understanding of the basic mechanisms of adaptation to hostile environments, pathogenicity, and virulence of this organism.

**Accession number(s).** The whole-genome shotgun project of *F. multimorphosa* CBS 980.96<sup>T</sup> has been deposited at DDBJ/EMBL/GenBank under the accession number [LVCI00000000](https://www.ncbi.nlm.nih.gov/nuclink/LVCI00000000).

## ACKNOWLEDGMENTS

We thank Roseli Prado and Marilza Doroti Lamour for technical assistance.

This project was supported by the INCT of Biological Nitrogen Fixation/CNPq/MCT, Fundação Araucária, Coordination for the Improvement of Higher Education Personnel (CAPES) and CBS-Fungal Biodiversity Center, Utrecht, the Netherlands.

## REFERENCES

- Vicente VA, Najafzadeh MJ, Jiufeng S, Gomes RR, Robl D, Marques SG, Azevedo CMPS, de Hoog GS. 2013. Environmental siblings of black agents of human chromoblastomycosis. *Fungal Divers* 65:1–17. <https://doi.org/10.1007/s13225-013-0246-5>.
- Doymaz MZ, Seyithanoglu MF, Hakyemez İ, Gulpepe BS, Cevik S, Aslan T. 2015. A case of cerebral phaeohyphomycosis caused by *Fonsecaea monophora*, a neurotropic dematiaceous fungus, and a review of the literature. *Mycoses* 58:187–192. <https://doi.org/10.1111/myc.12290>.
- Shinwari MW, Thomas AD, Orr JS. 1985. Feline cerebral phaeohyphomycosis associated with *Cladosporium bantianum*. *Aust Vet J* 62:383–384. <https://doi.org/10.1111/j.1751-0813.1985.tb14216.x>.
- Najafzadeh MJ, Vicente VA, Sun J, Meis JF, de Hoog GS. 2011. *Fonsecaea multimorphosa* sp. nov., a new species of *Chaetothyriales* isolated from a feline cerebral abscess. *Fungal Biol* 115:1066–1076. <https://doi.org/10.1016/j.funbio.2011.06.007>.
- Surash S, Tyagi A, de Hoog GS, Zeng JS, Barton RC, Hobson RP. 2005. Cerebral phaeohyphomycosis caused by *Fonsecaea monophora*. *Med Mycol* 43:465–472. <https://doi.org/10.1080/13693780500220373>.
- de Azevedo CMPS, Gomes RR, Vicente VA, Santos DWCL, Marques SG, do Nascimento MMF, Andrade CEW, Silva RR, Queiroz-Telles F, de Hoog GS. 2015. *Fonsecaea pugnacius*, a novel agent of disseminated chromoblastomycosis. *J Clin Microbiol* 53:2674–2685. <https://doi.org/10.1128/JCM.00637-15>.
- Andrews S. 2010. FastQC: A quality control tool for high throughput sequence data. <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Piro VC, Faoro H, Weiss VA, Steffens MBR, Pedrosa FO, Souza EM, Raittz RT. 2014. FGAP: an automated gap closing tool. *BMC Res Notes* 7:371. <https://doi.org/10.1186/1756-0500-7-371>.
- Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with bowtie 2. *Nat Methods* 9:357–359. <https://doi.org/10.1038/nmeth.1923>.
- Ter-Hovhannissyan V, Lomsadze A, Chernoff YO, Borodovsky M. 2008. Gene prediction in novel fungal genomes using an ab initio algorithm with unsupervised training. *Genome Res* 18:1979–1990. <https://doi.org/10.1101/gr.081612.108>.
- Viale RA, Pedrosa FO, Weiss VA, Guizelini D, Tibaes JH, Marchaukoski JN, Souza EM, Raittz RT. 31 May 2016. RAFTS3: rapid alignment-free tool for sequences similarity search. bioRxiv. <https://doi.org/10.1101/055269>.
- Quevillon E, Silventoinen V, Pillai S, Harte N, Mulder N, Apweiler R, Lopez R. 2005. InterProScan: protein domains identifier. *Nucleic Acids Res* 33:W116–W120. <https://doi.org/10.1093/nar/gki442>.
- Laslett D, Canback B. 2004. ARAGORN, a program to detect tRNA genes and tmRNA genes in nucleotide sequences. *Nucleic Acids Res* 32:11–16. <https://doi.org/10.1093/nar/gkh152>.