



Contents lists available at ScienceDirect

International Journal of Surgery Case Reports

journal homepage: www.casereports.com

Endovascular repair for an extracranial internal carotid aneurysm with cervical access: A case report



Ignacio J. Rivera-Chavarría*, Juan C. Alvarado-Marín

Division of Vascular Surgery, Rafael Ángel Calderón Guardia Hospital, Caja Costarricense del Seguro Social, San José, Costa Rica

ARTICLE INFO

Article history:

Received 11 September 2015

Received in revised form

29 November 2015

Accepted 29 November 2015

Available online 3 December 2015

Keywords:

Carotid aneurysm

Stentgraft

Endovascular therapy

Cervical access

ABSTRACT

BACKGROUND: Carotid aneurysms are a rare pathology. This vascular disorder can be asymptomatic or it can cause local compression. The disorder poses a high risk of embolization and rupture.

PRESENTATION OF CASE: A 79 years old female, presents with a right internal carotid fusiform aneurysm, approximately 3.8 cm in diameter, localized 3.30 cm from the common carotid artery bifurcation with an extremely tortuous common carotid artery.

DISCUSSION: Surgical management of the extracranial internal carotid artery remains varying and challenging, particularly with a distal internal carotid aneurysm and with anatomical difficulties.

CONCLUSION: Endovascular management of an internal carotid aneurysm with cervical access using an expanded polytetrafluoroethylene covered stent with Heparin Bioactive Surface in the carotid area, is safe and effective.

© 2015 The Authors. Published by Elsevier Ltd. on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Carotid aneurysms are a rare pathology, with an incidence of 0.4–4% of all aneurysms [1–3]. Occurrence of isolated aneurysms of the extracranial internal carotid artery (EICA) is even more rare. This vascular disorder can be asymptomatic or it can cause local compression. The disorder poses a high risk of embolization and rupture [4,5].

Surgical management of EICA remains varying and challenging [6]. Furthermore, when a distal internal carotid aneurysm and with an extremely tortuous common carotid artery, it is particularly difficult to use an open surgical or an endovascular approach.

2. Presenting concerns

The subject of this report is a 79-year-old white, married, non-drinking, non-smoking female, with hypertension, no history of trauma or surgery, who presents with a pulsatile cervical mass.

3. Clinical findings

A pulsatile retromandibular mass with systolic bruit in the right neck. The patient was evaluated by neurology department with




Fig. 1. Computed tomography: with right internal carotid fusiform aneurysm and the extremely tortuous of the common carotid artery.

* Corresponding author at: Servicio Vascular Periférico, Hospital Rafael Ángel Calderón Guardia, Barrio Aranjuez, San José 10101, Costa Rica.

E-mail address: riveraignacio@gmail.com (I.J. Rivera-Chavarría).

4. Diagnostic focus and assessment

The computed tomography shows a right internal carotid fusiform aneurysm of approximately 3.8 cm in diameter, localized 3.30 cm from the common carotid artery bifurcation with an extremely tortuous common carotid artery (Fig. 1).

5. Therapeutic focus and assessment

With an appropriate informed consent, we took the patient to a hybrid operating room. The procedure was performed with the patient under general anesthesia and endotracheal intubation.

The right common carotid artery was exposed through a 3 cm length incision in the anterior edge of the sternocleidomastoid muscle at the low neck. An 8 French introducer sheath (Avanti®) was inserted in the common carotid artery (Fig. 2), and 5000 Units of heparin was injected through the sheath. An angiography was performed (Fig. 3A and C). The internal carotid artery was canalized distal to the aneurysm with the cerebral emboli protection system (Filterwire EZ®) and was deployed just before the carotid siphon. The expanded polytetrafluoroethylene (ePTFE) covered stent with Heparin Bioactive Surface (Gore® VIABAHN® Endoprosthesis with Propaten® Bioactive Surface) was introduced and placed in the right position. A control angiography was performed with excellent results (Fig. 3B and D) and the cerebral protection system was removed.




Fig 2. Surgical approach with a 3 cm incision; 8 French introducer sheath in the common carotid artery.

The day after, the patient was evaluated by the same neurologist resulting in a NIHSS=0, and the patient was discharged the same day.

6. Discussion

Extracranial internal carotid aneurysms are extremely rare, most of them are asymptomatic, but have a potential in neuro-




Fig 3. A and C: angiography, right internal carotid fusiform aneurysm. B and D: control angiography after stenting.

logic morbimortality [6]. Historically the treatment is open surgical resection and reconstruction [7–9].

According to etiology, aneurysms can be classified as atherosclerotic, post-traumatic, dissecting or mycotic (infected) aneurysms or those associated with cystic medial necrosis, Marfan's syndrome and fibromuscular dysplasia. Other rare causes include tuberculosis, Takayasu's arteritis and human immunodeficiency virus-related arteritis [10].

The treatment of this kind of aneurysms presents a challenge for the vascular surgeon, even more when they are distal to the common carotid bifurcation that makes a difficult open surgery approach and with extremely tortuous common carotid artery or unfavorable aortic arc anatomy, that makes endovascular reach of the lesion impossible.

This case suggests that cervical access for the endovascular exclusion of an internal carotid aneurysm with ePTFE-covered stent with Heparin Bioactive Surface is an effective and safe procedure for the management of this pathology, especially with anatomical difficulty. This technique reduces operative time and makes a less hazardous dissection. This case represents an off-label use.

7. Conclusion

We report an endovascular management procedure of an internal carotid aneurysm with cervical access, and we demonstrate that the use ePTFE-covered stent with Heparin Bioactive Surface in carotid area is safe and effective.

Patient consent

The patient provided written permission for the publication of this case report.

Conflict of interest

Dr. Ignacio Rivera-Chavarría and Dr. Juan C. Alvarado-Marín have no conflict of interest.

Consent

We appropriate obtained written consent from the patient before the procedure.

Author contribution

Dr. Ignacio Rivera-Chavarría: Principal surgeon, writing the paper, final approval Dr. Juan C. Alvarado-Marín: assist the surgery, final approval.

Guarantor

Dr. Ignacio Rivera-Chavarría and Dr. Juan C. Alvarado-Marín.

References

- [1] K. Kaouel, S. Mehergui, I. Ben Mrad, M. Ben Mrad, F. Ghédira, H. Mizouni, et al., Traitement chirurgical des anévrismes carotidiens extracrâniens à propos de dix cas, *J. Mal. Vasc.* 37 (2012) 201–206.
- [2] K. Hosoda, S. Fujita, T. Kawaguchi, Y. Shibata, N. Tamaki, The use of an external-internal shunt in the treatment of extracranial internal carotid artery vascular aneurysms: technical case report, *Surg. Neurol.* 52 (1999) 153–155.
- [3] N.R. Hertzner, Extracranial carotid aneurysms: a new look at an old problem, *J. Vasc. Surg.* 31 (2000) 823–825.
- [4] R. Argenta, S. Braun, Surgical repair of an extracranial carotid aneurysm, *J. Vasc. Bras.* 14 (2015) 84–87.
- [5] J. Kakisis, T. Giannakopoulos, K. Moulakakis, C. Liapis, Extracranial internal carotid artery aneurysm, *J. Vasc. Surg.* 60 (2014) 1358.
- [6] J. Wu, Y. Chen, L. Qu, Y. He, K. Zhi, J. Bai, et al., Using PTFE covered stent-artery anastomosis in a new hybrid operation for giant juxta-skull internal carotid aneurysm with tortuous internal carotid artery, *Int. J. Cardiol.* 185 (2015) 25–28.
- [7] G.M. Longo, M.R. Kibbe, Aneurysms of the carotid artery, *Semin. Vasc. Surg.* 18 (2005) 178–183.
- [8] R. El-Sabrou, D.A. Cooley, Extracranial carotid artery aneurysms: Texas Heart Institute experience, *J. Vasc. Surg.* 31 (2000) 701–712.
- [9] N. Attigah, S. külkens, N. Zausig, J. Hansmann, P. Ringleb, M. Hakimi, et al., *Eur. J. Vasc. Endovasc. Surg.* 37 (2009) 127–133.
- [10] A.S. Mačina, V. Vidjak, L. Erdelez, S. Suknaic í, S. Gašparov, K. Karlo Novačić í, Open surgical and endovascular treatment of extracranial internal carotid artery, *Perspect. Vasc. Surg. Endovasc. Ther.* 21 (2009) 181–185.

Open Access

This article is published Open Access at sciedirect.com. It is distributed under the [IJSCR Supplemental terms and conditions](#), which permits unrestricted non commercial use, distribution, and reproduction in any medium, provided the original authors and source are credited.