



Prevalence and phylogenetic analysis of porcine deltacoronavirus in Sichuan province, China

Yu Feng¹ · Zhiwen Xu^{1,2} · Ling Zhu^{1,2}

Received: 23 October 2019 / Accepted: 4 August 2020 / Published online: 6 September 2020
© Springer-Verlag GmbH Austria, part of Springer Nature 2020

Abstract

In order to understand the prevalence and genetic diversity of porcine deltacoronavirus (PDCoV) in diarrhoeal pigs in Sichuan province, 634 clinical samples were collected from individual pigs with diarrhoea in 13 regions of Sichuan province, China, from January 2017 and June 2019. The detection results showed that the infection rate of PDCoV was relatively low in diarrhoeal pigs, 13.25% (84/634), but the infection rate of PEDV (porcine epidemic diarrhea virus) was high, 32.18% (204/634). Coinfection with PEDV was common (55.95%, 47/84) in PDCoV-infected diarrhoeal pigs. Additionally, the chance of PDCoV infection was 2.77 times higher in suckling piglets than in sows, and about 3.30 times higher in spring and winter than in summer. PDCoV/PEDV coinfection was 75% less likely in sows than in suckling piglets. The complete genomes of four Sichuan PDCoV strains were sequenced and analysed. There were some insertion-deletion signatures in the whole genome sequences of four strains, including a 6-nt deletion in the non-structural gene 2 region, a 9-nt insertion in the non-structural gene 3 region, a 3-nt deletion in the S gene region, and a distinguishing 11-nt deletion in the 3'UTR region. Phylogenetic analysis based on complete genome sequences revealed that the PDCoV Sichuan strains were closely related to other Chinese PDCoV reference strains; however, phylogenetic analysis based on S gene sequences showed that the CH/SC/2019 strain clustered in a large clade with strains from the USA, Japan, and Korea. These data advance our understanding of the genetic diversity and evolutionary characteristics of PDCoV in China and may contribute to vaccine development.

Introduction

Porcine deltacoronavirus (PDCoV), a member of the genus *Deltacoronavirus*, is a single-stranded, positive-sense enveloped RNA virus that causes enteritis and watery diarrhoea in pigs, especially nursing piglets [1]. The full genome of PDCoV is approximately 25.4 kb long, which is the smallest genome size among the porcine coronaviruses, arranged in the order 5' untranslated region (UTR), open reading frame 1a and b (ORF1ab), spike (S), envelope (E), membrane (M), non-structural gene 6 (NS6), nucleocapsid (N), non-structural gene 7 (NS7), and 3'UTR [2]. The PDCoV S gene is frequently used for studying genetic relationships among

different PDCoV strains and for performing epidemiological investigations [3].

PDCoV was first discovered in Hong Kong, and has since been reported in many countries, including the USA, Canada, Korea, China, Thailand, Laos, and Vietnam [4–9]. In China, PDCoV had been reported in many provinces, but little has been reported regarding the prevalence and epidemiology of PDCoV in Sichuan province, a major pig-raising province of China. Thus, in this study, we investigated the prevalence of PDCoV in diarrhoeal pigs and analysed the complete genome sequences of PDCoV isolates from Sichuan, China.

Materials and methods

Sampling

In this study, a total of 634 intestinal and faecal clinical samples (405 intestinal samples and 229 faecal samples) were collected from pigs with diarrhoea outbreaks diagnosed by a pig farm veterinarian from January 2017 and June 2019. All

Handling Editor: Tim Skern.

✉ Zhiwen Xu
abtctxw@126.com

¹ College of Veterinary Medicine, Sichuan Agricultural University, No. 211 Huiming Road, Chengdu 611130, China

² Animal Biology Technology Center, Sichuan Agricultural University, Chengdu, China

samples were collected from suckling piglets and sows and from pig farm in 13 regions in Sichuan province. The sampling locations are shown in Table 1. Soon after sampling, faeces and intestinal contents were prepared as 10% suspensions in phosphate-buffered saline (PBS). The suspensions were then vortexed and centrifuged at 8000 *g* at 4°C for 10 min. The clarified supernatants were stored at -80°C until use for RNA extraction. Total RNAs were extracted using TRIzol Reagent (Invitrogen, USA) according to the manufacturer's instructions. RT-qPCR was carried out to assess the frequency of PDCoV mono-infections and coinfections with PEDV and transmissible gastroenteritis virus (TGEV). We used the virotype PEDV/TGEV/PDCoV RT-PCR Rgt (QIAGEN) following the manufacturer's protocol.

Complete genome sequences and phylogenetic analysis

Representative PDCoV strains from four different pig farms where severe or moderate diarrhoea had been reported were selected for sequencing of the complete genome by RT-PCR amplification of 24 regions covering the PDCoV genome as described previously [10]. These representative PDCoV strains were named CH/SC/2017, CH/SC/2018/1, CH/SC/2018, and CH/SC/2019 (GenBank accession nos. MK211169, MK005882, MK330605, and MK993519). The PCR products were purified and cloned into the vector pMD18-T. Positive clones of each amplicon was sequenced twice (Sangon Biotech, Shanghai, China). At least three independent PCR amplicons were sequenced to obtain a consensus sequence. Complete sequences of four PDCoV strains were assembled by combining overlapping contigs and trimming off primer sequences using the DNASTAR software package (DNASTAR Inc., Madison, WI, USA). Sequence alignment analysis of these four strains and other PDCoV strains with sequences in the GenBank database was conducted using the Megalign program. Phylogenetic trees based on complete genome and S gene sequences were constructed by the maximum-likelihood method using MEGA software version 6 with bootstrap analysis of 1,000 replicates.

Statistical analysis

The rates of infection with PDCoV and coinfection with PDCoV and PEDV in diarrhoeal pigs were analysed according to age, year, and season. Significance was set at $p < 0.05$. Possible risk factors associated with virus infection were analysed by univariate and multivariate exact logistic regression models using Stata/SE 15.1 (Stata Corporation, USA). For each model the odds ratio (OR) and 95% confidence interval (CI) were calculated.

Results

Prevalence of PDCoV

Of the 634 clinical samples tested in our PDCoV prevalence survey, 84 (13.25%) were positive for PDCoV and 204 (32.18%) were PEDV positive. The positive rate for PDCoV in sows and piglets was 6.80% (14/206) and 16.36% (70/428), respectively. All of the samples were negative for TGEV, as shown in Table 2. The results indicate that the prevalence rate of PDCoV was relatively low in diarrhoeal pigs in Sichuan and that PEDV was the main pathogen causing diarrhoea in pigs. It is also noteworthy that 47 samples that were positive for PDCoV were also positive for PEDV, showing that mixed infections with these viruses are relatively common. The PDCoV infection rates in different regions of Sichuan province are shown in Table 1. Twelve out of the 13 regions were considered PDCoV positive, including the regions of Mianyang (14.93%), Deyang (16.67%), and Guan'an (22.73%), suggesting that PDCoV is widespread on pig farms in Sichuan province.

As shown in Table 3, The results show that the prevalence of PDCoV in diarrhoeal pigs increased from 10.53% in 2017 to 17.24% in 2019; however, based on statistical analysis, we could not conclude that PDCoV infection in diarrhoeal pigs has been increasing from year to year ($p > 0.05$). More clinical samples are needed to determine whether the PDCoV infection rate is correlated with the

Table 1 The number of samples and PDCoV positive rates in different regions in Sichuan, China

Region	Number of samples	PDCoV positive (%)	Region	Number of samples	PDCoV positive (%)
Mian Yang	67	10 (14.93%)	Ya An	55	5 (9.09%)
Guan Yuan	30	4 (13.33%)	De Yang	48	8 (16.67%)
Guan An	44	10 (22.73%)	Nan Chong	93	13 (13.98%)
Sui Ning	23	2 (8.70%)	Zi Yang	50	8 (16.00%)
Mei Shan	49	6 (12.24%)	Da Zhou	41	7 (17.07%)
Le Shan	31	0 (0%)	Nei Jiang	25	3 (12.00%)
Yi Bing	78	8 (10.26%)	Total	634	84 (13.25%)

Table 2 Detection of PDCoV, PEDV, and TGEV in diarrhoeal samples from pigs in Sichuan, China, 2017-2019

Year	Virus	Sows			Suckling piglets			Total		
		Number of samples	Positive	Positive rate, %	Number of samples	Positive	Positive rate, %	Number of samples	Positive	Positive rate, %
2017	PDCoV	76	5	6.58	152	19	12.50	228	24	10.53
	PEDV		17	22.37		53	34.87		70	30.70
	TGEV		0	0		0	0		0	0
	PDCoV+PEDV		1	1.32		13	8.55		14	6.14
2018	PDCoV	94	6	6.38	196	34	17.35	290	40	13.79
	PEDV		21	22.34		71	36.22		92	31.72
	TGEV		0	0		0	0		0	0
	PDCoV+PEDV		3	3.19		23	11.73		26	8.97
2019	PDCoV	36	3	8.33	80	17	21.25	116	20	17.24
	PEDV		10	27.78		32	40.00		42	36.21
	TGEV		0	0		0	0		0	0
	PDCoV+PEDV		0	0		7	8.75		7	6.03
2017-2019	PDCoV	206	14	6.80	428	70	16.36	634	84	13.25
	PEDV		48	23.30		156	36.45		204	32.18
2019	TGEV		0	0		0	0		0	0
	PDCoV+PEDV		4	1.94		43	10.05		47	7.41

Table 3 Analysis of univariate and multivariate exact logistic regression models of possible risk factors associated with PDCoV and PDCoV-PEDV coinfection

Variable	OR	95%CI	<i>p</i> -value
Univariate			
Age			
Sow	1	ref	
Piglet	2.77	1.52-5.04	0.001
Year			
2017	1	ref	
2018	1.36	0.79-2.33	0.263
2019	1.77	0.93-3.36	0.081
Season			
Summer	1	ref	
Spring	3.30	1.12-9.65	0.030
Autumn	1.85	0.57-5.94	0.304
Winter	3.32	1.19-9.13	0.022
Multiple model			
Age of piglet	2.67	1.46-4.87	0.001
Spring season	3.24	1.32-8.78	0.028
Winter season	3.30	1.40-9.08	0.022
Coinfection multivariate model			
Age of piglet	3.98	1.14-13.97	0.031

ref, reference variable; OR, odds ratio; CI, confidence interval

year of sample collection. Moreover, analysis of individual variables revealed that suckling piglets were 2.77 (OR) times more likely to be infected than sows. Additionally,

the PDCoV positive rates in spring (15.46%, 32/207) and winter (16.22%, 36/222) were clearly higher than in summer (5.26%, 4/76). Pigs had about 3.30 (OR) times more risk of PDCoV infection in spring and winter than in summer. PDCoV/PEDV coinfection was statistically correlated with age. Coinfection was 75% less likely in sows than suckling piglets.

Complete genomic characterisation of Sichuan PDCoV strains

The complete genomes of four novel PDCoV strains, representative of epidemic strains, were sequenced and analysed to perform a molecular characterisation of PDCoV in Sichuan. The results showed that the complete genomic sequences of strains CH/SC/2017, CH/SC/2018/1, CH/SC/2018, and CH/SC/2019 were 25,402 nt, 25,413 nt, 25,414 nt, and 25,392 nt in length, respectively, and all had a typical PDCoV genome organisation. Sequence alignment analysis showed that the CH/SC/2017 and CH/SC/2019 strains both had an 11-nt deletion in the 3'UTR region when compared with strains CH/SC/2018 and CH/SC/2018/1 (Fig. 1). All four sequenced strains had a 6-nt deletion in the non-structural gene (ns) 2 region (ORF1ab), which has been observed in some other Chinese strains. Additionally, CH/SC/2019 had a 9-nt deletion in the ns 3 region (ORF1ab) when compared with most Chinese strains. This deletion is also present in strain CH/Sichuan/S27/2012, detected previously in Sichuan, and in strains from Thailand

		ns 2		ns 3		S gene		3'UTR	
		1736	1750	2808	2822	19468	19482	25042	25055
China	HKU15-44	CAGTTTGAAGATCCG		GAGCCGGTTGGTAAG		GCTAATAATAATTTT		TGTCGT-TAAACCC	
	HKU15-155A..C..C..T	-C.....	
	CHN-HBC..C..T	-C.....	
	CHN-HNA..C..C..T		
	CH-HuNanC..C..T	-CT.....	
	▲ Sichuan/2017C..T	-C.....	
	▲ Sichuan/2018C..T	-C.....	
	▲ Sichuan/2018/1C..T	-C.....	
	▲ Sichuan/2019-G..	-C.....	
	CH-JSJGS02	...A.....	C..C..T	-C.....	
	CHN-SXD	...A.....	C..C..T	-C.....	
	CHJXNI2	...A.....	C..C..T	-C.....	
	SHJS/SLC..C..T	-C.....	
	CHN-HeB1C..C..T	-C.....	
	CHN-SDC..C..T	-C.....	
	Sichuan/S27-G..	-C.....	
	CHN-GD	...A.....	C..C..T	-C.....	
	CHN-AHC.....	
	CHN/GSCG.....	
	CHN/QHCG.....	
Thailand	TT_1115-G..		...C.....-C...	C..C..C..A..	
	S5011-G..		...C.....	C..C..C..A..	
	S5011L-G..		...C.....	C..C..C..A..	
Vietnam	Binh21-G..	C..CT.....	
	HaNoi6-G..	C..CT.....	
Japan	YMG/JPNC..C..T		
	HKD/JPNC..T		
Korea	KNU14-04C..C..T		
	DH2C..C..T		
	KNU16-11C..C..T	A..	
USA	Lowa459C..C..T		
	Nebraska145C..C..T		
	IL2768C..C..T		
	8734/USA-IAC..C..T		
	IN2847C..C..T		
	KY4813C..C..T		
	OH1987C..C..T		
	Ohio137C..C..T		

Fig. 1 The four main deletions or insertions in the complete genome sequence alignment. A multiple sequence alignment was constructed using ClustalW in the DNASTAR software. The four PDCoV strains sequenced in this study are indicated by black triangles. A dot (●)

indicates that the nucleotide exactly matches the reference sequence. A dash (-) indicates that the nucleotide is deleted relative to the reference sequence

and Vietnam. Moreover, compared with strains from other countries, a 3-nt (TAA) deletion was observed in the S gene of the four novel strains, which is also present in most Chinese PDCoV strains, except early Chinese strains (CHN-GD and CHN-AH).

Pairwise nucleotide sequence comparisons showed that the four sequenced strains shared 98.9%-99.3% nucleotide sequence identity with each other, and shared 97.6%-99.0% nucleotide and 98.0%-99.5% aa sequence identity with reference PDCoV sequences. Interestingly, the full genomes of the CH/SC/2017, CH/SC/2018, and CH/SC/2018/1 strains all had low nucleotide sequence identity (97.6%) to strains

from Vietnam and Thailand, while the CH/SC/2019 strain shared relatively high nucleotide sequence identity (98.9%) with these strains, suggesting that the Sichuan PDCoV strains might have undergone genetic changes and continued to evolve.

Phylogenetic analysis of PDCoV Sichuan strains

Phylogenetic analysis was carried out using the whole-genome and S gene sequences of the four sequenced strains and other reference PDCoV strains (Fig. 2A and B). The phylogenies based on the complete genome sequences and the S gene both




Fig. 2 Maximum-likelihood phylogenetic analysis based on (A) full-length genome sequences and (B) S gene sequences of PDCoV isolates. The scale bar indicates the number of nucleotide substitutions per site. The four strains sequenced in this study are indicated by black triangles

revealed that the PDCoV strains showed obvious regional characteristics, as has been reported previously [11]. Phylogenetic analysis based on whole genome sequences indicated that the PDCoV CH/SC/2017, CH/SC/2018/1, and CH/SC/2018 strains grouped within the Chinese clade and are more closely related to strains SHJS/SL/2016 and CH/HN/2014. However, the CH/SC/2019 strain is located in another subcluster and is similar to the CH/QH/2017 and CH/GS/2017 strains, which were isolated from piglets in recent years, suggesting that the origin of these isolates may be different.

A phylogenetic tree based on S gene sequences revealed that CH/SC/2017, CH/SC/2018/1, and CH/SC/2018 belong to the same group, while the CH/SC/2019 strain clustered in a large clade with strains from the USA, Japan, and Korea and demonstrated genetic differences when compared to the other Sichuan strains.

Discussion

In this study, RT-PCR analysis of 634 clinical samples from pigs with diarrhoea collected from 13 regions in Sichuan province from January 2017 and June 2019 revealed a positive rate of PDCoV of 13.25%, while that of PEDV was 32.18%, indicating that PEDV is still the primary pathogen causing porcine diarrhoea, which is consistent with previous studies [12]. The prevalence of PDCoV in diarrhoeal pigs in Sichuan was somewhat lower than in other provinces, such as Henan (23.49%) and Guangdong (21.8%), but a molecular-based detection study showed a similar low prevalence of PDCoV infection in Jiangsu (4.94%) and Hubei (6.45%) [13]. The prevalence of PDCoV in Qinghai and Tibet is also low, with a

positive rate of only 3.70% [14]. These data show that the prevalence of PDCoV in diarrhoeal pigs differs among the Chinese provinces, and more information regarding the molecular epidemiology of PDCoV in China is needed. In this study, the number of samples collected and regions investigated was limited, and therefore more samples need to be collected and tested, and PDCoV surveillance should be continued. Also, since the aim of this study was only to monitor the prevalence of PDCoV in diarrhoeal pigs more samples, including those from pigs without diarrhoea, are needed to evaluate the prevalence of PDCoV in Sichuan pig herds.

Although the overall prevalence of PDCoV in this study was relatively low, PDCoV was detected in most regions of Sichuan, implying that PDCoV prevention measures have not been very effective, and further study is needed to prevent and control this novel enteric virus. Statistical analysis demonstrated a higher chance of PDCoV infection in winter and spring, which is consistent with previous studies [15], suggesting that we should pay more attention to preventing infections during these seasons. Furthermore, we detected PDCoV in sows, as described previously [16]. However, unlike a previous study, our results showed that suckling piglets had a 2.77 times higher risk of PDCoV infection than sows. Previous reports have suggested that older pigs are more susceptible to PDCoV than piglets, which was not confirmed by our results [17]. Therefore, more clinical samples from older pigs are needed to test this conclusion. We plan to continue to monitor PDCoV infections in sows. Moreover, it has been reported that PDCoV/PEDV coinfection is more common in Chinese pigs than PDCoV/porcine rotavirus coinfection in US pig herds [18, 19]. Our detection data are consistent with this, also showing that suckling piglets have a higher chance of PDCoV/PEDV coinfection.

The results of sequence analysis showed that the four strains had some unique deletions/insertions compared to other PDCoV strains. The CH/SC/2019 strain, like the CH/Sichuan/S27/2012 strain, had a 6-nt deletion in the ns 2 region and a 9-nt deletion in the ns 3 region, respectively, while the others strains only had a 6-nt deletion in ns 2. Moreover, four strains had a 3-nt deletion in the deletion of the S gene, resulting in one amino acid. This deletion exists in most Chinese PDCoV isolates. The S1 region is responsible for virus receptor binding [20], suggesting that this aa deletion may have some biological significance regarding viral replication or pathogenesis. Importantly, the CH/SC/2017 and CH/SC/2019 strains had a unique 11-nt deletion in the 3'UTR region. This is the first report of a PDCoV strain with such a deletion. However, we do not know whether these naturally occurring deletions or insertions have biological significance in PDCoV biology and pathogenesis. The four sequenced PDCoV strains were all collected from suckling piglets, but the clinical signs in

these piglets were different. The piglets infected with the CH/SC/2017 and CH/SC/2019 strains suffered from severe watery diarrhoea and died after three days of diarrhoea, while the suckling piglet with the CH/SC/2018 and CH/SC/2018/1 strains had moderate diarrhoea and then recovered. Were these different clinical signs associated with the 11-nt deletion in the 3'UTR or another deletion/insertion? We will continue to isolate PDCoV strains for further investigation.

A phylogenetic tree constructed based on complete genome sequences indicated that these strains might have evolved from different Chinese strains. In the phylogenetic tree based on the S gene, the Sichuan strains clustered in a different clade. The CH/SC/2017, CH/SC/2018/1, and CH/SC/2018 strains are in the China lineage, like most Chinese PDCoVs, while the CH/SC/2019 strain is in the USA/Japan/South Korea lineage. This is possibly due to recombination or separate evolution. Additional PDCoV sequences are needed to identify possible recombination and evolution events.

In summary, we report for the first time the prevalence and genetic properties of PDCoV in diarrhoeal pigs in Sichuan, which could provide further insights into the epidemiology and evolution of PDCoV in China.

Acknowledgements This research was supported by the Science and Technology Program of Sichuan Province: Development of Bivalent Live Vaccine for Porcine Epidemic Diarrhea and Porcine Pseudorabies (2017NZ0038), the National Key Research Program During the 13th Five-Year Plan Period of China: Integrated Prevention and Control Technology for Major Diseases of Commercial Pigs and Integration of Safe Drug Use Technology (2018YFD051102), and the Program Sichuan Veterinary Medicine and Drug Innovation Group of China Agricultural Research System (CARS-SVDIP).

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval This project was approved by the appropriate ethical review committee. The “Guidelines for Experimental Animals” of the Ministry of Science and Technology (Beijing, China) were followed.

References

1. Jung K, Hu H, Bryan E, Lu Z, Chepngeno J, Saif L (2015) Pathogenicity of 2 porcine deltacoronavirus strains in gnotobiotic pigs. *Emerg Infect Dis* 21(4):650–654
2. Hu H, Jung K, Vlasova N, Chepngeno J, Lu Z, Wang Q, Saif J (2015) Isolation and characterisation of porcine deltacoronavirus from pigs with diarrhea in the United States. *J Clin Microbiol* 53(3):1537–1548
3. Shang J, Zheng Y, Yang Y, Liu C, Geng Q, Tai W, Du L, Zhou Y, Zhang W, Li F (2017) Cryo-EM structure of porcine delta coronavirus spike protein in the pre-fusion state. *J Virol* 92(4):e01556-17

4. Woo C, Lau K, Lam S, Lau C, Tsang K, Lau H, Bai R, Teng L, Tsang C, Wang M, Zheng J, Chan H, Yuen Y (2012) Discovery of seven novel mammalian and avian coronaviruses in the genus deltacoronavirus supports bat coronaviruses as the gene source of alphacoronavirus and betacoronavirus and avian coronaviruses as the gene source of gammacoronavirus and deltacoronavirus. *J Virol* 86(7):3995–4008
5. Marthaler D, Raymond L, Jiang Y, Collins J, Rossow K, Rovira A (2014) Rapid detection, complete genome sequencing, and phylogenetic analysis of porcine deltacoronavirus. *Emerg Infect Dis* 20(8):1347–1350
6. Ajayi T, Dara R, Misener M, Pasma T, Moser L, Poljak Z (2018) Herd-level prevalence and incidence of porcine epidemic diarrhoea virus (PEDV) and porcine deltacoronavirus (PDCoV) in swine herds in Ontario, Canada. *Transbound Emerg Dis* 65(5):1197–1207
7. Lee S, Lee C (2014) Complete genome characterization of Korean porcine deltacoronavirus strain KOR/KNU14-04/2014. *Genome Announc* 2(6):e01191-14
8. Liu J, Zuo Z, Gu Y, Luo X, Shi K, Hou S, Zhong F, Fan H (2018) Isolation and phylogenetic analysis of porcine deltacoronavirus from pigs with diarrhoea in Hebei province, China. *Transbound Emerg Dis* 65(3):874–882
9. Saeng-Chuto K, Stott J, Wegner M, Senasuthum R, Tantituvanont A, Nilubol D (2017) Retrospective investigation and evolutionary analysis of a novel porcine deltacoronavirus strain detected in Thailand from 2008 to 2015. *Arch Virol* 162(7):2103–2108
10. Wang L, Byrum B, Zhang Y (2014) Detection and genetic characterization of deltacoronavirus in pigs, Ohio, USA. *Emerg Infect Dis* 20(7):1227–1230
11. Zhang J (2016) Porcine Deltacoronavirus: Overview of infection dynamics, diagnostic methods, prevalence and genetic evolution. *Virus Res* 226(2):71–84
12. Jang G, Lee K, Kim H, Lee C (2017) Prevalence, complete genome sequencing and phylogenetic analysis of porcine deltacoronavirus in South Korea, 2014–2016. *Transbound Emerg Dis* 64(5):1364–1370
13. Dong N, Fang L, Zeng S, Sun Q, Chen H, Xiao S (2015) Porcine deltacoronavirus in mainland China. *Emerg Infect Dis* 21(12):2254–2255
14. Wang M, Wang Y, Baloch R, Pan Y, Tian L, Xu F, Shivaramu S, Chen S, Zeng Q (2018) Detection and genetic characterization of porcine deltacoronavirus in Tibetan pigs surrounding the Qinghai-Tibet Plateau of China. *Transbound Emerg Dis* 65(2):363–369
15. Zhang HL, Liang QQ, Li BX, Cui XG, Wei XL, Ding QW, Wang YB, Hu H (2019) Prevalence, phylogenetic and evolutionary analysis of porcine deltacoronavirus in Henan province, China. *Prev Vet Med* 166:8–15
16. Mai K, Feng J, Chen G, Li D, Zhou L, Bai Y, Wu Q, Ma J (2018) The detection and phylogenetic analysis of porcine deltacoronavirus from Guangdong Province in Southern China. *Transbound Emerg Dis* 65(1):166–173
17. Suzuki T, Shibahara T, Imai N, Yamamoto T, Ohashi S (2018) Genetic characterization and pathogenicity of Japanese porcine deltacoronavirus. *Infect Genet and Evol* 61:176–182
18. Song D, Zhou X, Peng Q, Chen Y, Zhang F, Huang T, Zhang T, Li A, Huang D, Wu Q, He H, Tang Y (2015) Newly emerged porcine deltacoronavirus associated with diarrhoea in swine in China: identification, prevalence and full-length genome sequence analysis. *Transbound Emerg Dis*. 62(6):575–580
19. Marthaler D, Raymond L, Jiang Y, Collins J, Rossow K, Rovira A (2014) Rapid detection, complete genome sequencing, and phylogenetic analysis of porcine Deltacoronavirus. *Emerg Infect Dis* 20:1347–1349
20. Li W, Hulswit G, Kenney P, Widjaja I, Jung K, Alhama A, Dieren B, Kuppeveld M, Saif J, Bosch J (2018) Broad receptor engagement of an emerging global coronavirus may potentiate its diverse cross-species transmissibility. *Proc Natl Acad Sci USA* 115(22):E5135–E5143

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.