

# A Comparison of Different Immune Activation Strategies to Reverse HIV-1 Latency

Caroline C. Garliss, Abena K. Kwaa, and Joel N. Blankson

Center for AIDS Research, Department of Medicine, Johns Hopkins Medicine, Baltimore, Maryland, USA

Resting CD4+ T cells are the best characterized component of the latent reservoir. Activation of these CD4+ T cells is needed to optimize transcription and viral replication, and this strategy has been used to measure the inducible reservoir. There are several methods that can be used to activate CD4+ T cells, and in this study, we compared 3 different strategies: the combination of the lectin phytohaemagglutinin (PHA) and irradiated allogeneic feeders, a combination of PHA and a superagonistic anti-CD28 antibody, and the combination of the protein kinase C agonist phorbol 12-myristate 13-acetate and the calcium ionophore ionomycin. We show that each strategy induces a different pattern of expression of activation markers on CD4+ T cells. However, the different activation strategies induced similar frequencies of latently infected CD4+ T cells from people living with HIV on suppressive antiretroviral therapy regimens to produce replication-competent virus. Furthermore, the frequency of infectious units per million induced by each regimen was positively correlated with the copies of intact proviral DNA per million CD4+ T cells. Our results suggest that no single pattern of activation marker expression is most associated with latency reversal and demonstrate that different immune activation strategies reverse latency in a low frequency of CD4+ T cells that harbor intact proviral DNA.

**Keywords:** culture assay; cure; HIV; latency.

Latently infected CD4+ T cells are a major barrier to HIV eradication [1, 2] but the true size of the inducible HIV reservoir has been challenging to measure [3]. Recent studies have shown that the number of CD4+ T cells with intact proviral DNA far exceeds the frequency of cells that produce infectious virus in the quantitative viral outgrowth assay (QVOA) [4, 5]. One potential reason for this discrepancy is that the activation strategy used in the conventional QVOA does not fully reverse latency. Evidence for this hypothesis comes from studies showing that repeated activation of CD4+ T cells leads to an increase in the number of latently infected CD4+ T cells that produce virus [4, 6]. In this study, we compared different immune activation strategies in order to determine whether these strategies were more effective at reversing latency than the phytohaemagglutinin (PHA) and irradiated feeder combination used in the standard QVOA. We also compared the frequency of cells that produced replication-competent virus with each strategy to the number of copies of intact and total proviral DNA using the newly described intact proviral DNA assay (IPDA) [7]. Our results have implications for HIV cure studies.

## METHODS

### Patient Cohort

We enrolled HIV-1-infected study subjects on suppressive ART regimens. This study was approved by the Johns Hopkins Institutional Review Board. All patients and HIV-1-negative healthy donors (HD) provided written consent before participating in this study. The clinical characteristics of these patients are shown in Table 1.

### Isolation of CD4+ T Cells

Peripheral blood mononuclear cells (PBMCs) from People living with HIV (PLWH) were isolated by Ficoll gradient centrifugation. Bulk CD4+ T cells were isolated from PBMCs by negative selection (Human CD4+ T Cell Isolation Kit, Miltenyi Biotec, Waltham, MA, USA).

### PHA/Feeders QVOA

The standard QVOA was performed on CD4+ T cells as described [8, 9]. Cells from each subject were cultured in either a 6-well plate with replicates of  $1 \times 10^6$  cells per well (CP21, CP25, VC20) or a 24-well plate with replicates of  $2 \times 10^5$  cells per well (all other subjects) with a 10-fold excess of irradiated PBMCs (feeders) from a healthy donor and cultured in RPMI 1640 (Thermo Fisher Scientific, Halethorpe, MD, USA) containing 10% fetal bovine serum (Thermo Fisher Scientific), 100 U/mL IL-2 (Chiron, Emeryville, CA, USA), 1% Pen Strep (Thermo Fisher Scientific), 1% T cell growth factor (activation media), and 0.5  $\mu$ g/mL of PHA (Remel, Thermo Fisher Scientific). PHA was washed off after 16 hours, and 2–4e6

Received 6 January 2020; editorial decision 25 February 2020; accepted 27 February 2020.

Correspondence: Joel N. Blankson, MD, PhD, Johns Hopkins Medicine, Baltimore, Maryland (jblanks@jhmi.edu).

Open Forum Infectious Diseases®

© The Author(s) 2020. Published by Oxford University Press on behalf of Infectious Diseases Society of America. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial reproduction and distribution of the work, in any medium, provided the original work is not altered or transformed in any way, and that the work is properly cited. For commercial re-use, please contact journals.permissions@oup.com  
 DOI: 10.1093/ofid/ofaa082

**Table 1. Clinical Characteristics of the Subjects in this Study; Total and Intact DNA Values Were Calculated by the Intact Proviral DNA Assay**

Subject	Current CD4 T-Cell Count, Cells/ $\mu$ L	HIV-1 RNA, Copies/mL	Time on Suppressive Regimen	Current Regimen	Intact DNA, Copies/Million Cells	Total DNA, Copies/Million Cells
CP11	1086	<20	8	FTC, TAF, EVG/c	82.82	441.07
CP21	525	<20	5	FTC, TAF, RPV	36.95	602.52
CP25	670	<20	1	FTC, TAF, RAL	23.67	532.48
CP39	829	<20	1	3TC, ABC, DTG	35.55	708.08
CP67	711	<20	6	FTC, TAF, EVG/c	13.66	518.33
CP68	599	<20	4	FTC, TAF, BTG	30.62	546.70
CP69	1395	<20	2	FTC, TAF, EVG/c	67.90	1348.72
CP70	805	<20	8	FTC, TAF, EVG/c	120.16	1004.87
CP71	561	<20	8	3TC, ABC, DTG	67.13	738.48
VC10	563	<20	3	TAF, FTC, DTG	3.84	129.18
VC20	575	<20	6	TAF, FTC, BTG	1.82	310.96

Abbreviations: 3TC, lamivudine; ABC, abacavir; BTG, bictegravir; CP, chronic progressor; DTG, dolutegravir; EVG/c, cobicistat boosted elvitegravir; FTC, emtricitabine; RAL, raltegravir; RPV, rilpivirine; TAF, tenofovir alafenamide; VC, viremic controller.

lymphoblasts from healthy donors were added to each 6-well plate well, whereas 0.5–1e6 lymphoblasts were added to each 24-well plate well, as previously described [8, 9]. A second round of lymphoblasts was then added on day 7 after the cultures were split. Supernatants from each well were tested for HIV-1 p24 protein by enzyme-linked immunosorbent assay (ELISA; Perkin Elmer, Waltham, MA, USA) on day 14 and repeated on day 21 of culture. The well size used for each subject was based on prior measurements of the subject's reservoir size.

#### Phorbol 12-Myristate 13-Acetate/Ionomycin QVOA

CD4+ T-cells from each subject were cultured in a T25 flask at 1–2e6 cells/mL in activation media with 50 ng/mL of phorbol 12-myristate 13-acetate (PMA; Sigma, St. Louis, MO, USA) and 1  $\mu$ m ionomycin (Sigma). After 12 hours of stimulation, the cells were washed twice and cultured in either a 12-well plate with replicates of 1  $\times$  10e6 cells per well (CP21, CP25, VC20) or a 48-well plate with replicates of 2  $\times$  10e5 cells per well (all other subjects); 2–4e6 lymphoblasts from healthy donors were added to each 12-well plate well, and 0.5–1e6 lymphoblasts were added to each 48-well plate well. The second round of lymphoblasts was added on day 7 after the cultures were split. Supernatants from each well were tested for HIV-1 p24 protein by ELISA (Perkin Elmer) on day 14; this was repeated on day 21 of culture. The well size used for each subject was based on prior measurements of the subject's reservoir size.

#### PHA/anti-CD28 QVOA

CD4+ T cells from each study subject were cultured in either a 96-well plate with replicates of 2  $\times$  10e5 cells per well (CP11, CP70, CP71) or a 48-well plate with replicates of 1  $\times$  10e6 cells per well (all other subjects) in activation media containing 1  $\mu$ g/mL of PHA (Remel) and 5  $\mu$ g/mL of the superagonistic anti-CD28 monoclonal antibody (Ansell, Bayport, MN, USA clone ANC28.1) [10]. Two days later, the PHA and anti-CD28 were washed off, and the replicates in 48-well plates were transferred into 12-well plates,

whereas those from 96-well plates were transferred into 48-well plates. Subsequently, 2–4e6 lymphoblasts from healthy donors were added to each well of a 12-well plate, and 0.5–1e6 lymphoblasts were added to each well of a 48-well plate. The second round of lymphoblasts was added on day 7 after the cultures were split. Supernatants from each well were tested for HIV-1 p24 protein by ELISA (Perkin Elmer) on days 14 and 21 of culture. The well size used for each subject was based on prior measurements of the subject's reservoir size.


**Flow Cytometry** To compare the CD4+ T-cell activation levels between the 3 assays, unfractionated CD4+ T cells were isolated from HLA-A2-positive individuals (3 CP and 7 HD). The cells were either untreated or stimulated with PHA/feeders, PMA/ionomycin, or PHA/anti-CD28 as above. On days 1 through 4, cells were stained with the following panel: CD4-PE (BD, Franklin Lakes, NJ, USA), CD25-FITC (BD), CD69-APC (BD), CD38-PE-Cy7 (BD), HLA-DR-PerCP-Cy5.5 (Biolegend, San Diego, CA, USA), HLA-A2-APC-H7 (BD), AnnexinV-PacBlue (BD). Stained cells were analyzed by flow cytometry on a FACS Canto (BD). Allogeneic feeders were obtained from HLA-A2-negative healthy donors for the PHA/feeders experiment, and we gated on HLA-A2-positive cells in order to distinguish between the subjects' cells and the allogeneic feeders [11].

#### Intact Proviral DNA Assay

The IPDA measures the HIV-1 reservoir by quantifying both defective and intact proviral DNA. The total provirus is the sum of intact, 5' defective, and 3' defective virus detected by the IPDA. Aliquots of 5 million unfractionated CD4+ T cells were assayed at Accelevir Diagnostics as previously described [7].

#### Data Analysis

Flow cytometry data were analyzed using FlowJo, version 10. All statistics were generated in GraphPad Prism 7. The Pearson



**Figure 1.** Activation marker profiles induced by different activation strategies. Bulk CD4+ T cells were either not treated (NT) or stimulated with phytohaemagglutinin (PHA)/feeders, PMA/ionomycin, or PHA/anti-CD28. Activation was measured by assessing the upregulation of 4 proteins: CD69, CD25, HLA-DR, and CD38. For each individual, the mean value from duplicates was obtained, and the mean values from 7 HDs (A) and 3 CPs (B) are shown. The error bars represent the standard error of means per treatment group per time point. To determine the differences between PHA/feeder stimulated cells and either treated or untreated cells, a 1-way repeated measures analysis of variance was performed.

test was run for all correlation statistics, and the Kruskal-Wallis test with Dunn's multiple comparison was performed for the frequency of infectious units per million cells (IUPM) values as well as the IUPM/intact provirus ratio. The IUPMs were based on the percentage of positive replicates and were calculated using the IUPMstats calculator [12]. Cultures that had no positive wells were assigned the median posterior estimate of infection frequency.


## RESULTS

### Different Stimulation Strategies Result in Different Patterns of Immune Activation

We compared the upregulation of activation markers at different time points after stimulation with PHA/feeders vs PHA and an anti-CD28 monoclonal antibody, PMA and ionomycin in CD4+ T cells from healthy donors and chronic progressors on ART. As shown in Figure 1, PMA/ionomycin stimulation resulted in higher and more sustained expression of the early activation marker CD69 than the other 2 strategies. In contrast, PHA/anti-CD28 stimulation resulted in the most efficient induction of CD25 expression. PHA/feeder stimulation resulted in the highest levels of HLA-DR expression, and stimulation with PHA/feeders and PHA/anti-CD28 was more effective at inducing CD38 expression than PMA/ionomycin stimulation. Interestingly, we found that cells stimulated with PHA/feeders had the highest levels of annexin V expression over the first 2 days of culture (Supplementary Figure 1).

### Different Stimulation Strategies Result in Similar Levels of Latency Reversal

To determine how effective each strategy was at latency reversal, we stimulated CD4+ T cells from 11 study subjects



**Figure 2.** Latency reversal by the different activation strategies. Statistical comparison of the infectious units per million (IUPM) generated with the 3 different assays by a nonparametric Kruskal-Wallis with Dunn's comparisons test. The open symbols indicate assays where there were no positive wells, and the IUPM value is dictated by the median posterior estimate.

on ART including 9 chronic progressors (CPs) and 2 viremic controllers (VCs) (Table 1). We compared the frequency of CD4+ T cells that produced replication-competent virus after stimulation with the 3 different strategies in viral outgrowth assays. Replication-competent virus was obtained from CD4+ T cells from 10, 8, and 9 subjects, after stimulation with PHA/feeders, PMA/ionomycin, and PHA/anti-CD28 strategies (Figure 2) respectively. There were no significant differences in the median frequency of IUPM generated with the different immune activation strategies (Figure 2).

### Positive Correlation Between Frequencies of Cells Induced to Produce Replication-Competent Virus by Different Activation Strategies

We asked whether there was any relationship between the frequencies of cells that produced replication-competent virus following stimulation of CD4+ T cells with the 3 different strategies. There was a significantly positive correlation of the IUPMs generated after stimulation with PMA/ionomycin and PHA/feeders (Figure 3A), PHA/anti-CD28 and PMA/ionomycin (Figure 3B), and PHA/anti-CD28 and PHA/feeders (Figure 3C).


### Relationship Between Intact DNA and Frequency of Cells Induced to Produce Replication-Competent Virus by Each Strategy

The IPDA is a recently developed digital droplet polymerase chain reaction assay that uses primers developed from near full-length proviral sequences to distinguish intact proviral DNA from most deleted and/or hypermutated proviral DNA [7]. The frequency of intact and total DNA for each study subject is shown in Table 1. As the frequency of intact DNA approximates the size of the inducible reservoir, we asked what fraction of cells with intact DNA produced replication-competent virus. As shown in Figure 4 each activation strategy induced replication-competent virus from only a fraction of the cells with intact DNA.

Finally, we asked whether there was any correlation between intact DNA and IUPM for each of the 3 activation strategies. We found a significant positive correlation between the IUPM generated by the 3 different activation strategies and intact DNA (Figure 5A-C) but not total DNA (Figure 5D-F).

## DISCUSSION

Reversal of HIV-1 latency will be required for most HIV cure strategies, but recent studies have suggested that only a small fraction of CD4+ T cells that harbor intact proviral DNA produce replication-competent virus upon activation with PHA and feeders [4, 5]. We asked whether other immune activation strategies would reverse latency more effectively. We used a combination of PMA and ionomycin that is often used to produce maximal T-cell activation. A recent study demonstrated that PHA alone is not as effective at reversing latency as the combination of PHA and allogeneic feeders [13], so for




**Figure 3.** Correlation of quantitative viral outgrowth assay infectious units per million generated by the different activation strategies. Comparisons of phorbol 12-myristate 13-acetate (PMA)/ionomycin vs phytohaemagglutinin (PHA)/feeders (A), PHA/anti-CD28 vs PMA/ionomycin (B), and PHA/anti-CD28 vs PHA/feeders (C) are shown with the Pearson's correlation coefficients. Values that were below the limit of detection were not included in the analysis.

this study we combined PHA with a superagonistic anti-CD28 antibody that provides a co-stimulatory signal and directly activates CD4+ T cells [10]. The pattern of immune activation produced by these different regimens was quite different with PMA/ionomycin inducing the highest levels of CD69, PHA/anti-CD28 inducing the highest levels of CD25, and PHA and feeders causing the highest expression of the late activation marker HLA-DR. Further studies will be needed to determine whether these differences are due to the activation of different subsets of cells and/or due to different mechanisms of action of the different activation strategies. However, in spite of these different patterns of activation, there was no significant difference in the frequency of cells that produced replication-competent virus when the activated cells were co-cultured with lymphoblasts from healthy donors. Furthermore, there was a positive


correlation between the frequency of cells producing virus in the 3 different assays.

We demonstrate that PMA/ionomycin and PHA/anti-CD28 stimulation can be used in place of the PHA/feeders activation step that is used in the standard viral outgrowth assay. Other studies have also shown comparable levels of latency reversal when different immune activation strategies are used in viral outgrowth assays [11, 13–16]. Our results extend these studies by also measuring the level of immune activation induced by each strategy. The PMA/ionomycin and PHA/anti-CD28 methods of immune activation do not require a 10:1 ratio of allogeneic feeders to patient CD4+ T cells and thus are less labor-intensive and less expensive. These methods will also be easier to use for the repeated stimulation of CD4+ T cells needed to trigger replication of non-induced viral clones that are refractory to a single round of T-cell activation [4, 6]. A recent study showed lower levels of HIV transcriptional initiation in infected rectal CD4+ T cells compared with infected peripheral CD4+ T cells in patients on ART, suggesting that cells in this compartment may be in a deeper state of latency [17]. Thus, it will be important to compare latency reversal in CD4+ T cells from different tissues with these alternate activation strategies. We also determined the frequency of CD4+ T cells that harbored intact proviral DNA and showed that there was a positive correlation between the frequency of intact proviral DNA and the IUPM induced with each activation method. A comparison of the 2 parameters suggested that <10% of CD4+ T cells with intact proviral DNA produced replication-competent virus with each activation method. This is consistent with a prior study that found that only a very small fraction of CD4+ T cells with proviral DNA could be induced to produce replication-competent virus following stimulation with antibodies to CD3 and CD28 [18]. These data suggest that different strategies that produce global immune activation



**Figure 4.** Ratio of quantitative viral outgrowth assay infectious units per million (IUPM)/intact proviral DNA for each activation strategy. The open symbols indicate assays where there were no positive wells, and the maximum possible IUPM value was divided by the intact proviral DNA frequency.





**Figure 5.** Correlations between quantitative viral outgrowth assay infectious units per million (IUPM) from each activation strategy and proviral DNA. IUPM derived from stimulation with phytohaemagglutinin (PHA)/feeders (A, D) phorbol 12-myristate 13-acetate (PMA)/ionomycin (B, E), and PHA/anti-CD28 (C, F) are plotted against intact (A–C) and total (D–F) proviruses per million CD4+ T cells. Pearson’s correlation coefficients are shown.

are inefficient at reversing latency. The Simoa ultrasensitive p24 assay is 1000-fold more sensitive than the standard ELISA we used for this study [19]; thus it is possible that we are underestimating the frequency of cells that produce replication-competent virus. However, in some studies, viral isolates that were positive by the ultrasensitive p24 assay alone were not always capable of replication in subsequent cultures, suggesting that some of these isolates may not be replication-competent [20, 21]. RNA induction assays are also more sensitive than QVOAs [22], but a recent study showed that a large percentage of transcription-competent virions produced after CD4+ T-cell activation are not replication-competent [23]. Further studies will therefore be required to determine what percentage of CD4+ T cells with intact DNA produce replication-competent viruses that are capable of initiating subsequent rounds of replication vs defective or attenuated viruses and/or transcription-competent viruses that are not replication-competent. In summary, our data suggest that different immune activation strategies induce different patterns of activation marker expression on CD4+ T cells and produce replication-competent virus from similar frequencies of latently infected CD4+ T cells. Novel latency-reversal strategies or repeated stimulation may be needed to fully reverse latency in viral outgrowth assays.

### Supplementary Data

Supplementary materials are available at *Open Forum Infectious Diseases* online. Consisting of data provided by the authors to benefit the reader, the posted materials are not copy-edited and are the sole responsibility of the authors, so questions or comments should be addressed to the corresponding author.

### Acknowledgments

**Financial support.** This work was supported by the Johns Hopkins University Center for AIDS Research (P30AI094189) and the National Institute of Allergies and Infectious Diseases (R01AI120024, J.N.B.).

**Potential conflicts of interest.** All authors: no reported conflicts of interest. All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

### References

- Siliciano JD, Kajdas J, Finzi D, et al. Long-term follow-up studies confirm the stability of the latent reservoir for HIV-1 in resting CD4+ T cells. *Nat Med* **2003**; 9:727–8.
- Crooks AM, Bateson R, Cope AB, et al. Precise quantitation of the latent HIV-1 reservoir: implications for eradication strategies. *J Infect Dis* **2015**; 212:1361–5.
- Wang Z, Simonetti FR, Siliciano RF, Laird GM. Measuring replication competent HIV-1: advances and challenges in defining the latent reservoir. *Retrovirology* **2018**; 15:21.
- Ho YC, Shan L, Hosmane NN, et al. Replication-competent noninduced proviruses in the latent reservoir increase barrier to HIV-1 cure. *Cell* **2013**; 155:540–51.
- Bruner KM, Murray AJ, Pollack RA, et al. Defective proviruses rapidly accumulate during acute HIV-1 infection. *Nat Med* **2016**; 22:1043–9.
- Hosmane NN, Kwon KJ, Bruner KM, et al. Proliferation of latently infected CD4+ T cells carrying replication-competent HIV-1: potential role in latent reservoir dynamics. *J Exp Med* **2017**; 214:959–72.

7. Bruner KM, Wang Z, Simonetti FR, et al. A quantitative approach for measuring the reservoir of latent HIV-1 proviruses. *Nature* **2019**; 566:120–5.
8. Siliciano JD, Siliciano RF. Enhanced culture assay for detection and quantitation of latently infected, resting CD4+ T-cells carrying replication-competent virus in HIV-1-infected individuals. *Methods Mol Biol* **2005**; 304:3–15.
9. Salgado M, Swanson MD, Pohlmeier CW, et al. HLA-B\*57 elite suppressor and chronic progressor HIV-1 isolates replicate vigorously and cause CD4+ T cell depletion in humanized BLT mice. *J Virol* **2014**; 88:3340–52.
10. Waibler Z, Sender LY, Merten C, et al. Signaling signatures and functional properties of anti-human CD28 superagonistic antibodies. *PLoS One* **2008**; 3:e1708.
11. Kuzmichev YV, Veenhuis RT, Pohlmeier CW, et al. A CD3/CD28 microbead-based HIV-1 viral outgrowth assay. *J Virus Erad* **2017**; 3:85–9.
12. Rosenbloom DI, Elliott O, Hill AL, et al. Designing and interpreting limiting dilution assays: general principles and applications to the latent reservoir for human immunodeficiency virus-1. *Open Forum Infect Dis* **2015**; 2(4).
13. Beliakova-Bethell N, Hezareh M, Wong JK, et al. Relative efficacy of T cell stimuli as inducers of productive HIV-1 replication in latently infected CD4 lymphocytes from patients on suppressive cART. *Virology* **2017**; 508:127–33.
14. Wong JK, Hezareh M, Günthard HF, et al. Recovery of replication-competent HIV despite prolonged suppression of plasma viremia. *Science* **1997**; 278:1291–5.
15. Tremblay CL, Giguel F, Merrill DP, et al. Marked differences in quantity of infectious human immunodeficiency virus type 1 detected in persons with controlled plasma viremia by a simple enhanced culture method. *J Clin Microbiol* **2000**; 38:4246–8.
16. Sanyal A, Mailliard RB, Rinaldo CR, et al. Novel assay reveals a large, inducible, replication-competent HIV-1 reservoir in resting CD4+ T cells. *Nat Med* **2017**; 23:885–9.
17. Telwate S, Lee S, Somsouk M, et al. Gut and blood differ in constitutive blocks to HIV transcription, suggesting tissue-specific differences in the mechanisms that govern HIV latency. *PLoS Pathog* **2018**; 14:e1007357.
18. Cillo AR, Sobolewski MD, Bosch RJ, et al. Quantification of HIV-1 latency reversal in resting CD4+ T cells from patients on suppressive antiretroviral therapy. *Proc Natl Acad Sci U S A* **2014**; 111:7078–83.
19. Passaes CPB, Bruel T, Decalf J, et al. Ultrasensitive HIV-1 p24 assay detects single infected cells and differences in reservoir induction by latency reversal agents. *J Virol* **2017**; 91:e02296–16.
20. Dhummakupt A, Siems LV, Singh D, et al. The latent human immunodeficiency virus (HIV) reservoir resides primarily in CD32-CD4+ T cells in perinatally HIV-infected adolescents with long-term virologic suppression. *J Infect Dis* **2019**; 219:80–8.
21. Bertagnolli LN, White JA, Simonetti FR, et al. The role of CD32 during HIV-1 infection. *Nature* **2018**; 561:E17–9.
22. Plantin J, Massanella M, Chomont N. Inducible HIV RNA transcription assays to measure HIV persistence: pros and cons of a compromise. *Retrovirology* **2018**; 15:9.
23. Massanella M, Yek C, Lada SM, et al. Improved assays to measure and characterize the inducible HIV reservoir. *EBioMedicine* **2018**; 36:113–21.