



GENOME ANNOUNCEMENT

Open Access

Genome sequencing and genomic characterization of a tigecycline-resistant *Klebsiella pneumoniae* strain isolated from the bile samples of a cholangiocarcinoma patient

Beiwen Zheng^{1,2†}, Ang Li^{1,2†}, Xiawei Jiang^{1,2}, Xinjun Hu^{1,2}, Jian Yao^{1,2}, Lina Zhao^{1,2}, Jinru Ji^{1,2}, Min Ye^{1,2}, Yonghong Xiao^{1,2*} and Lanjuan Li^{1,2*}

Abstract

Background: The relationship between *Klebsiella pneumoniae* and nosocomial and community-acquired infections is well known, and *K. pneumoniae* resistance to most antibiotics is increasing worldwide. In contrast, tigecycline remains active against many bacterial strains, and serves as a last resort for treating multi-drug resistant bacterial infections. That tigecycline nonsusceptibility among *K. pneumoniae* isolates has been reported worldwide is worrying. However, the mechanisms of tigecycline resistance in *K. pneumoniae* are less well known. We report the genome sequence and genomic characterization of tigecycline-resistant *K. pneumoniae* strain 5422 isolated from the bile samples of a patient with cholangiocarcinoma.

Results: We sequenced the *K. pneumoniae* strain 5422 genome using next-generation sequencing technologies. Sequence data assembly revealed a 5,432,440-bp draft genome and 57.1% G + C content, which contained 5397 coding sequences. The genome has extensive similarity to other sequenced *K. pneumoniae* genomes, but also has several resistance-nodulation-cell division (RND) efflux pump genes that may be related to tigecycline resistance.

Conclusions: *K. pneumoniae* strain 5422 is resistant to multiple antibiotics. The genome sequence of the isolate and comparative analysis with other *K. pneumoniae* strains presented in this paper are important for better understanding of *K. pneumoniae* multi-drug resistance. The RND efflux pump genes identified in the genome indicate the presence of an antibiotic resistance mechanism prior to antibiotics overuse. The availability of the genome sequence forms the basis for further comparative analyses and studies addressing the evolution of the *K. pneumoniae* drug resistance mechanism and the *K. pneumoniae* transcriptome.

Keywords: Tigecycline-resistant, *Klebsiella pneumoniae*, Cholangiocarcinoma, Next-generation sequencing, Comparative genomics

Background

Klebsiella pneumoniae is a Gram-negative opportunistic pathogen from the family *Enterobacteriaceae*. The increasing resistance detected in clinical isolates has become a matter of significant concern, and *K. pneumoniae* is an ESKAPE (*Enterococcus faecium*, *Staphylococcus aureus*,

K. pneumoniae, *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, and *Enterobacter* species) pathogen [1,2]. Infections caused by *K. pneumoniae* have been identified worldwide. It is worth noting that it is a major cause of nosocomial and community-acquired infections [3,4]. Moreover, a distinct invasive syndrome that causes liver abscesses has been detected in increasing numbers in Asia in the past two decades, and this syndrome is emerging as a global disease [5,6]. Furthermore, *K. pneumoniae* harboring extended-spectrum β-lactamases, and more recently, carbapenemase, which confers resistance to

* Correspondence: xiao-yonghong@163.com; lilj@zju.edu.cn

†Equal contributors

¹State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, School of Medicine, Zhejiang University, Hangzhou 310003, China

²Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, Zhejiang University, Hangzhou 310003, China

multiple antibiotics, has become a significant clinical concern worldwide [7,8]. Mortality among patients infected with extremely resistant *K. pneumoniae* is high, perhaps due to the limited therapeutic options remaining [9].

Tigecycline, a novel class of glycylcyclines, is a minocycline derivative synthesized to overcome the major tetracycline resistance mechanisms and to extend its spectrum of activity to multidrug-resistant (MDR) bacteria [10]. Tigecycline has enhanced antimicrobial activity compared to tetracycline, and can overcome efflux pump systems and ribosome protection mechanisms, retaining activity against a broad range of both Gram-positive and Gram-negative bacteria [11-13]. *K. pneumoniae* resistance to most antibiotics is increasing globally. Nevertheless, tigecycline remains active against many bacterial strains, and serves as last resort for treating MDR bacterial infections [14]. However, tigecycline resistance has emerged recently and been widely reported in *Enterobacteriaceae* isolates. It is worth noting that tigecycline non-susceptibility among *K. pneumoniae* isolates has been reported from different continents and ranges between 0% and 50% [15-17].

Previous studies have suggested that tigecycline resistance in *Enterobacteriaceae* is attributed to resistance-nodulation-cell division (RND)-type efflux pumps and transcriptional regulators of the efflux pump systems [15]. However, reports on the tigecycline resistance mechanisms in *K. pneumoniae* are rare. We hypothesized that a wide range of genes is involved in tigecycline resistance and contribute to decreased tigecycline susceptibility in *K. pneumoniae*, and performed whole-genome sequencing (WGS) to investigate this. We report the genome sequence of the tigecycline-resistant *K. pneumoniae* strain 5422 isolated from the bile samples of a patient with cholangiocarcinoma.

Methods

Strain information and growth conditions

Previously, we isolated strain 5422 from a bile sample obtained from a 54-year-old woman with bile duct cancer on February 25, 2012. After enrichment in Mueller-Hinton broth, the strain was identified as *K. pneumoniae* following the combination of its 16S rRNA gene sequencing and biochemical reaction results (VITEK 2 compact, bioMérieux, France). The strain exhibited high resistance to ciprofloxacin, ceftazidime, cefotaxime, cefoxitin, ampicillin/sulbactam, sulfamethoxazole, tigecycline, tetracycline, and piperacillin, and was susceptible to imipenem, meropenem, and gentamicin. Multi-locus sequence typing revealed that it belonged to ST37 (unpublished data). The strain we reported here is available in the State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, Zhejiang University.

Genomic DNA extraction

Late log-phase cells were harvested and lysed with EDTA, lysozyme, and detergent treatment, followed by proteinase K and RNase digestion. Genomic DNA was extracted using a DNeasy Blood & Tissue Kit (Qiagen, Germany) according to the manufacturer's recommended protocol. Genomic DNA yield, purity, and concentration was evaluated using 0.7% agarose gel electrophoresis with λ -Hind III digest DNA Marker and measured using a NanoDrop 1000 Spectrophotometer (Thermo Fisher Scientific, USA). The genomic DNA was stored at -20°C.


Genome sequencing and annotation

Whole-genome shotgun sequencing of *K. pneumoniae* strain 5422 was performed using a standard run of IlluminaHiSeq2000 sequencing by generating paired-end libraries (500-bp insert size) with a 2 × 100 pair-end sequencing strategy according to the manufacturer's instructions. Clean reads were assembled into scaffolds using Velvet version 1.2.07 [18], and then we used PAGIT (Post-Assembly Genome Improvement Toolkit) [19] to extend the initial contiguous sequences (contigs) and correct sequencing errors. We identified tRNAs and rRNAs using tRNAscan-SE [20] and RNAmmer [21], respectively. Open reading frames (ORFs) were identified using Glimmer version 3.0 [22]. The genome was annotated using the RAST (Rapid Annotation using Subsystem Technology) server [23]. The classification of some predicted genes and pathways was analyzed using the COGs (Clusters of Orthologous Groups of proteins) [24] and KEGG (Kyoto Encyclopedia of Genes and Genomes) [25] databases. Stretches of amino acids containing the efflux pump genes were searched using BLAST (Basic Local Alignment Search Tool, <http://blast.ncbi.nlm.nih.gov/Blast.cgi>); protein-coding sequences were further BLAST-searched against the Antibiotic Resistance Database (ARDB) [26]. To find genes with hypothetical or putative functions, we aligned genes against the National Center for Biotechnology Information (NCBI) nucleotide sequence database (downloaded September 20, 2013) using NCBI BLASTn: we accepted only hits with identity of ≥ 0.95 , coverage ≥ 0.9 , and putative or hypothetical reference gene annotation.

Initial comparative genomic and phylogenetic analysis

For comparative analysis, we downloaded the reference genome sequences of the closest genetic relatives of *K. pneumoniae* strain 5422 and representative strains from the NCBI website: *K. pneumoniae* LCT-KP182 (ATRN 00000000), *K. pneumoniae* LCT-KP289 (ATRO00000000), *K. pneumoniae* subsp. *pneumoniae* LZ (AJVY00000000), *K. pneumoniae* ATCC BAA-2146 (AOCV00000000), *K. pneumoniae* 12 3578 (PRJNA199972), *K. pneumoniae* NB60 (AZAP00000000), *K. pneumoniae* subsp. *pneumoniae* 1084 (CP003785), *K. pneumoniae* subsp. *pneumoniae*

A



B





Figure 1 (See legend on next page.)

(See figure on previous page.)

Figure 1 Genome map and phylogenetic analysis. A. Circular map of the genome generated using Circos. Circles from outside to inside: 1, contigs were arranged in clockwise direction from large to small; 2, CDS on forward strand; 3, CDS on reverse strand; 4, tRNA genes; 5, rRNA genes; 6, GC-skew (window size of 10 kb); and 7, blue indicates C content and yellow indicates G content (step size 500 bp). B. COGs distribution of *K. pneumoniae* strain 5422.

A



B




Figure 2 Subsystem distribution and phylogenetic analysis. A. Distribution of genes assigned to SEED subsystems (based on the RAST annotation server). B. Phylogenetic relationships (based on WGS and SNPs) of 12 *K. pneumoniae* strains and their genomic distance analysis. The snpTree server output used assembled genomes as input data.

NTUH-K2044 (AP006725), *K. pneumoniae* subsp. *pneumoniae* WGLW5 (AMLO00000000), *K. pneumoniae* ATCC 25955 (AQQH00000000), and *K. pneumoniae* G5-2 (AQQI00000000). Whole-genome alignments, single-nucleotide polymorphism (SNP) identification, and phylogenetic tree construction were performed using snpTree version 1.1, a server for online automatic SNP analysis of assembled genomes (<http://cge.cbs.dtu.dk/services/snpTree-1.1/>) [27].

Quality assurance

The 16S rDNA gene from the draft genome was used to check for contamination. Further VITEK biochemical identification data confirmed that the strain 5422 belonged to *K. pneumoniae*. Bioinformatics assessment of potential contamination of the genomic library by allochthonous microorganisms was achieved using the BLAST non-redundant database.

Initial findings

Genome characteristics and phylogenetic analysis

Filtered 520.8 M clean reads were assembled into scaffolds, and corresponding 99-fold coverage of the genome was generated. The draft genome sequence of *K. pneumoniae* strain 5422 was 5,432,440 bp in size and had a G + C content of 57.1% in 133 contigs, with N50 spanning 105,586 bp. Figure 1A depicts the overall genome profile.

Annotation of this assembly identified 5,397 coding sequences (CDSs), 65 tRNAs (excluding 0 pseudo tRNAs), and incomplete rRNA operons (three small subunit rRNAs, four large subunit rRNAs). We assigned putative function or hypothetical proteins to 1,478 protein-coding genes. We categorized 4,218 genes into COGs functional groups (including putative or hypothetical genes, Figure 1B). For COGs distribution, R (general function prediction only; 874 ORFs), E (amino acid transport and metabolism; 747 ORFs), G (carbohydrate metabolism and transport; 679 ORFs), and P (inorganic ion transport and metabolism; 525 ORFs) were abundant categories (>10% of total COGs matched counts).

Figure 2A illustrates the subsystem distribution and general information on the potential functional distribution of *K. pneumoniae* strain 5422. Genes responsible for carbohydrates (825 ORFs); amino acids and derivatives (520 ORFs); and cofactors, vitamins, prosthetic groups, and pigments (325 ORFs) were abundant among the SEED subsystem categories. Based on the raw reads and assembled genomes from published *K. pneumoniae* WGS data sets, we conducted phylogenetic analysis on tree topology and the SNP positions of the reference genome to identify the most closely related organism. The phylogenetic tree based on whole-genome SNPs showed that the closest ancestor to *K. pneumoniae* strain 5422 was *K. pneumoniae* ATCC BAA-2146 (Figure 2B),

Table 1 Summary of CDSs annotated to RND efflux pump genes

Aligned protein	Query sequence length	Coverage (%)	Hit length	Identity (%)	Description
AcrA	240	64.17	146	60.08	Probable RND efflux membrane fusion protein
OqxB	217	20.66	141	64.98	Probable RND efflux system inner membrane transporter CmeB
OqxB	181	18.53	110	60.77	Probable RND multi-drug efflux transporter
OqxB	160	15.27	105	65.62	Probable RND efflux system inner membrane transporter CmeB
AcrA	397	100	397	100	Membrane fusion protein of RND family multi-drug efflux pump
OqxA	483	100	480	99.38	Transcriptional regulator
OqxB	1050	100	1050	100	RND multi-drug efflux transporter
OqxA	391	100	391	100	RND multi-drug efflux transporter
RarA	369	100	366	99.19	Bacterial regulatory helix-turn-helix proteins, AraC family
OqxB	161	15.33	101	62.73	Probable RND multi-drug efflux transporter
AcrA	328	83.89	248	75.61	Probable RND efflux system membrane fusion protein CmeA
OqxB	217	20.67	143	65.9	Probable RND efflux system inner membrane transporter CmeB
TolC	1520	100	1474	96.97	Outer membrane efflux protein
OqxB	326	31.05	200	61.35	Probable RND efflux system inner membrane transporter CmeB

which is the first US isolate found to encode New Delhi metallo- β -lactamase 1 (NDM-1), eight β -lactamases, and 15 additional antibiotic-resistance enzymes [28,29].

RND efflux pumps in *K. pneumoniae* strain 5422

The RND family members are important mediators of MDR in Gram-negative bacteria. The AcrAB-TolC system in *Escherichia coli* and the MexAB-OprM complexes in *P. aeruginosa* are extremely well characterized, and the three-dimensional structures of various components have been resolved [30]. In the genome of *K. pneumoniae* strain 5422, 16 genes were indicated as probable efflux pumps or translational regulators based on their sequence similarity to known RND efflux pump genes (Table 1).

Future directions

The rapid progress of WGS has permitted detailed investigation of genetic differences between bacterial isolates with different phenotypic characteristics. Whole-genome studies of *K. pneumoniae* have mainly focused on comparing either distinct antibiotic-susceptible and MDR strains or related isolates from different patients. Therefore, large-scale genomic sequencing and comparative genome analysis of tigecycline-resistant, tigecycline non-susceptible, and tigecycline-susceptible clinical isolates will identify the differences in the genomic content of this species and yield evolutionary information on the development of tigecycline resistance through mutations. Moreover, further studies involving extensive high-throughput mRNA sequencing (RNA-Seq) experiments to significantly improve annotation and to provide exceptionally robust analysis of RNA expression under selective antibiotic pressure are warranted.

Ethics approval

This research was approved by the Research Ethics Committee of the First Affiliated Hospital, School of Medicine, Zhejiang University, and informed consent was obtained from the patient.

Availability of supporting data

This Whole Genome Shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession JGYH00000000. The version described in this paper is version JGYH01000000.

Competing interests

The authors have no competing interests.

Authors' contributions

BWZ, XJH, LNZ, JRJ and MY performed microbiology and molecular biology experiments. XWJ and JY generated and analyzed the sequencing data. BWZ and AL participated all discussions of data analysis and write the manuscript. BWZ, YHX and LJL were involved in overall experimental design. All authors read and approved the final manuscript.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (Grant No. 81361138021 and 81301461), Zhejiang Provincial Natural Science Foundation of China (Grant No. LQ13H190002) and the Scientific Research Foundation of Zhejiang Provincial Health Bureau (Grant No. 2012KYB083).

Received: 11 July 2014 Accepted: 2 September 2014

Published online: 27 September 2014

References

1. Boucher HW, Talbot GH, Bradley JS, Edwards JE, Gilbert D, Rice LB, Scheld M, Spellberg B, Bartlett J: **Bad bugs, no drugs: no ESKAPE! An update from the Infectious Diseases Society of America.** *Clin Infect Dis* 2009, 48:1–12.
2. Rice LB: **Federal funding for the study of antimicrobial resistance in nosocomial pathogens: no ESKAPE.** *J Infect Dis* 2008, 197:1079–1081.
3. Moore R, O'Shea D, Geoghegan T, Mallon PW, Sheehan G: **Community-acquired Klebsiella pneumoniae liver abscess: an emerging infection in Ireland and Europe.** *Infection* 2013, 41:681–686.
4. Henderson KL, Muller-Pebody B, Johnson AP, Wade A, Sharland M, Gilbert R: **Community-acquired, healthcare-associated and hospital-acquired bloodstream infection definitions in children: a systematic review demonstrating inconsistent criteria.** *J Hosp Infect* 2013, 85:94–105.
5. Siu LK, Yeh KM, Lin JC, Fung CP, Chang FY: **Klebsiella pneumoniae liver abscess: a new invasive syndrome.** *Lancet Infect Dis* 2012, 12:881–887.
6. Lin JC, Koh TH, Lee N, Fung CP, Chang FY, Tsai YK, Ip M, Siu LK: **Genotypes and virulence in serotype K2 Klebsiella pneumoniae from liver abscess and non-infectious carriers in Hong Kong, Singapore and Taiwan.** *Gut Pathog* 2014, 6:21.
7. Nordmann P, Cuzon G, Naas T: **The real threat of Klebsiella pneumoniae carbapenemase-producing bacteria.** *Lancet Infect Dis* 2009, 9:228–236.
8. Richter SN, Frasson I, Franchin E, Bergo C, Lavezzo E, Barzon L, Cavallaro A, Palu G: **KPC-mediated resistance in Klebsiella pneumoniae in two hospitals in Padua, Italy, June 2009–December 2011: massive spreading of a KPC-3-encoding plasmid and involvement of non-intensive care units.** *Gut Pathog* 2012, 4:7.
9. Munoz-Price LS, Poirel L, Bonomo RA, Schwaber MJ, Daikos GL, Cormican M, Cornaglia G, Garau J, Gniadkowski M, Hayden MK, Kumarasamy K, Livermore DM, Maya JJ, Nordmann P, Patel JB, Paterson DL, Pitout J, Vilegas MV, Wang H, Woodford N, Quinn JP: **Clinical epidemiology of the global expansion of Klebsiella pneumoniae carbapenemases.** *Lancet Infect Dis* 2013, 13:785–796.
10. Petersen PJ, Jacobus NV, Weiss WJ, Sum PE, Testa RT: **In vitro and in vivo antibacterial activities of a novel glycyclcycline, the 9-t-butylglycylamido derivative of minocycline (GAR-936).** *Antimicrob Agents Chemother* 1999, 43:738–744.
11. Jenner L, Starosta AL, Terry DS, Mikolajka A, Filonava L, Yusupov M, Blanchard SC, Wilson DN, Yusupova G: **Structural basis for potent inhibitory activity of the antibiotic tigecycline during protein synthesis.** *Proc Natl Acad Sci U S A* 2013, 110:3812–3816.
12. Brust K, Evans A, Plemmons R: **Tigecycline in treatment of multidrug-resistant Gram-negative bacillus urinary tract infections: a systematic review.** *J Antimicrob Chemother* 2014.
13. Kanj SS, Whitelaw A, Dowzicky MJ: **In vitro activity of tigecycline and comparators against Gram-positive and Gram-negative isolates collected from the Middle East and Africa between 2004 and 2011.** *Int J Antimicrob Agents* 2014, 43:170–178.
14. Lin YT, Wang FD, Chan YJ, Fu YC, Fung CP: **Clinical and microbiological characteristics of tigecycline non-susceptible Klebsiella pneumoniae bacteremia in Taiwan.** *BMC Infect Dis* 2014, 14:1.
15. Sun Y, Cai Y, Liu X, Bai N, Liang B, Wang R: **The emergence of clinical resistance to tigecycline.** *Int J Antimicrob Agents* 2013, 41:110–116.
16. Nigo M, Cevallos CS, Woods K, Flores VM, Francis G, Perlman DC, Revuelta M, Mildvan D, Waldron M, Gomez T, Koshy S, Jodłowski T, Riley W, Ruhe JJ: **Nested case-control study of the emergence of tigecycline resistance in multidrug-resistant Klebsiella pneumoniae.** *Antimicrob Agents Chemother* 2013, 57:5743–5746.
17. Huang TD, Berhan C, Bogaerts P, Glupczynski Y: **In vitro susceptibility of multidrug-resistant Enterobacteriaceae clinical isolates to tigecycline.** *J Antimicrob Chemother* 2012, 67:2696–2699.
18. Zerbino DR, Birney E: **Velvet: algorithms for de novo short read assembly using de Bruijn graphs.** *Genome Res* 2008, 18:821–829.

19. Swain MT, Tsai IJ, Assefa SA, Newbold C, Berriman M, Otto TD: A Post-Assembly Genome-Improvement Toolkit (PAGIT) to obtain annotated genomes from contigs. *Nat Protoc* 2012, **7**:1260–1284.
20. Lowe TM, Eddy SR: tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res* 1997, **25**:0955–0964.
21. Lagesen K, Hallin P, Rødland EA, Stærfeldt H-H, Rognes T, Ussery DW: RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res* 2007, **35**:3100–3108.
22. Delcher AL, Bratke KA, Powers EC, Salzberg SL: Identifying bacterial genes and endosymbiont DNA with Glimmer. *Bioinformatics* 2007, **23**:673–679.
23. Aziz RK, Bartels D, Best AA, De Jongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O: The RAST Server: rapid annotations using subsystems technology. *BMC Genomics* 2008, **9**:75.
24. Tatusov RL, Fedorova ND, Jackson JD, Jacobs AR, Kiryutin B, Koonin EV, Krylov DM, Mazumder R, Mekhedov SL, Nikolskaya AN, Rao BS, Smirnov S, Sverdlov AV, Vasudevan S, Wolf YI, Yin JJ, Natale DA: The COG database: an updated version includes eukaryotes. *BMC Bioinformatics* 2003, **4**:41.
25. Mitra S, Rupek P, Richter DC, Urich T, Gilbert JA, Meyer F, Wilke A, Huson DH: Functional analysis of metagenomes and metatranscriptomes using SEED and KEGG. *BMC Bioinformatics* 2011, **12 Suppl 1**:S21.
26. Liu B, Pop M: ARDB—Antibiotic Resistance Genes Database. *Nucleic Acids Res* 2009, **37**:D443–D447.
27. Leekitcharoenphon P, Kaas RS, Thomsen MC, Friis C, Rasmussen S, Aarestrup FM: snpTree—a web-server to identify and construct SNP trees from whole genome sequence data. *BMC Genomics* 2012, **13 Suppl 7**:S6.
28. Broberg CA, Palacios M, Miller VL: Whole-genome draft sequences of three multidrug-resistant *Klebsiella pneumoniae* strains available from the American type culture collection. *Genome Announc* 2013, **1**:e00312–e00313.
29. Hudson CM, Bent ZW, Meagher RJ, Williams KP: Resistance determinants and mobile genetic elements of an NDM-1-encoding *Klebsiella pneumoniae* strain. *PLoS One* 2014, **9**:e99209.
30. Alvarez-Ortega C, Olivares J, Martinez JL: RND multidrug efflux pumps: What are they good for? *Front Microbiol* 2013, **4**:7.

doi:10.1186/s13099-014-0040-2

Cite this article as: Zheng et al.: Genome sequencing and genomic characterization of a tigecycline-resistant *Klebsiella pneumoniae* strain isolated from the bile samples of a cholangiocarcinoma patient. *Gut Pathogens* 2014 **6**:40.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

