



Contents lists available at ScienceDirect

Animal Nutrition

journal homepage: <http://www.keaipublishing.com/en/journals/aninu/>

KeAi
CHINESE ROOTS
GLOBAL IMPACT



Original Research Article

Dietary lecithin improves feed efficiency without impacting meat quality in immunocastrated male pigs and gilts fed a summer ration containing added fat

Henny Akit ^{a,*}, Cherie Collins ^b, Fahri Fahri ^a, Alex Hung ^a, Darryl D'Souza ^c, Brian Leury ^a, Frank Dunshea ^a

^a Faculty of Veterinary and Agricultural Sciences, The University of Melbourne, Parkville, Victoria, 3010, Australia

^b Rivalea Australia Pty Ltd., Corowa, 2646, Australia

^c Australian Pork Ltd., Level 2, 2 Brisbane Avenue, Barton Capital Territory, 2600, Australia

ARTICLE INFO

Article history:

Received 23 August 2017

Received in revised form

11 January 2018

Accepted 23 January 2018

Available online 16 February 2018

Keywords:

Fat

Feed efficiency

Lecithin

Meat quality

Pigs

ABSTRACT

The aim of this study was to investigate the effects of sex and dietary lecithin on growth performance, meat quality, muscle collagen content and gene expression of key genes involved in collagen synthesis in finisher pigs. A total of 256 pigs (Large White × Landrace) were allotted to a 2 × 2 factorial arrangement involving sex (gilt or immunocastrated [IC] male) and dietary treatment (0 or 5 g/kg of dietary lecithin). All diets were formulated to contain 4.6% tallow with relatively high total fat of 6.3%. After 5 weeks of dietary treatment, pigs were slaughtered and *Longissimus dorsi* muscle was obtained for evaluation of meat quality and collagen content. *Rectus abdominis* muscle was analysed for gene expression of key genes involved in collagen synthesis namely, type I ($\alpha 1$) procollagen (*COL1A1*), type III ($\alpha 1$) procollagen (*COL3A1*), α -subunit of prolyl 4-hydroxylase (*P4H*), lysyl oxidase and metalloproteinase-1 (*MMP-1*). The results showed that lecithin improved feed efficiency of all pigs ($P < 0.05$) but it had no effect on feed intake, average daily gain and dressing percentage ($P > 0.05$). Lecithin also had no effect on meat compression, shear force, collagen content and gene expression ($P > 0.05$). Immunocastrated male had higher growth rate and increased *COL1A1* expression than gilts. However, sex had no effect on fat depth at the P2 site (65 mm from the midline over the last rib), collagen content and expression of other genes ($P > 0.05$). In conclusion, lecithin improved feed efficiency in finishing pigs without impacting pork quality. Thus, inclusion of lecithin in diets containing high amount of tallow during the summer period could be beneficial.

© 2018, Chinese Association of Animal Science and Veterinary Medicine. Production and hosting by Elsevier B.V. on behalf of KeAi Communications Co., Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Soy lecithin, a by-product from the processing of soybean oil, contains a mixture of various phospholipids. Lecithin is an emulsifier and it is generally used in various food applications, cosmetics and pharmaceutical industries. Lecithin also has been used in young pig starter diets to enhance the utilisation of dietary fat (Jones et al., 1992; Overland et al., 1993a; Reis de Souza et al., 1995; Soares and Lopez-Bote, 2002). The addition of high levels of fat in the pig finisher diet is particularly common in the summer period where the growth rate is slower compared with pigs reared in autumn (Myer and Bucklin, 2002). Fats and oils are used as energy sources in animal diets because of their high energy value and low heat increment of digestion. However, animal fats such as

* Corresponding author.

E-mail address: henny@upm.edu.my (H. Akit).

¹ Present address: Department of Animal Science, Faculty of Agriculture, Universiti Putra Malaysia, Serdang, Selangor, 43400, Malaysia.

Peer review under responsibility of Chinese Association of Animal Science and Veterinary Medicine.



Production and Hosting by Elsevier on behalf of KeAi

tallow and lard are rich in saturated fatty acids which are less digestible than vegetable oils (Cera et al., 1988; Soares and Lopez-Bote, 2002). A welfare friendly alternative to castration of male pigs is immunization against gonadotrophin releasing factor (GnRF) which allows producers to capitalize on the superior feed efficiency and carcass characteristics of boars without the risk of boar taint (Dunshea et al., 2001). It has been reported that immunocastrated (IC) males have higher feed intake and faster growth rate than gilts (Gispert et al., 2010; Morales et al., 2011; Font-i-Furnols et al., 2012). Rapid growth rate in animals can affect meat tenderness through increased muscle collagen synthesis and degradation (Fishell et al., 1985; Harper et al., 1999). Faster growing groups of cattle were reported to have more tender meat than slower growing groups of cattle (Harper et al., 1999; Zgur, 2003; Perry and Thompson, 2005). High growth rate before slaughter in IC males was suggested to be associated with lower pork shear force values in IC males compared with barrows and boars (Pauly et al., 2009). However, studies comparing meat tenderness between IC male pigs and other sex are limited (Channon et al., 2016). The aim was to investigate the impacts of sex and dietary lecithin on growth performance, meat quality and muscle collagen content of finisher pigs. In order to understand some of the mechanisms, the effects of dietary lecithin and sex on gene expression of key genes involved in intramuscular collagen synthesis and degradation were also investigated.

2. Materials and methods

2.1. Diets, animals and experimental design

A total of 256 pigs (Large White × Landrace) (PrimeGro Genetics, NSW, Australia) were identified at 17 weeks of age and transferred to a finisher accommodation with 32 pens. Each pen consisted of 8 pigs (same-sex). Pen weights (average start weight = 66.2 ± 0.70 kg) were recorded at entry to the finisher pens. Pens were randomly allocated to a 2 × 2 factorial experiment with the respective factors: 0 or 5 g/kg dietary lecithin (ADM Australia Pty Ltd., NSW, Australia) and sex (IC male or gilt). The priming dose of Improvac (CSL Pty Ltd., VIC, Australia) vaccination was administered to male pigs at 13 weeks of age with the second vaccination at 17 weeks of age. All diets were formulated to contain 0.52 g available lysine/MJ DE, 14.2 MJ DE/kg and 4.6% tallow (Table 1). Experimental diets were fed to pigs for 5 weeks starting from 17 weeks of age until slaughter at 22 weeks of age. Pen weights were recorded at the beginning of the experimental period at d 0 (17 weeks of age) and again at d 14. At d 35 (pre-slaughter), individual live weights were recorded. Pen feed intake was recorded over the intervals between d 0 to 14 and d 15 to 35 as measured by feed disappearance. Feed efficiency was subsequently calculated. At d 35, the pigs were slaughtered according to the standard procedures practised in commercial abattoirs. All procedures for animal care were approved by the Rivalea Animal Care and Use Committee.

2.2. Carcass quality measurements and sampling

Rectus abdominis muscles from a subset of 64 carcasses (2 median-weight pigs per pen) were collected. The tissues were immediately immersed in liquid nitrogen and then stored at -80 °C. This muscle was selected because of convenience of sampling at the slaughter house. The individual hot standard carcass weight was recorded. Fat depth at the P2 site (65 mm from the midline over the last rib) were measured using a PorkScan ultrasound system (PorkScan Pty Ltd., Canberra, Australia). Individual

Table 1
Ingredients and nutrient composition of experimental diets (as-fed basis).

Item	Experimental diets	
	Control	Lecithin
Ingredients, %		
Wheat	59.9	59.9
Barley	8.0	8.0
Millmix	13.6	13.6
Canola meal	7.0	7.0
Meat meal	3.8	3.8
Water	1.0	1.0
Tallow	4.6	4.6
Salt	0.2	0.2
Limestone	1.3	1.3
Lysine HCL	0.4	0.4
Threonine	0.1	0.1
Copper premix	0.1	0.1
Rivalea finisher premix	0.1	0.1
Rumensin	0.1	0.1
Lecithin	0.0	0.5
Estimated nutrient composition ¹ , %		
Digestible energy (DE), MJ/kg	14.2	14.2
Crude protein	15.2	15.2
Crude fat	6.3	6.3
Crude fibre	4.1	4.1
Total lysine	0.9	0.9
Available lysine to DE ratio, g/MJ DE	0.5	0.5

¹ Estimated from Rivalea Australia Pty Ltd. (NSW, Australia) composition data. This project was funded by Pork CRC. This was reported in Pork CRC report, 2011.

live weight and carcass weight measures were used to calculate dressing percentage. Twenty-four hours post-slaughter, *Longissimus dorsi* muscle (loin) from a subset of 64 carcasses (2 pigs per pen) was removed from the right side of each pig carcass. The loin samples for collagen content analysis were vacuum packed and stored at -20 °C until ready for use. Meanwhile, loin samples for texture analysis were vacuum packed and left to age for 5 days at 4 °C. Then, the muscle samples were stored at -20 °C for future analyses.

2.3. Meat pH and colour measurements

The pH of loin between 12th and 13th rib was measured at 45 min and 24 h post-slaughter using a portable pH/temperature metre Model 6009 (Jenco Electronic Ltd., Taipei, Taiwan, China) fitted with a polypropylene spear-type gel electrode (Ionode IJ42S, QLD, Australia) and a temperature probe.

Following a bloom time of 30 min, meat colour (L^* , a^* and b^*) was measured with a Minolta Chromameter CR-400 (Minolta, Osaka, Japan) using D65 lighting, a 2° standard observer, and an 8-mm aperture in the measuring head. L^* values correspond to lightness, a^* values to redness and b^* values to yellowness in the three-dimensional HunterLab colour space.

2.4. Cooking method and cooking loss

The cooking procedure was adopted from the method of Bouton et al. (1971). The loin was cut to a 70 ± 5 g cube (40 mm × 40 mm × 40 mm). Weight of each cube was recorded as weight before cooking. Then the samples were cooked in water bath at 70 °C for 35 min. After cooking, the samples were immersed in ice cold water for 20 min and then patted dry to remove excess moisture. Samples weight after cooking was recorded before being refrigerated overnight. Cooking loss was calculated as a percentage of weight loss before and after cooking.

2.5. Warner–Bratzler shear force and compression analyses

Texture of cooked meat was determined using Warner–Bratzler shear force and compression analyses (Bouton and Harris, 1972). For shear force, each sample was cut into 6 rectangular strips of 1 cm² cross-section, parallel to the muscle fibres. The shear force blade was driven to cut the samples at a crosshead speed of 300 mm/min and a 1 kN load cell was used. The meat tenderness was estimated using the mean of the peak shear force.

For compression analysis, each sample was cut into 6 cross-section samples (1-cm thick) with the fibres lying perpendicular on the face of the largest area. The sample was placed under a flat-ended plunger that moved downwards at about 80% through the sample. Hardness is defined as the peak force required for the first compression. Then, the plunger moved upwards and then returned to the same damaged area to measure the second compression. The ratio of the force required for the second compression to the force required for the first compression is defined as cohesiveness. Consequently, chewiness value was calculated as the product of hardness and cohesiveness (Bouton and Harris, 1972).

2.6. Meat collagen content analysis

Hydroxyproline content was determined using a colourimetric method developed by Kolar (1990) with minor modifications. Hydroxyproline was oxidised to pyrrole with chloramine-T reagent. Then, the complex formed by the liberated pyrrole and 4-dimethylaminobenzaldehyde was determined by the colourimetric method. The absorbance values of the sample solutions were measured on a plate reader (Thermo Multiskaner, Vantaa, Finland) at 558 nm wave length. The concentration of hydroxyproline (µg/mL) was determined from a standard curve. The total collagen content was determined using a factor of 7.46 (Colgrave et al., 2008) and was reported as milligramme of collagen per gram of sample.

2.7. Quantification of mRNA by real-time RT-PCR

Total RNA was extracted from 200 to 300 mg samples using Trizol reagent and PureLink Micro-to-Midi RNA isolation system kit (Invitrogen, San Diego, CA, USA) following the manufacturer's instructions. The RNA quality was evaluated using Experion automated electrophoresis system and Experion RNA StdSens analysis kit (BIORAD, Gladesville, NSW, Australia). Samples displaying 28S:18S ribosomal RNA ratio close to 1.5 were considered acceptable. Then, cDNA was synthesised using 1 µg of total RNA from each

sample using the SuperScript III First-Strand Synthesis System (Invitrogen) for RT-PCR and 50 ng/µL random hexamer primer (Invitrogen), following manufacturer's instructions. Primers matrix metalloproteinase-1 (*MMP-1*) and ribosomal 18s (*R18s*) were designed based on porcine sequences. Type I ($\alpha 1$) procollagen (*COL1A1*), type III ($\alpha 1$) procollagen (*COL3A1*), α -subunit of prolyl 4-hydroxylase (*P4H*) and lysyl oxidase primers were designed based on mouse (*Mus musculus*) sequences because complete genomes of the pig are not available (Table 2).

The reverse transcription PCR was performed on a MyiQ system (BIORAD) using SYBR green (BIORAD) in triplicate. The temperature profile consisted of initial hot start at 95 °C for 3 min as the initial denaturation step of one cycle, followed by 40 cycles at 95 °C for 10 s (denaturation), 51.0 to 61.2 °C for 45 s (annealing) and an extension step at 95.0 °C for 1 min. Melting curve analysis was performed to confirm the specificity of the primers. Non-template controls were included with each PCR run to validate that primers were not amplifying contaminating DNA. Relative gene expression was calculated using the 2^{-ΔΔCt} method, as described by Livak and Schmittgen (2001).

2.8. Statistical analysis

The main effects of dietary lecithin and sex on growth performance and carcass quality were analysed using an analysis of variance (ANOVA) for a randomised design. Replicate was used as a blocking factor. The experimental unit for all analyses was the pen of pigs. The meat colour, pH, collagen content, cooking loss, texture and gene expression data from a subset of pigs were analysed using restricted maximum likelihood (REML) to determine the main effects of dietary lecithin and sex. The pen number was included as a random model. All results were considered statistically significant when $P < 0.05$ and trends when $0.05 \leq P \leq 0.10$. All analyses were performed using the Genstat software 13th edition (VSN International Ltd, Hertfordshire, UK).

3. Results

3.1. Growth performance and carcass quality

Table 3 shows the results on the growth performance and carcass quality. During the initial d 0 to 14, lecithin tended to improve feed efficiency compared with pigs fed the control diet ($P = 0.068$). However, lecithin had no effect on feed intake and weight gain. Immunocastrated males had higher weight gain and better feed efficiency than gilts ($P < 0.001$). Feed intake was similar between sexes. Interactions between sex and diet

Table 2
Characteristics of the primers used for quantitative real-time PCR.

Gene	Accession number	Primers	Primer sequences (5' to 3')	Annealing temperature, °C	Amplicon size, bp
<i>COL1A1</i>	BC050014.1	Forward	GTCTGGTTTGAGAGAGCAT	60.9	189
		Reverse	CTTCTTGAGGTTGCCAGTCT		
<i>COL3A1</i>	NM_009930.2	Forward	TGATGTCAAGTCTGGAGTGG	53.4	223
		Reverse	TCCTGACTCTCCATCCCTTC		
<i>MMP-1</i>	EU722905.1	Forward	GTTCCACAAATGAGTGCTGA	60.9	212
		Reverse	ATAATAACGACGGCTCATCC		
Lysyl oxidase	M65142.1	Forward	CTGCTTGATGCCAACACA	58.7	156
		Reverse	TGCCGCATAGGTGTCATA		
α -subunit <i>P4H</i>	BC009654.1	Forward	CCCAGTCAGGTCTGCTATTC	51.0	204
		Reverse	GGAACAGTCTCTGGACAACC		
<i>R18s</i>	AY265350.1	Forward	GAACCCACTTGTCCCTCTA	61.2	219
		Reverse	GACTCAACACGGAAACCTC		

COL1A1 = type I ($\alpha 1$) procollagen; *COL3A1* = type III ($\alpha 1$) procollagen; *MMP-1* = matrix metalloproteinase-1; α -subunit *P4H* = α -subunit of prolyl 4-hydroxylase; *R18s* = ribosomal 18s.

Table 3
Effects of sex and dietary lecithin on average daily gain (ADG), average daily feed intake (ADFI), feed conversion ratio (FCR) and carcass quality.¹

Item	Gilt		Immunocastrated male		SED	P-value		
	Control	Lecithin	Control	Lecithin		Sex	Diet	Sex × Diet
Initial bodyweight, kg	65.1	65.1	67.3	67.3	0.70	<0.001	1.00	0.98
Final bodyweight ² , kg	96.8	98.5	105.3	104.7	1.31	<0.001	0.53	0.18
From 0 to 14 d on experiment								
ADG, kg/d	0.83	0.95	1.09	1.00	0.042	<0.001	0.65	0.002
ADFI, kg/d	2.48	2.53	2.61	2.45	0.066	0.60	0.26	0.028
FCR, kg/kg	3.04	2.68	2.40	2.46	0.114	<0.001	0.068	0.015
From 15 to 35 d on experiment								
ADG, kg/d	0.87	0.87	1.17	1.20	0.048	<0.001	0.64	0.66
ADFI, kg/d	2.76	2.64	3.54	3.39	0.120	<0.001	0.13	0.87
FCR, kg/kg	3.22	3.07	3.02	2.82	0.121	0.015	0.051	0.75
From 0 to 35 d on experiment								
ADG, kg/d	0.85	0.90	1.14	1.12	0.037	<0.001	0.56	0.21
ADFI, kg/d	2.65	2.60	3.17	3.02	0.091	<0.001	0.12	0.43
FCR, kg/kg	3.13	2.90	2.78	2.69	0.078	<0.001	0.006	0.21
Carcass quality								
Carcass weight ² , kg	75.5	76.8	80.3	80.1	1.04	<0.001	0.39	0.29
P2 backfat depth ³ , mm	10.8	10.2	10.5	10.8	0.40	0.49	0.40	0.067
Dressing, %	78.4	78.5	75.9	76.1	0.47	<0.001	0.70	0.79

SED = standard error of the difference; P2 = the site of 65 mm from the midline over the last rib.

¹ From Pork CRC report, 2011.

² Initial bodyweight was included as a covariate in the analysis.

³ Carcass weight was included as a covariate in the analysis.

Table 4
Effects of sex and dietary lecithin on meat quality of longissimus muscle.

Meat quality	Gilt		Immunocastrated male		SED	P-value		
	Control	Lecithin	Control	Lecithin		Sex	Diet	Sex × Diet
pH at 45 min	6.46	6.35	6.41	6.39	0.111	0.93	0.41	0.58
pH at 24 h	5.76	5.75	5.80	5.79	0.075	0.47	0.81	0.99
Lightness (L*)	49.41	48.59	48.50	49.16	0.593	0.68	0.85	0.085
Redness (a*)	5.66	6.04	5.81	5.97	0.307	0.86	0.22	0.63
Yellowness (b*)	2.09	1.94	2.10	2.22	0.234	0.37	0.91	0.41
Cooking loss, %	20.6	23.0	21.5	23.1	0.01	0.49	0.007	0.55
Warner–Bratzler shear force, kg	3.04	3.07	3.43	3.16	0.208	0.11	0.41	0.32
Chewiness	1.53	1.52	1.51	1.51	0.082	0.82	0.87	0.92
Cohesiveness	0.35	0.36	0.35	0.35	0.007	0.90	0.48	0.76
Hardness, kg	4.36	4.26	4.30	4.26	0.185	0.79	0.60	0.83
Collagen content, mg/g	0.65	0.64	0.67	0.69	0.055	0.46	0.94	0.79

SED = standard error of the difference.

showed that weight gain ($P = 0.002$) and feed intake ($P = 0.028$) were greater in gilts when offered the lecithin diets but not in IC males. As a result, gilts offered the lecithin diets had better feed efficiency ($P = 0.015$). During the subsequent period (d 15 to 35), there was a strong trend for improved feed efficiency in pigs fed lecithin diets ($P = 0.051$) but there was no effect on feed intake and weight gain. Immunocastrated males had higher weight gain and feed intake as well as better feed efficiency than gilts over this period ($P < 0.05$). There was no interaction between sex and diet on the growth performance during this period. Over the entire experimental period (d 0 to 35), lecithin improved pig feed efficiency ($P = 0.006$). However, lecithin had no effect on feed intake and weight gain. Immunocastrated males had higher weight gain and feed intake as well as better feed efficiency than gilts ($P < 0.001$). There was no interaction between sex and diet on the growth performance over the entire experiment.

Lecithin had no effect on final live weight, carcass weight, P2 backfat depth and dressing percentage whereas IC males had heavier final live weight and carcass weight than gilts ($P < 0.001$). Gilts had higher dressing percentage than IC males ($P < 0.001$). The P2 backfat depths were similar in both sexes. There was no interaction between sex and diet on the carcass quality.

3.2. Meat quality

Dietary lecithin or sex had no impact on meat pH, colour, collagen content, shear force, chewiness, cohesiveness and hardness (Table 4). Lecithin increased cooking loss ($P = 0.007$), but cooking loss was similar in both sexes. There was no interaction between sex and diet on the meat quality.

3.3. Skeletal muscle gene expression

Dietary lecithin or sex had no impact on the gene expression of the genes involved in muscle collagen synthesis and degradation (data on effect of lecithin was not shown). The exceptions were for *COL1A1* mRNA expression that was increased in IC males ($P = 0.051$) and *COL3A1* that tended to increase in IC males compared with gilts ($P = 0.093$) (Fig. 1). There was no interaction between sex and diet on the gene expression (data not shown).

4. Discussion

The major finding from the present study was that dietary lecithin improved feed efficiency in both gilts and IC male pigs fed a summer diet containing 4.6% tallow added. Similarly, Kim et al.




Fig. 1. Relative mRNA expression of type I ($\alpha 1$) procollagen (*COL1A1*), type III ($\alpha 1$) procollagen (*COL3A1*), matrix metalloproteinase-1 (*MMP-1*), lysyl oxidase and α -subunit of prolyl 4-hydroxylase (α -subunit *P4H*) in *rectus abdominis* muscle of finishing pigs. The expression levels were normalized against that of Ribosomal 18s (*R18s*) reference gene. Each column represents the means \pm SED of 32 pigs. The results were considered significant when $P < 0.05$ and as trends when $0.05 \leq P \leq 0.10$. SED = standard error of the difference; F = female; IC male = immunocastrated male.

(2008) also demonstrated improvement in feed efficiency in finisher pigs fed lecithin in diets containing 2.5% tallow. In contrast to the present study, lecithin had no effect on feed efficiency but it improved dressing percentage in finisher pigs supplemented lecithin in diets containing 4.5% tallow (Akit et al., 2014). Lecithin, as an emulsifier, was proposed to increase the capacity of bile salt micelles to solubilise long chain saturated fatty acids thus improving animal fat digestion and absorption (Freeman, 1969; Reynier et al., 1985). However, Kim et al. (2008) did not observe any improvements in nutrient digestibility. The degree of fatty acid saturation in the diet is an important factor in determining fat digestibility in pigs (Eusebio et al., 1965). Lecithin is high in unsaturated fatty acid, particularly linoleic acid. Therefore, it was hypothesised that rather than being due to lecithin's emulsifying effect, the benefits on digestibility might be due to an indirect increase in fatty acid unsaturation (Soares and Lopez-Bote, 2002). However, the same authors found that addition of lecithin increased the unsaturated to saturated fatty acid ratio of diets containing lard but lecithin had no effect on fat digestibility of weaning piglets. Effect of lecithin on feed efficiency of pigs fed diets containing vegetable oils was also inconsistent. A few studies showed no changes in the feed efficiency of pigs fed lecithin in diets containing canola oil (D'Souza et al., 2012, 2015). Another study on the other hand, showed lecithin improved feed efficiency in growing–finishing pigs fed diets containing soy oil (Overland et al., 1993b). They suggested that different sources of lecithin as well as varying type and level of dietary fats contribute to inconsistent responses of growth performance to lecithin.

Generally, muscles with higher collagen content possess higher cross-link concentrations per unit of collagen than muscles with lower collagen content (Shimokomaki et al., 1972; Light et al., 1985). A more tender meat was suggested to be as a consequence of reduced collagen content, and therefore reduced cross-links levels (Ngapo et al., 2002). Lecithin in the present experiment had no effect on meat texture and collagen content. In contrast, dietary lecithin decreased longissimus muscle chewiness and hardness (Akit et al., 2014; D'Souza et al., 2015) and this improvement was suggested to be due to decreased muscle collagen content (Akit et al., 2014). Dietary lecithin also decreased hardness and chewiness of semitendinosus muscle (D'Souza et al., 2012). Longissimus muscle had relatively lower collagen content compared with other muscles such as *Biceps femoris*, *Triceps bricii* and semitendinosus (Nold et al., 1999; Wheeler et al., 2000). Collagen content (0.64 to 0.69 mg/kg) was much lower in the present experiment than in a previous experiment (1.07 to 1.51 mg/kg) reported by Akit et al. (2014). Because of the already relatively low longissimus muscle collagen content in the present study, it was less likely to detect any

changes caused by dietary lecithin. In addition, the shear force values in the present study (3.04 to 3.43 kg) were lower compared with the values reported by Channon et al. (2001). Their study reported longissimus muscle shear force values were in a range of 3.72 to 5.13 kg. The meat from the present experiment was already tender which may have made it difficult to further improve its tenderness. Moreover, it was suggested to supplement lecithin in the diet of pigs as early as the grower stage through to the finisher stage (D'Souza et al., 2012) because meat becomes tougher as the animal grows in age due to the progressive maturation of collagen (Kauffman et al., 1964; Fang et al., 1999). Hence, lecithin in the present study could perhaps influence meat texture and collagen content if the pigs were fed lecithin for longer duration or if pigs with tougher muscle with higher collagen content was used.

Cooking loss was increased by lecithin in the present study. However, it should be noted that the cooking loss was much lower than the normal range of 25.8% to 27.3% (Akit et al., 2014). During cooking, denaturation and shrinkage of endomysial and perimysial collagen sheaths resulted in loss of water and reduced tenderness (Bailey, 1988; Avery et al., 1996). Cooking loss was shown to be positively correlated with meat total collagen and heat-insoluble collagen (Okeudo and Moss, 2005). This may explain the low values of cooking loss and total collagen in the present study.

As expected, IC males in the present experiment grew faster than the gilts and this was in agreement with previous studies (Gispert et al., 2010; Morales et al., 2011; Font-i-Furnols et al., 2012). Immunocastrated males also had more superior growth performance compared with boars and barrows (Dunshea et al., 2001, 2011). Immunocastration of male pigs decreased testosterone levels after the second immunisation against gonadotropin-releasing hormone (Dunshea et al., 2001; Zeng et al., 2002; Claus et al., 2007). Testosterone has a negative influence on appetite of pigs thus reduced levels of testosterone resulted in increased feed intake (Weiler et al., 1996). Reduced testosterone level also reduced aggression and mounting behaviours leading to increased growth rate because more feed was utilised for live weight gain rather than the non-beneficial behaviours in IC male pigs (Cronin et al., 2003; Dunshea et al., 2011).

Variation in meat characteristics due to sex effect can be explained by its influence on protein and body fat depositions in the carcass (Franco et al., 2008; Peinado et al., 2008; Serrano et al., 2009). However, carcass backfat depth at the P2 site was not affected by sex in the present study. When comparing IC male and gilt, backfat depth did not differ by much as reported by Gispert et al. (2010), and Fabrega et al. (2010) found backfat depth values of IC male and gilt were in between barrow and boar. Another study found that backfat depth did not differ between IC male and gilt but

was lower than barrow (Morales et al., 2011). Testosterone was suggested to contribute to high muscle collagen content in bulls because of its anabolic effects on collagen synthesis (Boccard et al., 1979). This coincides with a study that found muscle total collagen was greater in boars than barrows and gilts (Nold et al., 1999). A few studies comparing barrows and gilts found no effect of sex on muscle collagen content (Correa et al., 2006; Maiorano et al., 2007; Corino et al., 2008). However, one study found that collagen content and shear force values were lower in muscle of barrows than gilts (Kim et al., 2016). The lack of differences in backfat depth and collagen content correspond with the lack of differences in meat texture between IC male and gilt in the present study. Including boars and measuring the effect of testosterone on meat collagen synthesis could be worthy of further investigation.

Intramuscular collagen had less contribution to beef toughness from cattle slaughtered after a period of rapid growth (Fishell et al., 1985; Harper et al., 1999). Increased growth rate following a period of feed restriction showed variable responses on meat tenderness and collagen characteristics in pigs (Therkildsen et al., 2002; Kristensen et al., 2002, 2004). In the present experiment, IC male that had faster growth rate did not show any differences in the muscle collagen content and meat texture compared with the gilt. It is known that other than growth rate, type of muscle, animal age and management practices can also affect collagen synthesis and maturation (McCormick, 1994; Purslow, 2005). Apart from intramuscular collagen, growth rate prior to slaughter also affects post-mortem proteolytic degradation of myofibrillar proteins and thus meat texture (Kristensen et al., 2002; Sazili et al., 2004; Therkildsen et al., 2004). However, myofibrillar protein degradation may have been levelled out in all samples in the present experiment because the meat were left to age for 5 days prior to analysis. Moreover, muscle ultimate pH was similar in both IC male and gilt therefore any differences in the meat texture due to myofibrillar protein degradation may have been minimised.

Intramuscular connective tissues composed of extracellular matrix molecules such as collagens, proteoglycans, and glycoproteins. Collagen is synthesised as intracellular precursors known as procollagens. Type I and type III collagens make up the major components of intramuscular connective tissue. Extracellular matrix plays an important role in muscle cell growth and the organization of intramuscular connective tissue during skeletal muscle development (Nishimura, 2015). Faster growth rate was associated with greater rate of collagen synthesis (McCormick, 1994). The *COL3A1* and *COL5A3* (collagen type V, alpha 1) genes which have functions in collagen development were down-regulated in tissue of pigs fed low protein diets that had reduced growth rate (Hamill et al., 2013). The expression levels of several important genes involved in collagen synthesis including *COL3A1* and *COL1A1* were up-regulated in intact male cattle compared with the castrates (Zhang et al., 2011). They suggested that this was because intact males have a vigorous collagen anabolism and turnover rates compared with the castrates. The substantial difference in growth rate between IC male and gilt may explain the increase of *COL1A1* expression and the trend in increase of *COL3A1* expression in IC males in the present study. Although it should be kept in mind that this explanation should be supported by other genes involved in growth and development functions.

Lecithin was shown to prevent collagen accumulation as evidenced by down-regulation of *COL1A1* expression in hepatocyte cell culture (Poniachik et al., 1999; Cao et al., 2002a, 2002b) as well as to enhance breakdown of existing collagen (Lieber et al., 1994) in liver fibrosis. In a study using gilts, lecithin down-regulated skeletal muscle expression of *COL1A1*, *COL3A1* and *MMP-1* genes, and it tended to down-regulate the expression of α -subunit *P4H* (Akit et al., 2016). In contrast, lecithin did not show any changes in the

expression of genes involved in collagen synthesis in the present study. This lack of lecithin effect may partly be explained by the low level of muscle collagen content.

5. Conclusions

Dietary lecithin improved feed efficiency in both IC males and gilts without impacting pork quality. Hence, lecithin could be useful to improve feed efficiency in finisher pigs fed high amount of fat during the summer period. Lecithin and sex had no effect on meat texture, collagen content as well as gene expression of key genes involved in collagen synthesis and degradation. The lack of lecithin effect maybe because of the low collagen content and shear force values across treatments. Immunocastrated males had faster growth rate and higher *COL1A1* expression levels than gilts. Further investigation on the relationship between growth rate and collagen synthesis may be worth considering.

Conflicts of interest

The authors declare that they have no conflict of interest.

Acknowledgement

This research project was financed by Pork Cooperative Research Centers (CRC).

References

- Akit H, Collins CL, Fahri FT, Hung AT, D'Souza DN, Leury BJ, et al. Dietary lecithin decreases skeletal muscle *COL1A1* and *COL3A1* gene expression in finisher gilts. *Animals* 2016;6:38–46.
- Akit H, Collins CL, Fahri FT, Hung AT, D'Souza DN, Leury BJ, et al. Dietary lecithin improves dressing percentage and decreases chewiness in the longissimus muscle in finisher gilts. *Meat Sci* 2014;96:1147–51.
- Avery NC, Sims TJ, Warkup C, Bailey AJ. Collagen cross-linking in porcine *M.longissimus lumborum*: absence of a relationship with variation in texture at pork weight. *Meat Sci* 1996;42:355–69.
- Bailey AJ, Light ND. *Connective tissue in meat and meat products*. London: Elsevier Applied Science; 1988.
- Boccard RL, Naudi RT, Cronje DE, Smit MC, Venter HJ, Sossouw EJ. The influence of age, sex and breed of cattle on their muscle characteristics. *Meat Sci* 1979;3:261–80.
- Bouton PE, Harris PV. Comparison of some objective methods used to assess meat tenderness. *J Food Sci* 1972;37:218–21.
- Bouton PE, Harris PV, Shorthose WR. Effect of ultimate pH upon the water-holding capacity and tenderness of mutton. *J Food Sci* 1971;36:435–9.
- Cao Q, Mak KM, Lieber CS. Dilinoleoylphosphatidylcholine prevents transforming growth factor-beta1-mediated collagen accumulation in cultured rat hepatic stellate cells. *J Lab Clin Med* 2002a;139:202–10.
- Cao Q, Mak KM, Lieber CS. DLPC decreases TGF-beta1-induced collagen mRNA by inhibiting p38 MAPK in hepatic stellate cells. *Am J Physiol Gastrointest Liver Physiol* 2002b;283:1051–61.
- Cera KR, Mahan DC, Reinhart GA. Weekly digestibilities of diets supplemented with corn-oil, lard or tallow by weaning swine. *J Anim Sci* 1988;66:1430–7.
- Channon HA, Hamilton AJ, D'Souza DN, Dunshea FR. Estimating the impact of various pathway parameters on tenderness, flavour and juiciness of pork using Monte Carlo simulation methods. *Meat Sci* 2016;116:58–66.
- Channon HA, Reynolds J, Baud S. Identifying pathways to ensure acceptable eating quality of pork. Final report. Canberra: Australian Pork Ltd.; 2001.
- Claus R, Lacorn M, Danowski K, Pearce MC, Bauer A. Short-term endocrine and metabolic reactions before and after second immunization against GnRH in boars. *Vaccine* 2007;25:4689–96.
- Colgrave ML, Allingham PG, Jones A. Hydroxyproline quantification for the estimation of collagen in tissue using multiple reaction monitoring mass spectrometry. *J Chromatogr A* 2008;1212:150–3.
- Corino C, Musella M, Pastorelli G, Rossi R, Paolone K, Costanza L, et al. Influences of dietary conjugated linoleic acid (CLA) and total lysine content on growth, carcass characteristics and meat quality of heavy pigs. *Meat Sci* 2008;79:307–16.
- Correa JA, Faucitano L, Laforest JP, Rivest J, Marcoux M, Garipey C. Effects of slaughter weight on carcass composition and meat quality in pigs of two different growth rates. *Meat Sci* 2006;72:91–9.
- Cronin GM, Dunshea FR, Butler KL, McCauley I, Barnett JL, Hemsworth P. The effects of immuno- and surgical-castration on the behaviour and consequently growth of group-housed, male finisher pigs. *Appl Anim Behav Sci* 2003;81:111–26.

- D'Souza DN, Blake BL, Williams IH, Mullan BP, Pethick DW, Dunshea FR. Dietary lecithin supplementation can improve the quality of the *M. longissimus thoracis*. *Animals* 2015;5:1180–91.
- D'Souza DN, Mullan BP, Pethick DW, Pluske JR, Dunshea FR. Nutritional strategies affect carcass and pork quality but have no effect on intramuscular fat content of pork. *Anim Prod Sci* 2012;52:276–82.
- Dunshea FR, Colantoni C, Howard K, McCauley I, Jackson P, Long KA, et al. Vaccination of boars with a GnRH vaccine (Improvac) eliminates boar taint and increases growth performance. *J Anim Sci* 2001;79:2524–35.
- Dunshea FR, Cronin GM, Barnett JL, Hemsworth PH, Hennessy DP, Campbell RG, et al. Immunisation against gonadotrophin-releasing hormone (GnRH) increases growth and reduces variability in group-housed boars. *Anim Prod Sci* 2011;51:695–701.
- Eusebio JA, Hays VW, Speer VC, Call JT. Utilisation of fat by young pigs. *J Anim Sci* 1965;24:1001–7.
- Fabrega E, Velarde A, Cros J, Gispert M, Suarez P, Tibau J, et al. Effect of vaccination against gonadotrophin-releasing hormone, using Improvac on growth performance, body composition, behaviour and acute phase proteins. *Livest Sci* 2010;132:53–9.
- Fang SH, Nishimura T, Takahashi K. Relationship between development of intramuscular connective tissue and toughness of pork during growth of pigs. *J Anim Sci* 1999;77:120–30.
- Fishell VK, Aberle ED, Judge MD, Perry TW. Palatability and muscle properties of beef as influenced by pre-slaughter growth rate. *J Anim Sci* 1985;61:151–7.
- Font-i-Furnols M, Gispert M, Soler J, Diaz M, Garcia-Regueiro JA, Diaz I, et al. Effect of vaccination against gonadotrophin-releasing factor on growth performance, carcass, meat and fat quality of male Duroc pigs for dry-cured ham production. *Meat Sci* 2012;91:148–54.
- Franco MM, Antunes RC, Borges M, Melo EO, Goulart LR. Influence of breed, sex and growth hormone and halothane genotypes on carcass composition and meat quality traits in pigs. *J Muscle Foods* 2008;19:34–49.
- Freeman CP. Properties of fatty acids in dispersions of emulsified lipid and bile salt and the significance of these properties in fat absorption in the pig and the sheep. *Br J Nutr* 1969;23:249–63.
- Gispert M, Oliver MA, Velarde A, Suarez P, Perez J, Furnols MFI. Carcass and meat quality characteristics of immunocastrated male, surgically castrated male, entire male and female pigs. *Meat Sci* 2010;85:664–70.
- Hamil RM, Aslan O, Mullen AM, O'Doherty JV, McBryan J, Morris DG. Transcriptome analysis of porcine *M. semimembranosus* divergent in intramuscular fat as a consequence of dietary protein restriction. *BMC Genom* 2013;14:453–67.
- Harper GS, Allingham PG, Le Feuvre RP. Changes in connective tissue of *M. semitendinosus* as a response to different growth paths in steers. *Meat Sci* 1999;53:107–14.
- Jones DB, Hancock JD, Harmon DL, Walker CE. Effects of exogenous emulsifiers and fat sources on nutrient digestibility, serum lipids and growth performance in weaning pigs. *J Anim Sci* 1992;70:3473–82.
- Kauffman RG, Carpenter ZL, Bray RW, Hoekstra WG. Interrelations of tenderness, chronological age, connective tissue fractions of porcine musculature. *J Agric Food Chem* 1964;12:504–7.
- Kim TW, Kim CW, Kwon SG, Hwang JH, Park DH, Kang DG. pH as analytical indicator for managing pork meat quality. *Sains Malays* 2016;45:1097–103.
- Kim WT, Shinde P, Chae BJ. Effect of lecithin with or without chitoooligosaccharide on the growth performance, nutrient digestibility, blood metabolites and pork quality of finishing pigs. *Can J Anim Sci* 2008;88:283–92.
- Kolar K. Colorimetric determination of hydroxyproline as measure of collagen content in meat and meat-products. *J Assoc Off Anal Chem* 1990;73:54–7.
- Kristensen L, Therkildsen M, Riis B, Sorensen MT, Oksbjerg N, Purslow PP, et al. Dietary-induced changes of muscle growth rate in pigs: effects on *in vivo* and postmortem muscle proteolysis and meat quality. *J Anim Sci* 2002;80:2862–71.
- Kristensen L, Therkildsen M, Aaslyng MD, Oksbjerg N, Ertbjerg P. Compensatory growth improves meat tenderness in gilts but not in barrows. *J Anim Sci* 2004;82:3617–24.
- Lieber CS, Robins SJ, Li JJ, Decarli LM, Mak KM, Fasulo JM, et al. Phosphatidylcholine protects against fibrosis and cirrhosis in the baboon. *Gastroenterology* 1994;106:152–9.
- Light N, Champion AE, Voyle C, Bailey AJ. The role of epimysial, perimysial and endomysial collagen in determining texture in six bovine muscles. *Meat Sci* 1985;13:137–49.
- Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta C_T}$ Method. *Methods* 2001;5:402–8.
- Maiorano G, McCormick RJ, Field RA, Snowden G. Intramuscular collagen characteristics of ram, whether, and zeranol-implanted ram lambs. *J Anim Sci* 2007;71:1817–22.
- McCormick RJ. The flexibility of the collagen compartment of muscle. *Meat Sci* 1994;36:79–91.
- Morales JI, Camara L, Berrocoso JD, Lopez JP, Mateos GG, Serrano MP. Influence of sex and castration on growth performance and carcass quality of crossbred pigs from 2 Large White sire lines. *J Anim Sci* 2011;89:3481–9.
- Myer RO, Bucklin RA. Effect of season (summer vs. fall) and diet nutrient density on performance and carcass characteristics of growing-finishing swine. *Trans ASAE (Am Soc Agric Eng)* 2002;45:807–11.
- Ngapo, Berge P, Culioli J, Dransfield E, De Smet S, Claeys E. Perimysial collagen crosslinking and meat tenderness in Belgian Blue Double-muscling cattle. *Meat Sci* 2002;61:91–102.
- Nishimura N. Role of extracellular matrix in development of skeletal muscle and postmortem aging of meat. *Meat Sci* 2015;109:48–55.
- Nold RA, Romans JR, Costello WJ, Libal GW. Characterization of muscles from boars, barrows, and gilts slaughtered at 100 or 110 kilograms: differences in fat, moisture, color, water-holding capacity, and collagen. *J Anim Sci* 1999;77:1746–54.
- Okeudo NJ, Moss BW. Interrelationships amongst carcass and meat quality characteristics of sheep. *Meat Sci* 2005;69:1–8.
- Overland M, Tokach MD, Cornelius SG, Pettigrew JE, Rust JW. Lecithin in swine diets: I. weaning pigs. *J Anim Sci* 1993a;79:1187–93.
- Overland M, Tokach MD, Cornelius SG, Pettigrew JE, Wilson E. Lecithin in swine diets: II. growing-finishing pigs. *J Anim Sci* 1993b;71:194–1197.
- Pauly C, Spring P, O'Doherty JV, Kragten SA, Bee G. Growth performance, carcass characteristics and meat quality of group-penned surgically castrated, immunocastrated (Improvac) and entire male pigs and individually penned entire male pigs. *Animal* 2009;3:1057–66.
- Peinado J, Medel P, Fuentetaja A, Mateos GG. Influence of sex and castration of females on growth performance and carcass and meat quality of heavy pigs destined for the dry-cured industry. *J Anim Sci* 2008;86:1410–7.
- Perry D, Thompson JM. The effect of growth rate during backgrounding and finishing on meat quality traits in beef cattle. *Meat Sci* 2005;69:691–702.
- Poniachik J, Baraona E, Zhao JB, Lieber CS. Dilinoleoylphosphatidylcholine decreases hepatic stellate cell activation. *J Lab Clin Med* 1999;133:342–8.
- Purslow PP. Intramuscular connective tissue and its role in meat quality. *Meat Sci* 2005;70:435–47.
- Reis de Souza T, Peiniau J, Mounier A, Aumaitre A. Effect of addition of tallow and lecithin in the diet of weaning piglets on the apparent total tract and ileal digestibility of fat and fatty acids. *Anim Feed Sci Technol* 1995;52:77–91.
- Reynier MO, Lafont H, Crotte C, Sauve P, Gerolami A. Intestinal cholesterol uptake: comparisons between mixed micelles containing lecithin or lysolecithin. *Lipids* 1985;20:145–50.
- Sazili AQ, Lee GK, Parr T, Sensky PL, Bardsley RG, Buttery PJ. The effect of altered growth rates on the calpain proteolytic system and meat tenderness in cattle. *Meat Sci* 2004;66:195–201.
- Serrano MP, Valencia DG, Fuentetaja A, Lazaro R, Mateos GG. Effect of castration on productive performance, carcass characteristics and meat quality of Iberian pig females reared under intensive management systems. *Livest Sci* 2009;123:147–53.
- Shimokomaki M, Bailey AJ, Elsdon DF. Meat tenderness: age related changes in bovine intramuscular collagen. *J Food Sci* 1972;37:892–6.
- Soares M, Lopez-Bote CJ. Effects of dietary lecithin and fat unsaturation on nutrient utilisation in weaned piglets. *Anim Feed Sci Technol* 2002;95:169–77.
- Therkildsen M, Riis B, Karlsson A, Kristensen L, Ertbjerg P, Purslow PP, et al. Compensatory growth response in pigs, muscle protein turn-over and meat texture: effects of restriction/realimentation period. *Anim Sci* 2002;75:367–77.
- Therkildsen M, Vestergaard M, Busk H, Jensen MT, Riis B, Karlsson AH, et al. Compensatory growth in slaughter pigs-*in vitro* muscle protein turn-over at slaughter, circulating IGF-I, performance and carcass quality. *Livest Prod Sci* 2004;88:63–75.
- Weiler U, Claus R, Dehnhard M, Hofacker S. Influence of the photoperiod and a light reverse program on metabolically active hormones and food intake in domestic pigs compared with a wild boar. *Can J Anim Sci* 1996;76:531–9.
- Wheeler TL, Shackelford SD, Koohmaraie M. Variation in proteolysis, sarcomere length, collagen content, and tenderness among major pork muscles. *J Anim Sci* 2000;78:958–65.
- Zeng XY, Turkstra JA, Jongbloed AW, van Diepen J, Meloen RH, Oonk HB, et al. Performance and hormone levels of immunocastrated, surgically castrated and intact male pigs fed *ad libitum* high- and low-energy diets. *Livest Prod Sci* 2002;77:1–11.
- Zgur S. Influence of growth rate in 2 growth periods on intramuscular connective tissue and palatability traits of beef. *Czech J Anim Sci* 2003;48:113–9.
- Zhang YY, Zan LS, Wang HB. Screening candidate genes related to tenderness trait in Qinchuan cattle by genome array. *Mol Biol Rep* 2011;38:2007–14.