

REVIEW

Open Access



TGF- β /SMAD signaling regulation of mesenchymal stem cells in adipocyte commitment

Sheng-Nan Li^{1*}  and Jia-Fa Wu²

Abstract

Adipocytes arising from mesenchymal stem cells (MSCs) requires MSC adipocyte commitment and differentiation of preadipocytes to mature adipocytes. Several signaling and some cytokines affect the adipogenesis of MSCs. This review focuses on the roles of TGF- β /SMAD signaling in adipocyte commitment of MSCs. BMP4 and BMP7 signaling are sufficient to induce adipocyte lineage determination of MSCs. The roles of BMP2, TGF- β , and myostatin signaling in this process are unclear. Other TGF- β /SMAD signaling such as BMP3 and BMP6 signaling have almost no effect on commitment because of limited research available, while GDF11 signaling inhibits adipocyte commitment in human MSCs. In this review, we summarize the available information on TGF- β /SMAD signaling regulation of MSCs in adipocyte commitment. Deeper study of this commitment mechanism will offer new approaches in treating obesity, diabetes mellitus, and obesity-related metabolism syndrome.

Keywords: TGF- β /SMAD signaling, Commitment, Mesenchymal stem cells (MSCs)

Introduction

Mesenchymal stem cells (MSCs) are a group of cells with multilineage differentiation potential and self-renewal capacity, which were discovered and isolated from the bone marrow in the 1960s [1, 2]. Many studies have shown that MSCs can be isolated from almost all tissue types such as adipose tissue and muscle tissue because of the ease of their isolation and their extensive expansion rate [3, 4]. MSCs can differentiate into various specialized cell types such as osteoblasts, chondrocytes, myoblasts, and adipocytes under the stimuli of different conditions [5, 6]. Adipogenesis of MSCs is presently the research hotspot in the stem cell field. In recent years, some studies have shown that several signaling and some cytokines affect the adipogenesis of MSCs. Adipocytes arise from MSCs by two steps: MSCs commit to preadipocytes and preadipocytes differentiate into adipocytes. The first step adipocyte commitment is a stage that can be markedly changed, with hypomethylation of unknown specific genes [7–9]. The formation of

preadipocytes is an intermediate state between MSCs and adipocytes. While the adipocytes clearly contain fat droplets, the preadipocytes have none. 3T3-L1 cells and 3T3-F442A cells are some of the most common preadipocytes in use at present because of their high adipogenic potential. The second step adipocyte terminal differentiation is that preadipocytes proliferate by mitotic clonal expansion following the input of several hormones, resulting in terminal adipocyte differentiation. At present, the molecular mechanism of the preadipocytes differentiation into adipocytes is clear but little is known about the mechanism involved in the commitment process from MSCs to preadipocytes. Recent studies indicated that the TGF- β superfamily has an important role in regulating the adipocyte commitment of MSCs.

TGF- β superfamily signaling regulates the progress of many cellular processes such as apoptosis, inflammation, fibrosis, and adipocyte differentiation [10]. There are 33 human TGF- β superfamily ligands including activin, inhibin, bone morphogenetic proteins (BMPs), growth differentiation factors (GDFs), Lefty, transforming growth factor β (TGF- β), and Nodal [11, 12]. Structural and functional considerations separate the ligands into two subfamilies: TGF- β s and BMPs [12]. TGF- β , activin,

* Correspondence: lsn@hpu.edu.cn

¹School of Medicine, Henan Polytechnic University, Jiaozuo 454000, Henan, China

Full list of author information is available at the end of the article



myostatin, GDF11, and Nodal belong to the TGF- β s subfamily, while BMP2, BMP4, and BMP7 belong to the BMPs subfamily. Two pairs of transmembrane serine/threonine protein kinases constitute the TGF- β superfamily receptors: the type I receptors and the type II receptors. Ligand binding to the receptors activates SMAD-dependent pathways (TGF- β /SMAD signaling) or SMAD-independent pathways which mainly promote ERK, JNK, and p38 MAPK kinase signaling [13]. In this review, we focus on the roles of TGF- β /SMAD signaling (Fig. 1) in regulating the adipocyte commitment of MSCs to provide a reference point for future research.

TGF- β /SMAD signaling

Activin, inhibin, BMPs, GDFs, TGF- β , Nodal, and other ligands involved in TGF- β /SMAD signaling bind to cell membrane receptors to activate the intracellular downstream SMAD family proteins. The SMAD family proteins constitute three functional classes: the receptor-

regulated SMADs (R-SMADs), the common partner SMAD (Co-SMAD), and the inhibitory SMADs (I-SMADs) [14, 15]. The R-SMADs comprise SMAD1, SMAD2, SMAD3, SMAD5, and SMAD8. The Co-SMAD only refers to SMAD4. The I-SMADs consist of SMAD6 and SMAD7.

In the resting state, R-SMADs mainly exist in the cytoplasm, Co-SMAD predominantly localizes to both the cytoplasm and nucleus, and I-SMADs mainly tend to be in the nucleus. Upon TGF- β s ligand binding, the receptors recruit and phosphorylate SMAD2/3, and SMAD1/5/8 by BMPs ligands. After C-terminal phosphorylation of the R-SMADs, these receptor-activated R-SMADs (SMAD2/3 or SMAD1/5/8) are released from the receptor complex to form a trimeric complex consisting of two R-SMADs and a SMAD4 molecule, which then translocates to the nucleus to regulate the expression of specific genes [13, 16, 17]. After these genes respond to the TGF- β /SMAD signaling, the R-




Fig. 1 TGF- β /SMAD signaling. TGF- β s ligands such as TGF- β , Myostatin, and GDF11 in TGF- β /SMAD signaling bind to cell membrane receptors to phosphorylate the intracellular downstream SMAD2/3 (R-SMADs), and BMPs ligands such as BMP2, BMP4, and BMP7 phosphorylate the SMAD1/5/8 (R-SMADs). Activated R-SMADs form a complex with SMAD4 and then translocate to the nucleus to regulate the expression of specific genes. After the genes respond to the TGF- β /SMAD signaling, the R-SMADs-SMAD4 complex in the nucleus is depolymerized and they re-enter the cytoplasm. I-SMADs comprise SMAD6 and SMAD7, which negatively regulate TGF- β /SMAD signaling. In the resting state, I-SMADs mainly tend to be in the nucleus. Transcriptionally activated by TGF- β /SMAD signaling, SMAD7 shuttling from the nucleus to the cytoplasm prevents R-SMAD phosphorylation. SMAD6 competes with SMAD1 for binding to SMAD4

SMAD–SMAD4 complexes in the nucleus are dephosphorylated by dephosphorylase, and then are depolymerized and re-enter the cytoplasm through the nuclear pore, realizing the recycling of the R-SMADs and SMAD4 [18]. I-SMADs negatively regulate TGF-β/SMAD signaling. When the ligands bind to the receptors, SMAD7 shuttles from the nucleus to the cytoplasm, and binding to the type I receptor prevents R-SMAD phosphorylation to inhibit specific gene expression [19]. SMAD6 acts in a different manner to SMAD7. After the receptors are activated, SMAD6 undergoes shuttling from the nucleus to the cytoplasm and competes with SMAD1 for binding to SMAD4, serving as an inhibitor of TGF-β/SMAD signaling [20].

TGF-β/SMAD signaling regulation of MSCs in adipocyte commitment

Adipose tissue dysfunction can lead to obesity, fatty liver, dyslipidemia, and type II diabetes [8, 21, 22]. According to its color, function, and anatomy, adipose tissue is sub-classified as white adipose tissue (WAT), brown adipose tissue (BAT), and beige adipose tissue. WAT in humans is located abdominally and subcutaneously for lipid storage. BAT, which expresses uncoupling protein 1 (UCP1), is mostly located in the neck and supraclavicular regions of adult humans for thermogenesis [23]. When the body is under the stimulation of adrenalin or cold exposure for a long time, some cells in the white adipose tissue will express high levels of UCP1, stimulate thermogenesis, and undergo “browning” to adapt to the environmental change. These cells are called beige adipose cells [24, 25]. A lot of beige adipose cells compose the beige adipose tissue. Detailed molecular comparisons of white, brown, and beige adipose tissue have identified several specific markers (Table 1) [26, 27]. Adipose tissue is composed of plentiful adipocytes, which are derived from MSCs (MSCs involved in the study are listed in Table 2), cells that have multilineage differentiation capacity and reside in the vascular stroma of adipose tissue as well as in the bone marrow [50]. The Myf5⁻ lineage mainly gives rise to

Table 2 Cell type involved in the regulation of adipocyte commitment under TGF-β/SMAD signaling

Cell type (MSC)	Species	Signaling	References
C3H10T1/2	Mouse	BMP4 signaling	[8, 28–32]
		BMP2 signaling	[28–31, 33, 34]
		BMP7 signaling	[32, 35]
		Myostatin signaling	[36–38]
Human adipose stem cell (hASC)	Human	BMP4 signaling	[32, 39–41]
		BMP2 signaling	[42]
		BMP7 signaling	[41, 43]
Mouse bone marrow stromal stem cell (mouse BMSC)	Mouse	BMP7 signaling	[44]
		TGF-β signaling	[45]
Human bone marrow stromal stem cell (human BMSC)	Human	BMP7 signaling	[46]
		TGF-β signaling	[47, 48]
		Myostatin signaling	[49]


MSC, mesenchymal stem cell; *cell type (MSC)*, cell type involved in the study; *species*, species involved in study; *signaling*, TGF-β/SMAD signaling involved in the regulation of adipocyte commitment; *references*, references related to the study in this table

white preadipocytes, and the Myf5⁺ lineage gives rise to brown preadipocytes, but may not be absolute. Some studies confirmed that brown adipocytes and a subset of white adipocytes originated from Myf5⁺ lineage [51]. Most studies favor the brite adipocytes originate from white adipocytes, but some studies suggest brite adipocytes may originate from mesenchymal progenitor cells [52–55]. The formation of mature adipocytes requires adipocyte commitment of MSCs and adipocyte differentiation. It is now established that TGF-β/SMAD signaling has a dual role in the adipocyte differentiation process. BMPs signaling promotes the mature of adipocytes while TGF-βs signaling inhibits it. In recent years, several studies have shown that TGF-β/SMAD signaling has a critical role in regulating the adipocyte commitment of MSCs (Fig. 2).

Table 1 The comparisons of white, brown, and beige adipose tissue

Types of adipose tissue	Specific gene	Anatomy	Lipid droplet	Specific markers	Function
White adipose tissue	<i>UCP1</i> negative	Abdomen subcutis	A large unilocular lipid droplet	<i>Tcf21</i>	Lipid storage, secrete hormones and cytokines
Brown adipose tissue	<i>UCP1</i> positive	Neck supraclavicular regions	Multilocular lipid droplets	<i>Zic1</i>	Thermogenesis secrete hormones and cytokines
Beige adipose tissue	<i>UCP1</i> positive	WAT	Multilocular lipid droplets	<i>Tbx1, Epsti1, Cd137, Tmem26</i>	Thermogenesis

WAT, white adipose tissue; *UCP1*, uncoupling protein 1; *Tcf21*, transcription factor 21; *Zic1*, zinc finger protein of the cerebellum 1; *Tbx1*, T-box 1; *Epsti1*, epithelial stromal interaction 1; *Cd137*, tumor necrosis factor receptor superfamily, member 9; *Tmem26*, transmembrane protein 26



BMP4 signaling in the adipocyte commitment of MSCs

The role of BMP4 signaling has been validated in the commitment process of MSCs. C3H10T1/2 cells were isolated from C3H mice in 1973 and can differentiate into various cell types such as adipocytes, osteocytes, chondrocytes, and myoblasts [6, 56]. C3H10T1/2 cells can be faithfully considered as MSCs in terms of their multidirectional differentiation and strong proliferative ability. Their easy isolation, long lifetime in culture, and low culture cost, as well as the absence of ethical concerns, have resulted in the widespread use of these cells. Several studies indicate that BMP4 can induce pluripotent C3H10T1/2 cells to commit to the adipocyte lineage [28, 29]. Upon BMP4 ligand binding to the receptors BMP1A and

BMP2, the receptors phosphorylated SMAD1/5/8, which formed a complex with SMAD4. The complex translocated to the nucleus to regulate *LOX* (lysyl oxidase), *TPT1* (translationally controlled tumor protein 1), and *α B-crystallin* which are target genes of BMP4 signaling in the adipocyte commitment of C3H10T1/2 cells [30, 31]. Treatment of C3H10T1/2 cells with BMP4 revealed their capacity to differentiate into adipocytes, and the expression of adipocyte markers CCAAT/enhancer binding protein- α (C/EBP α), peroxisome proliferator-activated receptor- γ (PPAR γ), and adipocyte protein 2 (AP2) were detected. C3H10T1/2 cells pretreated with BMP4 were subcutaneously implanted into athymic mice, and the cells developed into tissue that was undistinguishable from

adipose tissue [28]. The research of Bowers et al. also confirmed that BMP4 signaling was sufficient and necessary to induce adipocyte lineage determination in C3H10T1/2 cells [8]. A33 cells are a committed preadipocyte cell line derived from C3H10T1/2 cells after 5-azacytidine treatment [8]. Exposure of A33 cells with noggin, a BMP4 binding antagonist, blocked the subsequent differentiation, indicating BMP4 is necessary to maintain commitment [8]. BMP4 also plays a crucial role in regulating human adipose tissue stromal cell commitment. Human adipose tissue stromal cells are also referred to as human adipose stem cells (hASCs), which are a subset of stromal vascular fraction (SVF) cells of adipose tissue [57]. The hASCs are multipotent cells that can differentiate into several different lineages of cells, such as adipocytes, chondrocytes, and osteoblasts. Because hASCs exhibit properties similar to MSCs, some researchers advise that the two cell lines are identical [58]. Inhibition of BMP4 signaling decreased adipogenesis in hASCs [39]. SMAD1/5/8 is phosphorylated by activated BMP receptors. It then associates with SMAD4 and the complex translocates to the nucleus where it regulates gene expression. Overexpression of constitutively active BMP receptors induced adipocyte commitment without exogenous BMP4, whereas overexpression of a dominant-negative BMP receptor suppressed the commitment that was induced by exogenous BMP4 [30]. Also, knockdown of *SMAD4* impeded adipocyte commitment by exogenous BMP4. Treatment with 40 ng/mL BMP4 triggered the adipocyte commitment of hASCs. The committed cells differentiated into adipocytes was associated with increased activation of PPAR γ , APM1, AP2, and GLUT4 [40]. BMP4 has been shown to promote the commitment of the early precursor cells. WISP2, as a secreted protein of adipose precursor cells, can form a complex with ZFP423 without BMP4 stimulation. The addition of BMP4 activated the phosphorylation of SMAD1/5/8, dissociated the WISP2/ZFP423 complex, allowed ZFP423 to enter the nucleus for PPAR γ activation, and committed the cells to the adipose lineage [59]. To date, there have been few studies of BMP4 in BAT. Some research had found that BMP4 combined with BMP7 induced expression of the terminal BAT-specific marker UCP1 in hASCs [32]. BMP4 also induced the formation of brown adipocytes in C3H10T1/2 cells [32]. The expression of UCP1 and the early regulator of brown fat fate protein PRDM16 were induced when C3H10T1/2 cells were pretreated with 20 ng/ml recombinant human BMP4. Implantation of C3H10T1/2 cells pretreated with BMP4 into nude mice resulted in the development of UCP1-positive brown adipocytes. BMP4 increased the white-to-brown transition and the expression of UCP1 and decreased the expression of the white specific marker transcription factor 21 (TCF21) in hASCs [41]. It also enhanced insulin sensitivity, adipocyte cell number, and

whole-body energy expenditure by browning subcutaneous adipose tissue in mature mice [60, 61].

BMP2 signaling in the adipocyte commitment of MSCs

BMP2 signaling in the adipocyte commitment of MSCs remains unclear. Several studies found that BMP2 signaling can induce pluripotent C3H10T1/2 cells to commit to the adipocyte lineage [29–31, 33]. There was a certain degree of plasticity between adipogenesis, chondrogenesis, and osteogenesis, and low-level addition of BMP2 to C3H10T1/2 cells favored adipogenesis [29, 33]. BMP2, just like BMP4, activated the expression and phosphorylation of SMAD1/5/8, which formed a complex with SMAD4. The complex translocated to the nucleus to regulate *LOX*, *TPT1*, and α *B-crystallin* in the adipocyte commitment of C3H10T1/2 cells [30, 31]. Knockdown of *SMAD4* expression prevented the adipocyte commitment of C3H10T1/2 cells by BMP2 [30]. BMP2 induced the adipocyte commitment of C3H10T1/2 cells by induction of PPAR γ expression through activation of SMAD1 and p38 kinase, suggesting a central role of PPAR γ in this commitment event [34]. SMAD6 blocked PPAR γ expression and adipocyte commitment of C3H10T1/2 cells induced by BMP2. BMP2 (100 ng/ml) regulated SMAD signaling, decreased proliferation, and exhibited a donor-dependent dual role, inducing osteogenic or adipogenic differentiation in hASCs [42]. The expression of alkaline phosphatase, an osteoblast-specific marker, as well as the mineralization levels were notably enhanced in some donor cell lines by BMP2 stimulation, while diminishing lipid formation in other cell lines. The hASCs derived from older donors had higher potential to demonstrate BMP2-induced adipogenesis than cells from younger individuals. BMP2 induced SMAD1/5 phosphorylation, activation, and translocation to the nucleus to regulate adipocyte commitment, and the expression of adipogenic marker AP2 was remarkably detected [42]. The relevance of the origin of BMP2 production is that it influences the adipocyte commitment of hASCs. BMP2 produced in mammalian cells, Chinese hamster ovary (CHO) cells, induced higher levels of lipid formation compared with BMP2 of *Escherichia coli* origin [42]. Conversely, some studies have shown that BMP2 inhibits adipocyte commitment [28, 62, 63]. Tang et al. found that adding 50 ng/ml BMP2 to C3H10T1/2 cells until post-confluency followed by standard adipocyte differentiation failed to cause an accumulation of triglyceride or the expression of the adipocyte-specific marker AP2 [28]. Thus, it can be concluded that BMP2 inhibits adipocyte commitment in C3H10T1/2 cells. However, the resulting research of Tang et al. considered that BMP2 can induce pluripotent C3H10T1/2 cells to commit to the adipocyte lineage, as shown previously [30, 31]. Despite some common ground,

we identified some disagreements through a comparative study of these papers and analyses. C3H10T1/2 cells in these studies were all plated at low density with purified recombinant BMP2 at the same concentration of 50 ng/ml. C3H10T1/2 cells were induced to differentiate using the adipocyte differentiation protocol after the cells reached post-confluency. The difference was that they added different concentrations of an adipogenic induction cocktail consisting of insulin, dexamethasone, and 3-isobutyl-1-methylxanthine. The later research of Tang et al. changed DMEM containing 10% calf serum (CS) for DMEM containing 10% fetal bovine serum (FBS) in the process of differentiation because FBS may have some effects on adipocyte commitment. Future research should focus on the effect of the different components between CS and FBS in adipocyte commitment. Other studies have shown that BMP2 enhanced osteoblast commitment in a bone marrow stromal cell line by increasing the expression of osteoblast-specific markers OSF2/CBFA1 and mineralized nodule formation, concomitantly inhibiting adipocyte maturation [62, 63]. The effects of BMP2 on adipocyte differentiation reduced adipogenesis and decreased the levels of leptin and the formation of cytoplasmic lipid droplets. It should be noted that in the article, the multipotent murine bone marrow stromal cell line BMS2 cannot be regarded as MSCs, because up to 40% or more of the BMS2 population could spontaneously differentiate into adipocytes only in the presence of adipogenic induction cocktail. In conclusion, BMP2 may guide the commitment destiny of MSCs depending on the donor line, the concentrations of an adipogenic induction cocktail, and the type of serum added to DMEM. Further studies are required to clarify how BMP2 regulates the commitment of MSCs.

BMP7 signaling in the adipocyte commitment of MSCs

Many studies have demonstrated that BMP7 has an important role in brown adipocyte lineage determination [32, 35, 36, 41, 44, 46]. BMP7 triggered the commitment of C3H10T1/2 cells to a brown adipocyte lineage with a significant increase in lipid accumulation and expression of UCP1 [32, 35]. BMP7 stimulated cell proliferation and differentiation in mouse bone marrow stromal stem cells (BMSCs) and human adult MSCs [44, 46]. Low concentrations stimulated adipocyte differentiation, whereas it was inhibited at higher concentrations under BMP7 treatment in mouse BMSCs [44]. BMP7 promoted the adipogenic differentiation of human BMSCs but not the osteogenic or chondrogenic lineage development in a high-density micro-mass culture [46]. BMP7 or BMP4 also induced adipogenic differentiation by lipid accumulation and PPAR γ expression, while inducing the white-to-brown transition by increasing UCP1 expression and decreasing the expression of white specific marker TCF21 in

hASCs dependent on the donors [41]. Implantation of BMP7-treated C3H10T1/2 cells subcutaneously into the sternal region of athymic nude mice resulted in the development of adipose tissue containing brown adipocytes in vivo [35]. Although *BMP7* knockout mice died shortly after birth, BAT mass was reduced and there was an almost complete absence of UCP1 in the knockout newborns [35]. Mice injected with adenovirus expressing BMP7 showed a significant increase in brown fat mass and energy expenditure [35]. BMP7 also drove human adipogenic stem cells to form metabolically active beige adipocytes [43]. Human adipogenic stem cells are also referred to hASCs. Exposure of hASCs to human BMP7 was associated with a significant increase in UCP1 expression and beige specific marker gene expression as well as metabolic activity [43].

TGF- β signaling in the adipocyte commitment of MSCs

TGF- β is another important molecule that regulates the adipocyte commitment of MSCs. TGF- β signaling inhibited adipocyte commitment via SMAD3 signaling [64]. TGF- β inhibited C/EBPs and PPAR γ expression and induced PPAR γ phosphorylation, resulting in inhibition of the adipocyte commitment of BMSCs [45]. Deleting TGF- β in mesenchymal cells resulted in a marked expansion of adipocytes in the bone marrow of mice and significantly increased the expression of PPAR γ and FABP4 associated with adipocyte commitment, while osteoblasts were significantly reduced [65]. Since PPAR γ is a master regulator in adipocyte commitment, TGF- β suppresses the adipocyte commitment of MSCs through phosphorylation of PPAR γ . Several articles focused on the modulation of TGF- β signaling in adipogenic differentiation of hASCs and the multipotent mesenchymal stromal cells misleading the MSC origin. In these studies, TGF- β inhibited adipogenic differentiation in these cells [66, 67]. However, it should be noted that the hASCs and the multipotent mesenchymal stromal cells in these articles cannot be regarded as MSCs because more of the population could spontaneously differentiate to adipocytes with exposure to only adipogenic induction cocktail. In a sense, the cells can be regarded as pre-adipocytes. We should not only focus on the surface but also think carefully about what MSCs are, perhaps according to their property. Although some studies have reported that TGF- β suppresses the adipocyte commitment of MSCs, recent studies showed that TGF- β promotes adipocyte commitment. A single pulse dose treatment with TGF- β during the commitment phase led to increased adipogenesis of human BMSCs, while continuous treatment during the whole phase (commitment phase and differentiation phase) inhibited adipogenesis [47]. The reason could be TGF- β promoted the first step adipocyte commitment, but inhibited the

second step adipogenic differentiation of BMSCs by repressing the function of C/EBP β and C/EBP δ during the differentiation phase [68]. TGF- β promoted osteogenic and adipogenic differentiation of a clone isolated from human BMSC lines, which have limited differentiation capacity, suggesting an epigenetic change rather than a gene mutation [48].

Myostatin signaling in the adipocyte commitment of MSCs

Myostatin, also called GDF8, is a well-known regulator of myogenesis inhibition. Some researchers found that myostatin can promote adipogenesis in MSCs. C3H10T1/2 cells were incubated in regular medium with graded concentrations of myostatin and found that there were many oil droplets present after a 2-week incubation, as visualized by Oil Red O staining [36]. Myostatin failed to trigger adipogenesis in the 3T3-L1 preadipocyte cell line, but it was also a substitute for dexamethasone, which can trigger adipogenesis and induce small and apparently immature adipocytes of C3H10T1/2 cells [37]. Myostatin gene (*Mstn*) knockout in mice (*Mstn*^{-/-} mice) increased myogenesis and decreased adipogenesis [69, 70]. *Mstn*^{-/-} mice had reduced adipogenesis and consequently decreased leptin secretion and C/EBP α and PPAR γ levels [69]. *Mstn*^{-/-} mice also had suppressed obesity and glucose metabolism than control mice [70]. The molecular mechanism of the phenomenon by which *Mstn*^{-/-} mice had reduced adipogenesis was that the absence of *Mstn* resulted in enhanced peripheral tissue fatty acid oxidation and brown adipose formation in white adipose tissue [71]. Some studies found that myostatin inhibited adipogenesis in MSCs. The presence of myostatin downregulated the expression of adipocyte markers, induced SMAD3 phosphorylation, and activated β -catenin in human BMSCs [49]. As previously mentioned, BMP7 triggered the adipocyte commitment of C3H10T1/2 cells. Myostatin prevented BMP7 signaling by blocking BMP7 receptor binding, resulting in reducing adipogenesis in C3H10T1/2 cells [38]. Myostatin is an important factor in brown adipocyte lineage determination. *Mstn*^{-/-} mouse embryonic fibroblasts (MEFs) had enhanced expression levels of brown adipocyte markers including UCP1, PRDM16, and PGC1 compared with the control group undergoing adipogenic differentiation [72]. From the above statement, we can conclude that myostatin plays an extremely complicated role in the adipocyte commitment of MSCs.

Conclusion

TGF- β superfamily signaling has an important function in regulating the adipocyte commitment of MSCs. TGF- β s ligands such as TGF- β and myostatin use SMAD2/3

signaling to regulate adipocyte commitment. However, BMPs ligands such as BMP2/4/7 use SMAD1/5/8 signaling to regulate adipocyte commitment. BMP4 signaling is explicitly sufficient to induce adipocyte lineage determination and the formation of brown adipocytes in MSCs. BMP2 and myostatin signaling have an effect on the adipocyte commitment of MSCs. Some studies found that BMP2 or myostatin signaling can promote adipocyte commitment in MSCs, while animal experiments showed the same results. However, some studies found that BMP2 or myostatin signaling can inhibit this commitment. It is unclear whether they promote adipocyte commitment or inhibit it. It is well known that BMP2 and myostatin signaling were originally identified based on their ability to induce osteogenesis and myogenesis inhibition. BMP2 and myostatin signaling may play a small part in adipocyte commitment, perhaps due to different donor lines, the concentrations of an adipogenic induction cocktail, or serum types in culture medium. Previous articles have demonstrated that TGF- β signaling promoted the proliferation of MSCs and suppressed the adipocyte commitment of MSCs by inhibiting C/EBPs and PPAR γ expression. Recent studies have showed that TGF- β promoted adipocyte commitment. The difference was that they studied the adipocyte commitment of BMSCs isolated from different species: mice and humans [45, 47]. MSC origin may affect adipocyte commitment under TGF- β signaling. Various clones isolated from human BMSC lines have different differentiation capacities, which may be the factor affecting the commitment event. More studies are required to address the effect of BMP2, TGF- β , and myostatin on adipocyte commitment in MSCs. BMP7 mainly promotes the commitment of MSCs to a brown adipocyte lineage. Other TGF- β superfamily members such as BMP3, BMP6, and GDF11 have been evaluated as to whether they affect commitment. BMP3 stimulated the proliferation rather than the commitment of MSCs through TGF- β /activin signaling [73]. BMP6 is highly homologous with BMP7, but BMP6 mainly affects the adipocyte differentiation process of MSCs rather than the commitment process. BMP6 governs the lineage adipocyte commitment of other cell types. BMP6 stimulation in muscle precursor cells C2C12 was sufficient to induce brown fat determination [74]. Little research has been performed on GDF11, but a recent article discussed the effect of GDF11 on adipocyte commitment. The data indicated that GDF11 inhibited adipogenic differentiation in human MSCs by activating SMAD2/3-dependent TGF- β signaling [75]. It is becoming clear that R-SMADs have an impact on the commitment of MSCs, but we still know very little about Co-SMAD (SMAD4). SMAD4 is common to all receptor pathways, and its mechanism is complicated. A recent article reported that SMAD4 depletion in C3H10T1/2 cells and hASCs reduced the nuclear retention of TAZ, a

transcriptional coactivator with a PDZ-binding motif, resulting in enhanced adipogenesis of the MSCs through dissociating TAZ–PPAR γ interactions [76]. However, this study runs counter to the above conclusion [30]. Both studies used RNA interference to silence *SMAD4* in C3H10T1/2 cells. One study added BMP2 protein but the other did not, causing a different outcome because the function of *SMAD4* was independent of the TGF- β signaling in some cases, and sometimes BMP2 had a role in adipocyte commitment independent of TGF- β /SMAD signaling. PPAR γ is a master transcription factor in adipocyte commitment. BMP2/4 signaling induces adipocyte commitment of MSCs by activating PPAR γ expression. A recent novel study reported the epigenetics of adipogenic commitment under TGF- β /SMAD signaling. *miR-20a* promoted C3H10T1/2 cells to commit, and this function may depend upon its direct inhibitory effect on lysine-specific demethylase 6B (KDM6B) and TGF- β signaling [77]. Thus far, the research on adipocyte commitment in MSCs has had some limitations: (1) the disunified source and unclear features of MSCs, (2) the limited research available on the role of the entire TGF- β superfamily, and (3) the unclear relationships between the TGF- β superfamily members. Perhaps future studies should focus more on these limitations.

The TGF- β superfamily members have an effect on adipocytokines secreted from WAT and BAT as well as beige adipose tissue, influencing white-to-brown transition and disease. Future studies of TGF- β superfamily signaling will offer new approaches in treating obesity, diabetes mellitus, and obesity-related metabolism syndrome.

Abbreviations

AP2: Adipocyte protein 2; BAT: Brown adipose tissue; BMPs: Bone morphogenetic proteins; BMSCs: Bone marrow stromal stem cells; C/EBP α : CCAAT/enhancer binding protein- α ; CS: Calf serum; FBS: Fetal bovine serum; GDFs: Growth differentiation factors; hASCs: Human adipose stem cells; KDM6B: Lysine-specific demethylase 6B; LOX: Lysyl oxidase; MEFs: Mouse embryonic fibroblasts; MSCs: Mesenchymal stem cells; PPAR γ : Peroxisome proliferator-activated receptor- γ ; SVF: Stromal vascular fraction; TCF21: Transcription factor 21; TGF- β : Transforming growth factor- β ; TPT1: Translationally controlled tumor protein 1; UCP1: Uncoupling protein 1; WAT: White adipose tissue

Acknowledgements

We thank the National Natural Science Foundation of China and the Ph.D. Programs Foundation of Henan Polytechnic University.

Authors' contributions

SNL drafted the manuscript. SNL and JFW participated in the paper modification. Both authors read and approved the final manuscript.

Funding

This work was supported by the National Natural Science Foundation of China (grant number 81700775) and the Ph.D. Programs Foundation of Henan Polytechnic University (grant number B2017-61).

Availability of data and materials

Not applicable.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹School of Medicine, Henan Polytechnic University, Jiaozuo 454000, Henan, China. ²School of Food and Bioengineering, Henan University of Science and Technology, Luoyang, Henan, China.

Received: 14 August 2019 Revised: 18 December 2019

Accepted: 5 January 2020 Published online: 29 January 2020

References

- Friedenstein AJ, Petrakova KV, Kurolesova AI, Frolova GP. Heterotopic of bone marrow. Analysis of precursor cells for osteogenic and hematopoietic tissues. *Transplantation*. 1968;6:230–47.
- Friedenstein AJ, Deriglasova UF, Kulagina NN, Panasuk AF, Rudakowa SF, Luria EA, et al. Precursors for fibroblasts in different populations of hematopoietic cells as detected by the in vitro colony assay method. *Exp Hematol*. 1974;2:83–92.
- Mushahary D, Spittler A, Kasper C, Weber V, Charwat V. Isolation, cultivation, and characterization of human mesenchymal stem cells. *Cytometry A*. 2018; 93:19–31.
- Compte M, Cuesta AM, Sanchez-Martin D, Alonso-Camino V, Vicario JL, Sanz L, et al. Tumor immunotherapy using gene-modified human mesenchymal stem cells loaded into synthetic extracellular matrix scaffolds. *Stem Cells*. 2009;27:753–60.
- Pittenger MF, Mackay AM, Beck SC, Jaiswal RK, Douglas R, Mosca JD, et al. Multilineage potential of adult human mesenchymal stem cells. *Science*. 1999;284:143–7.
- Lin CS, Xin ZC, Deng CH, Ning H, Lin G, Lue TF. Defining adipose tissue-derived stem cells in tissue and in culture. *Histol Histopathol*. 2010;25:807–15.
- McBeath R, Pirone DM, Nelson CM, Bhadriraju K, Chen CS. Cell shape, cytoskeletal tension, and RhoA regulate stem cell lineage commitment. *Dev Cell*. 2004;6:483–95.
- Bowers RR, Kim JW, Otto TC, Lane MD. Stable stem cell commitment to the adipocyte lineage by inhibition of DNA methylation: role of the BMP-4 gene. *Proc Natl Acad Sci U S A*. 2006;103:13022–7.
- Konieczny SF, Emerson CP Jr. 5-Azacytidine induction of stable mesodermal stem cell lineages from 10T1/2 cells: evidence for regulatory genes controlling determination. *Cell*. 1984;38:791–800.
- Massague J, Wotton D. Transcriptional control by the TGF-beta/Smad signaling system. *EMBO J*. 2000;19:1745–54.
- Zhang YE. Mechanistic insight into contextual TGF-beta signaling. *Curr Opin Cell Biol*. 2018;51:1–7.
- David CJ, Massague J. Contextual determinants of TGFbeta action in development, immunity and cancer. *Nat Rev Mol Cell Biol*. 2018;19:419–35.
- Derynck R, Zhang YE. Smad-dependent and Smad-independent pathways in TGF-beta family signalling. *Nature*. 2003;425:577–84.
- Shi Y, Massague J. Mechanisms of TGF-beta signaling from cell membrane to the nucleus. *Cell*. 2003;113:685–700.
- Miyazawa K, Miyazono K. Regulation of TGF-beta family signaling by inhibitory Smads. *Cold Spring Harb Perspect Biol*. 2017;9:a022095.
- Hill CS. Nucleocytoplasmic shuttling of Smad proteins. *Cell Res*. 2009;19:36–46.
- Feng XH, Derynck R. Specificity and versatility in tgfbeta signaling through Smads. *Annu Rev Cell Dev Biol*. 2005;21:659–93.
- Lin X, Duan X, Liang YY, Su Y, Wrighton KH, Long J, et al. PPM1A functions as a Smad phosphatase to terminate TGFbeta signaling. *Cell*. 2006;125:915–28.
- Kavsak P, Rasmussen RK, Causing CG, Bonni S, Zhu H, Thomsen GH, et al. Smad7 binds to Smurf2 to form an E3 ubiquitin ligase that targets the TGF beta receptor for degradation. *Mol Cell*. 2000;6:1365–75.
- Hata A, Lagna G, Massague J, Hemmati-Brivanlou A. Smad6 inhibits BMP/Smad1 signaling by specifically competing with the Smad4 tumor suppressor. *Genes Dev*. 1998;12:186–97.

21. Covas DT, Panepucci RA, Fontes AM, Silva WA Jr, Orellana MD, Freitas MC, et al. Multipotent mesenchymal stromal cells obtained from diverse human tissues share functional properties and gene-expression profile with CD146+ perivascular cells and fibroblasts. *Exp Hematol*. 2008;36:642–54.
22. Wolfgang MJ, Lane MD. Control of energy homeostasis: role of enzymes and intermediates of fatty acid metabolism in the central nervous system. *Annu Rev Nutr*. 2006;26:23–44.
23. Hilton C, Karpe F, Pinnick KE. Role of developmental transcription factors in white, brown and beige adipose tissues. *Biochim Biophys Acta*. 2015;1851:686–96.
24. Wu J, Bostrom P, Sparks LM, Ye L, Choi JH, Giang AH, et al. Beige adipocytes are a distinct type of thermogenic fat cell in mouse and human. *Cell*. 2012;150:366–76.
25. Seale P, Bjork B, Yang W, Kajimura S, Chin S, Kuang S, et al. PRDM16 controls a brown fat/skeletal muscle switch. *Nature*. 2008;454:961–7.
26. de Jong JM, Larsson O, Cannon B, Nedergaard J. A stringent validation of mouse adipose tissue identity markers. *Am J Physiol Endocrinol Metab*. 2015;308:E1085–105.
27. Peirce V, Carobbio S, Vidal-Puig A. The different shades of fat. *Nature*. 2014;510:76–83.
28. Tang QQ, Otto TC, Lane MD. Commitment of C3H10T1/2 pluripotent stem cells to the adipocyte lineage. *Proc Natl Acad Sci U S A*. 2004;101:9607–11.
29. Ahrens M, Ankenbauer T, Schroder D, Hollnagel A, Mayer H, Gross G. Expression of human bone morphogenetic proteins-2 or -4 in murine mesenchymal progenitor C3H10T1/2 cells induces differentiation into distinct mesenchymal cell lineages. *DNA Cell Biol*. 1993;12:871–80.
30. Huang H, Song TJ, Li X, Hu L, He Q, Liu M, et al. BMP signaling pathway is required for commitment of C3H10T1/2 pluripotent stem cells to the adipocyte lineage. *Proc Natl Acad Sci U S A*. 2009;106:12670–5.
31. Huang HY, LL Hu, TJ Song, X Li, Q He, X Sun, et al. Involvement of cytoskeleton-associated proteins in the commitment of C3H10T1/2 pluripotent stem cells to adipocyte lineage induced by BMP2/4. *Mol Cell Proteomics*. 2011;10:M110 002691.
32. Xue R, Wan Y, Zhang S, Zhang Q, Ye H, Li Y. Role of bone morphogenetic protein 4 in the differentiation of brown fat-like adipocytes. *Am J Physiol Endocrinol Metab*. 2014;306:E363–72.
33. Wang EA, Israel DI, Kelly S, Luxenberg DP. Bone morphogenetic protein-2 causes commitment and differentiation in C3H10T1/2 and 3T3 cells. *Growth Factors*. 1993;9:57–71.
34. Hata K, Nishimura R, Ikeda F, Yamashita K, Matsubara T, Nokubi T, et al. Differential roles of Smad1 and p38 kinase in regulation of peroxisome proliferator-activating receptor gamma during bone morphogenetic protein 2-induced adipogenesis. *Mol Biol Cell*. 2003;14:545–55.
35. Tseng YH, Kokkotou E, Schulz TJ, Huang TL, Winnay JN, Taniguchi CM, et al. New role of bone morphogenetic protein 7 in brown adipogenesis and energy expenditure. *Nature*. 2008;454:1000–4.
36. Artaza JN, Bhasin S, Magee TR, Reisz-Porszasz S, Shen R, Groome NP, et al. Myostatin inhibits myogenesis and promotes adipogenesis in C3H 10T1/2 mesenchymal multipotent cells. *Endocrinology*. 2005;146:3547–57.
37. Feldman BJ, Streeper RS, Farese RV Jr, Yamamoto KR. Myostatin modulates adipogenesis to generate adipocytes with favorable metabolic effects. *Proc Natl Acad Sci U S A*. 2006;103:15675–80.
38. Rebbapragada A, Benchabane H, Wrana JL, Celeste AJ, Attisano L. Myostatin signals through a transforming growth factor beta-like signaling pathway to block adipogenesis. *Mol Cell Biol*. 2003;23:7230–42.
39. Gustafson B, Hammarstedt A, Hedjazifar S, Hoffmann JM, Svensson PA, Grimsby J, et al. BMP4 and BMP antagonists regulate human white and beige adipogenesis. *Diabetes*. 2015;64:1670–81.
40. Gustafson B, Smith U. The WNT inhibitor Dickkopf 1 and bone morphogenetic protein 4 rescue adipogenesis in hypertrophic obesity in humans. *Diabetes*. 2012;61:1217–24.
41. Elsen M, Raschke S, Tennagels N, Schwahn U, Jelenik T, Roden M, et al. BMP4 and BMP7 induce the white-to-brown transition of primary human adipose stem cells. *Am J Physiol Cell Physiol*. 2014;306:C431–40.
42. Vanhatupa S, Ojansivu M, Autio R, Juntunen M, Miettinen S. Bone morphogenetic protein-2 induces donor-dependent osteogenic and adipogenic differentiation in human adipose stem cells. *Stem Cells Transl Med*. 2015;4:1391–402.
43. Okla M, Ha JH, Temel RE, Chung S. BMP7 drives human adipogenic stem cells into metabolically active beige adipocytes. *Lipids*. 2015;50:111–20.
44. Chen TL, Shen WJ, Kraemer FB. Human BMP-7/OP-1 induces the growth and differentiation of adipocytes and osteoblasts in bone marrow stromal cell cultures. *J Cell Biochem*. 2001;82:187–99.
45. Ahdjoudj S, Kaabeche K, Holy X, Fromiguet O, Modrowski D, Zerath E, et al. Transforming growth factor-beta inhibits CCAAT/enhancer-binding protein expression and PPARgamma activity in unloaded bone marrow stromal cells. *Exp Cell Res*. 2005;303:138–47.
46. Neumann K, Endres M, Ringe J, Flath B, Manz R, Haupt T, et al. BMP7 promotes adipogenic but not osteo-/chondrogenic differentiation of adult human bone marrow-derived stem cells in high-density micro-mass culture. *J Cell Biochem*. 2007;102:626–37.
47. Elsafadi M, Manikandan M, Attaya M, Dawud RA, Almalki S, Kaimkhani ZA, et al. SERPINB2 is a novel TGFbeta-responsive lineage fate determinant of human bone marrow stromal cells. *Sci Rep*. 2017;7:10797.
48. Elsafadi M, Shinwari T, Al-Malki S, Manikandan M, Mahmood A, Aldahmash A, et al. Convergence of TGFbeta and BMP signaling in regulating human bone marrow stromal cell differentiation. *Sci Rep*. 2019;9:4977.
49. Guo W, Flanagan J, Jasuja R, Kirkland J, Jiang L, Bhasin S. The effects of myostatin on adipogenic differentiation of human bone marrow-derived mesenchymal stem cells are mediated through cross-communication between Smad3 and Wnt/beta-catenin signaling pathways. *J Biol Chem*. 2008;283:9136–45.
50. Dani C, Smith AG, Dessolin S, Leroy P, Staccini L, Villageois P, et al. Differentiation of embryonic stem cells into adipocytes in vitro. *J Cell Sci*. 1997;110(Pt 11):1279–85.
51. Sanchez-Gurmaches J, Guertin DA. Adipocytes arise from multiple lineages that are heterogeneously and dynamically distributed. *Nat Commun*. 2014;5:4099.
52. Barbatelli G, Murano I, Madsen L, Hao Q, Jimenez M, Kristiansen K, et al. The emergence of cold-induced brown adipocytes in mouse white fat depots is determined predominantly by white to brown adipocyte transdifferentiation. *Am J Physiol Endocrinol Metab*. 2010;298:E1244–53.
53. Sanchez-Gurmaches J, Hung CM, Guertin DA. Emerging complexities in adipocyte origins and identity. *Trends Cell Biol*. 2016;26:313–26.
54. Qian SW, Wu MY, Wang YN, Zhao YX, Zou Y, Pan JB, et al. BMP4 facilitates beige fat biogenesis via regulating adipose tissue macrophages. *J Mol Cell Biol*. 2019;11:14–25.
55. Lee YH, Petkova AP, Konkara AA, Granneman JG. Cellular origins of cold-induced brown adipocytes in adult mice. *FASEB J*. 2015;29:286–99.
56. Reznikoff CA, Brankow DW, Heidelberger C. Establishment and characterization of a cloned line of C3H mouse embryo cells sensitive to postconfluence inhibition of division. *Cancer Res*. 1973;33:3231–8.
57. Bourin P, Bunnell BA, Casteilla L, Dominici M, Katz AJ, March KL, et al. Stromal cells from the adipose tissue-derived stromal vascular fraction and culture expanded adipose tissue-derived stromal/stem cells: a joint statement of the International Federation for Adipose Therapeutics and Science (IFATS) and the International Society for Cellular Therapy (ISCT). *Cytotherapy*. 2013;15:641–8.
58. Baer PC. Adipose-derived mesenchymal stromal/stem cells: an update on their phenotype in vivo and in vitro. *World J Stem Cells*. 2014;6:256–65.
59. Hammarstedt A, Hedjazifar S, Jenndahl L, Gogg S, Grunberg J, Gustafson B, et al. WISP2 regulates preadipocyte commitment and PPARgamma activation by BMP4. *Proc Natl Acad Sci U S A*. 2013;110:2563–8.
60. Hoffmann JM, Grunberg JR, Church C, Elias I, Palsdottir V, Jansson JO, et al. BMP4 gene therapy in mature mice reduces BAT activation but protects from obesity by browning subcutaneous adipose tissue. *Cell Rep*. 2017;20:1038–49.
61. Tang Y, Qian SW, Wu MY, Wang J, Lu P, Li X, et al. BMP4 mediates the interplay between adipogenesis and angiogenesis during expansion of subcutaneous white adipose tissue. *J Mol Cell Biol*. 2016;8:302–12.
62. Gimble JM, Morgan C, Kelly K, Wu X, Dandapani V, Wang CS, et al. Bone morphogenetic proteins inhibit adipocyte differentiation by bone marrow stromal cells. *J Cell Biochem*. 1995;58:393–402.
63. Gori F, Thomas T, Hicok KC, Spelsberg TC, Riggs BL. Differentiation of human marrow stromal precursor cells: bone morphogenetic protein-2 increases OSF2/CBFA1, enhances osteoblast commitment, and inhibits late adipocyte maturation. *J Bone Miner Res*. 1999;14:1522–35.
64. Ng F, Boucher S, Koh S, Sastry KS, Chase L, Lakshminath U, et al. PDGF, TGF-beta, and FGF signaling is important for differentiation and growth of mesenchymal stem cells (MSCs): transcriptional profiling can identify markers and signaling pathways important in differentiation of MSCs into adipogenic, chondrogenic, and osteogenic lineages. *Blood*. 2008;112:295–307.
65. Abou-Ezzi G, Supakorndej T, Zhang J, Anthony B, Kramps J, Celik H, et al. TGF-beta signaling plays an essential role in the lineage specification of

- mesenchymal stem/progenitor cells in fetal bone marrow. *Stem Cell Reports*. 2019;13:48–60.
66. Kim YJ, Hwang SJ, Bae YC, Jung JS. MiR-21 regulates adipogenic differentiation through the modulation of TGF-beta signaling in mesenchymal stem cells derived from human adipose tissue. *Stem Cells*. 2009;27:3093–102.
 67. Walenda G, Abnaof K, Joussem S, Meurer S, Smeets H, Rath B, et al. TGF-beta1 does not induce senescence of multipotent mesenchymal stromal cells and has similar effects in early and late passages. *PLoS One*. 2013;8:e77656.
 68. Choy L, Derynck R. Transforming growth factor-beta inhibits adipocyte differentiation by Smad3 interacting with CCAAT/enhancer-binding protein (C/EBP) and repressing C/EBP transactivation function. *J Biol Chem*. 2003;278:9609–19.
 69. Lin J, Arnold HB, Della-Fera MA, Azain MJ, Hartzell DL, Baile CA. Myostatin knockout in mice increases myogenesis and decreases adipogenesis. *Biochem Biophys Res Commun*. 2002;291:701–6.
 70. McPherron AC, Lee SJ. Suppression of body fat accumulation in myostatin-deficient mice. *J Clin Invest*. 2002;109:595–601.
 71. Zhang C, McFarlane C, Lokireddy S, Masuda S, Ge X, Gluckman PD, et al. Inhibition of myostatin protects against diet-induced obesity by enhancing fatty acid oxidation and promoting a brown adipose phenotype in mice. *Diabetologia*. 2012;55:183–93.
 72. Braga M, Pervin S, Norris K, Bhasin S, Singh R. Inhibition of in vitro and in vivo brown fat differentiation program by myostatin. *Obesity (Silver Spring)*. 2013;21:1180–8.
 73. Stewart A, Guan H, Yang K. BMP-3 promotes mesenchymal stem cell proliferation through the TGF-beta/activin signaling pathway. *J Cell Physiol*. 2010;223:658–66.
 74. Sharma A, Huard C, Vernochet C, Ziemek D, Knowlton KM, Tyminski E, et al. Brown fat determination and development from muscle precursor cells by novel action of bone morphogenetic protein 6. *PLoS One*. 2014;9:e92608.
 75. Luo H, Guo Y, Liu Y, Wang Y, Zheng R, Ban Y, et al. Growth differentiation factor 11 inhibits adipogenic differentiation by activating TGF-beta/Smad signalling pathway. *Cell Prolif*. 2019;52:e12631.
 76. Park JS, Kim M, Song NJ, Kim JH, Seo D, Lee JH, et al. A reciprocal role of the Smad4-Taz axis in osteogenesis and adipogenesis of mesenchymal stem cells. *Stem Cells*. 2019;37:368–81.
 77. Zhou J, Guo F, Wang G, Wang J, Zheng F, Guan X, et al. miR-20a regulates adipocyte differentiation by targeting lysine-specific demethylase 6b and transforming growth factor-beta signaling. *Int J Obes*. 2015;39:1282–91.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.