Could Helicobacter pylori cause an abnormal immune response?

Genel Sur¹, Lucia Sur², Anamaria Girbovan³, Gabriel Samasca⁴, Giulia Alexandru⁵, Rahela Carpa⁵, Iulia Lupan⁶

To the Editor

Rostami-Nejad et al and Samasca et al. found that the screening of H. pylori infection is necessary in patients with CD and iron deficiency (1,2). In a recent our study, the prevalence of H. pylori infection in our pediatric lot was 12%. 38 patients were positive for IgA H. Pylori antibodies. We performed IgA tissue transglutaminase antibodies to all 38 patients with positive results for IgA H. pylori antibodies, however, the results were negative. But, clinical diagnosis of children with H. pylori infection was diversified, moreover, nutritional diseases were the most frequent. H. pylori was involved in the following autoimmune diseases: type 1 diabetes mellitus (T1DM), thyroiditis and arthritis (Table 1). CD may be associated with H. pylori gastritis infection (3) however, few studies revealed this aspect. The association between H. pylori infection and thyroiditis in patients with T1DM was observed (4). H. pylori and patterns of gut microbiome have associated with low risk of T1DM (5). We also observed an association between H. pylori and T1DM in our study. Therefore, could autoimmunity be triggered by a decreased exposure to bacterial antigens? In our study, H. Pylori was associated with autoimmune diseases such as: type 1 diabetes mellitus, thyroiditis or arthritis but further studies are necessary for this etiology.

E-mail: Gabriel.Samasca@umfcluj.ro

Conflict of interests

The authors declare that they have no conflict of interest.

Table 1. Clinical diagnosis in patients with positive *H. pylori* infection

Diagnostic	Absolute
	frequency
Mild or moderate protein-energy	5
malnutrition	
Diabetes mellitus type 1	4
Gastritis/duodenitis	4
Nutritional anemia	3
Urticaria and erythema	2
Diarrhoea and gastroenteritis of presumed	1
infectious origin	
Bacterial infection	1
Benign tumor of the breast	1
Hereditary deficiency through missing	1
factor VIII	
Other coagulation abnormalities	1
Other bleeding disorders	1
Thyroiditis	1
Obesity	1
Pulmonary valve disorders	1
Other chronic obstructive pulmonary	1
disease	
Asthma	1
Acute appendicitis	1
Erythema nodosum	1
Other abnormalities in of pigmentation	1
Infectious arthropathy	1
Other arthritis	1
Back pain	1
Abnormal heart rhythm	1
Bleeding from respiratory system	1
Nausea and vomiting	1
TOTAL	38

¹Department of Pediatrics II, Iuliu Hatieganu University of Medicine, Romania

² Department of Pediatrics I, Iuliu Hatieganu University of Medicine, Romania

³ Department of Immunology II, Iuliu Hatieganu University of Medicine, Romania

⁴ Emergency Hospital for Children, Romania

⁵ Department of Microbiology, Babeş-Bolyai University, Romania

⁶ Department of Molecular Biology and Biotechnology, Babeş-Bolyai University, Romania

Received: 15 January 2017 Accepted: 27 February 2017 Reprint or Correspondence: Gabriel Samasca, MD. Emergency Hospital for Children, Romania.

References

- 1.Rostami-Nejad M, Aldulaimi D, Livett H, Rostami K. H.pylori associated with iron deficiency anemia even in celiac disease patients; strongly evidence based but weakly reflected in practice. Gastroenterol Hepatol Bed Bench 2015;8:178-82.
- 2. Samasca G, Deleanu D, Sur G, Lupan I, Alexandru GD, Carpa R. Is it necessary to screen Helicobacter pylori infection in patients with celiac disease and iron deficiency? Gastroenterol Hepatol Bed Bench 2016;9:345.
- 3. Aydogdu S, Cakir M, Yuksekkaya HA, Tumgor G, Baran M, Arikan C, et al. Helicobacter pylori infection in children with celiac disease. Scand J Gastroenterol 2008;43:1088-93.
- 4. Zekry OA, Abd Elwahid HA. The association between Helicobacter pylori infection, type 1 diabetes mellitus, and autoimmune thyroiditis. J Egypt Public Health Assoc 2013;88:143-7.
- 5. Kondrashova A, Hyöty H. Role of viruses and other microbes in the pathogenesis of type 1 diabetes. Int Rev Immunol 2014;33:284-95.