



OPEN

Perioperative hemodynamic instability in pheochromocytoma and sympathetic paraganglioma patients

Jung Hee Kim^{1,9}, Hyung-Chul Lee^{2,9}, Su-jin Kim^{3,4,5,6}✉, Soo Bin Yoon², Sung Hye Kong¹, Hyeong Won Yu⁷, Young Jun Chai⁸, June Young Choi⁷, Kyu Eun Lee^{3,4,5,6}, Kwang-Woong Lee³, Seung-Kee Min³, Chan Soo Shin¹ & Kyu Joo Park³

For pheochromocytoma and sympathetic paraganglioma (PPGL), surgery can be used as a curative treatment; however, the life-threatening risk of perioperative hemodynamic instability (HI) presents challenges. This study aimed to analyze the incidence and predictive factors of perioperative HI. The electronic medical records of 114 consecutive patients who underwent surgery for PPGLs at our institution were retrospectively reviewed. HI was defined as one or more episodes of systolic blood pressure > 200 mmHg or mean blood pressure < 60 mmHg during surgery. The factors predictive of perioperative HI were determined using both univariate and multivariate analyses. Intraoperative HI occurred in 79 (69.3%) patients. In multivariate analysis, α -adrenergic receptor blocker duration (days) (odds ratio, 1.015; 95% confidence interval, 1.001–1.029) was a predictor for intraoperative HI. Postoperative hypotension occurred in 36 (31.6%) patients. Higher urine epinephrine levels, and greater preoperative highest heart rate (HR) were predictive factors for postoperative hypotension in PPGL patients. Caution should be taken in perioperative management for PPGL, especially with long duration of α -adrenergic receptor blocker use, higher urine epinephrine levels, and greater preoperative highest HR.

Pheochromocytoma and sympathetic paraganglioma (PPGL) are rare neuroendocrine tumors arising from catecholamine-producing chromaffin cells in the adrenal medulla and extraadrenal sympathetic paraganglia. The overall prevalence of PPGL was 2.13 per 100,000 persons, and the overall age-standardized incidence rate was 0.18 per 100,000 person years¹. Excessive catecholamine release from PPGLs can induce life-threatening cardiovascular manifestations, including hypertensive crisis, myocardial infarction, brady- and tachyarrhythmias, Takotsubo cardiomyopathy, and acute heart failure^{2,3}.

Surgery can be used as a curative treatment for these tumors to prevent cardiovascular and other organ system complications associated with catecholamine excess⁴. However, surgery is considered challenging due to the life-threatening risk of perioperative hemodynamic instability (HI). Perioperative HI, characterized by arrhythmias, abrupt increases in blood pressure (BP) during manipulation of the tumor, and decreases in BP after ligation of the adrenal vein can be major challenges for perioperative management of PPGL patients^{5,6}.

To avoid HI, routine use of a blockade and volume expansion has been incorporated into the perioperative optimization of the patient's condition before surgery^{4,7}. Despite proper preoperative preparation for PPGL, HI

¹Department of Internal Medicine, Seoul National University College of Medicine, Seoul National University Hospital, Seoul, Republic of Korea. ²Department of Anesthesiology and Pain Medicine, Seoul National University College of Medicine, Seoul National University Hospital, Seoul, Republic of Korea. ³Department of Surgery, Seoul National University College of Medicine, Seoul National University Hospital, Seoul, Republic of Korea. ⁴Cancer Research Institute, Seoul National University College of Medicine, Seoul, Republic of Korea. ⁵Division of Surgery, Thyroid Center, Seoul National University Cancer Hospital, Seoul, Republic of Korea. ⁶Medical Big Data Research Center, Institute of Medical and Biological Engineering, Seoul National University, Seoul, Republic of Korea. ⁷Department of Surgery, Seoul National University Bundang Hospital, Seongnam, Republic of Korea. ⁸Department of Surgery, Seoul National University Boramae Medical Center, Seoul, Republic of Korea. ⁹These authors contributed equally: Jung Hee Kim and Hyung-Chul Lee. ✉email: su.jin.kim.md@snu.ac.kr

Variables	HI (-)	HI (+)	Total	P-value
	(N = 35)	(N = 79)	(N = 114)	
Age (years)	49.1 ± 14.4	49.9 ± 16.0	49.6 ± 15.5	0.800
Female	16 (45.7%)	39 (49.4%)	55 (48.2%)	0.875
Body weight (kg)	63.9 ± 11.4	62.9 ± 11.3	63.2 ± 11.3	0.671
BMI (kg/m ²)	23.6 ± 3.3	23.7 ± 3.8	23.6 ± 3.7	0.764
Diabetes mellitus	6 (17.1%)	22 (27.8%)	28 (24.6%)	0.323
Hypertension	17 (48.6%)	43 (54.4%)	60 (52.6%)	0.708
Coronary artery disease	7 (20.0%)	9 (11.4%)	16 (14.0%)	0.249
Cerebrovascular disease	3 (8.6%)	6 (7.6%)	9 (7.9%)	1.000
History of abdominal surgery	5 (14.3%)	17 (21.5%)	22 (19.3%)	0.519
Symptomatic presentation	18 (51.4%)	43 (54.4%)	61 (53.5%)	0.926
Maximal size (cm)	4.3 ± 2.1	5.1 ± 2.8	4.8 ± 2.6	0.212
Side				
Right	7 (25.9%)	29 (44.6%)	36 (39.1%)	0.194
Left	17 (63.0%)	28 (43.1%)	45 (48.9%)	
Bilateral	3 (11.1%)	8 (12.3%)	11 (12.0%)	
Metastasis	1 (2.9%)	7 (8.9%)	8 (7.0%)	0.606
Multifocal	1 (2.9%)	5 (6.3%)	6 (5.3%)	0.326
Preoperative vital signs				
Preblockade SBP (mmHg)	134.1 ± 15.1	129.2 ± 20.7	130.7 ± 19.2	0.196
Lowest SBP (mmHg)	106.2 ± 10.9	101.9 ± 10.7	103.3 ± 10.9	0.058
Highest SBP (mmHg)	150.4 ± 23.1	144.2 ± 24.4	146.2 ± 24.1	0.118
Lowest HR (bpm)	60.0 ± 10.9	62.3 ± 12.5	61.6 ± 12.0	0.231
Highest HR (bpm)	91.4 ± 18.0	93.5 ± 19.5	92.9 ± 19.0	0.866
ASA				0.351
1	5 (14.3%)	10 (13.2%)	15 (13.5%)	
2	20 (57.1%)	52 (68.4%)	72 (64.9%)	
3	9 (25.7%)	14 (18.4%)	23 (20.7%)	
4	1 (2.9%)	0 (0.0%)	1 (0.9%)	
Surgical approach				0.896
Open*	13 (37.1%)	32 (40.5%)	45 (39.5%)	
Laparoscopic	22 (62.9%)	47 (59.5%)	69 (60.5%)	
Duration of anesthesia (min)	178 (157–224)	193 (147–252)	185 (148–247)	0.398
Duration of surgery (min)	110 (83–172)	125 (80–185)	123 (80–185)	0.341

Table 1. Clinical characteristics of study subjects according to hemodynamic instability (HI). Data are presented as the mean ± standard deviation, median (interquartile range) or n (%). HI, hemodynamic instability; SBP, systolic blood pressure; ASA, American Society of Anesthesiologists; *12 cases of open conversion were included in the open approach.

is unpredictable and poorly defined. Recent studies have analyzed the predictive factors for HI, but the results are conflicting^{8–10}. There are several potential predictive factors for HI: a large tumor size^{9,10}, the type of operation (open vs. laparoscopic)¹⁰, the type of a blockade¹⁰, symptomatic preoperative high BP⁸, and high levels of catecholamine⁸. In addition, the incidence of HI is highly variable in previous studies, such as in reports by Gajoux et al.⁸ (8.7%, 13/149) and Jiang et al.⁹ (67%, 44/134). This discrepancy is thought to be due to various definitions of HI in these studies, the lack of any prospective studies reducing the selection bias, and the small study populations.

The aim of this study was to evaluate the incidence of HI during surgery and to analyze the predictive factors for intraoperative HI and postoperative hypotension in PPGL patients.

Results

Among 114 subjects, HI occurred in 79 (69.3%) subjects. There are 10 (12.7%, 10/79) patients with SBP > 200 mmHg, and 76 (96.2%, 76/79) patients with MBP < 60 mmHg. The clinical and biochemical characteristics of the study subjects, according to HI, are compared in Tables 1 and 2. The demographic, clinical, and biochemical variables were similar in subjects with and without HI. The type of α -adrenergic receptor blocker was not different between the two groups. However, the duration of α -adrenergic receptor blocker use was longer in subjects with HI than in subjects without HI (63.8 ± 57.3 vs. 40.7 ± 33.0 days, $P = 0.008$) (Supplemental Fig. 1). American Society of Anesthesiologists (ASA) class, surgical approach, anesthesia, and duration of surgery were not different between the two groups.

Variables	HI (-)	HI (+)	Total	P-value
	(N = 35)	(N = 79)	(N = 114)	
Hormone type				0.076
Nonfunctioning	5 (14.3%)	2 (2.5%)	7 (6.1%)	
Epinephrine	19 (54.3%)	50 (63.3%)	69 (60.5%)	
Norepinephrine	11 (31.4%)	27 (34.2%)	38 (33.3%)	
Serum metanephrines (pg/mL)	2350.0 ± 3116.0	5593.5 ± 7226.2	4603.4 ± 6423.0	0.089
Serum normetanephrines (pg/mL)	8486.2 ± 7492.1	8647.4 ± 7363.5	8597.1 ± 7363.4	0.930
Urine epinephrine (µg/day)	45.0 ± 88.1	126.2 ± 376.4	102.1 ± 320.7	0.147
Urine norepinephrine (µg/day)	458.3 ± 512.8	500.4 ± 799.9	487.9 ± 724.4	0.824
Urine dopamine (µg/day)	247.3 ± 79.2	342.7 ± 310.7	314.4 ± 267.0	0.223
Urine metanephrines (µg/day)	912.4 ± 1597.2	2275.8 ± 5972.0	1870.5 ± 5108.7	0.197
Urine normetanephrines (µg/day)	3049.0 ± 3212.6	5110.5 ± 12,925.3	4497.6 ± 10,992.9	0.854
Medications				
α-blocker use	32 (91.4%)	73 (92.4%)	105 (92.1%)	1.000
α-blocker use				
Phenoxybenzamine	9 (28.1%)	16 (21.9%)	25 (23.8%)	1.000
Doxazosin	23 (65.7%)	49 (62.0%)	72 (63.2%)	
Terazosin	0 (0%)	8 (0.10%)	8 (7.0%)	
Duration of α-blocker use (days)	40.7 ± 33.0	63.8 ± 57.3	56.7 ± 52.0	0.008
α-blocker dose (DDD)	0.8 ± 0.5	0.9 ± 0.6	0.8 ± 0.6	0.576
β-blocker use	22 (62.9%)	45 (57.0%)	67 (58.8%)	0.701
Duration of β-blocker use (days)	15.6 ± 55.7	27.8 ± 80.9	67.3 ± 31.8	0.183
β-blocker dose (DDD)	0.3 ± 0.6	0.3 ± 0.8	0.3 ± 0.7	0.257
Calcium channel blocker (DDD)	0.5 ± 0.6	0.4 ± 0.6	0.4 ± 0.6	0.393
RAS inhibitor (DDD)	0.3 ± 0.5	0.5 ± 0.9	0.4 ± 0.8	0.527
Anti-HTN drug (DDD)	1.3 (0.9–2.3)	1.6 (0.8–2.7)	1.3 (0.9–2.7)	0.725

Table 2. Biochemical characteristics, and medications of study subjects according to hemodynamic instability (HI). Data are presented as the mean ± standard deviation, median (interquartile range) or n (%). HI, hemodynamic instability; SBP, systolic blood pressure; DDD, daily defined dose; HTN, hypertension.

Variables	Univariate		Multivariate	
	OR	P-value	OR	P-value
α-blocker duration	1.012 (1.001–1.023)	0.033	1.015 (1.001–1.029)	0.041
Serum metanephrines (pg/mL)	1.0004 (1.0001–1.0007)	0.036	1.000 (1.000–1.000)	0.060
Urine dopamine (µg/day)	1.004 (1.000–1.008)	0.075		
Preoperative lowest SBP	0.964 (0.928–1.002)	0.062		

Table 3. Logistic regression analysis for predicting intraoperative hemodynamic instability in PPGL patients. Variables with a *P*-value < 0.1 in univariate analysis were included in the multivariate logistic regression models. Data are shown as odds ratios (ORs) (95% confidence intervals). PPGL, pheochromocytoma and sympathetic paraganglioma.

We conducted logistic regression models of preoperative risk factors for HI (Table 3). Among the variables shown in Tables 1 and 2, the variables with a *P*-value less than 0.1, including the duration of α-adrenergic receptor blocker use, serum metanephrine level, urine dopamine level, and lowest preoperative systolic blood pressure (SBP), were included in the logistic regression models. In the multivariate analysis, the duration of α-adrenergic receptor blocker use (odds ratio [OR], 1.015; 95% confidence interval [CI], 1.001–1.029) was a predictive factor for intraoperative HI.

Table 4 demonstrates the intraoperative characteristics of the study subjects according to the HI. Subjects with HI exhibited lower mean SBP and mean blood pressure (MBP) but a longer time when SBP was higher than 200 mmHg and a longer time when MBP was less than 60 mmHg than those without HI. RBC transfusion and the administered doses of phenylephrine and norepinephrine were greater in subjects with HI than in those without HI. However, the dose of vasodilators was not different between the two groups.

Supplemental Table 1 demonstrates intraoperative HI according to phenoxybenzamine use. Phenoxybenzamine was not superior to other α-blockers, such as doxazosin or prazosin.

Postoperative hypotension (one or more episodes of MBP < 60 mmHg) occurred in 36 (31.6%) subjects. We further analyzed the clinical predictors for postoperative hypotension in PPGL subjects. Female sex, lower BMI,

	HI (-) (n = 35)	HI (+) (n = 79)	Total (N = 114)	P-value
Hemodynamic variables				
Mean SBP	119.3 ± 8.9	112.9 ± 10.6	114.8 ± 10.5	< 0.001
Maximum SBP	188 (169–206)	190 (164–211)	189 (164.5–209.5)	0.847
Minimum SBP	84 (74–90)	68 (55–76)	72 (62–80)	< 0.001
Mean MBP	86.9 ± 6.1	79.7 ± 6.1	81.9 ± 7.0	< 0.001
Maximum MBP	138 (124–148)	132 (118–160)	134 (120–151.5)	0.511
Minimum MBP	60 (57–67)	48 (43–52)	50 (46–58)	< 0.001
Mean HR	69.4 ± 11.3	70.1 ± 13.8	69.9 ± 13.0	0.770
Maximum HR	102.1 ± 16.1	104.6 ± 21	103.8 ± 19.5	0.490
Minimum HR	49.3 ± 9	50.1 ± 11.4	49.8 ± 10.7	0.706
Hemodynamic instabilities				
Time SBP > 200 (min)	0.0 ± 0.0	0.9 ± 3.2	0.6 ± 2.7	0.028
Time MBP < 60 (min)	0.0 ± 0.0	16.8 ± 16.2	11.7 ± 15.6	< 0.001
Area SBP > 200 (mmHg × min)	0.0 ± 0.0	24.3 ± 105.4	16.8 ± 88.3	0.028
Area MBP < 60 (mmHg × min)	0.0 ± 0.0	97.1 ± 121.7	67.3 ± 110.6	< 0.001
Std of SBP	19.4 ± 5.1	22.5 ± 5.5	21.5 ± 5.6	0.004
HIS	20.0 (14.0–28.5)	30.0 (22.0–44.0)	28.0 (19.2–36.8)	< 0.001
ARV	15.7 (10.7–19.4)	16.4 (12.7–20.4)	16.2 (12.1–20.0)	0.260
MDAPE	13.1 (9.0–16.5)	15.7 (12.8–20.5)	14.6 (11.5–18.9)	0.002
Fluid management				
Urine output (mL)	300 (125–350)	205 (95–458)	240 (100–400)	0.773
Estimated blood loss (mL)	100 (0–325)	100 (25–425)	100 (0–350)	0.386
Total infused fluid (mL)	1350 (725–1750)	1400 (825–2250)	1375 (800–2000)	0.163
Tidal transfused RBC (pack)	0.0 (0.0–0.0)	0.0 (0.0–400.0)	0.0 (0.0–0.0)	0.030
Vasopressors				
Use of vasopressor	18 (75.0%)	53 (84.1%)	71 (81.6%)	0.361
Number of vasopressors	1.3 ± 0.9	1.8 ± 1.3	1.7 ± 1.2	0.120
Phenylephrine (µg)	30 (0–105)	100 (15–325)	60 (0–200)	0.028
Norepinephrine (µg)	0.0 (0.0–37.4)	0.0 (0.0–0.0)	0.0 (0.0–60.3)	0.014
Epinephrine (µg)	0.0 (0.0–0.0)	0.0 (0.0–0.0)	0.0 (0.0–0.0)	0.177
Ephedrine (mg)	5.0 (0.0–11.2)	10.0 (0.0–20.0)	5.0 (0.0,15.0)	0.295
Vasopressin (U)	0.0 (0.0–0.0)	0.0 (0.0–0.0)	0.0 (0.0–0.0)	0.245
Vasodilators				
Use of vasodilator	15 (62.5%)	46 (73.0%)	61 (70.1%)	0.487
Number of vasodilators	1.1 ± 1.1	1.2 ± 1.1	1.2 ± 1.1	0.670

Table 4. Intraoperative characteristics of the study subjects. Data are presented as the mean ± standard deviation or median (interquartile range). HI, hemodynamic instability; RBC, red blood cell; SBP, systolic blood pressure; MBP, mean blood pressure; Std, standard deviation; HIS, hemodynamic instability score; ARV, average real variability; MDAPE, median absolute performance error.

symptoms, higher normetanephrine and norepinephrine levels, greater highest heart rates (HR), and higher anti-hypertensive medication dose were related to postoperative hypotension (Table 5). In the multivariate analysis, subjects with higher urine epinephrine levels (OR, 1.003; 95% CI, 1.000–1.007) and greater preoperative highest HR (OR, 1.046; 95% CI, 1.014–1.080) had a higher risk of postoperative hypotension (Table 6). There was two readmissions due to pleural effusion, and one readmission due to ileus. There was no mortality within 30 days of surgery among the subjects.

Discussion

The cornerstone treatment of PPGL is surgical resection. However, the surgery itself is accompanied by intraoperative HI and postoperative hypotension, which can lead to life-threatening complications. In the present study, intraoperative HI was present in 69.3% of subjects, and postoperative hypotension was present in 31.6% of PPGL patients after preoperative medical preparation. We demonstrated that the duration of α -adrenergic receptor blocker use was the only predictive factor for intraoperative HI and that a higher urine epinephrine level and greater preoperative highest HR were predictive factors for postoperative hypotension.

Previous researchers have focused on intraoperative HI and the related risk factors to improve the surgical outcome and reduce perioperative morbidity/mortality^{11,12}. The definition of intraoperative HI was heterogeneous according to the studies, and there was no agreement regarding the definition of HI. Intraoperative hypertension

Variables	Postoperative hypotension (-)	Postoperative hypotension (+)	Total	P-value
	(N = 78)	(N = 36)	(N = 114)	
Age (years)	50.2 ± 14.9	48.4 ± 16.8	49.6 ± 15.5	0.582
Female	32 (41.0%)	23 (63.9%)	55 (48.2%)	0.039
Body weight (kg)	65.0 (57.3–71.0)	58.6 (52.0–64.0)	62.9 (55.0–69.4)	0.004
BMI (kg/m ²)	24.0 (21.8–26.0)	22.4 (20.8–23.7)	23.1 (21.2–25.4)	0.041
Symptomatic presentation	36 (46.2%)	25 (69.4%)	61 (53.5%)	0.034
Maximal size (cm)	3.8 (2.6–5.0)	5.5 (3.5–6.7)	4.0 (3.0–5.5)	0.005
Metastasis	4 (5.1%)	4 (11.1%)	8 (7.0%)	0.259
Multifocal	2 (2.6%)	4 (11.1%)	6 (5.3%)	0.078
Hormone type				
Nonfunctioning	6 (7.7%)	1 (2.8%)	7 (6.1%)	0.492
Epinephrine	45 (57.7%)	24 (66.7%)	69 (60.5%)	
Norepinephrine	27 (34.6%)	11 (30.6%)	38 (33.3%)	
Serum metanephrines (pg/mL)	3949.1 ± 5376.3	6168.9 ± 8328.8	4603.4 ± 6423.0	0.201
Serum normetanephrines (pg/mL)	7233.4 ± 7057.0	11,930.7 ± 7147.6	8597.1 ± 7363.4	0.006
Urine epinephrine (µg/day)	51.7 ± 99.4	207.0 ± 534.3	102.0 ± 320.7	0.092
Urine norepinephrine (µg/day)	323.6 ± 456.7	830.1 ± 1014.8	487.9 ± 724.4	0.007
Urine dopamine (µg/day)	290.0 ± 219.5	365.0 ± 344.1	314.3 ± 267.1	0.238
Urine metanephrines (µg/day)	1198.8 ± 1983.5	3269.8 ± 8410.0	1870.5 ± 5108.8	0.153
Urine normetanephrines (µg/day)	4398.7 ± 12,976.8	4703.6 ± 4866.8	4497.6 ± 10,992.9	0.858
Preoperative vital signs				
Preblockade SBP (mmHg)	129.7 ± 19.8	133.7 ± 17.4	130.7 ± 19.2	0.375
Lowest SBP (mmHg)	104.6 ± 10.7	100.5 ± 10.9	103.3 ± 10.9	0.073
Highest SBP (mmHg)	146.8 ± 23.4	145.0 ± 25.7	146.2 ± 24.1	0.730
Lowest HR (bpm)	60.7 ± 11.8	63.5 ± 12.4	61.6 ± 12.0	0.273
Highest HR (bpm)	89.1 ± 15.7	100.8 ± 22.7	92.9 ± 19.0	0.008
Medications				
α-blocker use	70 (89.7%)	35 (97.2%)	105 (92.1%)	0.268
α-blocker type				
Phenoxybenzamine	20 (25.6%)	5 (13.9%)	25 (21.9%)	0.244
Others	58 (74.4%)	31 (86.1%)	89 (78.1%)	
Duration of α-blocker use	44.5 (25.2–80.0)	42.0 (19.0–65.2)	44.0 (22.2–75.5)	0.567
α-blocker dose (DDD)	0.7 (0.5–1.0)	0.9 (0.6–1.1)	0.8 (0.5–1.0)	0.108
β-blocker use	46 (59.0%)	21 (58.3%)	67 (58.8%)	0.889
Duration of β-blocker use	2.0 (0.0–19.8)	2.5 (0.0–14.2)	2.0 (0.0–19.8)	0.982
β-blocker dose (DDD)	0.1 (0.0–0.3)	0.1 (0.0–0.5)	0.1 (0.0–0.3)	0.366
Anti-HTN drug (DDD)	1.1 (0.6–2.3)	1.9 (1.2–3.4)	1.3 (0.9–2.7)	0.029

Table 5. Clinical predictors for postoperative hypotension in PPGL patients.

Variables	Univariate		Multivariate	
	OR	P-value	OR	P-value
Female sex	2.543 (1.124–5.752)	0.025		
Maximal size (cm)	1.194 (1.013–1.408)	0.035		
Symptomatic presentation	2.652 (1.148–6.125)	0.022		
Multifocal	5.516 (1.096–27.774)	0.038		
Urine epinephrine (µg/day)	1.003 (1.000–1.006)	0.049	1.003 (1.000–1.007)	0.026
Urine norepinephrine (µg/day)	1.001 (1.000–1.002)	0.004		
Preoperative highest HR (/min)	1.035 (1.011–1.061)	0.005	1.046 (1.014–1.080)	0.005

Table 6. Logistic regression analysis for predicting postoperative hypotension in PPGL patients. Variables with a *P*-value < 0.05 in univariate analysis were included in the multivariate logistic regression models, and forward stepwise selection based on the Wald test was performed. Data are shown as odds ratios (ORs) (95% confidence intervals).

may be related to intubation for general anesthesia, pneumoperitoneum during laparoscopic surgery, and PPGL manipulation. Intraoperative hypertension can lead to subsequent hypertensive crisis, consisting of various cardiac arrhythmias, left ventricular failure, myocardial ischemia, and stroke^{2,3}. On the other hand, after surgical resection of the tumor, a sudden decline in catecholamine excess and the downregulation of adrenergic receptors due to catecholamine excess result in intraoperative hypotension. Since intraoperative hypotension is also related to postoperative complications, including acute kidney injury, myocardial injury¹³, stroke¹⁴, and even death¹⁵ we combined intraoperative hypertension and hypotension under the intraoperative HI definition. Therefore, we reported that the incidence of HI was 69.3% (n = 79), and only 8.8% (n = 10) of the subjects exhibited intraoperative hypertension. In our center, HI was caused mostly by intraoperative hypotension.

Median absolute performance error (MDAPE), average real variability (ARV), and hemodynamic instability score (HIS) were used to evaluate HI in the different operation fields. MDAPE was calculated through preoperative and postoperative SBP in noncardiac surgery, ARV was calculated through BP, and the HIS was calculated through hemodynamic variables, including BP, HR, volume therapy, and cardiovascular medication in high-risk abdominal surgery. In the present study, we found that there was a significant association of HI, MDAPE, and HIS.

Although some researchers have suggested that α -adrenergic receptor blockers might be harmful, causing orthostatic hypotension and intra- or postoperative hypotension^{16–18}, preoperative use of α -adrenergic receptor blockers has been advocated by several guidelines^{2,4,7}. Among several α -adrenergic receptor blockers, phenoxybenzamine was preferred. Phenoxybenzamine is a nonselective and noncompetitive α 1- and α 2-adrenergic receptor blocker with irreversible and long-duration effects. Thus, phenoxybenzamine is known to be superior to prevent intraoperative hypertension, but prolonged hypotension after surgery and reflex tachycardia are other concerns¹⁹. Selective and competitive α 1-adrenergic receptor blockers, such as doxazosin, have a shorter duration effect and less associated reflex tachycardia¹⁹. Previous retrospective studies have yielded conflicting results^{20–23}. The recent first randomized controlled trial comparing the efficacies of phenoxybenzamine and doxazosin demonstrated that the total duration of BP during surgery outside a predefined target range was not different between the two drugs²⁴. Instead, the HIS was lower in the phenoxybenzamine group than in the doxazosin group, which was not linked with the cardiovascular complication rate²⁴. However, we failed to show a significant difference between phenoxybenzamine users and nonusers in terms of intraoperative hypertension, postoperative hypotension, and even the HIS.

The present study demonstrated that the duration of α -adrenergic receptor blocker use was the only predictive factor for intraoperative HI. Tian et al.²⁵ showed that the use of phenoxybenzamine for more than two weeks after the final dose adjustment could not further reduce the HI rate. In addition, Kong et al.²⁶ demonstrated that the use of doxazosin for more than 30 days after the final dose adjustment might increase the risk of intraoperative bradycardia and postoperative hypotension. In the present study, duration of preoperative α -adrenergic receptor blocker use was defined as period from date of beginning of α -blockade to ending of α -blockade. Data collected from the electronic medical records, long duration of α -blockade was not associated with the difficulty to obtain target BP below 130/80 mmHg in supine position. Some patients with the long duration of preoperative α -blockade had been prescribed before diagnosis of PPGLs due to benign prostate hyperplasia or voiding problem. Without uncontrolled HI with appropriate medications for PPGL, elective operation was performed. Nine patients did not use α -adrenergic receptor blockers due to low BP or side effects, and 12 patients were prescribed less than 14 days. However, we cannot figure out the significant difference of incidence of HI between patients with a duration of α -blocker use ≥ 14 days and < 14 days. Patients who used α -blocker use ≥ 14 days were present in 66/79 (83.5%) without HI, and 28/35 (80.0%) with HI ($P = 0.848$). As the other authors acknowledged (Tian et al.²⁵, Kong et al.²⁶), the reason why the prolonged use of α -blocker is harmful to HI remained to be elucidated. But presumably, the prolonged use of α -blocker may suppress the vascular tone response to a sudden decrease of catecholamine after resection of PPGLs. However, to date, there was no clear evidence to support this. In our cohort, most patients (79 patients, 96.2%) with HI exhibited a MBP < 60 mmHg. Therefore, we suggest that prolonged use of alpha blocker was a predictor of only MBP < 60 mmHg. However, the underlying mechanism required further research.

Based on the results of the present study, we recognized the long-duration was not beneficial to prevent intraoperative HI, and needed further study to obtain the optimal duration of α -blockade for PPGL patients.

Contrary to previous data^{9,10}, a larger tumor size was not a risk factor for HI in our study since the tumor size was relatively small compared with that in the other studies. In the present study, incidentally detected PPGLs during chest or abdominal computed tomography (CT) for the purpose of routine health check-ups were included.

There is controversy in previous studies regarding the effect of the surgical approach on HI. Kiernan et al.¹⁰ reported that subjects undergoing open surgery had a higher risk of HI than those undergoing laparoscopic surgery, but other studies reported that there was no increased risk of HI in open surgery groups^{11,20}. Although open surgery was preferred in subjects with large PPGLs and the possibility of malignancy in our center, the surgical approach (open vs. laparoscopic) had no influence on HI.

The genetic testing results were available only in 63 of 114 patients, and only 10 of 63 patients harbored the genetic mutation: RET (n = 1), VHL (n = 4), SDHB (n = 1), SDHD (n = 3), and MAX (n = 1). Therefore, we could not analyze the HI between cluster 1 and cluster 2 tumors.

In the present study, postoperative hypotension occurred in 31.6% of subjects. Hypotension develops intraoperatively after tumor removal and may last for 24–48 h postoperatively, which requires fluid therapy and vasopressors. Premedication with phenoxybenzamine is known to be accompanied by a higher risk of postoperative hypotension compared to that with doxazosin premedication²². However, in our study, phenoxybenzamine was not a determinant for postoperative hypotension, in agreement with the PRESCRIPT study²⁴. The high urine epinephrine level and elevated preoperative highest HR may reflect the extent of catecholamine secretion.

Elevation in epinephrine levels downregulates β -adrenergic receptors in the heart, which depresses myocardial contractility²⁷. After tumor removal during surgery, depressed myocardial function does not compensate for peripheral vasodilation, which might lead to intraoperative and postoperative hypotension^{27,28}.

The major strength of the present study is detailed clinical and biochemical information, including the intraoperative and postoperative hemodynamic parameters, fluid management, and medication for PPGL patients undergoing surgical treatment. Intraoperative hemodynamic parameters were retrieved from the anesthetic records, which contain BP measurements at a 1-min resolution to evaluate HI. To evaluate postoperative hypotension, postoperative hemodynamic parameters were collected from electronic medical records obtained at the surgical intensive care unit for 24 h after surgery. We calculated the DDD of all antihypertensive medications. Thus, we could analyze the effect of the dosage of other antihypertensive medications as well as the use of α -adrenergic receptor blockers on HI.

This study has several limitations. First, this study was retrospective in a single institution and limited by selective and recall bias. Second, the sample size was relatively small, even though our institution is a large tertiary referral hospital. PPGLs are rare endocrine tumors with an overall prevalence of 2.13 per 100,000 in the Republic of Korea.

In conclusion, the present study revealed that intraoperative HI and postoperative hypotension were present in 69.3% and 31.6% of subjects who underwent surgery for PPGLs, respectively. Long-term use of α -adrenergic receptor blocker before surgery was significantly associated with intraoperative HI, and a higher urine epinephrine level and greater highest HR were predictive factors for postoperative hypotension. Therefore, we believe that the present study would be helpful to identify at-risk patients for intraoperative HI and postoperative hypotension and to guide delicate and intensive management in PPGL intraoperatively and postoperatively to improve clinical outcomes.

Methods

Study subjects. We recruited a total of 114 consecutive patients with PPGLs who underwent surgery at Seoul National University Hospital (Seoul, Korea) from March 2012 to April 2019. Patients who met the following criteria were excluded: (1) surgery due to carotid body tumor and (2) age less than 18 years. The present study was approved by the Institutional Review Board of the Seoul National University Hospital (No. H-1801-010-911), and performed in accordance with the Declaration of Helsinki. Informed consent was waived because of the retrospective design in accordance with requirement of the Institutional Review Board of the Seoul National University Hospital.

Preoperative variables. All clinical data were obtained from the medical records. Clinical information included i) age at initial diagnosis, sex, and body mass index; ii) comorbidities, such as diabetes mellitus, hypertension, coronary artery diseases, and cerebrovascular diseases, and history of abdominal surgery; iii) clinical symptoms/signs, including headache, sweating, palpitation, pain (neck, chest, abdomen, or bone), palpable mass, and hypertension (new onset, paroxysmal, or uncontrolled); iv) tumor characteristics, such as tumor size, side, metastasis, location, and serum/24-h urine catecholamine production, including epinephrine, norepinephrine, metanephrine, normetanephrine, and dopamine; v) preoperative hemodynamic parameters, including BP and HR before and after use of α -adrenergic receptor blockers; vi) preoperative medications, including α - and β -adrenergic receptor blockers and other antihypertensive drugs (antihypertensive medication regimens were calculated as the daily defined dose, DDD), which is the assumed average maintenance dose per day for a drug used for its main indication in adults, according to World Health Organization ATC/DDD Index 2019 (<https://www.whooc.no/ddd>); and vii) American Society of Anesthesiologists physical status classification (ASA class).

PPGLs were diagnosed based on catecholamine excess before surgery and/or pathology after surgery. Catecholamine secretion was evaluated with 24-h urine catecholamine/fractionated metanephrines or serum fractionated metanephrines. The catecholamine type was classified as epinephrine, norepinephrine, or nonfunctioning type²⁹. If urinary or serum epinephrine levels were elevated with or without high norepinephrine levels, the type was designated “epinephrine type”. If urinary or serum norepinephrine levels were elevated without high epinephrine levels regardless of dopamine levels, the type was classified as “norepinephrine type”. If both urinary or serum epinephrine and norepinephrine levels were within the normal range, the type was classified as “nonfunctioning type”. Serum fractionated metanephrines and urinary catecholamines were measured by liquid chromatography-tandem mass spectrometry (LC-MS/MS), and urinary fractionated metanephrines were assessed by high-performance liquid chromatography-electrochemical detection (HPLC-ECD).

Anatomical imaging, such as CT or magnetic resonance imaging (MRI), was used to locate PPGLs. Functional imaging, such as MIBG and positron emission tomography/computed tomography (PET/CT) with ⁶⁸Ga-labeled DOTA-Tyr3 octreotide (DOTATOC), was performed to evaluate metastatic lesions. Metastasis was defined as the presence of pheochromocytomas in nonchromaffin organs, such as the lung, kidney, liver, bone, mediastinum, and lymph node, at diagnosis.

The ASA class was recorded from the patients’ electronic medical records.

Preoperative medications. α -Adrenergic receptor blockers were administered to reach BP < 130/80 mmHg in the supine position and SBP > 90 mmHg in the upright position⁴. If several α -adrenergic receptor blockers were used, the final type and dose were included in the analysis. β -adrenergic receptor blockers were added in subjects with a HR at upright position > 100 bpm⁴. A high-salt diet and fluid intake were recommended. At our institution, we routinely performed preoperative infusion of saline (1–2 L) for several days (5–6 days) in all subjects except in cases with contraindication for volume expansion, to restore the catecholamine-induced volume depletion and prevent severe hypotension after tumor removal.

Surgical techniques. Operations were performed under general anesthesia, and an arterial line was inserted into the patient's radial artery after anesthesia induction for continuous arterial pressure monitoring. The surgical approach (open vs. laparoscopic) was determined based on the possibility of malignancy, tumor size, relationship to adjacent organs, and previous history of abdominal surgery. For most benign tumors < 7 cm on preoperative CT imaging, laparoscopic adrenalectomy was preferred.

Definition of hemodynamic instability (HI). HI was defined as one or more episodes of SBP > 200 mmHg or MBP < 60 mmHg during the surgery.

Intraoperative variables. The hemodynamic variables, including BP and HR, were collected from the electronic anesthetic chart, which included vital signs with a 1-min interval during the operation. Fluid management, use of vasopressors and vasodilators, duration of surgery, and duration of anesthesia were also collected from electronic medical records.

We also analyzed HI using the HIS, ARV, and MDAPE. The HIS was calculated using the method reported by Buitenwerf et al.³⁰. The ARV and MDAPE were calculated using the following formulas^{31–33}.

$$ARV = \frac{1}{N-1} \sum_{k=1}^{N-1} |SBP_{k+1} - SBP_k|$$

$$MDAPE = \text{median} \left(\left| \frac{SBP_k - SBP_{\text{ward}}}{SBP_{\text{ward}}} \right| \right) \times 100\%,$$

where SBP_{ward} represents the median ward BP measured one day before operation.

Postoperative variables. After surgery, PPGL patients were routinely transferred to the surgical intensive care unit for close monitoring of hemodynamic status and moved to the ward on postoperative day 1. Postoperative clinical information included postoperative hemodynamic variables, fluid management, use of vasopressors and vasodilators, and postoperative complications, including postoperative mortality within 30 days.

Postoperative hemodynamic variables were collected from the electronic vital sign sheet in the intensive care unit for 24 h after surgery. Postoperative hypotension was defined as one or more episodes of MBP < 60 mmHg.

Statistical analysis. All data are expressed as the mean ± SD, median (interquartile range) or absolute numbers (percentage) and were analyzed with SPSS software (version 25.0; IBM Corp., Armonk, NY, USA). In all analyses, $P < 0.05$ was considered to indicate statistical significance.

Continuous variables were compared with Student's t-test or the Mann–Whitney U-test based on the results of the Shapiro–Wilk test. Categorical variables were compared using the Pearson chi-square test. Univariate logistic regression analysis was performed to identify predictive variables for intraoperative HI and postoperative hypotension. Multivariate logistic regression was performed using forward stepwise selection based on the Wald test. Cox proportional hazard ratio modeling for intraoperative HI was performed using the Python Lifelines library. Receiver operating characteristic (ROC) curve analysis was used to assess the prediction variables for intraoperative HI and postoperative hypotension.

Received: 6 October 2020; Accepted: 2 September 2021

Published online: 17 September 2021

References

- Kim, J. H., Moon, H., Noh, J., Lee, J. & Kim, S. G. Epidemiology and prognosis of pheochromocytoma/paraganglioma in Korea: A nationwide study based on the national health insurance service. *Endocrinol. Metab. (Seoul)* **35**, 157–164. <https://doi.org/10.3803/EnM.2020.35.1.157> (2020).
- Lenders, J. W. M. *et al.* Genetics, diagnosis, management and future directions of research of phaeochromocytoma and paraganglioma: A position statement and consensus of the working group on endocrine hypertension of the European society of hypertension. *J. Hypertens* **38**, 1443–1456. <https://doi.org/10.1097/HJH.0000000000002438> (2020).
- Prejbisz, A., Lenders, J. W., Eisenhofer, G. & Januszewicz, A. Cardiovascular manifestations of phaeochromocytoma. *J. Hypertens* **29**, 2049–2060. <https://doi.org/10.1097/HJH.0b013e32834a4ce9> (2011).
- Lenders, J. W. *et al.* Pheochromocytoma and paraganglioma: An endocrine society clinical practice guideline. *J. Clin. Endocrinol. Metab.* **99**, 1915–1942. <https://doi.org/10.1210/jc.2014-1498> (2014).
- Fernandez-Cruz, L., Taura, P., Saenz, A., Benarroch, G. & Sabater, L. Laparoscopic approach to pheochromocytoma: Hemodynamic changes and catecholamine secretion. *World J. Surg.* **20**, 762–768; discussion 768, doi:<https://doi.org/10.1007/s002689900116> (1996).
- Flavio Rocha, M. *et al.* Laparoscopic surgery for pheochromocytoma. *Eur. Urol.* **45**, 226–232. <https://doi.org/10.1016/j.eururo.2003.09.016> (2004).
- Pacak, K. Preoperative management of the pheochromocytoma patient. *J. Clin. Endocrinol. Metab.* **92**, 4069–4079. <https://doi.org/10.1210/jc.2007-1720> (2007).
- Gaujoux, S. *et al.* Preoperative risk factors of hemodynamic instability during laparoscopic adrenalectomy for pheochromocytoma. *Surg. Endosc.* **30**, 2984–2993. <https://doi.org/10.1007/s00464-015-4587-x> (2016).
- Jiang, M. *et al.* Preoperative risk factors for haemodynamic instability during pheochromocytoma surgery in Chinese patients. *Clin. Endocrinol.* **88**, 498–505. <https://doi.org/10.1111/cen.13544> (2018).
- Kiernan, C. M. *et al.* Predictors of hemodynamic instability during surgery for pheochromocytoma. *Ann. Surg. Oncol.* **21**, 3865–3871. <https://doi.org/10.1245/s10434-014-3847-7> (2014).

11. Ma, L., Shen, L., Zhang, X. & Huang, Y. Predictors of hemodynamic instability in patients with pheochromocytoma and paraganglioma. *J Surg Oncol.* <https://doi.org/10.1002/jso.26079> (2020).
12. Li, N. *et al.* Intraoperative hypotension is associated with increased postoperative complications in patients undergoing surgery for pheochromocytoma-paraganglioma: A retrospective cohort study. *BMC Anesthesiol.* **20**, 147. <https://doi.org/10.1186/s12871-020-01066-y> (2020).
13. Salmasi, V. *et al.* Relationship between Intraoperative hypotension, defined by either reduction from baseline or absolute thresholds, and acute kidney and myocardial injury after noncardiac surgery: A retrospective cohort analysis. *Anesthesiology* **126**, 47–65. <https://doi.org/10.1097/ALN.0000000000001432> (2017).
14. Sun, L. Y. *et al.* Defining an intraoperative hypotension threshold in association with stroke in cardiac surgery. *Anesthesiology* **129**, 440–447. <https://doi.org/10.1097/ALN.0000000000002298> (2018).
15. Monk, T. G. *et al.* Association between intraoperative Hypotension and hypertension and 30-day Postoperative mortality in noncardiac Surgery. *Anesthesiology* **123**, 307–319. <https://doi.org/10.1097/ALN.0000000000000756> (2015).
16. Groeben, H. *et al.* Perioperative alpha-receptor blockade in pheochromocytoma surgery: An observational case series. *Br. J. Anaesth.* **118**, 182–189. <https://doi.org/10.1093/bja/aew392> (2017).
17. Groeben, H. *et al.* International multicentre review of perioperative management and outcome for catecholamine-producing tumours. *Br. J. Surg.* **107**, e170–e178. <https://doi.org/10.1002/bjs.11378> (2020).
18. Buisset, C. *et al.* Pheochromocytoma surgery without systematic preoperative pharmacological preparation: Insights from a referral tertiary center experience. *Surg. Endosc.* <https://doi.org/10.1007/s00464-020-07439-1> (2020).
19. Naranjo, J., Dodd, S. & Martin, Y. N. Perioperative management of pheochromocytoma. *J. Cardiothorac. Vasc. Anesth.* **31**, 1427–1439. <https://doi.org/10.1053/j.jvca.2017.02.023> (2017).
20. Bruynzeel, H. *et al.* Risk factors for hemodynamic instability during surgery for pheochromocytoma. *J. Clin. Endocrinol. Metab.* **95**, 678–685. <https://doi.org/10.1210/jc.2009-1051> (2010).
21. Weingarten, T. N. *et al.* Comparison of two preoperative medical management strategies for laparoscopic resection of pheochromocytoma. *Urology* **76**(508), e506–511. <https://doi.org/10.1016/j.urology.2010.03.032> (2010).
22. Prys-Roberts, C. & Farndon, J. R. Efficacy and safety of doxazosin for perioperative management of patients with pheochromocytoma. *World J. Surg.* **26**, 1037–1042. <https://doi.org/10.1007/s00268-002-6667-z> (2002).
23. Kocak, S., Aydintug, S. & Canakci, N. Alpha blockade in preoperative preparation of patients with pheochromocytomas. *Int. Surg.* **87**, 191–194 (2002).
24. Buitenwerf, E. *et al.* Efficacy of alpha-Blockers on hemodynamic control during pheochromocytoma resection: A randomized controlled trial. *J. Clin. Endocrinol. Metab.* **105**, 2381–2391. <https://doi.org/10.1210/clinem/dgz188> (2020).
25. Tian, J. *et al.* The duration of preoperative administration of single alpha-receptor blocker phenoxybenzamine before adrenalectomy for pheochromocytoma: 18 years of clinical experience from nationwide high-volume center. *Biomed. Res. Int.* **2019**, 2613137. <https://doi.org/10.1155/2019/2613137> (2019).
26. Kong, H. *et al.* The use of doxazosin before adrenalectomy for pheochromocytoma: Is the duration related to intraoperative hemodynamics and postoperative complications?. *Int. Urol. Nephrol.* <https://doi.org/10.1007/s11255-020-02539-2> (2020).
27. Olson, S. W., Deal, L. E. & Piesman, M. Epinephrine-secreting pheochromocytoma presenting with cardiogenic shock and profound hypocalcemia. *Ann. Intern. Med.* **140**, 849–851. <https://doi.org/10.7326/0003-4819-140-10-200405180-00033> (2004).
28. Namekawa, T. *et al.* Clinical predictors of prolonged postresection hypotension after laparoscopic adrenalectomy for pheochromocytoma. *Surgery* **159**, 763–770. <https://doi.org/10.1016/j.surg.2015.09.016> (2016).
29. Kimura, N. *et al.* Pathological grading for predicting metastasis in pheochromocytoma and paraganglioma. *Endocr. Relat. Cancer* **21**, 405–414. <https://doi.org/10.1530/ERC-13-0494> (2014).
30. Buitenwerf, E. *et al.* The haemodynamic instability score: Development and internal validation of a new rating method of intraoperative haemodynamic instability. *Eur. J. Anaesthesiol.* **36**, 290–296. <https://doi.org/10.1097/EJA.0000000000000941> (2019).
31. Mena, L. *et al.* A reliable index for the prognostic significance of blood pressure variability. *J. Hypertens* **23**, 505–511. <https://doi.org/10.1097/01.hjh.0000160205.81652.5a> (2005).
32. Lee, H. C., Ryu, H. G. & Jung, C. W. Performance measurement of intraoperative systolic arterial pressure to predict in-hospital mortality in adult liver transplantation. *Sci. Rep.* **7**, 7030. <https://doi.org/10.1038/s41598-017-07664-0> (2017).
33. Mascha, E. J., Yang, D., Weiss, S. & Sessler, D. I. Intraoperative mean arterial pressure variability and 30-day mortality in patients having noncardiac surgery. *Anesthesiology* **123**, 79–91. <https://doi.org/10.1097/ALN.0000000000000686> (2015).

Acknowledgements

None.

Author contributions

JH. K: Conceptualization; Visualization; Writing—original draft; HC. L: Formal analysis; Writing—original draft; S. K: Conceptualization; Writing—original draft; Supervision; SB. Y: Formal analysis; Investigation; SH. K: Investigation, Writing—review & editing; HW Y: Investigation, Writing—review & editing; YJ. C: Investigation, Writing—review & editing; JY. C: Investigation, Writing—review & editing; KE. L: Investigation, Writing—review & editing; KW. L: Investigation, Writing—review & editing; SK. M: Investigation, Writing—review & editing; CS. S: Investigation, Writing—review & editing; KJ. P: Investigation, Writing—review & editing.

Funding

This research was supported by the MSIT (Ministry of Science and ICT), Republic of Korea, under the ITRC (Information Technology Research Center) support program (IITP-2021–2018-0–01833) supervised by the IITP (Institute for Information & Communications Technology Planning & Evaluation).

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-021-97964-3>.

Correspondence and requests for materials should be addressed to S.K.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021