

RESEARCH ARTICLE

Mocetinostat combined with gemcitabine for the treatment of leiomyosarcoma: Preclinical correlates

Gonzalo Lopez¹, Danielle Braggio¹, Abeba Zewdu¹, Lucia Casadei¹, Kara Batte¹, Hemant Kumar Bid², David Koller¹, Peter Yu¹, Obiajulu Hans Iwenofu³, Anne Strohecker¹, Edwin Choy⁴, Dina Lev⁵, Raphael Pollock^{6*}

1 Comprehensive Cancer Center, The Ohio State University, Columbus, OH, United States of America, **2** Life Science Institute, University of Michigan, Ann Arbor, MI, United States of America, **3** Department of Pathology, The Ohio State University, Columbus, OH, United States of America, **4** Division of Hematology Oncology, Massachusetts General Hospital, Boston, MA, United States of America, **5** Surgery B, Sheba Medical Center, Tel Aviv, Israel, **6** Department of Surgery, The Ohio State University, Columbus, OH, United States of America

* raphael.pollock@osumc.edu



OPEN ACCESS

Citation: Lopez G, Braggio D, Zewdu A, Casadei L, Batte K, Bid HK, et al. (2017) Mocetinostat combined with gemcitabine for the treatment of leiomyosarcoma: Preclinical correlates. PLoS ONE 12(11): e0188859. <https://doi.org/10.1371/journal.pone.0188859>

Editor: Irina V. Lebedeva, Columbia University, UNITED STATES

Received: August 10, 2017

Accepted: November 14, 2017

Published: November 29, 2017

Copyright: © 2017 Lopez et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was supported by NIH SARC SP0RE grant (U54CA168512), funding recipient RP; and National Leiomyosarcoma Foundation grant (GRT00041181/60052311), funding recipient GL. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Abstract

Leiomyosarcoma (LMS) is a malignant soft tissue sarcoma (STS) with a dismal prognosis following metastatic disease. Chemotherapeutic intervention has demonstrated to have modest clinical efficacy with no curative potential in LMS patients. Previously, we demonstrated pan-HDAC inhibition to have a superior effect in various complex karyotypic sarcomas. In this study, our goal is to evaluate the therapeutic efficacy of mocetinostat alone and in combination with gemcitabine in LMS. Human leiomyosarcoma (LMS) cell lines were used for *in vitro* and *in vivo* studies. Compounds tested included the class I HDAC inhibitor, mocetinostat, and nucleoside analog, gemcitabine. MTS and clonogenic assays were used to evaluate the effect of mocetinostat on LMS cell growth. Cleaved caspase 3/7 analysis was used to determine the effects of mocetinostat on apoptosis. Compusyn software was used to determine *in vitro* synergy studies for the combination of mocetinostat plus gemcitabine. A LMS xenograft model in SCID mice was used to test the impact of mocetinostat alone, gemcitabine alone and the combination of mocetinostat plus gemcitabine. Mocetinostat abrogated LMS cell growth and clonogenic potential, and enhanced apoptosis in LMS cell lines. The combination of mocetinostat plus gemcitabine exhibited a synergistic effect in LMS cells *in vitro*. Similarly, mocetinostat combined with gemcitabine resulted in superior anti-LMS effects *in vivo*. Mocetinostat reduced the expression of gemcitabine-resistance markers RRM1, RRM2, and increased the expression of gemcitabine-sensitivity marker, hENT1, in LMS cells. LMS are aggressive, metastatic tumors with poor prognosis where effective therapeutic interventions are wanting. Our studies demonstrate the potential utility of mocetinostat combined with gemcitabine for the treatment of LMS.

Competing interests: The authors have declared that no competing interests exist.

Introduction

Leiomyosarcoma (LMS) is a malignant tumor of mesenchymal origin, and is one of the most common soft tissue sarcoma subtypes. LMS can present in various locations including the uterus, skin, blood vessels, retroperitoneum, gastrointestinal tract, trunk, and extremities [1]. The pathogenesis of LMS remains largely unknown, with complex karyotype and both intra- and intertumor heterogeneity [2]. Advanced-stage LMS is usually incurable with current systemic therapies, suggesting the need for novel antitumor strategies [1].

A common LMS systemic therapeutic option includes gemcitabine (2',2'-difluorodeoxycytidine) combined with docetaxel [3]. Gemcitabine is a fluorinated analogue of the nucleoside deoxycytidine. Intracellular phosphorylation of gemcitabine leads to an active diphosphate form which inhibits ribonucleotide reductase and an active triphosphate form which is incorporated into DNA [4]. Administration of gemcitabine alone was reported to have a response rate of 21% in second-line therapy in uterine LMS based on Gynecological Oncology Group (GOG) response criteria [5]. In a phase II study, gemcitabine combined with docetaxel showed a response rate of 25% by RECIST criteria [6]. Even in the first-line setting, a phase II trial of locally advanced/metastatic LMS demonstrated that gemcitabine combined with docetaxel was active [7]. The combination of gemcitabine and docetaxel are synergistic [8], raising the possibility that gemcitabine be synergistic with other anticancer therapeutics as well.

One antitumor strategy of growing interest are histone deacetylases (HDAC) inhibitors [9]. HDACs play a crucial role in chromatin remodeling and gene regulation; histone acetylation of lysines in histone tails is associated with a condensed chromatin state and gene silencing [10]. Inhibition of this process leads to a more relaxed chromatin state and active gene transcription [11]. There are 11 isoforms of HDACs which are grouped into 4 different classes, and many inhibitors have been developed which can inhibit various HDAC isoforms [12]. Mocetinostat is a class I and IV selective HDAC inhibitor which has shown potent and selective antiproliferative effects in various human cancer cells preclinically [13]. Clinical trials show that mocetinostat is well-tolerated, with favorable pharmacokinetics and pharmacodynamics and promising antitumor activity in several hematological diseases [14]. Combining mocetinostat with other antitumor agents for the treatment of solid tumors is an additional promising approach [15].

In this study we examined a combination of mocetinostat and gemcitabine in the treatment of LMS. In addition to elucidating the *in vitro* anticancer effects, we used an established LMS xenograft model to study the combination effects of gemcitabine and mocetinostat *in vivo*. Our results demonstrate synergistic anti-LMS effects of combined gemcitabine and Mocetinostat, an approach that warrants further study.

Materials and methods

Cell lines and reagents

The human LMS cell lines SKLMS1 (ATCC) and LMS1 (Dominique Broccoli, Mercer University, Savannah, GA) was used in this study. LMS cell strains Leio-012 and Leio-196A were developed at the University of Texas MD Anderson Cancer Center/MDACC (Houston, TX) under IRB approval from MDACC and with patient written informed consent. LMS-117 was developed in the Sarcoma Research Lab (The Ohio State University, Columbus, OH) under IRB approval from The Ohio State University and with patient written informed consent. LMS1 was developed in Dr. Dominique Broccoli's lab (Mercer University, Savannah, GA) and acquired via collaboration; no identifying patient information was provided when acquiring LMS1. To establish LMS cell strains, patient tumor tissue was acquired at surgery and tumor

tissue was digested using 3% collagenase Type I, 0.02% DNase I Type II, and 1.5 mg/mL hyaluronidase. The digested tumor was strained and centrifuged at 1500 RPM for 5 minutes at room temperature, the cell pellet was washed once with sterile PBS and centrifuged again. The cell pellet was resuspended and maintained in DMEM 1X media supplemented with 10% FBS and Primocin. Short Tandem Repeat (STR) analysis was used to confirm the cell origin to the corresponding patient tumor (S1 Table). Control cell lines, human aortic smooth muscle cells (HASMC) and human colonic smooth muscle cells (HCSMC) were used and maintained per supplier protocol (ScienCell). Antibodies used for Western blot analysis: acetylated histone 3 (EMD Millipore), acetylated histone 4 (EMD Millipore), acetylated tubulin (Sigma), RRM1 (Abcam), RRM2 (Abcam), hENT1 (Santa Cruz Biotechnology), β -actin (Santa Cruz Biotechnology), GAPDH (Santa Cruz Biotechnology).

Cell growth assays

Cell growth conducted via MTS analysis (CellTiter96 Aqueous Non-Radioactive Cell Proliferation Assay kit, Promega) per manufacturer protocol. Cells were plated at 5,000 cells per well and allowed to adhere overnight. Growth rates were assessed 96 hr after treatment with DMSO (control), or 0.1, 0.5, 1 μ M of mocetinostat and absorbances measured at 490 nm wavelength. Clonogenic analysis was conducted by plating 400 cells per well. On the following day, cells were treated with DMSO or mocetinostat for 24 hr. Cells were grown for 10 days. On day 10, colonies were stained with 0.5% crystal violet solution for 30 min at RT. Crystal violet solution was removed, wells washed 5x with deionized H₂O, stained colonies were scanned and counted. Drug synergy studies were conducted using MTS and analysis was conducted using Compusyn software [16].

Apoptosis assays

Apoptosis was determined by quantifying cleaved caspase 3 and cleaved caspase 7 activity using Cell Event Caspase 3/7 Green Detection Reagent (Life Technologies) analyzed in the IncuCyte Zoom (Essen BioScience). Cells were plated at 2000 cells per well and treated with the Caspase 3/7 reagent together with DMSO or varying concentrations of mocetinostat for 96 hr. Endpoint analysis was performed in each well using Vybrant DyeCycle Green Stain (Life Technologies) in complete DMEM media.

In vivo experiment

Six week old female SCID mice ($n = 40$) weighing approximately (AVG \pm SD) 20 \pm 0.9 grams (Taconic Biosciences, Hudson, NY) were injected s.c. with SKLMS1 (1×10^6) into the right flank. Once tumors reached approximately 0.5 cm, mice were randomized into four treatment arms ($n = 9$ per treatment arm) and treatment was initiated: Vehicle (PEG400/0.2 N HCl), mocetinostat (50 mg/kg PO QD) (Mirati Therapeutics, Inc.), gemcitabine (20 mg/kg, i.p. BID) (Selleck Chemicals), mocetinostat combined with gemcitabine. Mice in the 'mocetinostat combined with gemcitabine' treatment arm were given mocetinostat 24 hr prior to combining with gemcitabine. Four mice were excluded from experimentation due to low tumor take. Drug doses were used per manufacture recommendation. Mice were monitored for well being, weighed, and tumors were measured twice weekly. Animals appeared to be in good health with no significant weight loss or death observed during the course of the experiment. Ulceration occurred in some mice at the tumor site nearing the 1.5 cm endpoint; mice were provided with buprenorphine (0.05–0.1 mg/kg) as analgesic. Mice were humanely euthanized (euthanized by CO₂ followed by cervical dislocation to ensure death according to IACUC guidelines) once tumors in control mice grew to approximately 1.5 cm. Final tumor volumes

and weights were measured. All mice were maintained under barrier conditions at a temperature of $72^{\circ}\text{F}\pm 4^{\circ}\text{F}$ and 12:12 hr light:dark cycle. Mice ($n = 5/\text{cage}$) were housed in $194\times 178\times 397$ mm cages (NexGen caging, Allentown Inc, Allentown, NJ) given feed (Harlan Teklad Irradiated diet 7912, Envigo, Huntingdon, UK) and water *ad libitum*. Environmental enrichment included bedding (1/4" corn cob) and autoclaved nesting material. All procedures were conducted under The Ohio State University's Institutional Animal Care and Use Committee (IACUC) approval (number 2014A00000085) and in accordance with The Ohio State University's Animal Welfare Assurance (number A3261-01). All sections of this report adhere to the ARRIVE Guidelines for reporting animal research [17]. A completed ARRIVE guidelines checklist is included in [S1 Checklist](#).

Statistical analysis

Statistical analysis was performed using GraphPad Prism6. Mean \pm SEM was calculated for all *in vitro* and *in vivo* assays using ANOVA. Significance was set at * ($p\leq 0.05$), ** ($p\leq 0.01$), *** ($p\leq 0.001$), **** ($p\leq 0.0001$). Median values and confidence interval (95%) for all data are located in the [S2 Table](#).

Results

Mocetinostat abrogates cell growth and induces apoptosis in LMS cells

The class I HDAC inhibitor, mocetinostat was used in all experiments. We previously demonstrated the efficacy of the pan-HDAC inhibitor abexinostat/PCI-24781 on a subset of STS cell lines, including the leiomyosarcoma cell line SKLMS1 [18, 19]. Pan-HDAC inhibitors, per abexinostat, demonstrate a high affinity to class I HDAC isoforms; e.g., HDAC1 and HDAC2 [20]. Pan-HDAC inhibitors yield numerous undesirable side effects, whereas the inhibition of various HDAC isoforms results in improved therapeutic windows and less toxicity [21], thus providing an impetus to target a smaller set, i.e., class I HDACs in LMS. Mocetinostat increased LMS cell line/strain histone 3 and 4 acetylation in a dose- and time-dependent manner (Fig 1A). Tubulin acetylation, a HDAC6 substrate, remained unacetylated upon treatment with mocetinostat, demonstrating the compound's affinity to class I vs class II HDAC substrates. The effect of mocetinostat on LMS cell growth was tested. Mocetinostat abrogated cell growth in a time- and dose-dependent manner (Fig 1B and S3 Table). LMS1 and Leio-196A demonstrated sensitivity to mocetinostat followed by LMS-117; SKLMS1 and Leio-012 exhibited the highest tolerance among the LMS cells. Mocetinostat had a modest impact on normal cells (HASMC and HCSMC). Mocetinostat significantly reduced LMS clonogenic potential in LMS1 and SKLMS1 cells (Fig 1C). Again, LMS1 exhibited sensitivity whereas SKLMS1 was more tolerant to mocetinostat anti-cancer effects.

To further assess the impact of mocetinostat on LMS cell proliferation, we evaluated the compound's effect on apoptosis. Cleaved caspase 3/7 activity in LMS cells treated with mocetinostat was measured using the IncuCyte Zoom system. A similar trend in drug sensitivity was observed in LMS cells in response to caspase 3/7 activity (Fig 1D); LMS1 displaying a significant increase in cleaved caspase 3/7 and SKLMS1 displaying a modest cleaved caspase 3/7 increase in response to mocetinostat.

Mocetinostat combined with gemcitabine exhibits a synergistic anti-LMS effect in vitro

Previously, we identified superior anti-STs effects when combining pan-HDAC inhibition (abexinostat) with doxorubicin or cisplatin [18] *in vitro* and *in vivo*. In this study we evaluated




Fig 1. Mocetinostat inhibits LMS cell growth and induces apoptosis. A, Mocetinostat increased acetylated histone 3 and 4 in a time- and dose-dependent manner in LMS cells. Mocetinostat did not increase acetylated tubulin expression. B, Mocetinostat-induced growth inhibition was determined using MTS assays. C, Colony formation assays recapitulate the sensitivity and tolerant dichotomy between LMS1 and SKLMS1 to mocetinostat treatment. D, Mocetinostat induced a significant increase in cleaved caspase 3/7 in LMS1 cells and a modest increase in SKLMS1 cells.

<https://doi.org/10.1371/journal.pone.0188859.g001>

the efficacy of mocetinostat combined with gemcitabine. SKLMS1 and LMS1 were used as representative cell lines that were tolerant/sensitive in response to mocetinostat. Cells were pretreated with mocetinostat 24 hr prior to combining with gemcitabine. The combination of mocetinostat and gemcitabine in this dosage sequence exhibited a synergistic effect in both cell lines (Fig 2A and 2B). No synergy was identified when the cells were pretreated with gemcitabine prior to combining with mocetinostat or when both drugs were administered concurrently (data not shown). The combination effects on apoptosis were evaluated next. SKLMS1 cells were pretreated with mocetinostat then combined with gemcitabine and analyzed for caspase 3/7 activity using the Incucyte Zoom. The data demonstrates a significant pro-apoptotic effect in cells treated with the combination versus either drug alone (Fig 2C). These *in vitro* findings lead us to test this combination *in vivo*.

Mocetinostat combined with gemcitabine versus either drug alone demonstrates a superior anti-LMS effect *in vivo*

SKLMS1 xenografts were used for *in vivo* testing of mocetinostat combined with gemcitabine in that LMS1 cells failed to grow as xenografts in SCID mice. Once tumors reached approximately 100 mm³, mocetinostat was administered i.p. daily at a dose of 50 mg/kg to mice in the mocetinostat alone and mocetinostat plus gemcitabine groups. The treatments were well tolerated without significant weight loss. Treatment with mocetinostat alone did not significantly affect SKLMS1 tumor growth, whereas gemcitabine alone induced significant tumor growth inhibition ($p \leq 0.0001$). Mocetinostat combined with gemcitabine significantly inhibited SKLMS1 tumor growth compared to either treatment alone (Fig 3; $p \leq 0.0001$). The average tumor weights at the end of the study were: 0.29 g \pm 0.03 for vehicle, 0.27 g \pm 0.03 for mocetinostat alone, 0.12 g \pm 0.02 for gemcitabine alone, and 0.05 g \pm 0.01 for the combination (Fig 3). Mocetinostat alone modestly reduced tumor weight, while gemcitabine alone ($p \leq 0.0001$) and the drug combination significantly ($p \leq 0.0001$) reduced SKLMS1 tumor weight compared to vehicle.

Mocetinostat reduces gemcitabine-resistance associated target expression *in vitro* and *in vivo*

To identify potential mechanisms of mocetinostat-induced sensitization to gemcitabine, LMS cells were treated with mocetinostat and immunoblotted for the key gemcitabine-resistant targets, RRM1, RRM2, and hENT1 [22, 23], demonstrating mocetinostat-induced down-regulation of RRM1, RRM2, and an increase in hENT1 protein expression in LMS cells (Fig 4). Mocetinostat-induced regulation of these proteins may contribute to the observed synergistic sensitization to gemcitabine *in vitro* and *in vivo*, and this direction is being evaluated by us at this time.

Discussion

Our investigations demonstrate anti-LMS synergy using the combination of mocetinostat and gemcitabine. Our preclinical data explores a novel therapeutic approach for the treatment of LMS. The role of gemcitabine in sarcoma is limited. Initially, gemcitabine was shown to have




Fig 2. Mocetinostat synergizes with gemcitabine in LMS cells *in vitro*. A and B, MTS assays demonstrating synergistic combination of mocetinostat and gemcitabine in SKLMS1 and LMS cells, respectively (cells were pretreated with mocetinostat prior to combining with gemcitabine). C, Mocetinostat combined with gemcitabine enhances caspase 3/7 positive SKLMS1 cells. Combination index (CI) values of the drug combinations were calculated using Compusyn software. Synergy, additivity, and antagonism are defined as $CI < 1$, $CI = 1$, and $CI > 1$, respectively.

<https://doi.org/10.1371/journal.pone.0188859.g002>

benefit for the treatment of chemo-resistant osteosarcoma [24], leading the way for various clinical trials in sarcomas of different histologies. The trials demonstrate uterine LMS (ULMS) to respond best to gemcitabine compared to non-uterine LMS and STS of different histology [8]. Recently, markers associated with gemcitabine resistance were shown to be upregulated in a cohort of ULMS patient samples [25]. Studies focusing on the role of HDAC inhibition in STS warrants further investigation, especially in the context of LMS [18, 26, 27].

The efficacy of HDAC inhibition combined with gemcitabine has been previously described in various tumor models [28–41]. Extensive examination of gemcitabine resistance in pancreatic cancer has yielded strategies to overcome gemcitabine resistance using rational combination therapies, including several classes of HDAC inhibitor compounds. Similar to our study, Sung et al. demonstrated the synergistic efficacy of mocetinostat in combination with gemcitabine in human pancreatic cancer cells [42]. Gemcitabine resistant cells have been shown to be sensitive to HDAC inhibition [43]. Recently, Lee et al., demonstrated that the novel HDAC inhibitor, CG200745, reduced the expression of multidrug-resistant protein MRP4 in pancreatic cells, suggesting a potential synergistic mechanism of HDAC inhibition combined with gemcitabine [44].

Pan-HDAC inhibitors exhibit side effects prompting the development of isoform-specific HDAC inhibitors, with the aim of reducing toxicities while enhancing the therapeutic window. Gong et al. demonstrated the effective combination of sirtuin 1 (SIRT1) inhibition and gemcitabine in pancreatic carcinoma. Sirtuin 1 is one of seven NAD-dependent deacetylase isoforms [45], suggesting that the combination of HDAC/SIRT isoform-specific inhibition and gemcitabine may warrant further investigation.

Epigenetic regulation of acquired gemcitabine resistance has been described [46–49]. HDAC inhibition may be crucial to help establish potential epigenetic markers of LMS gemcitabine resistance that can be targeted for therapy. Essential proteins that have a role in gemcitabine resistance include RRM1, RRM2, and hENT1. We have demonstrated mocetinostat-induced regulation of these proteins, suggesting a potential mechanism of mocetinostat-




Fig 3. Mocetinostat combined with gemcitabine exhibits significant anti-LMS effect *in vivo*. Mocetinostat combined with gemcitabine significantly reduced tumor growth and tumor weight.

<https://doi.org/10.1371/journal.pone.0188859.g003>




Fig 4. Mocetinostat regulates gemcitabine-resistant markers expression. Mocetinostat reduces RRM1 and RRM2, and increases hENT1 expression in LMS cell lines.

<https://doi.org/10.1371/journal.pone.0188859.g004>

induced sensitivity to gemcitabine. In LMS, RRM1 is more highly expressed in uterine LMS versus extrauterine LMS patients [25], a possible biological dichotomy relevant to therapeutic intervention. The roles of RRM2 and hENT1 in the context of gemcitabine resistance in leiomyosarcoma have yet to be identified, and are currently under study in our laboratory.

Our work expands the knowledge base needed to identify potentially novel therapeutic options for the treatment of leiomyosarcoma, especially in the context of chemotherapeutic resistance. This work also identifies novel LMS cell lines; there is a major lack of available LMS cell lines at this time. This study presents a platform to further study the mechanistic role of HDAC inhibition and gemcitabine resistance in LMS. Accordingly, we currently investigating isoform-specific HDAC inhibition combined with gemcitabine to determine the utility of specific HDAC isoforms in anti-LMS therapy, including their potential synergism with gemcitabine as per above. These studies also utilize several LMS patient-derived xenograft (PDX) models we have recently developed as a preclinical bridge to future clinical trials to evaluate these severely needed and promising new approach to LMS patient therapeutics.

Supporting information

S1 Checklist. The ARRIVE guidelines checklist.
(PDF)

S1 Table. STR analysis of LMS cell lines compared to their corresponding patient tumor.
(XLSX)

S2 Table. Median values and confidence intervals (95%).
(XLSX)

S3 Table. EC50 of mocetinostat on cell lines.
(XLSX)

Acknowledgments

We would like to acknowledge Dr. Nyla Heerema and Jadwiga Labanowska for karyotyping analysis, the OSU Plant-Microbe Genomics Facility for STR analysis, the National Leiomyosarcoma Foundation/NLMSF (GRT00041181/60052311), and the SARC SPORE (U54CA168512).

Author Contributions

Conceptualization: Gonzalo Lopez, Edwin Choy, Dina Lev, Raphael Pollock.

Data curation: Gonzalo Lopez, Danielle Braggio, Abeba Zewdu, Lucia Casadei, Kara Batte, Hemant Kumar Bid.

Formal analysis: Gonzalo Lopez, Abeba Zewdu, David Koller, Peter Yu, Obiajulu Hans Iwenofu.

Investigation: Gonzalo Lopez, Danielle Braggio, Abeba Zewdu.

Methodology: Gonzalo Lopez, Hemant Kumar Bid.

Project administration: Kara Batte.

Resources: Gonzalo Lopez.

Software: Gonzalo Lopez, Abeba Zewdu.

Supervision: Gonzalo Lopez, Anne Strohecker, Edwin Choy, Dina Lev, Raphael Pollock.

Validation: Gonzalo Lopez.

Visualization: Gonzalo Lopez, Obiajulu Hans Iwenofu.

Writing – original draft: Gonzalo Lopez, Raphael Pollock.

Writing – review & editing: Gonzalo Lopez, Anne Strohecker, Raphael Pollock.

References

1. Wang Z, Shi N, Naing A, Janku F, Subbiah V, Araujo DM, et al. Survival of patients with metastatic leiomyosarcoma: the MD Anderson Clinical Center for targeted therapy experience. *Cancer Med.* 2016; 5: 3437–3444. <https://doi.org/10.1002/cam4.956> PMID: 27882721
2. Wang R, Lu YJ, Fisher C, Bridge JA, Shipley J. Characterization of chromosome aberrations associated with soft-tissue leiomyosarcomas by twenty-four-color karyotyping and comparative genomic hybridization analysis. *Genes Chromosomes Cancer.* 2001; 31: 54–64. <https://doi.org/10.1002/gcc.1118> PMID: 11284036
3. Hensley ML, Maki R, Venkatraman E, Geller G, Lovegren M, Aghajanian C, et al. Gemcitabine and docetaxel in patients with unresectable leiomyosarcoma: results of a phase II trial. *J Clin Oncol.* 2002; 20: 2824–2831. <https://doi.org/10.1200/JCO.2002.11.050> PMID: 12065559
4. Heinemann V, Hertel L W, Grindey G B, Plunkett W. Comparison of the cellular pharmacokinetics and toxicity of 2',2'-difluorodeoxycytidine and 1-beta-D-arabinofuranosylcytosine. *Cancer Res.* 1988; 48: 4024–4031. PMID: 3383195
5. Look KY, Sandler A, Blessing JA, Lucci JA 3rd, Rose PG, Gynecologic Oncology Group S. Phase II trial of gemcitabine as second-line chemotherapy of uterine leiomyosarcoma: a Gynecologic Oncology Group (GOG) Study. *Gynec Oncol.* 2004; 92: 644–647.
6. Seddon B, Scurr M, Jones RL, Wood Z, Probert-Lewis C, Fisher C, et al. A phase II trial to assess the activity of gemcitabine and docetaxel as first line chemotherapy treatment in patients with unresectable leiomyosarcoma. *Clin Sarcoma Res.* 2015; 5: 13. <https://doi.org/10.1186/s13569-015-0029-8> PMID: 25987978
7. Hensley ML, Blessing JA, Mannel R, Rose PG. Fixed-dose rate gemcitabine plus docetaxel as first-line therapy for metastatic uterine leiomyosarcoma: a Gynecologic Oncology Group phase II trial. *Gynecol Oncol.* 2008; 109: 329–34. <https://doi.org/10.1016/j.ygyno.2008.03.010> PMID: 18534250

8. Maki RG. Gemcitabine and docetaxel in metastatic sarcoma: past, present, and future. *Oncologist*. 2007; 12: 999–1006. <https://doi.org/10.1634/theoncologist.12-8-999> PMID: 17766660
9. West AC, Johnstone RW. New and emerging HDAC inhibitors for cancer treatment. *J Clin Invest*. 2014; 124: 30–9. <https://doi.org/10.1172/JCI69738> PMID: 24382387
10. Iizuka M, Smith MM. Functional consequences of histone modifications. *Curr Opin Genet Dev*. 2003; 13: 154–160. PMID: 12672492
11. Johnstone RW. Histone-deacetylase inhibitors: novel drugs for the treatment of cancer. *Nat Rev Drug Disc*. 2002; 1: 287–299.
12. Haberland M, Montgomery RL, Olson EN. The many roles of histone deacetylases in development and physiology: implications for disease and therapy. *Nat Rev Genet*. 2009; 10: 32–42. <https://doi.org/10.1038/nrg2485> PMID: 19065135
13. Fournel M, Bonfils C, Hou Y, Yan PT, Trachy-Bourget MC, Kalita A, et al. MGCD0103, a novel isotype-selective histone deacetylase inhibitor, has broad spectrum antitumor activity in vitro and in vivo. *Mol Cancer Ther*. 2008; 7: 759–68. <https://doi.org/10.1158/1535-7163.MCT-07-2026> PMID: 18413790
14. Boumber Y, Younes A, Garcia-Manero G. Mocetinostat (MGCD0103): a review of an isotype-specific histone deacetylase inhibitor. *Expert Opin Investig Drugs*. 2011; 20: 823–9. <https://doi.org/10.1517/13543784.2011.577737> PMID: 21554162
15. Siu LL, Pili R, Duran I, Messersmith WA, Chen EX, Sullivan R, et al. Phase I study of MGCD0103 given as a three-times-per-week oral dose in patients with advanced solid tumors. *J Clin Oncol*. 2008; 26: 1940–7. <https://doi.org/10.1200/JCO.2007.14.5730> PMID: 18421048
16. Chou T. Drug combination studies and their synergy quantification using the Chou-Talalay method. *Cancer Res*. 2010; 70: 440–46. <https://doi.org/10.1158/0008-5472.CAN-09-1947> PMID: 20068163
17. Kilkeny C, Browne WJ, Cuthill IC, Emerson M, Altman DG. Improving bioscience research reporting: the ARRIVE guidelines for reporting animal research. *PLoS Biol*. 2010; 8: e1000412 <https://doi.org/10.1371/journal.pbio.1000412> PMID: 20613859
18. Lopez G, Liu J, Ren W, Wei W, Wang S, Lahat G, et al. Combining PCI-24781, a novel histone deacetylase inhibitor, with chemotherapy for the treatment of soft tissue sarcoma. *Clin Cancer Res*. 2009; 15: 3472–83. <https://doi.org/10.1158/1078-0432.CCR-08-2714> PMID: 19417021
19. Lopez G, Torres K, Liu J, Hernandez B, Young E, Belousov R, et al. Autophagic survival in resistance to histone deacetylase inhibitors: novel strategies to treat malignant peripheral nerve sheath tumors. *Cancer Res*. 2011; 71: 185–96. <https://doi.org/10.1158/0008-5472.CAN-10-2799> PMID: 21084276
20. Buggy JJ, Cao ZA, Bass KE, Verner E, Balasubramanian S, Liu L, et al. CRA-024781: a novel synthetic inhibitor of histone deacetylase enzymes with antitumor activity in vitro and in vivo. *Mol Cancer Ther*. 2006; 5: 1309–17. <https://doi.org/10.1158/1535-7163.MCT-05-0442> PMID: 16731764
21. Wagner JM, Hackanson B, Lubbert M, Jung M. Histone deacetylase (HDAC) inhibitors in recent clinical trials for cancer therapy. *Clin Epigenetics*. 2010; 1: 117–36. <https://doi.org/10.1007/s13148-010-0012-4> PMID: 21258646
22. Aye Y, Li M, Long MJC, Weiss RS. Ribonucleotide reductase and cancer: biological mechanisms and targeted therapies. *Oncogene*. 2015; 34: 2011–21. <https://doi.org/10.1038/onc.2014.155> PMID: 24909171
23. Pastor-Anglada M, Pérez-Torras S. Nucleoside transporter proteins as biomarkers of drug responsiveness and drug targets. *Front Pharmacol*. 2015; 6: 13. <https://doi.org/10.3389/fphar.2015.00013> PMID: 25713533
24. Merimsky O, Meller I, Kollender Y, Inbar M. Palliative effect of gemcitabine in osteosarcoma resistant to standard chemotherapy. *Eur J Cancer*. 1998; 34: 1296–7. PMID: 9849496
25. Chatterjee S, Xiu J, Reddy SK, Jones NL, Buckley de Meritens A, Tergas AI, et al. Uterine leiomyosarcomas exhibit distinct drug resistance molecular profiles compared to extrauterine leiomyosarcomas: A comprehensive analysis of 1,023 leiomyosarcomas. *Gynecol Oncol*. 2016; 141(Suppl 1):51–2.
26. Sampson ER, Amin V, Schwarz EM, O’Keefe RJ, Rosier RN. The histone deacetylase inhibitor vorinostat selectively sensitizes fibrosarcoma cells to chemotherapy. *J Orthop Res*. 2011; 29: 623–32. <https://doi.org/10.1002/jor.21274> PMID: 20957741
27. Lee J, McGuire C. Clinical efficacy of Vorinostat in a patient with leiomyosarcoma. *Clin Med Insights Oncol*. 2012; 6: 101–5. <https://doi.org/10.4137/CMO.S7194> PMID: 22408374
28. Rundall BK, Denlinger CE, Jones DR. Suberoylanilide hydroxamic acid combined with gemcitabine enhances apoptosis in non-small cell lung cancer. *Surgery*. 2005; 138: 360–7. <https://doi.org/10.1016/j.surg.2005.06.016> PMID: 16153448
29. Piacentini P, Donadelli M, Costanzo C, Moore PS, Palmieri M, Scarpa A. Trichostatin A enhances the response of chemotherapeutic agents in inhibiting pancreatic cancer cell proliferation. *Virchows Arch*. 2006; 448: 797–804. <https://doi.org/10.1007/s00428-006-0173-x> PMID: 16568310

30. Schniewind B, Heintz K, Kurdow R, Ammerpohl O, Trauzold A, Emme D, et al. Combination phenylbutyrate/gemcitabine therapy effectively inhibits in vitro and in vivo growth of NSCLC by intrinsic apoptotic pathways. *J Carcinog*. 2006; 5: 25. <https://doi.org/10.1186/1477-3163-5-25> PMID: 17123441
31. Ammerpohl O, Trauzold A, Schniewind B, Griep U, Pilarsky C, Grutzmann R, et al. Complementary effects of HDAC inhibitor 4-PB on gap junction communication and cellular export mechanisms support restoration of chemosensitivity of PDAC cells. *Br J Cancer*. 2007; 96: 73–81. <https://doi.org/10.1038/sj.bjc.6603511> PMID: 17164759
32. Donadelli M, Costanzo C, Beghelli S, Scupoli MT, Dandrea M, Bonora A, Piacentini P, et al. Synergistic inhibition of pancreatic adenocarcinoma cell growth by trichostatin A and gemcitabine. *Biochim Biophys Acta*. 2007; 1773: 1095–106. <https://doi.org/10.1016/j.bbamcr.2007.05.002> PMID: 17555830
33. Iwahashi S, Shimada M, Utsunomiya T, Morine Y, Imura S, Ikemoto T, Mori H, et al. Histone deacetylase inhibitor augments anti-tumor effect of gemcitabine and pegylated interferon- α on pancreatic cancer cells. *Int J Clin Oncol*. 2011; 16: 671–8. <https://doi.org/10.1007/s10147-011-0246-y> PMID: 21556798
34. Iwahashi S, Shimada M, Utsunomiya T, Morine Y, Imura S, Ikemoto T, et al. Histone deacetylase inhibitor enhances the anti-tumor effect of gemcitabine: a special reference to gene-expression microarray analysis. *Oncol Rep*. 2011; 26: 1057–62. <https://doi.org/10.3892/or.2011.1407> PMID: 21805043
35. Dovzhanskiy DI, Arnold SM, Hackert T, Oehme I, Witt O, Felix K, et al. Experimental in vivo and in vitro treatment with a new histone deacetylase inhibitor belinostat inhibits the growth of pancreatic cancer. *BMC Cancer*. 2012; 812: 226.
36. Klein JM, Henke A, Sauer M, Bessler M, Reiners KS, Engert A, et al. The histone deacetylase inhibitor LBH589 (panobinostat) modulates the crosstalk of lymphocytes with Hodgkin lymphoma cell lines. *PLoS One*. 2013; 8: e79502. <https://doi.org/10.1371/journal.pone.0079502> PMID: 24278143
37. Minjie S, Defei H, Zhimin H, Weiding W, Yuhua Z. Targeting pancreatic cancer cells by a novel hydroxamate-based histone deacetylase (HDAC) inhibitor ST-3595. *Tumour Biol*. 2015; 36: 9015–22. <https://doi.org/10.1007/s13277-015-3537-5> PMID: 26084607
38. Xue K, Gu JJ, Zhang Q, Mavis C, Hernandez-Ilizaliturri FJ, Czuczman MS, et al. Vorinostat, a histone deacetylase (HDAC) inhibitor, promotes cell cycle arrest and re-sensitizes rituximab- and chemo-resistant lymphoma cells to chemotherapy agents. *J Cancer Res Clin Oncol*. 2016; 142: 379–87. <https://doi.org/10.1007/s00432-015-2026-y> PMID: 26314218
39. Sakamoto T, Kobayashi S, Yamada D, Nagano H, Tomokuni A, Tomimaru Y, et al. A histone deacetylase inhibitor suppresses epithelial-mesenchymal transition and attenuates chemoresistance in biliary tract cancer. *PLoS One*. 2016; 11: e0145985. <https://doi.org/10.1371/journal.pone.0145985> PMID: 26726879
40. Henderson SE, Ding LY, Mo X, Bekaii-Saab T, Kulp SK, Chen CS, et al. Suppression of tumor growth and muscle wasting in a transgenic mouse model of pancreatic cancer by the novel histone deacetylase inhibitor AR-42. *Neoplasia*. 2016; 18: 765–774. <https://doi.org/10.1016/j.neo.2016.10.003> PMID: 27889645
41. Lee HS, Park SB, Kim SA, Kwon SK, Cha H, Lee DY, et al. A novel HDAC inhibitor, CG200745, inhibits pancreatic cancer cell growth and overcomes gemcitabine resistance. *Sci Rep*. 2017; 7: 41615. <https://doi.org/10.1038/srep41615> PMID: 28134290
42. Sung V, Richard N, Brady H, Maier A, Kelter G, Heise C. Histone deacetylase inhibitor MGCD0103 synergizes with gemcitabine in human pancreatic cells. *Cancer Sci*. 2011; 102: 1201–7. <https://doi.org/10.1111/j.1349-7006.2011.01921.x> PMID: 21375679
43. Samulitis BK, Pond KW, Pond E, Cress AE, Patel H, Wisner L, et al. Gemcitabine resistant pancreatic cancer cell lines acquire an invasive phenotype with collateral hypersensitivity to histone deacetylase inhibitors. *Cancer Biol Ther*. 2015; 16: 43–51. <https://doi.org/10.4161/15384047.2014.986967> PMID: 25485960
44. Lee HS, Park SB, Kim SA, Kwon SK, Cha H, Lee DY, et al. A novel HDAC inhibitor, CG200745, inhibits pancreatic cancer cell growth and overcomes gemcitabine resistance. *Sci Rep*. 2017; 30: 41615.
45. Cen Y, Youn DY, Sauve AA. Advances in characterization of human sirtuin isoforms: chemistries, targets and therapeutic applications. *Curr Med Chem*. 2011; 18: 1919–35. PMID: 21517779
46. Candelaria M, de la Cruz-Hernández E, Pérez-Cárdenas E, Trejo-Becerril C, Gutiérrez-Hernández O, Dueñas-González A. Pharmacogenetics and pharmacoeugenetics of gemcitabine. *Med Oncol*. 2010; 27: 1133–43. <https://doi.org/10.1007/s12032-009-9349-y> PMID: 19902390
47. Candelaria M, de la Cruz-Hernandez E, Taja-Chayeb L, Perez-Cardenas E, Trejo-Becerril C, Gonzalez-Fierro A, et al. DNA methylation-independent reversion of gemcitabine resistance by hydralazine in cervical cancer cells. *PLoS One*. 2012; 7: e29181. <https://doi.org/10.1371/journal.pone.0029181> PMID: 22427797

48. Ansari D, Urey C, Hilmersson KS, Bauden MP, Ek F, Olsson R, et al. Apicidin sensitizes pancreatic cancer cells to gemcitabine by epigenetically regulating MUC4 expression. *Anticancer Res.* 2014; 34: 5269–76. PMID: [25275019](https://pubmed.ncbi.nlm.nih.gov/25275019/)
49. Hosokawa M, Saito M, Nakano A, Iwashita S, Ishizaka A, Ueda K, et al. Acquired resistance to decitabine and cross-resistance to gemcitabine during the long-term treatment of human HCT116 colorectal cancer cells with decitabine. *Oncol Lett.* 2015; 10: 761–767. <https://doi.org/10.3892/ol.2015.3253> PMID: [26622566](https://pubmed.ncbi.nlm.nih.gov/26622566/)