



POSTER PRESENTATION

Open Access

Pulse wave velocity in real-time cardiac magnetic resonance

Martin Fasshauer^{1,5*}, Johannes T Kowallick^{1,5}, Arun Joseph^{2,5}, Christina Unterberg-Buchwald^{3,5}, Klaus-Dietmar Merboldt^{2,5}, Michael Steinmetz^{4,5}, Jan M Sohns^{1,5}, Wieland Staab^{1,5}, Andreas Schuster^{3,5}, Dirk Voit^{2,5}, Sebastian Schaetz^{2,5}, Shoun Zhang^{2,5}, Jens Frahm^{2,5}, Joachim Lotz^{1,5}

From 17th Annual SCMR Scientific Sessions
New Orleans, LA, USA. 16-19 January 2014

Background

Atherosclerosis and its associated diseases are constantly increasing in developed countries. Aortic stiffness is an indicator for atherosclerosis and is associated with mortality and morbidity especially in aortic abdominal aneurysms. For further evaluation of aortic stiffness, we examined n = 13 healthy volunteers using real-time magnetic-resonance imaging (RT-MRI) with highly undersampled radial fast low-angle shot (FLASH) acquisitions, phase-sensitive image reconstructions and regularized nonlinear inversion (NLINV). We hypothesized that RT-MRI is able to determine pulse wave velocity (PWV) and flow data using just one transverse view of the ascending and descending aorta. This method could be superior to complex known MRI methods using velocity projection today.

Methods

We assessed PWV as surrogate parameter for aortic stiffness by velocity-encoded RT-MRI. Time lag between the ascending and descending aortic pulse wave was calculated and divided by the mean length of the aortic arch for each individual in detail. RT-MRI can determine PWV during normal breathing, physical strain and recovery from strain using the Valsalva (VM) and Mueller maneuvers (MM). During strain/maneuvers volunteers had visual feedback of intra-thoracic pressure via a mouthpiece to ensure adequate strain performance. We calculated PWV in normal breathing, at the end of strain and at the end of the recovery phase. Due to the advantage of a single heart beat-to-beat variability with

RT-MRI, the average PWV of 4 heartbeats could be calculated in each phase with Standard deviation (SD).

Results

While changes during MM were very subtle (normal breathing $3.23 \text{ m/s} \pm 0.05 \text{ SD}$ vs. MM $2.99 \text{ m/s} \pm 0.04 \text{ SD}$ vs. recovery $3.24 \text{ m/s} \pm 0.05$), increase and especially decrease during recovery from VM was more pronounced (normal breathing $3.55 \text{ m/s} \pm 0.06$ vs. VM $4.12 \text{ m/s} \pm 0.08$ vs. recovery $3.13 \text{ m/s} \pm 0.06$, VM vs. recovery $p < 0.05$, values shown as mean \pm SEM).

Conclusions

Successful measurements of PWV in one transverse view during physical strain have not been shown before to our knowledge and prove to be an easy alternative to previous MR attempts determining PWV. In addition, the significant PWV decrease measured in RT-MRI from VM to recovery resembles the normal variability of aortic stiffness in healthy subjects due to change in intra-thoracic pressure. Real time MRI is able to measure these data under stress maneuvers such as the VM or MM. A combined score of PWV and its variability might be helpful for risk assessment in the development of aortic aneurysms

Funding

This work was funded by the Deutsche Forschungsgemeinschaft (LO 1773/1-1).

Authors' details

¹Institute for Diagnostic and Interventional Radiology, University Medical Center Göttingen, Göttingen, Germany. ²Biomedizinische NMR ForschungsGmbH, Max-Planck-Institut für biophysikalische Chemie, Göttingen, Germany.

³Clinic for Cardiology and Pneumology, Heart Center, University Medical Center Göttingen, Göttingen, Germany. ⁴Clinic for Pediatric Cardiology and

¹Institute for Diagnostic and Interventional Radiology, University Medical Center Göttingen, Göttingen, Germany
Full list of author information is available at the end of the article

© 2014 Fasshauer et al.; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Intensive Care Medicine, University Medical Center Göttingen, Göttingen, Germany. ⁵partner site Göttingen, DZHK (German Cardiovascular Research Center), Göttingen, Germany.

Published: 16 January 2014

doi:10.1186/1532-429X-16-S1-P382

Cite this article as: Fasshauer et al.: Pulse wave velocity in real-time cardiac magnetic resonance. *Journal of Cardiovascular Magnetic Resonance* 2014 **16**(Suppl 1):P382.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

