



ARTICLE

Antimicrobial Activity of Propolis Extract and Their Application as a Natural Preservative in Livestock Products: A Meta-Analysis

Andre^{1,*}, Irma Isnafia Arief¹, Astari Apriantini¹, Anuraga Jayanegara², and Cahyo Budiman¹

¹Department of Animal Production Science and Technology, Faculty of Animal Science, IPB University, Bogor 16680, Indonesia

²Department of Nutrition and Feed Science and Technology, Faculty of Animal Science, IPB University, Bogor 16680, Indonesia

OPEN ACCESS

Received September 14, 2021

Revised November 17, 2021

Accepted January 18, 2022

*Corresponding author : Andre
Department of Animal Production Science
and Technology, Faculty of Animal Science,
IPB University, Bogor 16680, Indonesia
Tel: +6282128295872
E-mail: andreandre@apps.ipb.ac.id

*ORCID
Andre
<https://orcid.org/0000-0003-2484-506X>
Irma Isnafia Arief
<https://orcid.org/0000-0001-8193-0194>
Astari Apriantini
<https://orcid.org/0000-0002-4811-5101>
Anuraga Jayanegara
<https://orcid.org/0000-0001-7529-9770>
Cahyo Budiman
<https://orcid.org/0000-0002-2052-9572>

Abstract This study aimed to evaluate the effectiveness of propolis extract as a natural preservative for livestock products in term of chemical and microbiological characteristics by meta-analysis. The stages carried out in this study were identification, selection, checking suitability, and the resulting selected articles were used in the meta-analysis. The selection results obtained a total of 22 selected journal articles consisting of 9 articles for analysis of the antimicrobial activity of propolis extract and 13 articles for analysis of the chemical and microbiological characteristics of livestock products. The articles were obtained from electronic databases, namely Science Direct and Google Scholar. The model used in this study is the random-effect model involving two groups, control and experimental. Heterogeneity and effect size values were carried out in this study using Hedge's obtained through openMEE software. Forest plot tests and data validation on publication bias was obtained using Kendall's test through JASP 0.14.1 software. The results showed that there is a significant relationship between propolis extract with the results of the antimicrobial activity ($p < 0.05$). In addition, the results of the application of propolis extract on the livestock products for the test microbes and the value of thiobarbituric acid reactive substances (TBARs) showed significant results ($p < 0.05$). Conclusion based on the random-effect model on the effectiveness of antimicrobial activity of propolis extract and their application as a natural preservative of the chemical and microbiological characteristics of livestock products is valid by Kendall's test ($p > 0.05$). Propolis in this case effectively used as natural preservatives in livestock products.

Keywords antimicrobial, livestock products, natural preservative, propolis

Introduction

Propolis is a product produced by honey bees from a mixture of wax compounds, β -glucosidase enzymes, and resins from plant parts, shoots, and exudates derived from bee saliva (Rocha et al., 2012). For decades, the chemical composition and properties

of propolis have been extensively studied and research data have also been published in various scientific papers around the world. From 1998 until now, more than 5,000 articles on propolis have been published in Science Direct. The chemical composition of propolis varies according to geographic area, climate, environmental conditions, and harvest season (López et al., 2014). More than 420 chemical compounds have been identified in propolis from various geographic regions of the world (Bankova, 2005; Milojković Opsenica et al., 2016). Propolis has a reputation as a natural product in the world. In recent decades, it has been widely accepted in many countries as a dietary supplement for promoting health and preventing disease. In general, propolis influences human health (Azemin et al., 2018; Farooqui and Farooqui, 2012).

The health-promoting properties of propolis come from its chemical composition, including antimicrobial and antiviral (Bankova et al., 2014; das Neves et al., 2016; Nolkemper et al., 2010), antioxidant properties (Azemin et al., 2018; Mello and Hubinger, 2012; Sun et al., 2015), anticancer (Markiewicz-Zukowska et al., 2013; Xuan et al., 2014), anti-inflammatory and cytostatic (Corrêa et al., 2017; Kismet et al., 2017), immunostimulants (Nassar et al., 2012), and anti-allergic (Yasar et al., 2016). The rich bioactive components are useful in their application in various fields such as medicine and dentistry, pharmaceuticals, cosmetics, and the food industry.

Research data related to antimicrobial properties and the application of propolis extract to livestock products are quite widely published at the national and international levels. The results of research on the antimicrobial properties of propolis extract have been widely published and quite a lot related to its application. The antimicrobial properties of propolis are effective against the type of bacteria, both gram-positive and negative, molds, and fungi (gram-positive bacteria: *Staphylococcus aureus*, *Bacillus cereus*, *Listeria monocytogenes*, *Enterococcus faecalis*; gram-negative bacteria: *Salmonella enteritidis*, *Shigella sonnei*, *Klebsiella pneumoniae*, *Escherichia coli* O157, *Proteus mirabilis*, *Enterobacter aerogenes*, *Pseudomonas aeruginosa*; molds: *Rhodotorula mucilaginosa*, *Candida albicans*, *Candida krusei*, *Saccharomyces cerevisiae*; and fungi: *Colletotrichum gloeosporoides*, *Alternaria solani*, *Fusarium solani*, *Rhizopus stolonifer*, *Botrytis cinerea*, *Cladosporium cladosporoides*, *Aspergillus niger*, *Aspergillus ochraceus*, *Mucor mucedo*, *Penicillium expansum*, *Penicillium chrysogenum*; Pobiega et al., 2019a). Meanwhile, several studies stated that the antimicrobial properties of propolis extract were applied to livestock products (such as fermented meat sausage, beef patties, fresh oriental sausage, sausage, Tuscan sausage, milk, and ice cream) with various concentrations of propolis extract added (Pobiega et al., 2019b). Preliminary research was also carried out related to the use of propolis extract as food preservation in beef products stored for 24 hours. The result data shows that the higher the concentration used, the better the quality given (Andre et al., 2021). However, in this study, there is not enough data to produce comprehensive information. Thus, it is necessary to conduct a study using meta-analysis to produce information related to problems in the field of livestock products.

Regulation of propolis on the production in the food sector are grouped under the category of health products. However, the regulation depends on region on each with the requirements of registration are different. Now, the legal regulations regarding the use of propolis has been introduced in various areas, such as Brazil, the USA, European Union, Australia, Canada, China, Japan, and Korea (Berretta et al., 2017). Propolis in it is classified as a dietary supplement and has been listed on the group Directive 2002/46/EC, as a source of concentrated nutrients with physiological effects (European Parliament and Council of the European Union, 2002). However, everything depends on the stage of production and the method of extraction used [European Food Safety Authority (EFSA), 2010]. Aspects of the commercialization of propolis, especially in my region it was reported that as many as 3,957 kg/year production propolis is produced from a total of four farmer groups (Utama et al., 2021). Given the magnitude of the results of such production, along with groups from different areas and regions, propolis enough potential to be commercialized synergy with the resulting production.

A meta-analysis, according to Anwar (2005), is a statistical technique for quantitatively combining two or more original studies (a statistical procedure for combining data from various studies). Meta-analyses have been carried out in the fields of medicine, pharmacy, education, psychology, criminology, business, marketing, economics, management, nutrition, and food (Donker et al., 2014). This study aims to obtain comprehensive information through quantitative data analysis using meta-analysis. In addition to providing information related to a meta-analysis on food, especially in the livestock sector, this study aims to evaluate the effectiveness of the potential of propolis as food preservation by considering the results of the analysis of the antimicrobial compound content of propolis extract and its application to the chemical and microbiological characteristics of livestock products. Therefore, this study is expected to provide information related to natural preservatives as a solution to problems in the field of livestock products.

Materials and Methods

The meta-analysis in this study was divided into two groups which were analyzed separately using the same method. This is due to the limitations of several studies. The division of these groups, the antimicrobial activity of propolis extract, and the application of propolis extract to chemical and microbiological properties are presented in Table 1.

Research stages

The stages carried out in this study were identification, selection, suitability check, and the final included articles were selected to be used in the meta-analysis. The identification stage included the number of journals produced by scientific database search engines such as Science Direct, Springer, and others using certain keywords and search results on other sources. The selection of articles was done using reference software such as Zotero and Mendeley to read the titles and abstracts of the journals and then eliminated the duplicate journals. The conformity assessment stage was carried out by viewing the full article and selecting it based on its suitability with the topic under study. The data required included complete control and experimental treatment with the average value, number of samples, and SD or SE. Articles that meet these requirements were then used for meta-analysis. The meta-analysis data processing was carried out using Microsoft Excel 2019 on the recap data of the selected article data extraction results. The articles were obtained from electronic databases, namely Science Direct and Google Scholar. The effect size value used in this study used Hedge's d .

Data analysis

The data obtained were analyzed using OpenMME software, calculated effect size to obtain information related to

Table 1. Group division in the meta-analysis study

Group 1	Group 2
(antimicrobial activity of propolis extract)	(application of propolis extract on chemical and microbiological characteristics)
Subgroup	
Pathogenic microbes	Chemical characteristics
	TBARs
	Microbiology
	Microbial test

TBARs, thiobarbituric acid reactive substances.

heterogeneity test and JASP 0.14.1 to obtain information related to standardized mean difference and publication bias.

Results and Discussion

Initial search results on antimicrobial activity from databases of international journals and Google search engine obtained 522 articles. After removing the articles with the same content and selecting the title and abstract contents, 9 complete articles were selected which were assessed for their suitability for meta-analysis. Complete articles were excluded from 101 articles as many as 92 due to inappropriate substances such as incomplete and inappropriate information. The following 9 articles that were used are presented in Table 2. Furthermore, the search results for the application of propolis extract on livestock products from the database of international journals and the google search engine are 216 articles. After removing articles with the same content and selecting titles and abstracts, 12 complete articles were selected which were assessed for their suitability for meta-analysis. Thirty eight complete articles were excluded due to inappropriate content; 109 articles were excluded with 50 selected articles from 159 duplicated articles. The following 12 articles that were used in this study are presented in Table 3 and diagram of the meta-analysis of screening, inclusion, and exclusion of articles are presented in Fig. 1.

Antimicrobial activity

The number of study data used in the meta-analysis of the difference in antimicrobial activity through the inhibition zone diffusion method at various concentrations of propolis extract was 22 with a range of 1998 to 2020 (Fig. 2). The total number of samples/ replications in the bacterial inhibition zone of propolis extract was 182. Based on the results of the calculation with random-effect model values obtained SMD/d+ overall by 4.05 with CI95% (2.96, 5.13; Table 4), because the confidence interval does not contain 0 (zero), then the treatment given to the experimental group different from the control group in terms of inhibiting the microbial activity. The results of the analysis showed (Table 4), then in this case the true effect size is

Table 2. Description of studies used in the meta-analysis of antimicrobial activity of propolis extract

Source	Study location	Extract method	N	EPc	Control	Bacterial species
Abdullah et al. (2020)	Brunei Darussalam	Ethanol	55	2 g/L	Negative	<i>Basillus subtilis</i> and <i>Staphylococcus aureus</i>
Abdullah et al. (2019)	Brunei Darussalam	Ethanol	18	2 mg/mL; 8 mg/mL	Negative	<i>Staphylococcus aureus</i> , and <i>Pseudomonas aeruginosa</i>
Seibert et al. (2019)	Brazil	Ethanol, hexane, ethyl acetate	36	50 mg/mL	Negative	<i>Staphylococcus saprophyticus</i> , <i>Listeria monocytogenes</i> , and <i>Enterococcus faecalis</i>
Khodayari et al. (2019)	Iran	Ethanol	4	2%	Negative	<i>Escherichia coli</i>
Rezaeigolestani et al. (2017)	Iran	Ethanol	12	2%	Negative	<i>Staphylococcus aureus</i> , and <i>V. parahaemolyticus</i>
Kujumgiev et al. (1999)	Bulgaria	Ethanol	6	0.1%	Negative	<i>Streptococcus aureus</i>
Oliveira et al. (2017)	Portugal	Ethanol	12	20 uL	Negative	<i>Staphylococcus aureus</i> , and <i>Salmonella Typhimurium</i>
Airen et al. (2018)	India	Ethanol	60	5%; 20%	Negative	<i>L. acidophilus</i> , and <i>Streptococcus mutans</i>
Tosi et al. (2007)	Argentina	Ethanol	10	1.4 mg	Negative	<i>Escherichia coli</i>

N, sample size; EPc, extract of propolis concentrations.

Table 3. Description of the studies used in the meta-analysis of the application of propolis extract to the value of TBARs and microbial testing of livestock products

Source	Study location	Product	N	EPc	Treatment	Output
Mehdizadeh and Mojaddar Langroodi (2019)	Iran	Chicken breast meat	18	1%	Storage at 4°C for 16 days	Inhibits oxidative activity and prolongs the shelf life.
Pedonese et al. (2019)	Italia	Milk and whey cheese	48	2% and 5%	Cultivation of pasteurized milk products at 37°C for 24 hours and storage for 28 days on cheese products	Inhibits the growth of <i>Bacillus cereus</i> , <i>Pseudomonas fluorescens</i> , and <i>Staphylococcus aureus</i> and prolongs shelf life.
Vargas-Sánchez et al. (2019)	Mexico	Beef patties	4	2%	Storage at 2°C for 16 days	Inhibits the growth of <i>Staphylococcus aureus</i> and prolongs shelf life.
Kisa et al. (2018)	Turkey	Turkish dry-fermented sausage	96	1% and 2%	Storage at 4°C for 30 days	Inhibits the occurrence of fat oxidation and the growth of mesophilic bacteria and extends the shelf life.
dos Reis et al. (2017)	Brazil	Burger meat	6	0.3 g/kg	Storage at -16°C for 28 days	Inhibits fat oxidation reactions and prolongs shelf life.
Viera et al. (2016)	Brazil	Tuscan sausage	24	0.5%	Storage at 4°C for 56 days	Inhibits the growth of mesophilic, psychotropic, and <i>Staphylococcus</i> bacteria and prolongs the shelf life.
Gutiérrez-Cortés and Suarez Mahecha (2014)	Colombia	Sausage	6	0.8 mg/mL	Storage for 24 days	Inhibits fat oxidation.
Vargas-Sánchez et al. (2014)	Mexico	Beef patties	6	2%	Refrigerator temperature storage for 8 days	Inhibits the growth of psychotropic bacteria.
El-Mossalami and Abdel-Hakeim (2013)	Egypt	Fresh Egyptian sausage	12	400 and 600 mg/kg	Storage at 5°C for 21 days	Inhibits fat oxidation reactions and prolongs shelf life.
Ali et al. (2010)	Egypt	Fresh oriental sausage	6	0.6%	Storage at 5°C for 21 days	Inhibits fat oxidation reactions and prolongs shelf life.
Kunrath et al. (2017)	Brazil	Salami Italian	12	0.01 and 0.05%	Storage at 18°C for 35 days	Inhibit oxidation reactions and prolongs shelf life.
Coró et al. (2020)	Brazil	Jerked beef	12	200 and 400 ppm	Storage at 25°C for 60 days	Inhibits oxidation reaction and prolongs shelf life.
Andre et al. (2021)	Indonesia	Beef slice	18	1% and 2%	Storage at 25°C for 24 hours	Inhibits oxidation reaction.

TBARs, thiobarbituric acid reactive substances; N, sample size; EPc, extract of propolis concentrations.

not equal to 0. It indicates that there is a significant relationship between propolis extract with the results of the antimicrobial activity ($p < 0.05$). However, before drawing any conclusions based on the random-effect model this would be accurate if it is proved that all research results in true effect size that is different in the population so that the need for the test of heterogeneity. Analysis of the heterogeneity of the impact on antimicrobial activity of propolis extract showed the presence of variability that occurs in all of the research ($p < 0.05$; Table 4). Then the assumption of homogeneity needs to be rejected and accept the assumption of heterogeneity, namely the variability that occurs not more caused by sampling error ($Y_i = \mu + \tau_i + \varepsilon_i$). It was also evidenced by the high percentage of I^2 (inconsistency) that is equal to 97.38% (Table 4). Thus it can be concluded that propolis extract is effective in inhibiting the growth of microbes.

Further evaluation of the validation of the effectiveness of propolis extract in inhibiting the growth of microbes test is required validation against the bias of the publication using the test provided by Kendall's test. The results of the analysis




Fig. 1. Diagram of the meta-analysis of selection, inclusion, and exclusion of articles. n1, group 1 (antimicrobial activity of propolis extract articles); n2, group 2 (application of propolis extract on chemical and microbiological characteristics articles).




Fig. 2. Forest plot meta-analysis of the differences in antimicrobial activity through the inhibition zone using the diffusion method at various concentrations of PE (propolis extract) using the Random Effect model.

Table 4. The results of the meta-analysis of the antimicrobial activity parameters of propolis extract microbial test and TBARs value for their application to livestock products

No	Parameter	N	SMD/d+ (RE 95% CI)	p-value	I ²	p-value	Kendall's τ	p-value
1	Antimicrobial activity	182	4.05 (2.96, 5.13)	<0.05	95.32%	<0.05	0.025	0.862
2	Microbial test	188	-1.20 (-1.69, -0.72)	<0.05	84.10%	<0.05	-0.179	0.138
3	TBARs	132	-1.62 (-2.27, -0.98)	<0.05	99.32%	<0.05	0.048	0.781

TBARs, thiobarbituric acid reactive substances; N, sample size; SMD/d+ (RE 95% CI), standardized mean difference (true effect size) by random effect model with 95% of confident interval; I², Inconsistency (percentage).

show that the p-value on the method (rank correlation test for Funnel plot Asymmetry) is symmetrical or in other words did not happen or finding evidence of a bias publication ($p > 0.05$). Negative rank correlation (-0.096) in Table 4 indicates that research with a big sample isn't included in the study sample meta-analysis. So the conclusion made based on the random-effect model on the effectiveness of antimicrobial activity of propolis extract is valid is indicated by the inconsistency is high and the bias publication of the resulting low through the test provided by Kendall's ($p < 0.05$; Table 4).

The potency of propolis extract in various studies shows that there is an activity from the influence of antimicrobial compound content with different concentrations. This is evidenced by the variability that occurs in all the studies analyzed. The variability that occurs is no longer caused by sampling error. The results showed that all propolis extracts showed strong antimicrobial activity against, namely *B. subtilis*, *S. aureus*, and *P. aeruginosa* (Kharsany et al., 2019; Okińczyc et al., 2020). Several alternatives can complement or can be used as a substitute for the use of synthetic preservatives, propolis provides antimicrobial effects in several studies that have been carried out (Aga et al., 1994; Farnesi et al., 2009). Furthermore, propolis has been tested as a food preservative because of its activity that can inhibit various bacteria and is safe (Tosi et al., 2007). The potential of propolis can show that propolis is economically feasible by introducing safe additive compounds as preservatives in food technology.

Microbial test

The number of study data used in the meta-analysis of differences in product microbial test results with the addition of different concentrations of propolis extract was 34 with the range from 2014 to 2019 (Fig. 3). The total number of samples/replications in the microbial test was 188. Based on the results of the calculation with random-effect model values obtained SMD/d+ overall by -1.20 with CI95% ($-1.69, -0.72$; Table 4), because the confidence interval does not contain 0 (zero), then the treatment given to the experimental group different from the control group in terms of inhibiting pathogenic microbes on products. These values indicate that the test microbes with the treatment of the addition of propolis extract on various products lower than the control. It is shown through the summary effect (the difference value is negative). The results of calculations showed (Table 4), then in this case the true effect size is not equal to 0. It indicates that there is a significant relationship between the addition of propolis extract with the results of the test microbes on the farm ($p < 0.05$). However, before drawing any conclusions based on the random-effect model this would be accurate if it is proved that all research results in true effect size that is different in the population so that the need for the test of heterogeneity. Analysis of the heterogeneity against test microbes of the product indicates the presence of variability that occurs in all of the research ($p < 0.05$; Table 4). Then the assumption of homogeneity needs to be rejected and accept the assumption of heterogeneity, namely the variability that occurs not more caused by sampling error. It was also evidenced by the high percentage of I² (inconsistency) that is equal to 84.10% (Table 4). Thus it can be concluded that the addition of propolis extract is effective in




Fig. 3. Forest plot of microbial analysis at various concentrations of PE (propolis extract) on various livestock products using the Random Effect model.

inhibiting the growth of microbial pathogens in livestock products.

Further evaluation of the validation of the effectiveness of propolis extract in inhibiting the growth of pathogenic microbes necessary validation test against the bias of the publication using the test provided by Kendall's test. The results of the analysis show that the p-value on the method (rank correlation test for Funnel plot Asymmetry) is symmetrical or in other words did not happen or finding evidence of a bias publication ($p > 0.05$). The rank correlation value is negative (-0.179) in Table 4 indicate that research with a large sample is not included in the sample of the study meta-analysis, more dominant research with a small sample size. So the conclusion made based on the random-effect model about the effectiveness of the extract of propolis in its application as a natural preservative in products of livestock is valid is indicated by the inconsistency is high and the bias publication of the resulting low through the test provided by Kendall's test ($p > 0.05$; Table 4).

Propolis in terms of its use has been quite promising as a natural preservative in various food products, such as juices, fruit, and vegetables even in various livestock products (meat and milk) due to its antimicrobial and antioxidant properties (Bankova et al., 2016). The antimicrobial effect of propolis has been extensively studied, as it was reported that the use of propolis as a preservative was able to inhibit mesophilic and psychotropic bacteria in beef patties (Vargas-Sánchez et al., 2014), inhibit micrococcaceae, molds, and yeasts on sausage surfaces (Ozturk, 2015), inhibit the activity of pathogenic microbes (Gutiérrez-Cortés and Suarez Mahecha, 2014), inhibit *L. monocytogenes* (Thamnopoulos et al., 2018), inhibit *S. aureus* (El-Bassiony et al., 2012), and increase the shelf life of yogurt (Özer, 2020). In this case, propolis can be used as an enrichment in food products, both as a natural additive, improving food quality, and as a natural preservative (Pobiega et al., 2019b; Seibert et al., 2019).

Thiobarbituric acid reactive substances (TBARs) value

The number of study data used in the meta-analysis of the difference in TBARs values in products with the addition of

various concentrations of propolis extract was 22 with the range from 2010 to 2021 (Fig. 4). The total number of samples/replicates on the TBARs value of the product with the addition of propolis extract was 132. Based on the results of the calculation with a random-effect model value obtained SMD/d+ overall by -1.62 with CI95% ($-2.27, -0.98$; Table 4), because the confidence interval does not contain 0 (zero), then the treatment given to the experimental group different from the control group in terms of inhibiting the oxidation of lipids on the farm with the addition of propolis extract as a natural preservative. These values indicate that the TBARs value with the addition of propolis extract on various products lower than the control treatment. It is shown through the summary effect (the difference value is negative). The results of the analysis showed (Table 4), then in this case the true effect size is not equal to 0. It indicates there is a significant relationship between the addition of propolis extract with the TBARs value produced on products ($p < 0.05$). However, before drawing any conclusions based on the random-effect model this would be accurate if it is proved that all the research produces true effect size differences in the population so that the need for the test of heterogeneity. The analysis of heterogeneity on the value of TBARs product indicates the presence of variability that occurs in all of the research ($p < 0.05$; Table 4). Then the assumption of homogeneity needs to be rejected and accept the assumption of heterogeneity, namely the variability that occurs not more caused by sampling error. This was evidenced by the high percentage of I² (inconsistency) that is equal to 99.32% (Table 4). Thus it can be concluded that the addition of propolis extract is effective in inhibiting the oxidation reaction on the product as indicated by TBARs value.

Further evaluation of the validation of the effectiveness of propolis extract in inhibiting the oxidation reaction required a validation test against the bias of the publication using the test provided by Kendall's test. The results of the analysis show that the p-value on the method (rank correlation test for Funnel plot Asymmetry) is symmetrical or in other words did not happen or finding evidence of a bias publication ($p > 0.05$). The positive rank correlation (0.048) in Table 4 indicates that research with a large sample is included in the study sample meta-analysis. So the conclusion made based on the random-




Fig. 4. Forest plot analysis of TBARs values at various concentrations of PE (propolis extract) on various livestock products using the Random Effect model. TBARs, thiobarbituric acid reactive substances.

effect model about the effectiveness of the extract of propolis in its application as a natural preservative to inhibit the oxidation reaction on the product is the result of breeding is valid is indicated by the inconsistency is high and the bias publication of the resulting low through the test provided by Kendall's test ($p > 0.05$; Table 4).

Fat oxidation is one of the main causes of food spoilage. This is indicated by the resulting TBARs value. Several cases reported that propolis can inhibit the reduction of fat oxidation in sausage products (Ali et al., 2010), beef patties (Vargas-Sánchez et al., 2014), and research articles used in this study. Preliminary research also reported that giving propolis extract to beef products stored at room temperature for 24 hours was able to inhibit fat oxidation, the greater the concentration of propolis extract added, the smaller the TBARs value produced (Andre et al., 2021).

Active compounds and mechanism to inhibit of bacteria

The chemical composition of propolis is composed of resin (flavonoids and phenolic compounds) by 42% to 58%, candles and oil (oleic acid and palmitic acid fiber of essential oils and aromatic) by 33% to 47%, polen (protein, free amino acids, vitamins, and minerals) of 3% to 5%, and other components (ketones, lactones, steroids, and sugars) by 2% to 5% (Barlak, 2009; Burdock, 1998; Değirmencioğlu, 2018).

The difference of opinion related to the mechanism of action of flavonoids in inhibiting the growth of bacteria. Flavonoids cause damage to the permeability of the bacterial cell wall, microsomes, and lysosomes as a result of the interaction between flavonoids with DNA of bacteria (Bryan, 1982; Wilson, 1982). Flavonoids are able to release energy tranduksi against the cytoplasmic membrane of bacteria it also inhibits the motility of bacteria (Mirzoeva et al., 1997). A different mechanism is also reported that the hydroxyl groups contained in the structure of the flavonoid compounds cause changes in the organic component and the transport of nutrients which will eventually lead to the onset of toxic effects against bacteria (Di Carlo et al., 1999; Estrela et al., 1995). Although the mechanism of the detail of the antibacterial activity of propolis is still unknown (Santos et al., 2002), the possibility it is related to the compound polar and phenolic lipophilic namely flavonoid compounds. The compound has a carbonyl electronegative, amen, imina, sulfid, thiol, metoksil, and hydroxyl groups are very polar and lipophilic, and is responsible for contact with the bacterial cells and induce damage to the structure of the cell wall and membrane, causing cell lysis and death (Cushnie and Lamb, 2005; Cushnie et al., 2003; Echeverría et al., 2017; Kim and Chung, 2011; Sanpa et al., 2015).

Conclusion

The effectiveness of propolis extract as a natural preservative in products of livestock indicates the presence of a significant relationship between the addition of propolis extract at various concentrations of the antimicrobial activity as well as test microbes and the value of TBARs in its application in a variety of storage. Based on the random-effect model on the effectiveness of antimicrobial activity of propolis extract and its application as a natural preservative against the characteristics of the chemical and microbiological analysis on products of livestock is valid and not the discovery of bias publication that is produced through the test provided by kendall's. Propolis in this case effectively used as natural preservatives in the products of livestock.

Conflicts of Interest

The authors declare no potential conflicts of interest.

Author Contributions

Conceptualization: Andre. Data curation: Andre, Jayanegara A. Formal analysis: Andre, Jayanegara A. Methodology: Andre, Jayanegara A. Software: Jayanegara A. Validation: Arief II, Apriantini A, Jayanegara A. Investigation: Andre. Writing - original draft: Andre. Writing - review & editing: Andre, Arief II, Apriantini A, Jayanegara A, Budiman C.

Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

References

- Abdullah NA, Ja'afar F, Yasin HM, Taha H, Petalcorin MIR, Mamit MH, Kusriani E, Usman A. 2019. Physicochemical analyses, antioxidant, antibacterial, and toxicity of propolis particles produced by stingless bee *Heterotrigona itama* found in Brunei Darussalam. *Heliyon* 5:e02476.
- Abdullah NA, Zulkiflee N, Zaini SNZ, Taha H, Hashim F, Usman A. 2020. Phytochemicals, mineral contents, antioxidants, and antimicrobial activities of propolis produced by Brunei stingless bees *Geniotrigona thoracica*, *Heterotrigona itama*, and *Tetrigona binghami*. *Saudi J Biol Sci* 27:2902-2911.
- Aga H, Shibuya T, Sugimoto T, Kurimoto M, Nakajima S. 1994. Isolation and identification of antimicrobial compounds in Brazilian propolis. *Biosci Biotechnol Biochem* 58:945-946.
- Airen B, Sarkar PA, Tomar U, Bishen KA. 2018. Antibacterial effect of propolis derived from tribal region on *Streptococcus mutans* and *Lactobacillus acidophilus*: An *in vitro* study. *J Indian Soc Pedod Prev Dent* 36:48-52.
- Ali FH, Kassem GM, Atta-Alla OA. 2010. Propolis as a natural decontaminant and antioxidant in fresh oriental sausage. *Vet Ital* 46:167-172.
- Andre, Apriantini A, Budiman C. 2021. Pengaruh ekstrak propolis sebagai edible coating terhadap karakteristik kimia dan aktifitas antioksidan daging sapi pada penyimpanan suhu ruang. *J Ilmu Produksi Dan Teknologi Hasil Peternakan* 9:72-78.
- Anwar R. 2005. Meta-analisis. Universitas Padjajaran Press, West Java, Indonesia.
- Azemin A, Md-Zin NB, Mohd-Rodi MM, Kim-Chee AS, Zakaria AJ, Mohd KS. 2018. Application of metabolite profiling and antioxidant activity in assessing the quality of processed and unprocessed stingless bee's propolis. *J Fundam Appl Sci* 9:637-660.
- Bankova V. 2005. Chemical diversity of propolis and the problem of standardization. *J Ethnopharmacol* 100:114-117.
- Bankova V, Galabov AS, Antonova D, Vilhelmova N, Di Perri B. 2014. Chemical composition of propolis extract ACF® and activity against herpes simplex virus. *Phytomedicine* 21:1432-1438.
- Bankova V, Popova M, Trusheva B. 2016. New emerging fields of application of propolis. *Maced J Chem Chem Eng* 35:1-11.
- Barlak Y. 2009. Türk propolisiekstraktlarının prostat kanser hücre serilerinin proteomisine etkisi. Ph.D. dissertation, Karadeniz Technical Univ., Trabzon, Turkey.
- Berretta AA, Arruda C, Miguel FG, Baptista N, Piacezzi Nascimento A, Marquele-Oliveira F, Hori JI, da Silva Barud H, Damaso B, Ramos C, Ferreira R, Bastos JK. 2017. Functional properties of Brazilian propolis: From chemical composition until the market. In *Superfood and functional food: An overview of their processing and utilization*.

- Waisundara VY, Shiomi N (ed). IntechOpen, London, UK. pp 55-98.
- Bryan LE. 1982. Bacterial resistance and susceptibility to chemotherapeutic agents. Cambridge University Press, Cambridge, UK.
- Burdock GA. 1998. Review of the biological properties and toxicity of bee propolis (propolis). *Food Chem Toxicol* 36:347-363.
- Coró FAG, Gaino VO, Carneiro J, Coelho AR, Pedrão MR. 2020. Control of lipid oxidation in jerked beef through the replacement of sodium nitrite by natural extracts of yerba mate and propolis as antioxidant agent. *Braz J Dev* 6:4834-4850.
- Corrêa FRS, Schanuel FS, Moura-Nunes N, Monte-Alto-Costa A, Daleprane JB. 2017. Brazilian red propolis improves cutaneous wound healing suppressing inflammation-associated transcription factor NF κ B. *Biomed Pharmacother* 86:162-171.
- Cushnie TPT, Hamilton VES, Lamb AJ. 2003. Assessment of the antibacterial activity of selected flavonoids and consideration of discrepancies between previous reports. *Microbial Res* 158:281-289.
- Cushnie TPT, Lamb AJ. 2005. Antimicrobial activity of flavonoids. *Int J Antimicrob Agents* 26:343-356.
- das Neves MVM, da Silva TMS, de Oliveira Lima E, da Cunha EVL, Oliveira EJ. 2016. Isoflavone formononetin from red propolis acts as a fungicide against *Candida* sp. *Braz J Microbiol* 47:159-166.
- Değirmencioglu HT. 2018. Evaluation of phenolic profile, botanical origin, antioxidant and antimicrobial activities of Turkish propolis. M.S. thesis, Yeditepe Univ., İstanbul, Turkey.
- Di Carlo G, Mascolo N, Izzo AA, Capasso F. 1999. Flavonoids: Old and new aspects of a class of natural therapeutic drugs. *Life Sci* 65:337-353.
- Donker AS, de Boer H, Kostons D, Dignath van Ewijk CC, van der Werf MPC. 2014. Effectiveness of learning strategy instruction on academic performance: A meta-analysis. *Educ Res Rev* 11:1-26.
- dos Reis AS, Diedrich C, de Moura C, Pereira D, de Florio Almeida J, da Silva LD, Oviedo MSVP, Tavares RAW, Carpes ST. 2017. Physico-chemical characteristics of microencapsulated propolis co-product extract and its effect on storage stability of burger meat during storage at -15°C . *LWT-Food Sci Technol* 76:306-313.
- Echeverría J, Opazo J, Mendoza L, Urzúa A, Wilkens M. 2017. Structure-activity and lipophilicity relationships of selected antibacterial natural flavones and flavanones of Chilean flora. *Molecules* 22:608.
- El-Bassiony TA, Saad NM, El-Zamkan MA. 2012. Study on the antimicrobial activity of ethanol extract of propolis against enterotoxigenic methicillin-resistant *Staphylococcus aureus* in lab prepared ice-cream. *Vet World* 5:155-159.
- El-Mossalami HHA, Abdel-Hakeim YA. 2013. Using of propolis extract as a trial to extend the shelf-life and improving the quality criteria of fresh Egyptian sausage. *Assiut Vet Med J* 59:23-33.
- Estrela C, Sydney GB, Bammann LL, Felipe O Jr. 1995. Mechanism of action of calcium and hydroxyl ions of calcium hydroxide on tissue and bacteria. *Braz Dent J* 6:85-90.
- European Food Safety Authority [EFSA]. 2010. Scientific opinion on the substantiation of health claims related to propolis (ID 1242, 1245, 1246, 1247, 1248, 3184) and flavonoids in propolis (ID 1244, 1644, 1645, 3526, 3527, 3798, 3799) pursuant to Article 13(1) of Regulation (EC) No 1924/2006. *EFSA J* 8:1810.
- European Parliament and Council of the European Union. 2002. Directive 2002/46/EC of the European Parliament and of the council of 10 June 2002 on the approximation of the laws of the Member States relating to food supplements (Text with EEA relevance). *Off J Eur Communities Legis* 183:51-57.

- Farnesi AP, Aquino-Ferreira R, De Jong D, Bastos JK, Soares AEE. 2009. Effects of stingless bee and honey bee propolis on four species of bacteria. *Genet Mol Res* 8:635-640.
- Farooqui T, Farooqui AA. 2012. Beneficial effects of propolis on human health and neurological diseases. *Front Biosci* 4:779-793.
- Gutiérrez-Cortés C, Suarez Mahecha H. 2014. Antimicrobial activity of propolis and its effect on the physicochemical and sensorial characteristics in sausages. *Vitae* 21:90-96.
- Kharsany K, Viljoen A, Leonard C, van Vuuren S. 2019. The new buzz: Investigating the antimicrobial interactions between bioactive compounds found in South African propolis. *J Ethnopharmacol* 238:111867.
- Khodayari M, Basti AA, Khanjari A, Misaghi A, Kamkar A, Shotorbani PM, Hamed H. 2019. Effect of poly (lactic acid) films incorporated with different concentrations of *Tanacetum balsamita* essential oil, propolis ethanolic extract and cellulose nanocrystals on shelf life extension of vacuum-packed cooked sausages. *Food Packag Shelf Life* 19:200-209.
- Kim YH, Chung HJ. 2011. The effects of Korean propolis against foodborne pathogens and transmission electron microscopic examination. *N Biotechnol* 28:713-718.
- Kisa Ç, Karagöz E, Cici G, Köker Ö, Kiliç B, Şimşek A, Bilecen D, Soyucok A. 2018. Effects of pomegranate peel and propolis powders and their combinations on physico-chemical and microbiological properties of Turkish dry-fermented sausage (Sucuk) with various nitrite levels. *Sci Pap D Anim Sci* 61:275-282.
- Kismet K, Ozcan C, Kuru S, Gencay Celemlı O, Celepli P, Senes M, Guclu T, Sorkun K, Hucumenoglu S, Besler T. 2017. Does propolis have any effect on non-alcoholic fatty liver disease? *Biomed Pharmacother* 90:863-871.
- Kujungiev A, Tsvetkova I, Serkedjieva Y, Bankova V, Christov R, Popov S. 1999. Antibacterial, antifungal and antiviral activity of propolis of different geographic origin. *J Ethnopharmacol* 64:235-240.
- Kunrath CA, Savoldi DC, Mileski JPF, Novello CR, da Trindade Alfaro A, Marchi JF, Tonial IB. 2017. Application and evaluation of propolis, the natural antioxidant in Italian-type salami. *Braz J Food Technol* 20:e2016035.
- López BGC, Schmidt EM, Eberlin MN, Sawaya ACHF. 2014. Phytochemical markers of different types of red propolis. *Food Chem* 146:174-180.
- Markiewicz-Żukowska R, Borawska MH, Fiedorowicz A, Naliwajko SK, Sawicka D, Car H. 2013. Propolis changes the anticancer activity of temozolomide in U87MG human glioblastoma cell line. *BMC Complement Altern Med* 13:50.
- Mehdizadeh T, Mojaddar Langroodi A. 2019. Chitosan coatings incorporated with propolis extract and *Zataria multiflora* Boiss oil for active packaging of chicken breast meat. *Int J Biol Macromol* 141:401-409.
- Mello BCBS, Hubinger MD. 2012. Antioxidant activity and polyphenol contents in Brazilian green propolis extracts prepared with the use of ethanol and water as solvents in different pH values. *Int J Food Sci Technol* 47:2510-2518.
- Milojković Opsenica D, Ristivojević P, Trifković J, Vovk I, Lušić D, Tešić Ž. 2016. TLC fingerprinting and pattern recognition methods in the assessment of authenticity of poplar-type propolis. *J Chromatogr Sci* 54:1077-1083.
- Mirzoeva OK, Grishanin RN, Calder PC. 1997. Antimicrobial action of propolis and some of its components: The effects on growth, membrane potential and motility of bacteria. *Microbiol Res* 152:239-246.
- Nassar SA, Mohamed AH, Soufy H, Nasr SM, Mahran KM. 2012. Immunostimulant effect of Egyptian propolis in rabbits. *Sci World J* 2012:901516.
- Nolkemper S, Reichling J, Sensch KH, Schnitzler P. 2010. Mechanism of herpes simplex virus type 2 suppression by propolis extracts. *Phytomedicine* 17:132-138.
- Okińczyc P, Paluch E, Franiczek R, Widelski J, Wojtanowski KK, Mroczek T, Krzyżanowska B, Skalicka-Woźniak K, Sroka

- Z. 2020. Antimicrobial activity of *Apis mellifera* L. and *Trigona* sp. propolis from Nepal and its phytochemical analysis. *Biomed Pharmacother* 129:110435.
- Oliveira AV, Ferreira AL, Nunes S, Dandlen SA, Miguel MDG, Faleiro ML. 2017. Antibacterial activity of propolis extracts from the south of Portugal. *Pak J Pharm Sci* 30:1-9.
- Özer ED. 2020. Propolis and potential use in food products. *Turk J Agric Food Sci Technol* 8:1139-1144.
- Ozturk I. 2015. Antifungal activity of propolis, thyme essential oil and hydrosol on natural mycobiota of sucuk, a Turkish fermented sausage: Monitoring of their effects on microbiological, color and aroma properties. *J Food Process Preserv* 39:1148-1158.
- Pedonese F, Verani G, Torracca B, Turchi B, Felicioli A, Nuvoloni R. 2019. Effect of an Italian propolis on the growth of *Listeria monocytogenes*, *Staphylococcus aureus* and *Bacillus cereus* in milk and whey cheese. *Ital J Food Saf* 8:8036.
- Pobiega K, Kraśniewska K, Gniewosz M. 2019a. Application of propolis in antimicrobial and antioxidative protection of food quality: A review. *Trends Food Sci Technol* 83:53-62.
- Pobiega K, Kraśniewska K, Przybył JL, Bączek K, Żubernik J, Witrowa-Rajchert D, Gniewosz M. 2019b. Growth biocontrol of foodborne pathogens and spoilage microorganisms of food by Polish propolis extracts. *Molecules* 24:2965.
- Rezaeigolestani M, Misaghi A, Khanjari A, Basti AA, Abdulkhani A, Fayazfar S. 2017. Antimicrobial evaluation of novel poly-lactic acid based nanocomposites incorporated with bioactive compounds *in-vitro* and in refrigerated vacuum-packed cooked sausages. *Int J Food Microbiol* 260:1-10.
- Rocha BA, Rodrigues MR, Bueno PCP, de Mello Costa-Machado AR, de Oliveira Lima Leite Vaz MM, Nascimento AP, Barud HS, Berretta-Silva AA. 2012. Preparation and thermal characterization of inclusion complex of Brazilian green propolis and hydroxypropyl- β -cyclodextrin: Increased water solubility of the chemical constituents and antioxidant activity. *J Therm Anal Calorim* 108:87-94.
- Sanpa S, Popova M, Bankova V, Tunkasiri T, Eitssayeam S, Chantawannakul P. 2015. Antibacterial compounds from propolis of *Tetragonula laeviceps* and *Tetrigona melanoleuca* (Hymenoptera:Apidae) from Thailand. *PLOS ONE* 10:e0126886.
- Santos FA, Bastos EMA, Uzeda M, Carvalho MAR, Farias LM, Moreira ESA, Braga FC. 2002. Antibacterial activity of Brazilian propolis and fractions against oral anaerobic bacteria. *J Ethnopharmacol* 80:1-7.
- Seibert JB, Bautista-Silva JP, Amparo TR, Petit A, Pervier P, dos Santos Almeida JC, Azevedo MC, Silveira BM, Brandão GC, de Souza GHB, de Medeiros Teixeira LF, dos Santos ODH. 2019. Development of propolis nanoemulsion with antioxidant and antimicrobial activity for use as a potential natural preservative. *Food Chem* 287:61-67.
- Sun C, Wu Z, Wang Z, Zhang H. 2015. Effect of ethanol/water solvents on phenolic profiles and antioxidant properties of Beijing propolis extracts. *Evid Based Complement Alternat Med* 2015:595393.
- Thamnopoulos IAI, Michailidis GF, Fletouris DJ, Badeka A, Kontominas MG, Angelidis AS. 2018. Inhibitory activity of propolis against *Listeria monocytogenes* in milk stored under refrigeration. *Food Microbiol* 73:168-176.
- Tosi EA, Ré E, Ortega ME, Cazzoli AF. 2007. Food preservative based on propolis: Bacteriostatic activity of propolis polyphenols and flavonoids upon *Escherichia coli*. *Food Chem* 104:1025-1029.
- Utama GL, Mahani M, Djali M. 2021. Skill and knowledge improvement training on honey and propolis commercialization in Central Bangka Regency. *J Pengabdian Kepada Masyarakat* 5:59-73.
- Vargas-Sánchez RD, Acedo-Félix E, Pérez-Morales R, Sánchez-Escalante A, Torrescano-Urrutia GR. 2019. Propolis ethanolic extract against *S. aureus* growth in beef patties. *J Food Process Preserv* 2:8-9.
- Vargas-Sánchez RD, Torrescano-Urrutia GR, Acedo-Félix E, Carvajal-Millán E, González-Córdova AF, Vallejo-Galland B,

- Torres-Llanaez MJ, Sánchez-Escalante A. 2014. Antioxidant and antimicrobial activity of commercial propolis extract in beef patties. *J Food Sci* 79:C1499-C1504.
- Viera VB, Piovesan N, Moro KIB, Rodrigues AS, Scapin G, da Rosa CS, Kubota EH. 2016. Preparation and microbiological analysis of Tuscan sausage with added propolis extract. *Food Sci Technol* 36:37-41.
- Wilson G. 1982. *Kimia farmasi dan medisinal organik*. 8th ed. Dirjen Dikti dan Kebudayaan, Jakarta, Indonesia. pp 10-12.
- Xuan H, Li Z, Yan H, Sang Q, Wang K, He Q, Wang Y, Hu F. 2014. Antitumor activity of Chinese propolis in human breast cancer MCF-7 and MDA-MB-231 cells. *Evid Based Complement Alternat Med* 2014:280120.
- Yasar M, Savranlar Y, Karaman H, Sagit M, Silici S, Ozcan I. 2016. Effects of propolis in an experimental rat model of allergic rhinitis. *Am J Otolaryngol* 37:287-293.