




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Case Reports and Series

Higher mortality of hospitalized haematologic patients with COVID-19 compared to non-haematologic is driven by thrombotic complications and development of ARDS: An age-matched cohorts study[☆]

Ana Fernández-Cruz ^{a,g,*}, Alba Puyuelo ^b, Lucía Núñez Martín-Buitrago ^b, Enrique Sánchez-Chica ^c, Carmen Díaz-Pedroche ^{d,h,j}, Rosa Ayala ^{e,h}, Manuel Lizasoain ^{f,h}, Rafael Duarte ^{b,g,i}, Carlos Lumbreiras ^{f,h,j}, Juan Antonio Vargas ^{c,g,i}

^a Infectious Diseases Department, Hospital Puerta de Hierro-Majadahonda, Madrid, Spain

^b Haematology Department, Hospital Puerta de Hierro-Majadahonda, Madrid, Spain

^c Internal Medicine Department, Hospital Puerta de Hierro-Majadahonda, Madrid, Spain

^d Internal Medicine Department, Hospital Doce de Octubre, Madrid, Spain

^e Haematology Department, Hospital Doce de Octubre, Madrid, Spain

^f Infectious Diseases Department, Hospital Doce de Octubre, Madrid, Spain

^g Fundación de Investigación Sanitaria Puerta de Hierro - Segovia de Arana, Spain

^h Fundación para la Investigación Biomédica del Hospital Universitario, 12 de Octubre, Spain

ⁱ Universidad Autónoma de Madrid, Spain

^j Universidad Complutense de Madrid, Spain

ARTICLE INFO

ABSTRACT

Keywords
COVID-19
Haematologic
Outcome
Mortality

Background and Objectives: The characteristics of COVID-19 in haematologic patients compared to non-haematologic patients have seldom been analyzed. Our aim was to analyze whether there are differences in clinical characteristics and outcome of haematologic patients with COVID-19 as compared to non-haematologic.

Patients and methods: Retrospective cohort study in 2 University hospitals of patients admitted with laboratory-confirmed COVID-19 included in the SEMICOVID19 database. The cohort with underlying haematologic disease was compared to a cohort of age and date-of-COVID-19-matched controls without haematologic disease (1:2).

Results: 71 cases and 142 controls were included from March-May 2020.

Twenty (28.1%) had received recent chemotherapy. Twelve (16.9%) were stem cell transplant recipients (SCT). Eleven (15.5%) were neutropenic concurrently with COVID-19 diagnosis.

Haematologic patients presented ARDS (58.5 vs 20.7%, p = 0.0001), thrombotic complications (15.7 vs 2.1%, p = 0.002), DIC (5.7 vs 0.0%, p = 0.011), heart failure (14.3 vs 4.9%, p = 0.029) and required ICU admission (15.5 vs 2.8%, p = 0.001), MV (14.1% vs 2.1%, p = 0.001), steroid (64.8 vs 33.1%, p = 0.0001), tocilizumab (33.8 vs 8.5%, p = 0.0001) or anakinra treatment (9.9% vs 0%, p = 0.0001) more often. In-hospital mortality was significantly higher (38.0% vs 18.3%, p = 0.002).

Conclusions: Our results suggest COVID-19 has worse outcomes in haematologic patients than in non-haematologic, independently of age, and that the development of ARDS and thrombotic complications drive the higher in-hospital mortality.

Abbreviations: ARDS, acute respiratory distress syndrome; C-RP, C-reactive protein; CI, confidence interval; CLL, chronic lymphocytic leukemia; COVID-19, Coronavirus disease 2019; DIC, disseminated intravascular coagulation; ECOG scale, Eastern Cooperative Oncology Group scale; G-CSF, granulocyte stimulating factor; HFNC, high flow nasal cannula; ICU, Intensive Care Unit; IL6, Interleukin 6; IQR, interquartile range; LDH, Lactate dehydrogenase; MDS, myelodysplastic syndrome; MM, multiple myeloma; MV, mechanical ventilation; NHL, non-Hodgkin lymphoma; NIMV, non-invasive mechanical ventilation; OR, Odds ratio; PaO2/FiO2, arterial oxygen tension/inspiratory oxygen fraction; PEEP, positive end expiratory pressure; SARS-CoV-2, Severe acute respiratory syndrome coronavirus 2; SCT, stem cell transplantation; SD, standard deviation.

* This study was previously presented at the 31st ECCMID as an e-Poster (Session: 12c. Clinical features, case management, outcome) 9–12 July online.

* Corresponding autor at: Unidad de Enfermedades Infecciosas, Servicio de Medicina Interna, Hospital Universitario Puerta de Hierro, Instituto de Investigación Sanitaria Puerta de Hierro - Segovia de Arana, C/ Joaquín Rodrigo n° 2, 28222 Majadahonda, Madrid, Spain.

E-mail address: afcruz@salud.madrid.org (A. Fernández-Cruz).

Introduction

Haematologic patients present a high risk for infection, due to immune-compromise secondary to underlying disease and subsequent therapy. Viral infections such as RSV or Influenza that are considered mild in immunocompetent hosts can become life-threatening in certain haematologic patients (Kmeid et al., 2016; Sheshadri et al., 2019).

The characteristics of SARS-CoV-2 infection and COVID-19 in haematologic patients are not yet well known. In the context of COVID-19 pandemic, the underlying haematologic disease could influence the inflammatory response and viral clearance, and modify manifestations and outcome of the disease (Chamilos et al., 2021).

Studies published so far suggest haematologic patients with COVID-19 present a higher mortality as compared with general population data (García-Suárez et al., 2020). However, the characteristics of COVID-19 in haematologic patients as directly compared to non-haematologic patients have seldom been analyzed. There is a lack of information regarding differences in clinical presentation, incidence of different complications and management of patients with haematologic malignancy and COVID-19, compared to non-haematologic cases. The scarce prior published series that compare to the general population (Passamonti et al., 2020) present mainly population-based data and lack detailed information of cases.

We present a cohort of haematologic patients with COVID-19, and compare them to non-haematologic patients with COVID-19.

Material/Patients and methods

Setting and study design

We performed a retrospective cohort study in 2 University hospitals in Madrid, Spain, of admitted patients with SARS-CoV-2 laboratory-confirmed pneumonia included in the SEMICOVID19 Registry (compiled by the Spanish Society of Internal Medicine) from March to May 2020. Both centres are tertiary teaching hospitals, with reference Haematology Departments, that possess stem cell transplantation units and treat complex Haematology patients.

The SEMICOVID19 is an ongoing, nationwide multicentre anonymized online database of consecutive adult patients admitted with SARS-CoV-2 laboratory-confirmed pneumonia from 131 different Spanish hospitals. Inclusion criteria for the registry were age ≥ 18 years and first hospital discharge with a confirmed diagnosis of COVID-19; exclusion criteria were subsequent admissions of the same patient and denial or withdrawal of informed consent, as described elsewhere (Casas-Rojo et al., 2020). Patients were cared for according to local protocols and clinical judgment of their attending physician.

For the present study, only 2 of these hospitals were selected that had included all their haematologic hospitalized patients with COVID-19 in the Registry database. A retrospective cohort study was designed to compare the differences between patients with underlying haematologic disease and patients without underlying haematologic disease. All patients with underlying haematological disease were selected and two controls without haematologic disease were selected for each haematologic patient, matched by age and date of COVID-19. To ensure a standard process of choosing controls, an algorithm was used to select those of the same age among the possible controls diagnosed at the nearest date of COVID diagnosis.

Data collection

The SEMICOVID Registry includes epidemiological, clinical, laboratory and radiologic data extracted from electronic medical records. For more comprehensive information on the registry, see previously published works (Casas-Rojo et al., 2020).

A complementary standardized form was fulfilled for haematologic patients that included specific data about haematologic disease:

underlying haematologic disease, ECOG, status, therapy, stem cell transplantation.

Definitions

We considered SARS-CoV-2 infected patients those with a microbiological confirmation by reverse transcription polymerase chain reaction (RT-PCR) testing of a respiratory sample. All patients admitted with symptomatic COVID-19 infection were included, with or without pneumonia.

We included in the "Haematologic disease cohort" patients (or "Haematologic patients") admitted with SARS-CoV-2 infection who had an underlying active haematological malignancy, or were stem cell-transplantation (SCT) recipients (as treatment for haematological malignancy). Patients were considered as having active onco-hematologic disease when they were under treatment (chemotherapy, or targeted therapy) or being still immunocompromised due to their underlying hematological condition or treatment. The "Non-haematologic disease cohort" (or "Non-haematologic patients") included patients admitted with SARS-CoV-2 infection and without onco-haematologic disease or SCT. Patients with active solid tumours were excluded from both cohorts and will be analysed separately.

Disease status at the time of SARS-CoV-2 detection was defined according to each specific disease's revised criteria for leukemia, myeloproliferative neoplasm, multiple myeloma and lymphoma (Döhner et al., 2017; Cheson et al., 2014; Kumar et al., 2016).

Performance status at the diagnosis of COVID-19 was graded according to the Eastern Cooperative Oncology Group (ECOG) (Oken et al., 1982).

The diagnosis and grading of ARDS was determined according to modified Berlin criteria (Force et al., 2012) (in non-ventilated patients, the PEEP value in the modified criteria was not taken into consideration): Mild: PaO₂/FiO₂ 200–300 mmHg (PEEP or CPAP ≥ 5 cmH₂O, or non-ventilated); Moderate: PaO₂/FiO₂ 100–200 (PEEP ≥ 5 cmH₂O, or non-ventilated); Severe: PaO₂/FiO₂ ≤ 100 mmHg (PEEP ≥ 5 cmH₂O, or non-ventilated).

The main outcome variable was in-hospital mortality.

Statistical analysis

Quantitative variables were expressed as means and standard deviations (SD) and/or medians and interquartile ranges, and qualitative variables as frequencies and percentages.

To compare differences between haematologic and non-haematologic cohorts, the Mann-Whitney *U* test, χ^2 test, Fisher's exact test or Student *t* test were used where appropriate.

To explore risk factors associated with in-hospital death among haematologic patients, univariable and multivariable logistic regression models were used. Variables with a *p* < 0.05 in univariable analyses were selected into the multivariable.

All statistical analyses were performed using SPSS system (version 26.0 for Windows, SPSS Inc., Chicago, IL, USA). The statistical significance level was set at a two-sided *p* value of < 0.05. An odds ratio (OR) was reported along with 95% confidence interval (CI).

Results

From March to May 2020, 5592 patients with COVID-19 were admitted to the 2 hospitals. Among them, 71 (1.3%) cases had an underlying haematologic disease. One-hundred and forty-two patients with COVID-19 but without haematologic disease admitted to the hospital during the study period were selected as the control cohort.

Patients were followed until discharge, and if readmitted, until subsequent discharge (median follow-up of 7 days (IQR 4–14)).

Characteristics of underlying haematologic disease

Characteristics of haematologic diseases are summarized in Table 1 and Fig. 1. The most common was NHL, followed by MM, CLL and MDS. In 14.1% the haematologic disease was at an initial stage, whereas in 16.9% it was refractory or relapsed. Performance status measured by ECOG scale was > 1 in 12.9 %.

Twelve patients (16.9%) were stem cell transplantation recipients (SCT). In the majority of cases SCT was autologous (58.3% autologous versus 41.7% allogenic). Median time from SCT to COVID-19 diagnosis was 607 days (IQR 259–1291 days).

Forty-five cases (63.4%) were under oncologic therapy at COVID-19 diagnosis, which in 25 (55.6%) of cases was considered as palliative/symptomatic care.

Twenty patients (28.1%) had received chemotherapy in the prior month and 11 (15.5%) were neutropenic concurrently with COVID-19 diagnosis. Among these, 6 (54.5%, 8.5% of the whole series) received G-CSF. In seven cases (33.3%) chemotherapy was administered in combination with targeted therapy.

Thirty-two patients (45%) were receiving targeted therapy, including 15 (21.1%) under rituximab, 10 (14.1%) treated with ibrutinib (7) or other BTK inhibitors (3), 6 (8.5%) with bortezomib (4) or other proteasome inhibitors (2), 2 with imatinib, 2 with obinutuzumab (one of them combined with zanabrutinib) and 1 with ofatumumab (combined with ibrutinib), 1 with daratumumab (combined with carfilzomib) and 1 with ruxolitinib. Treatment with ibrutinib or other BTK inhibitors was maintained throughout the COVID episode in all cases.

Comparison of haematologic and non-haematologic patients with COVID-19

Differential characteristics of patients with and without haematologic disease are shown in Table 2 (baseline characteristics) and Table 3 (management and outcome characteristics).

Baseline significant differences included a higher age-adjusted Charlson comorbidity index (5.4 vs 3.4, p = 0.0001), creatinine (1.26 vs 0.95 mg/dl, p = 0.027), C-reactive protein (111 vs 79 mg/dl, p = 0.023) and D-dimer levels (2893 vs 1077 ng/mL, p = 0.047), increased prevalence of neutropenia (15.5% vs 1.4%, p = 0.0001) and anemia (hemoglobin 11.9 vs 14.2 g/dl, p = 0.0001) and more frequent receipt of prior anticoagulants (21.1% vs 7.7 %, p = 0.003) and prior steroids (18.3% vs 3.5%, p = 0.024) or immunosuppressive therapy (26.7% vs




Fig. 1. Type of haematological malignancy.

2.1%, p = 0.0001) in haematologic patients. Both cohorts had similar median age (70.7 vs 69.6 y, p = 0.734). Of note, there were no differences either in prevalence of baseline lymphopenia, thrombocytopenia or other alteration of coagulation parameters (different from D-dimer) between haematologic cases and controls, or in LDH, ferritin or IL6 levels.

Regarding outcome, haematologic patients presented ARDS (58.5 vs 20.7%, p = 0.0001), thrombotic complications (15.7 vs 2.1%, p = 0.002), disseminated intravascular coagulation (5.7 vs 0.0%, p = 0.011) and heart failure (14.3 vs 4.9%, p = 0.029) more often and required ICU admission (15.5 vs 2.8%, p = 0.001), mechanical ventilation (14.1% vs 2.1%, p = 0.001), and steroid (64.8 vs 33.1%, p = 0.0001), tocilizumab (33.8 vs 8.5%, p = 0.0001) or anakinra treatment (9.9% vs 0%, p = 0.0001) in a higher proportion than controls. Length of stay after COVID-19 diagnosis was longer for haematologic patients (18 days (SD17.1) versus 7 (7.4) days, p = 0.0001). Interestingly, haematologic patients did not present more infectious complications than non-haematologic. In 67 (31.5%) cases that had a follow-up SARS-CoV-2 PCR, there was a non-significant trend to a longer duration of positivity in haematologic patients (11 vs 4.5 days, p = 0.861).

In-hospital mortality was significantly higher among haematologic cases versus controls (38.0% vs 18.3%, p = 0.002). However, there were no significant differences in in-hospital mortality in patients who developed ARDS, or required ICU admission or ventilation, according to the presence of haematologic disease (Table 3).

Risk factors for in-hospital mortality among haematologic patients

Univariable analysis of factors associated with in-hospital mortality are displayed in Table 4.

When considering only haematologic patients, only development of ARDS (96.3 vs 36.4, p 0.0001) (OR 37.635 (4.583–309.060) p 0.001) was independently associated with a higher probability of in-hospital death in the multivariable analysis. Other factors such as recent chemotherapy, neutropenia, targeted therapies or uncontrolled haematologic disease were not predictors of in-hospital mortality in the multivariable analysis.

In-hospital mortality in recipients of SCT was similar to that of non-recipients (33.3% vs 39% (p = 0.999), however, there were no preengraftment cases of SCT.

Administration of G-CSF in haematologic patients was not associated with development of ARDS (p = 0.417).

Table 1
Underlying haematologic malignancy.

Patients with underlying hematologic disease	n = 71
Type of hematological malignancy	
Hodgkin lymphoma	1 (1.4%)
Non-Hodgkin lymphoma	21 (29.6%)
Chronic lymphocytic leukemia	13 (18.3%)
Acute lymphoblastic leukemia	1 (1.4%)
Acute myeloid leukemia	3 (4.2%)
Multiple myeloma	15 (21.1%)
Myelodysplastic syndrome	10 (14.1%)
Myeloproliferative neoplasm	6 (8.5%)
Rosai-Dorfman disease	1 (1.4%)
Hematological malignancy status	
Initial diagnosis	10 (14.1%)
Stable, no remission	16 (22.5%)
Relapsed or refractory	12 (16.9%)
Complete/partial response	33 (46.5%)
ECOG > 1	9 (12.9%)
Oncologic treatment	45 (63.4%)
Curative	20 (44.4%)
Palliative	25 (55.6%)
G-CSF treatment	6 (8.5%)
Stem cell transplant type	
Allogeneic	5 (7%)
Autologous	7 (9.9%)

Table 2
Baseline characteristics.

Demographic and Clinical Characteristics (n, %)	Haematological malignancy (71)	Non Haematological malignancy (142)	p
Demographic Characteristics			
Sex			0.378
Male	44 (61.9%)	77 (54.2%)	
Female	27 (38.0%)	64 (45.1%)	
Age (median-IQR)	70.7 (59.6–80.7)	69.6 (59.5–80.3)	0.734
(mean-SD)	68.6 (13.9)	68.4 (13.8)	0.915
Age-adjusted Charlson comorbidity index (mean-SD)	5.4 (2.6)	3.4 (2.4)	0.0001
Ethnicity			0.983
Caucasian	63 (88.7%)	124 (87.3%)	
Latin American	6 (8.5%)	13 (9.2%)	
Other	1 (1.4%)	2 (1.4%)	
COVID-19 close contact			0.088
No	58 (81.7%)	96 (67.6%)	
Yes	12 (17.1%)	39 (27.5%)	
Acquisition			0.174
Community-acquired	59 (83.1%)	129 (90.8%)	
Hospital-acquired	7 (9.9%)	6 (4.2%)	
Nursing home	5 (7.0%)	6 (4.2%)	
Comorbidities and prior therapies			
Enolism	2 (2.8%)	5 (3.5%)	0.999
Smoking status			0.133
No	45 (63.4%)	106 (74.6%)	
Ex-smoker	22 (31%)	27 (19%)	
Smoker	3 (4.2%)	4 (2.8%)	
Hypertension			0.246
Yes	35 (42.3%)	58 (40.8%)	
ACE inhibitors	11 (31.4%)	18 (31%)	0.673
ARBs	14 (40%)	29 (50%)	0.999
Dyslipidemia			0.223
Yes	29 (40.8%)	45 (31.7%)	
Statins	26 (89.7%)	43 (30.3%)	0.356
Atrial fibrillation	11 (15.5%)	11 (7.7%)	0.097
Obesity (Body Mass Index > 30)	14 (19.7%)	33 (23.2%)	0.721
Degenerative neurological disease	3 (4.2%)	11 (7.7%)	0.394
Chronic kidney disease			0.131
Moderate/Severe	7 (9.9%)	6 (4.2%)	
Dialysis			0.346
Hemodialysis	2 (2.8%)	3 (2.1%)	
Peritoneal dialysis	1 (1.4%)	0 (0.0%)	
Solid-organ transplantation	1 (1.4%)	1 (0.7%)	0.687
Liver	0 (0.0%)	1 (0.7%)	
Kidney			
Immunosuppression			0.0001
Azathioprine	0 (0.0%)	1 (0.7%)	
Methotrexate	0 (0.0%)	1 (0.7%)	
Cyclophosphamide	0 (0.0%)	2 (1.4%)	
Others	17 (24%)	1 (0.7%)	
Diabetes mellitus without end-organ damage	11 (15.5%)	23 (16.2%)	0.999
6 (8.5%)	6 (4.2%)	0.220	
Diabetes mellitus with end-organ damage	13 (18.3%)	25 (17.6%)	0.851
5 (7%)	13 (9.2%)	0.795	
Metformin	2 (2.8%)	0 (0.0%)	0.110
DPP-4 inhibitors	3 (4.2%)	5 (3.5%)	0.999
GLP-1 agonists	5 (7%)	5 (3.5%)	0.307
SGLT2 inhibitors			
Insulin			
Prior systemic corticosteroids	13 (18.3%)	5 (3.5%)	0.001
Hydroxychloroquine	1 (1.4%)	0 (0.0%)	0.335
Biological agents	7 (9.9%)	2 (1.4%)	0.007
Anticoagulants			0.003
VKAs	6 (8.5%)	3 (2.1%)	
DOACs	5 (7.0%)	8 (5.6%)	
LMWH	4 (5.6%)	0 (0.0%)	
	8 (11.3%)	6 (4.2%)	0.075

Table 2 (continued)

Demographic and Clinical Characteristics (n, %)	Haematological malignancy (71)	Non Haematological malignancy (142)	p
Acute myocardial infarction			
Angina	6 (8.5%)	7 (4.9%)	0.366
Acetylsalicylic acid	19 (26.8%)	25 (17.6%)	0.151
Heart failure	6 (8.5%)	8 (5.6%)	0.559
Chronic obstructive pulmonary disease	1 (1.4%)	9 (6.3%)	0.170
Asthma	3 (4.2%)	12 (8.5%)	0.395
Inhaled steroids	3 (4.2%)	10 (7.0%)	0.430
Cerebrovascular disease	2 (2.8%)	4 (2.8%)	0.999
Dementia	2 (2.8%)	11 (7.7%)	0.227
Peripheral vascular disease	1 (1.4%)	4 (2.8%)	0.667
Liver disease			
Mild	1 (1.4%)	8 (5.6%)	0.277
Moderate/Severe	1 (1.4%)	2 (1.4%)	0.999
Connective tissue disease	1 (1.4%)	4 (2.8%)	0.667
Gastroduodenal ulcer	1 (1.4%)	3 (2.1%)	0.999
HIV	1 (1.4%)	1 (0.7%)	0.999
Obstructive sleep apnea	5 (7.0%)	0 (0.0%)	0.536
Clinical findings at admission			
Cough			0.682
Dry	37 (52.1%)	82 (57.7%)	
Expectoration	14 (19.7%)	29 (20.4%)	
Arthromyalgia	15 (21.1%)	50 (35.2%)	0.040
Ageusia	1 (1.4%)	8 (5.6%)	0.277
Anosmia	2 (2.8%)	8 (5.6%)	0.503
Asthenia	37 (52.1%)	62 (43.7%)	0.381
Anorexia	14 (19.7%)	14 (9.9%)	0.057
Pharyngeal ache	11 (15.5%)	14 (9.9%)	0.264
Headache	9 (12.7%)	15 (10.6%)	0.652
Low grade fever	15 (21.1%)	24 (16.9%)	0.525
Fever	44 (62.0%)	99 (69.7%)	
Dyspnea	28 (39.4%)	81 (57.0%)	0.020
Diarrhea	14 (19.7%)	21 (14.8%)	0.433
Nausea	7 (9.9%)	15 (10.6%)	0.999
Vomit	3 (4.2%)	9 (6.3%)	0.755
Abdominal pain	4 (5.6%)	10 (7.0%)	0.778
Confusion	5 (7.0%)	13 (9.2%)	0.795
Tachypnea	17 (23.9%)	47 (33.1%)	0.201
Lung auscultation			
Crackling	44 (62%)	76 (53.5%)	0.362
Wheezing	0 (0.0%)	8 (5.6%)	0.053
Ronchi	2 (2.8%)	10 (7.0%)	0.345
Laboratory and radiology parameters			
Positive PCR sample			0.363
Nasopharyngeal	65 (91.5%)	133 (93.7%)	
Sputum	4 (5.6%)	8 (5.6%)	
Broncho-alveolar lavage	1 (1.4%)	0 (0.0%)	
Neutropenia			
Neutrophils < 500 (cells/µL)	5 (7.0%)	1 (0.7%)	0.009
Neutrophils < 1000 (cells/µL)	11 (15.5%)	2 (1.4%)	0.000
Blood cell counts (cells/µL) (mean-SD)			
Absolute white blood cell count	10,245 (14006)	7827 (8733)	0.187
Absolute lymphocyte count	4001 (9880)	1583 (5443)	0.058
Absolute neutrophil count	4509 (3750)	4948 (2361)	0.370
Hemoglobin (mean-SD)	11.9 (2.5)	14.2 (1.7)	0.0001
C-RP (mg/L) (mean-SD)	111 (105)	79 (79)	0.023
Serum creatinine (mg/dL) (mean-SD)	1.26 (1.07)	0.95 (0.62)	0.027
LDH (U/L) (mean-SD)	461 (665)	327 (126)	0.110
Serum ferritin (µg/L) (mean-SD)	1280 (1708)	1217 (1768)	0.875
IL-6 (pg/mL) (mean-SD)	184 (291)	79 (121)	0.152
D-dimer (ng/mL) (mean-SD)	2893 (6431)	1077 (1376)	0.047

(continued on next page)

Table 2 (continued)

Demographic and Clinical Characteristics (n, %)	Haematological malignancy (71)	Non Haematological malignancy (142)	p
Alveolar infiltrates			0.805
Unilateral	10 (14.1%)	26 (18.3%)	
Bilateral	27 (38.0%)	53 (37.3%)	
Ground-glass infiltrates			0.016
Unilateral	4 (5.6%)	16 (11.3%)	
Bilateral	36 (50.7%)	89 (62.7%)	
Pleural effusion			0.019
Unilateral	3 (4.2%)	1 (0.7%)	
Bilateral	5 (7.0%)	2 (1.4%)	
Influenza PCR test			0.029
Negative	6 (8.5%)	25 (17.6%)	
Positive	2 (2.8%)	0 (0.0%)	
Not performed	62 (87.3%)	112 (78.9%)	
Urinary antigens			0.114
Negative	19 (26.8%)	56 (39.4%)	
S. pneumoniae	1 (1.4%)	2 (1.4%)	
Not performed	49 (69%)	74 (52.1%)	

Discussion

Our results show that, despite immunosuppression, haematologic patients with COVID-19 present significantly more respiratory and thrombotic complications as compared to non-haematologic patients, and a higher in-hospital mortality.

Some of the cancer-associated factors that have been advocated to contribute to worse outcomes of SARS-CoV-2 infection (lymphopenia and lymphocyte dysfunction, hypercoagulability, immuno-metabolic deregulation related to myeloid cell dysfunction) converge in patients with haematological malignancy. On the other hand, chemotherapy-induced neutropenia and monocytopenia might attenuate the hyper-inflammatory response to the virus, whereas neutrophil recovery, treatment with G-CSF or immunotherapy could enhance it (Chamilos et al., 2021).

Several series attribute the increased mortality of haematologic patients to their higher age (García-Suárez et al., 2020). Age is an important prognostic factor in COVID-19 (Moreno-Torres et al., 2021). However, in our age-matched cohort, we still observed a significant difference in mortality between haematologic and non-haematologic cohorts. Nevertheless, patients with ARDS and patients requiring ICU had a similar mortality, regardless of the presence underlying haematologic disease. In the present series, both the development of ARDS and thrombotic complications were more frequent and could account for the increased mortality, in the haematologic cohort.

In COVID-19 patients, ARDS is driven by the inflammatory response to SARS-CoV-2, rather than by direct viral damage (Osuchowski et al., 2021). However, despite immunosuppression, haematologic patients in this series presented ARDS more often than non-haematologic. Haematologic patients with pneumonia are at risk of developing ARDS during neutropenia recovery (Rhee et al., 2009; Malek et al., 2021). It is a matter of controversy whether G-CSF could exacerbate the effect of neutrophil recovery contributing to ARDS (Rhee et al., 2009; Mignard et al., 2019). G-CSF upregulates the production of cytokines that increase alveolar permeability and neutrophil influx, and may enhance secretion of pro-inflammatory cytokines by alveolar macrophages (Rhee et al., 2009). In the present series, G-CSF was not a risk factor for ARDS, although it was administered to only a minority of patients.

Nevertheless, endothelitis and coagulopathy leading to in-situ thrombosis is increasingly gaining consideration in the pathogenesis of respiratory failure in COVID-19 (Bonaventura et al., 2021). Micro-thrombosis seems to be involved in the physiopathology of acute respiratory distress syndrome (ARDS) (Bonaventura et al., 2021; O'Donnell et al., 2021). The development of a pro-thrombotic state is an important feature of COVID-19. In the present series, thrombotic events were strikingly more frequent in haematologic patients. Cancer is a well-known risk factor for thrombosis and, in particular, patients with

Table 3

Management and outcome.

Complications and outcome	Haematological malignancy (71)	Non Haematological malignancy (142)	p
COMPLICATIONS (n, %)			
Heart failure	10 (14.3%)	7 (4.9%)	0.029
Cardiac arrhythmia	2 (2.8%)	3 (2.1%)	0.116
Atrial cardiac arrhythmia	2 (2.8%)	0 (0.0%)	
Ventricular cardiac arrhythmia	2 (2.8%)	0 (0.0%)	
Acute myocardial infarction	2 (2.8%)	0 (0.0%)	0.108
Myocarditis	3 (4.2%)	0 (0.0%)	0.035
Stroke			0.283
Ischemic stroke	0 (0.0%)	1 (0.7%)	
Hemorrhagic stroke	1 (1.4%)	0 (0.0%)	
Acute kidney failure	15 (21.1%)	19 (13.4%)	0.164
Sepsis	4 (5.7%)	2 (1.4%)	0.094
Bacterial pneumonia	3 (4.2%)	12 (8.5%)	0.395
Persistence of positive SARS-CoV-2 PCR (in 67 cases with available control PCR)			
(mean-SD)	16.8 (18.3)	10.1 (13.8)	0.303
(median-IQR)	11 (0.0–26.5)	4.5 (0.0–16.5)	0.861
Disseminated intravascular coagulation (DIC)	4 (5.7%)	0 (0.0%)	0.011
Thromboembolic disease	11 (15.5%)	3 (2.1%)	0.002
Deep vein thrombosis (DVT)	2 (2.8%)	1 (0.7%)	
Pulmonary thromboembolism (PE)	8 (11.3%)	2 (1.4%)	
DVT + PE	1 (1.4%)	0 (0.0%)	
Acute peripheral arterial disease	1 (1.4%)	1 (0.7%)	0.552
Acute respiratory distress syndrome (ARDS)	41 (58.5%)	29 (20.7%)	0.0001
Mild ARDS	6 (8.5%)	9 (6.3%)	
Moderate ARDS	10 (14.1%)	3 (2.1%)	
Severe ARDS	25 (35.3%)	17 (12%)	
Shock	3 (4.2%)	2 (1.4%)	0.335
Multiorgan failure	4 (5.6%)	4 (5.6%)	0.444
MANAGEMENT			
Systemic corticosteroids	46 (64.8%)	47 (33.1%)	0.0001
Systemic corticosteroids maximum daily dose (mg prednisone or equivalent) (mean-SD)	220 (214)	104 (70)	0.001
Days of treatment with systemic corticosteroids (mean-SD)	11.3 (9.6)	5.6 (5.6)	0.001
Cumulative dose of systemic corticosteroids during admission (mean-SD)	997 (1015)	319 (214)	0.002
Number of days with systemic corticosteroid pulses (>150 mg prednisone equivalent) (mean-SD)	2 (2.3)	0.5 (1.0)	0.000
Lopinavir/ritonavir	40 (56.3%)	94 (66.2%)	0.177
Beta Interferon	9 (12.7%)	34 (24%)	0.070
Remdesivir	1 (1.4%)	0 (0.0%)	0.340
Hydroxichloroquine	63 (44.4%)	135 (95.1%)	0.097
Colchicine	0 (0.0%)	1 (0.7%)	0.999
Tocilizumab	24 (33.8%)	12 (8.5%)	0.0001
Inmunoglobulins	1 (1.4%)	0 (0.0%)	0.330
Anakinra	7 (9.9%)	0 (0.0%)	0.0001
Baricitinib	0 (0.0%)	0 (0.0%)	NA
Intensive care unit (ICU) admission	11 (15.5%)	4 (2.8%)	0.001
Ventilation			
High flow nasal cannula	14 (19.7%)	19 (13.4%)	0.232
Non-invasive mechanical ventilation (NIMV)	7 (9.9%)	13 (9.2%)	0.999
Invasive mechanical ventilation (IMV)	10 (14.1%)	3 (2.1%)	0.001
Prone positioning therapy	12 (17%)	8 (5.6%)	0.012

(continued on next page)

Table 3 (continued)

Complications and outcome	Haematological malignancy (71)	Non Haematological malignancy (142)	p
Hospital mortality (overall)	27 (39%)	26 (18.3%)	0.002
-in patients who developed ARDS	24 (64.9%)	16 (57.1%)	0.610
-in patients who required ICU admission	7 (63.6%)	2 (50%)	0.999
-in patients who required MV	6 (60%)	1 (33.3%)	0.559
Outcome	5 (71.4%)	9 (69.2%)	0.999
Death during admission or readmission	30 (42.3%)	26 (18.3%)	0.002
Length of stay after COVID diagnosis (mean-SD)	18 (17.1)	7 (7.4)	0.0001
Re-admission	7 (9.9%)	5 (3.5%)	0.110
Reason for discharge	39 (55%)	114 (89.3%)	0.0001
Improvement: Home	5 (7%)	2 (1.4%)	
Convalescence: support centre	27 (38%)	26 (18.3%)	
Death			

active onco-haematologic conditions are known to be at higher risk for thromboembolism ([Kekre and Connors, 2019](#)). The baseline predisposition for thrombotic events seems to place haematologic patients more at risk for developing COVID-19 complications, both at the macro and at the microvascular level.

In immune-compromised patients there is a trend to longer persistence of viral shedding ([Taramasso et al., 2021](#)) that could contribute to a greater direct damage and mortality. Persistence of positive PCR could not be adequately evaluated in this retrospective series, as only 35% of cases had at least one control test after the diagnosis of COVID. Among those, there was a non-significant trend to a longer persistence of SARS-CoV-2 positivity.

Several studies report an inferior mortality of stem cell transplantation recipients as compared to other haematologic patients ([Piñana et al., 2020](#)). In published series, median time from SCT was, in general, long, and patients had had time enough to recover before presenting COVID-19. In addition, therapies typically used for graft versus host disease could mitigate the inflammatory response ([Saraceni et al., 2021](#)). On the contrary, outcome of recently transplanted patients that suffer from SARS-CoV-2 infection during the pre-engraftment period is not well known, and cases of ARDS at the moment of neutrophil recovery have been described ([Malek et al., 2021](#)). In our series, median time from transplantation to COVID-19 was close to 2 years and no pre-engraftment cases were detected. Nevertheless, we were not able to find differences in in-hospital mortality as compared to non-transplant recipients.

Patients receiving small molecule kinase inhibitors (such as JAK or BTK inhibitors) might be protected from hyper-inflammation, and it has been speculated that discontinuation of such therapies in cancer patients with COVID-19 could unleash the hyper-inflammatory response to SARS-CoV-2, in addition of adversely affecting the outcome of the underlying malignancy ([Wijaya et al., 2021; Stack et al., 2021](#)). In our series, all patients who were under ibrutinib or other BTK inhibitors maintained it throughout the COVID-19 episode. There were no significant differences in outcome according to BTK inhibitor treatment, though the sample is too small to draw any conclusion.

The major strength of the present study is the direct comparison of a cohort of haematologic patients with COVID-19 with a non-haematologic cohort matched by age and date of diagnosis of COVID. This allows to avoid bias secondary to different age range of patients with haematologic disease, and bias secondary to presentation in different moments of the COVID-19 learning curve at the beginning of the pandemic. Patients included were only those admitted to two

Table 4

In-hospital mortality in patients with haematologic disease and COVID-19.			
Haematologic patients mortality (n, %)	Non-survivors (27)	Survivors (44)	p
Age (mean, SD)	72.5 (16.4)	66.2 (11.6)	0.065
Sex			0.318
Male	19 (70.4%)	25 (56.8%)	
Female	8 (29.6%)	19 (43.2%)	
Race			0.694
Caucasian	25 (92.6%)	38 (86.4%)	
Latin	2 (7.4%)	4 (9.1%)	
Haematological malignancy status			0.278
Initial diagnosis	5 (18.5%)	5 (11.4%)	
Stable, no remission	5 (18.5%)	11 (25.0%)	
Relapsed or refractory	7 (25.9%)	5 (11.4%)	
Complete/partial response	10 (37.0%)	23 (52.3%)	
ECOG > 1	7 (26.9%)	2 (4.5%)	0.011
Oncologic treatment			0.372
Curative	6 (22.2%)	14 (31.8%)	
Palliative	11 (40.7%)	14 (31.8%)	
Chemotherapy during the last month	7 (25.9%)	13 (29.5%)	0.563
Rituximab	8 (29.6%)	7 (15.9%)	0.232
Ibrutinib	2 (7.4%)	5 (11.4%)	0.701
G-CSF	2 (7.4%)	4 (9.1%)	0.999
Stem cell transplant recipient	4 (14.8%)	8 (18.2%)	0.999
Allogeneic	2 (7.4%)	3 (6.8%)	
Autologous	2 (7.4%)	5 (11.4%)	
Type of haematological malignancy			0.447
Hodgkin lymphoma	0 (0.0%)	1 (2.3%)	
Non-Hodgkin lymphoma	11 (40.7%)	10 (22.7%)	
Chronic lymphocytic leukemia	3 (11.1%)	10 (22.7%)	
Acute lymphoblastic leukemia	1 (3.7%)	0 (0.0%)	
Acute myeloid leukemia	1 (3.7%)	2 (4.5%)	
Multiple myeloma	4 (14.8%)	11 (25.0%)	
Myelodysplastic syndrome	4 (14.8%)	6 (13.6%)	
Myeloproliferative neoplasm	2 (7.4%)	4 (9.1%)	
Smoker	9 (33.3%)	16 (36.4%)	0.373
Hypertension	16 (59.3%)	19 (43.2%)	0.227
Dyslipidemia	11 (40.7%)	18 (40.9%)	0.999
Obesity (Body Mass Index > 30)	3 (11.1%)	11 (25.0%)	0.215
Chronic kidney disease	2 (7.4%)	5 (11.4%)	0.701
Diabetes	7 (25.9%)	10 (22.7%)	0.086
Heart failure	2 (7.4%)	4 (9.1%)	0.999
COPD	1 (3.7%)	0 (0.0%)	0.380
Asthma	1 (3.7%)	2 (4.5%)	0.999
Laboratory parameters (mean, SD)			
Neutrophils (<500 cells/µL)	4 (14.8%)	1 (2.3%)	0.066
Lymphocytes (<1000 cells/µL)	16 (59.3 %)	23 (52.3%)	0.243
C-RP (mg/dL)	139.3 (25.8)	93.7 (11.9)	0.117
Ferritin (mg/L)	13,989 (1600)	11,807 (18097)	0.680
LDH (U/L)	663.9 (219.3)	345.2 (20.0)	0.161
DD (ng/mL)	3505 (8812)	2521 (4551)	0.594
DIC	3 (11.1%)	1 (2.3%)	0.141
ARDS	25 (92.5%)	16 (36.3%)	0.0001
COVID-19 therapy			
Lopinavir/ritonavir	14 (51.9%)	26 (59.1%)	0.626
Beta Interferon	2 (7.4%)	7 (15.9%)	0.693
Remdesivir	1 (3.7%)	0 (0.0%)	0.380
Hidroxichloroquina	23 (85.2%)	40 (90.9%)	0.469
Tocilizumab	13 (48.1%)	11 (25.0%)	0.070
Inmunoglobulin	1 (3.7%)	0 (0.0%)	0.371
Anakinra	6 (22.2%)	1 (2.3%)	0.010
Systemic corticoids	23 (85.2%)	23 (52.3%)	0.005
ICU admission	7 (25.9%)	4 (9.1%)	0.090
HFNC	10 (37.0%)	4 (9.1%)	0.005
NIMV	5 (18.5%)	2 (4.5%)	0.097
MV	6 (22.2%)	4 (9.1%)	0.164

reference hospitals in Madrid, which ensures the homogeneity in management and therapeutic options, and increases the probability of the observed differences being attributable to differences in the response to SARS-CoV-2 in haematologic patients.

Limitations of our study include the small sample size and the

heterogeneous haematologic population, which prevents from drawing any conclusion about specific types of haematologic disease or specific haematologic therapies in relation to COVID-19 outcome. During the study period, at the beginning of the pandemic, patients who received steroids, remdesivir and tocilizumab did so in the setting of clinical trials, or off-label, as compassionate use. In the case of steroids and tocilizumab, their use was significantly inferior than in patients without hematological malignancy, probably for fear of increasing immunocompromise without the certainty of a beneficial effect. Only 1 case had access to remdesivir. In this respect the results may not be applicable to the current management of COVID-19. Patients included belong only to the first COVID-19 wave, when antibody detection during recovery was not systematically addressed, and consequently it was not possible to analyze. Our series includes only hospitalized patients with COVID-19 and the results cannot be generalizable to a wider population of non-admitted haematologic patients.

Our results suggest COVID-19 has worse outcomes in haematologic patients than in non-haematologic, independently of age, and that the development of ARDS and thrombotic complications drive the higher in-hospital mortality. Immune-compromise does not prevent inflammatory complications but may in addition impede viral elimination. Maximal stress in preventive measures in haematologic patients is warranted (Malek et al., 2021), and, if unfortunately infected, close surveillance with antiviral, anti-inflammatory and anticoagulant treatment before decompensation as well as prompt consideration of intensive care management in those deteriorating (Giesen et al., 2021).

Funding

There was no funding granted for this article.

Availability of data and materials

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

Ethics

This study was carried out in accordance with the Declaration of Helsinki and was approved by the Institutional Research Ethics Committee of Málaga on March 27, 2020 (Ethics Committee code: SEMI-COVID-19 27-03-20), as per the guidelines of the Spanish Agency of Medicines and Medical Products.

Consent for publication

Only patients who had previously given consent for their medical records to be used for medical research were included in this registry. Data confidentiality and patient anonymity were maintained at all times, in accordance with Spanish regulations on observational studies.

CRediT authorship contribution statement

Ana Fernández-Cruz: Conceptualization, Formal analysis, Writing – original draft, Writing – review & editing. **Alba Puyuelo:** Data curation, Formal analysis, Writing – review & editing. **Lucía Núñez Martín-Buitrago:** Data curation, Formal analysis, Writing – review & editing. **Enrique Sánchez-Chica:** Data curation, Formal analysis, Writing – review & editing. **Carmen Díaz-Pedroche:** Conceptualization, Data curation, Formal analysis, Writing – review & editing. **Rosa Ayala:** Data curation, Writing – review & editing. **Manuel Lizasoain:** Data curation, Writing – review & editing. **Rafael Duarte:** Conceptualization, Writing – review & editing. **Carlos Lumbreras:** Conceptualization, Writing – review & editing, Project administration. **Juan Antonio Vargas:** Conceptualization, Writing – review & editing, Project administration.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We gratefully acknowledge all the investigators who participate in the SEMI-COVID-19 Registry (appendix), as well as all the clinicians of all specialties that cared for COVID-19 patients during the first wave.

Appendix

List of the SEMI-COVID-19 network members

Coordinator of the SEMI-COVID-19 Registry: José Manuel Casas Rojo.

SEMI-COVID-19 Scientific Committee Members: José Manuel Casas Rojo, José Manuel Ramos Rincón, Carlos Lumbreras Bermejo, Jesús Millán Núñez-Cortés, Juan Miguel Antón Santos, Ricardo Gómez Huelgas.

Members of the SEMI-COVID-19 Group

H. Univ. de Bellvitge. L'Hospitalet de Llobregat (Barcelona)

Xavier Corbella, Francesc Formiga Pérez, Narcís Homs, Abelardo Montero, Jose María Mora-Luján, Manuel Rubio-Rivas

H. U. 12 de Octubre. Madrid

Paloma Agudo de Blas, Coral Arévalo Cañas, Blanca Ayuso, José Bascuñana Morejón, Samara Campos Escudero, María Carnevali Frías, Santiago Cossío Tejido, Borja de Miguel Campo, Carmen Díaz Pedroche, Raquel Diaz Simon, Ana García Reyne, Laura Ibarra Veganzones, Lucia Jorge Huerta, Antonio Lalueza Blanco, Jaime Laureiro Gonzalo, Jaime Lora-Tamayo, Carlos Lumbreras Bermejo, Guillermo Maestro de la Calle, Rodrigo Miranda Godoy, Barbara Otero Perpiña, Diana Paredes Ruiz, Marcos Sánchez Fernández, Javier Tejada Montes

H. Costa del Sol. Marbella (Málaga)

Victoria Augustín Bandera, Javier García Alegria, Nicolás Jiménez-García, Jairo Luque del Pino, María Dolores Martín Escalante, Francisco Navarro Romero, Victoria Núñez Rodriguez, Julián Olalla Sierra

H. U. Gregorio Marañón. Madrid

Laura Abarca Casas, Álvaro Alejandro de Oña, Rubén Alonso Beato, Leyre Alonso Gonzalo, Jaime Alonso Muñoz, Christian Mario Amodeo Oblitas, Cristina Ausín García, Marta Bacete Cebrián, Jesús Baltasar Corral, María Barrientos Guerrero, Alejandro D. Bendala Estrada, María Calderón Moreno, Paula Carrascosa Fernández, Raquel Carrillo, Sabela Castañeda Pérez, Eva Cervilla Muñoz, Agustín Diego Chacón Moreno, María Carmen Cuenca Carvajal, Sergio de Santos, Andrés Enríquez Gómez, Eduardo Fernández Carracedo, María Mercedes Ferreiro-Mazón Jenaro, Francisco Galeano Valle, Alejandra García, Irene García Fernandez-Bravo, María Eugenia García Leoni, María Gómez Antúnez, Candela González San Narciso, Anthony Alexander Gurjian, Lorena Jiménez Ibáñez, Cristina Lavilla Olleros, Cristina Llamazares Mendo, Sara Luis García, Víctor Mato Jimeno, Clara Millán Nohales, Jesús Millán Núñez-Cortés, Sergio Moragón Ledesma, Antonio Muiño Míguez, Cecilia Muñoz Delgado, Lucía Ordieres Ortega, Susana Pardo Sánchez, Alejandro Parra Virto, María Teresa Pérez Sanz, Blanca Pinilla Llorente, Sandra Piqueras Ruiz, Guillermo Soria Fernández-Llamazares, María Toledo Macías, Neera Toledo Samaniego, Ana Torres do Rego, María Victoria Villalba García, Gracia Villarreal, María Zurita Etayo

H. de Cabueñas. Gijón (Asturias)

Ana María Álvarez Suárez, Carlos Delgado Vergés, Rosa Fernandez-Madera Martínez, Eva Mª Fonseca Aizpurua, Alejandro Gómez Carrasco, Cristina Helguera Amezua, Juan Francisco López Caleya, Diego López Martínez, María del Mar Martínez López, Aleida Martínez Zapico, Carmen Olabuenaga Iscar, Lucía Pérez Casado, María Luisa Taboada Martínez, Lara María Tamargo Chamorro

H. Royo Villanova. Zaragoza

Nicolás Alcalá Rivera, Anxela Crestelo Vieitez, Esther del Corral Beamonte, Jesús Díez Manglano, Isabel Fiteni Mera, María del Mar García Andreu, Martín Gericó Aseguinolaza, Cristina Gallego Lezaun, Claudia Josa Laorden, Raul Martínez Murgui, Marta Teresa Matía Sanz

H. Reg. Univ. de Málaga

M^a Mar Ayala-Gutiérrez, Rosa Bernal López, José Bueno Fonseca, Verónica Andrea Buonaiuto, Luis Francisco Caballero Martínez, Lidia Cobos Palacios, Clara Costo Muriel, Francis de Windt, Ana Teresa Fernandez-Truchaud Christophe, Paula García Ocaña, Ricardo Gómez Huelgas, Javier Gorospe García, José Antonio Hurtado Oliver, Sergio Jansen-Chaparro, María Dolores López-Carmona, Pablo López Quirantes, Almudena López Sampalo, Elizabeth Lorenzo-Hernández, Juan José Mancebo Sevilla, Jesica Martín Carmona, Luis Miguel Pérez-Belmonte, Iván Pérez de Pedro, Araceli Pineda-Cantero, Carlos Romero Gómez, Michele Ricci, Jaime Sanz Cánovas

H. U. La Paz. Madrid

Jorge Álvarez Troncoso, Francisco Arnalich Fernández, Francisco Blanco Quintana, Carmen Busca Arenzana, Sergio Carrasco Molina, Aranzazu Castellano Candalija, Germán Daroca Bengoa, Alejandro de Gea Grela, Alicia de Lorenzo Hernández, Alejandro Díez Vidal, Carmen Fernández Capitán, María Francisca García Iglesias, Borja González Muñoz, Carmen Rosario Herrero Gil, Juan María Herrero Martínez, Víctor Hontañón, María Jesús Jaras Hernández, Carlos Lahoz, Cristina Marcelo Calvo, Juan Carlos Martín Gutiérrez, Monica Martínez Prieto, Elena Martínez Robles, Araceli Menéndez Saldaña, Alberto Moreno Fernández, Jose María Mostaza Prieto, Ana Noblejas Mozo, Carlos Manuel Oñoro López, Esmeralda Palmier Peláez, Marina Palomar Pampyn, María Angustias Quesada Simón, Juan Carlos Ramos Ramos, Luis Ramos Ruperto, Aquilino Sánchez Purificación, Teresa Sancho Bueso, Raquel Sorrigüeta Torre, Clara Itziar Soto Abanedes, Yeray Untoria Tabares, Marta Varas Mayoral, Julia Vásquez Manau

H. U. Reina Sofía. Córdoba

Antonio Pablo Arenas de Larriva, Pilar Calero Espinal, Javier Delgado Lista, Francisco Fuentes-Jiménez, María del Carmen Guerrero Martínez, María Jesús Gómez Vázquez, Jose Jiménez Torres, Laura Limia Pérez, José López-Miranda, Laura Martín Piedra, Marta Millán Orge, Javier Pascual Vinagre, Pablo Pérez-Martínez, María Elena Revilles Vilchez, Angela Rodrigo Martínez, Juan Luis Romero Cabrera, José David Torres-Peña.

C. H. U. de Badajoz

Rafael Aragón Lara, Inmaculada Cimadevilla Fernandez, Juan Carlos Cira García, Gema María García García, Julia Gonzalez Granados, Beatriz Guerrero Sánchez, Francisco Javier Monreal Periénez, María Josefina Pascual Perez

H. Clínico de Santiago de Compostela (A Coruña)

Maria del Carmen Beceiro Abad, María Aurora Freire Romero, Sonia Molinos Castro, Emilio Manuel Paez Guillan, María Pazo Nuñez, Paula María Pesqueira Fontan

H. U. Puerta de Hierro. Madrid

Ane Andrés Eisenhofer, Ana Arias Milla, Isolina Baños Pérez, Laura Benítez Gutiérrez, Javier Bilbao Garay, Jorge Calderón Parra, Alejandro Callejas Díaz, Erika Camacho Da Silva, M^a Cruz Carreño Hernández, Raquel Castejón Díaz, María Jesús Cidores Sánchez, Carmen Cubero Gozalo, Valentín Cuervas-Mons Martínez, Laura Dorado Doblado, Sara de la Fuente Moral, Alberto Díaz de Santiago, Itziar Diego Yagüe, Ignacio Donate Velasco, Ana María Duca, Pedro Durán del Campo, Gabriela Escudero López, Esther Expósito Palomo, Ana Fernández Cruz, Amy Galán Gómez, Sonia García Prieto, Beatriz García Revilla, Miguel Ángel García Viejo, Javier Gómez Irusta, Patricia González Merino, Edith Vanessa Gutiérrez Abreu, Isabel Gutiérrez Martín, Ángela Gutiérrez Rojas, Andrea Gutiérrez Villanueva, Jesús Herráiz Jiménez, Fátima Ibáñez Estélez, Pedro Laguna del Estal, M^a Carmen Máñez Saiz, Carmen de Mendoza Fernández, María Martínez Urbistondo, Fernando Martínez Vera, María Mateos Seirul-lo, Susana Mellor Pita, Patricia A. Mills Sánchez, Esther Montero Hernández, Alberto Mora Vargas, Victor Moreno-Torres Concha, Ignacio Morras De La Torre, Elena Múñez Rubio, Rosa Muñoz de Benito, Alejandro Muñoz Serrano, Pablo Navarro Palomo, Ilduara Pintos Pascual, Arturo José Ramos Martín-Vegue, Antonio Ramos Martínez, Celia Rodríguez Olleros, Alberto Roldán Montaud, Yolanda Romero Pizarro, Silvia Rosado García, Diana Ruiz de Domingo,

David Sánchez Ortiz, Enrique Sánchez Chica, Irene Solano Almena, Elena Suanzes Martín, Yale Tung Chen, Pablo Tutor de Ureta, Ángela Valencia Alijo, Jose Manuel Vázquez Comendador, Juan Antonio Vargas Núñez.

H. Universitario Dr. Peset. Valencia

Juan Alberto Aguilera Ayllón, Arturo Artero, María del Mar Carmona Martín, María José Fabiá Valls, María de Mar Fernández Garcés, Ana Belén Gómez Belda, Ian López Cruz, Manuel Madrazo López, Elisabeth Mateo Sanchis, Jaume Micó Gandia, Laura Piles Roger, Adela María Pina Belmonte, Alba Viana García

H. Clínico San Carlos. Madrid

Inés Armenteros Yeguas, Javier Azaña Gómez, Julia Barrado Cuchillo, Irene Burruzeo López, Noemí Cabello Clotet, Alberto E. Calvo Elías, Elpidio Calvo Manuel, Carmen María Cano de Luque, Cynthia Chocron Benbunan, Laura Dans Vilan, Claudia Dorta Hernández, Ester Emilia Dubon Peralta, Vicente Estrada Pérez, Santiago Fernández-Castelao, Marcos Oliver Fragié Saavedra, José Luis García Klepzig, María del Rosario Iguarán Bermúdez, Esther Jaén Ferrer, Alejandro Maceín Rodríguez, Alejandro Marcelles de Pedro, Rubén Ángel Martín Sánchez, Manuel Méndez Bailón, Sara Miguel Álvarez, María José Nuñez Orantos, Carolina Olmos Mata, Eva Orviz García, David Oteo Mata, Cristina Oton González, Juncal Perez-Somarriba, Pablo Pérez Mateos, María Esther Ramos Muñoz, Xabier Rivas Regaira, Laura M^a Rodríguez Gallardo, Iñigo Sagastagoitia Fornie, Alejandro Salinas Botrán, Miguel Suárez Robles, Maddalena Elena Urbano, Andrea María Vellisca González, Miguel Villar Martínez

H. U. La Princesa. Madrid

María Aguilera García, Ester Alonso Monge, Jesús Álvarez Rodríguez, Claudia Alvarez Varela, Miquel Berniz Gòdia, Marta Briega Molina, Marta Bustamante Vega, Jose Curbelo, Alicia de las Heras Moreno, Ignacio Descalzo Godoy, Alexia Constanza Espiño Alvarez, Ignacio Fernández Martín-Caro, Alejandra Franquet López-Mosteiro, Gonzalo Galvez Marquez, María José García Blanco, Yaiza García del Álamo Hernández, Clara García-Rayó Encina, Noemí Gilabert González, Carolina Guillermo Rodríguez, Nicolás Labrador San Martín, Manuel Molina Báez, Carmen Muñoz Delgado, Pedro Parra Caballero, Javier Pérez Serrano, Laura Rabes Rodríguez, Pablo Rodríguez Cortés, Carlos Rodriguez Franco, Emilia Roy-Vallejo, Monica Rueda Vega, Aresio Sancha Lloret, Beatriz Sánchez Moreno, Marta Sanz Alba, Jorge Serrano Ballesteros, Alba Somovilla, Carmen Suarez Fernández, Macarena Vargas Tirado, Almudena Villa Martí

H. U. de A Coruña

Alicia Alonso Álvarez, Olaya Alonso Juarros, Ariadna Arévalo López, Carmen Casariego Castiñeira, Ana Cerezales Calviño, Marta Contreras Sánchez, Ramón Fernández Varela, Santiago J. Freire Castro, Ana Padín Trigo, Rafael Prieto Jarel, Fátima Raad Varea, Ignacio Ramil Freán, Laura Ramos Alonso, Francisco Javier Sanmartín Pensado, David Vieito Porto

H. Moisès Broggi. Sant Joan Despí (Barcelona)

Judit Aranda Lobo, Lucía Feria Casanovas, Jose Loureiro Amigo, Miguel Martín Fernández, Isabel Oriol Bermúdez, Melani Pestaña Fernández, Nicolas Rhyman, Nuria Vázquez Piqueras

H. U. S. Juan de Alicante (Alicante)

Marisa Asensio Tomás, David Balaz, David Bonet Tur, Ruth Cañizares Navarro, Paloma Chazarra Pérez, Jesús Corbacho Redondo, Eliana Damonte White, María Escamilla Espínola, Leticia Espinosa Del Barrio, Pedro Jesús Esteve Atiénzar, Carles García Cervera, David Francisco García Núñez, Francisco Garrido Navarro, Vicente Giner Galván, Angie Gómez Uranga, Javier Guzmán Martínez, Isidro Hernández Isasi, Lourdes Lajara Villar, Verónica Martínez Sempere, Juan Manuel Núñez Cruz, Sergio Palacios Fernández, Juan Jorge Peris García, Rafael Piñol Pleguezuelos, Andrea Riaño Pérez, José Miguel Seguí Ripoll, Azucena Sempere Mira, Philip Wikman-Jorgensen

H. U. Infanta Cristina. Parla (Madrid)

Juan Miguel Antón Santos, Ana Belén Barbero Barrera, Blanca Beaumonte Vela, Coralía Bueno Muñoz, Charo Burón Fernández, Ruth

Calderón Hernáiz, Irene Casado López, José Manuel Casas Rojo, Andrés Cortés Troncoso, Pilar Cubo Romano, Francesco Deodati, Alejandro Estrada Santiago, Gonzalo García Casasola Sánchez, Elena García Guijarro, Francisco Javier García Sánchez, Pilar García de la Torre, Mayte de Guzmán García-Monge, Davide Luordo, María Mateos González, José A. Melero Bermejo, Cruz Pastor Valverde, José Luis Pérez Quero, Fernando Roque Rojas, Lorea Roteta García, Elena Sierra Gonzalo, Francisco Javier Teigell Muñoz, Juan Vicente de la Sota, Javier Villanueva Martínez

H. G. U. de Elda (Alicante)

Carmen Cortés Saavedra, Jennifer Fernández Gómez, Borja González López, María Soledad Hernández Garrido, Ana Isabel López Amorós, Santiago López Gil, María de los Reyes Pascual Pérez, Nuria Ramírez Perea, Andrea Torregrosa García

H. U. Son Llàtzer. Palma de Mallorca

Andrés de la Peña Fernández, Almudena Hernández Milián

C. A. U. de Salamanca

Gloria María Alonso Claudio, Víctor Barreales Rodríguez, Cristina Carbonell Muñoz, Adela Carpio Pérez, María Victoria Coral Orbes, Daniel Encinas Sánchez, Sandra Inés Revuelta, Miguel Marcos Martín, José Ignacio Martín González, José Ángel Martín Oterino, Leticia Moralejo Alonso, Sonia Peña Balbuena, María Luisa Pérez García, Ana Ramón Prados, Beatriz Rodríguez-Alonso, Ángela Romero Alegría, María Sanchez Ledesma, Rosa Juana Tejera Pérez

C. H. U. de Ferrol (A Coruña)

Hortensia Alvarez Diaz, Tamara Dalama Lopez, Estefania Martul Pego, Carmen Mella Pérez, Ana Pazos Ferro, Sabela Sánchez Trigo, Dolores Suarez Sambade, María Trigas Ferrin, María del Carmen Vázquez Friol, Laura Vilarino Maneiro

H. U. Virgen del Rocío. Sevilla

Reyes Aparicio Santos, Máximo Bernabeu-Wittel, Santiago Rodríguez Suárez, María Nieto, Luis Giménez Miranda, Rosa María Gámez Mancera, Fátima Espinosa Torre, Carlos Hernandez Quiles, Concepción Conde Guzmán, Juan Delgado de la Cuesta, Jara Eloisa Ternero Vega, María del Carmen López Ríos, Pablo Díaz Jiménez, Bosco Baron Franco, Carlos Jiménez de Juan, Sonia Gutiérrez Rivero, Julia Lanseros Tenlaldo, Verónica Alfaro Lara, Aurora González Estrada

H. Público de Monforte de Lemos (Lugo)

José López Castro, Manuel Lorenzo López Reboiro, Cristina Sardiña González

H. Marina Baixa. Villajoyosa (Alicante)

Javier Ena, José Enrique Gómez Segado

H. General Defensa. Zaragoza

Anyuli Gracia Gutiérrez, Leticia Esther Royo Trallero

H. U. Quironsalud Madrid

Pablo Guisado Vasco, Ana Roda Santacruz, Ana Valverde Muñoz

H. U. HM Montepriñce

José F. Varona Arche

References

- Kmeid, J., Vanichanan, J., Shah, D.P., El Chaer, F., Azzi, J., Ariza-Heredia, E.J., Hosing, C., Mulanovich, V., Chemaly, R.F., 2016. Outcomes of Influenza Infections in Hematopoietic Cell Transplant Recipients: Application of an Immunodeficiency Scoring Index. *Biol Blood Marrow Transplant* 22 (3), 542–548.
- Sheshadri, A., Karimipour, M., Vakil, E., Bashoura, L., Godoy, M., Arain, M.H., Evans, S. E., Dickey, B.F., Ost, D.E., Chemaly, R.F., Faiz, S.A., 2019. Refinement of estimates of mortality risk using the Radiologic Severity Index in hematologic malignancy patients with respiratory syncytial virus infection. *Transpl Infect Dis* 21 (4).
- Chamilos, G., Lionakis, M.S., Kontoyiannis, D.P., 2021. Are All Patients with Cancer at Heightened Risk for Severe Coronavirus Disease 2019 (COVID-19)? *Clin Infect Dis* 72 (2), 351–356. <https://doi.org/10.1093/cid/ciaa1079> [published Online First: 2021/01/28].
- García-Suárez, J., de la Cruz, J., Cedillo, Á., Llamas, P., Duarte, R., Jiménez-Yuste, V., Hernández-Rivas, J.Á., Gil-Manso, R., Kwon, M.i., Sánchez-Godoy, P., Martínez-Barranco, P., Colás-Lahuerta, B., Herrera, P., Benito-Parra, L., Alegre, A., Velasco, A., Matilla, A., Aláez-Usón, M.C., Martos-Martínez, R., Martínez-Chamorro, C., Susana Quiroz, K., Del Campo, J.F., de la Fuente, A., Herráez, R., Pascual, A., Gómez, E., Pérez-Oteyza, J., Ruiz, E., Alonso, A., González-Medina, J., Martín-Buitrago, L.N., Canales, M., González-Gascón, I., Vicente-Ayuso, M.C., Valenciano, S., Roa, M.G., Monteliu, P.E., López-Jiménez, J., Escobar, C.E., Ortiz-Martín, J., Diez-Martin, J.L., Martínez-Lopez, J., Seri-Merino, C., Queiroz-Cervantes, K., Fernandez, M.E., Peñalvo-Moreno, M.-J., Naya-Errea, D., Bermejo-Martínez, L., Llorente-González, L., 2020. Impact of hematologic malignancy and type of cancer therapy on COVID-19 severity and mortality: lessons from a large population-based registry study. *J Hematol Oncol* 13 (1).
- Passamonti, F., Cattaneo, C., Arcaini, L., Bruna, R., Cavo, M., Merli, F., Angelucci, E., Krampéra, M., Cairoli, R., Della Porta, M.G., Fracchiolla, N., Ladetto, M., Gambacorti Passerini, C., Salvini, M., Marchetti, M., Lemoli, R., Molteni, A., Busca, A., Cuneo, A., Romano, A., Giuliani, N., Galimberti, S., Corso, A., Morotti, A., Falini, B., Billio, A., Gherlinzoni, F., Visani, G., Tisi, M.C., Tafuri, A., Tosi, P., Lanza, F., Massaia, M., Turrini, M., Ferrara, F., Gurrieri, C., Vallisa, D., Martelli, M., Derenzini, E., Guarini, A., Conconi, A., Cuccaro, A., Cudillo, L., Russo, D., Ciambelli, F., Scattolin, A.M., Luppi, M., Selleri, C., Ortu La Barbera, E., Ferrandina, C., Di Renzo, N., Olivieri, A., Bocchia, M., Gentile, M., Marchesi, F., Musto, P., Federici, A. B., Candoni, A., Venditti, A., Fava, C., Pinto, A., Galieni, P., Rigacci, L., Armiento, D., Pane, F., Oberti, M., Zappasodi, P., Visco, C., Franchi, M., Grossi, P.A., Bertù, L., Corrao, G., Pagano, L., Corradini, P., 2020. Clinical characteristics and risk factors associated with COVID-19 severity in patients with haematological malignancies in Italy: a retrospective, multicentre, cohort study. *Lancet Haematol* 7 (10), e737–e745.
- Casas-Rojo, J.M., Anton-Santos, J.M., Millan-Nunez-Cortes, J., et al., 2020. Clinical characteristics of patients hospitalized with COVID-19 in Spain: Results from the SEMI-COVID-19 Registry. *Rev Clin Esp* 220 (8), 480–494. <https://doi.org/10.1016/j.rce.2020.07.003> [published Online First: 2021/05/18].
- Döhner, H., Estey, E., Grimwade, D., Amadori, S., Appelbaum, F.R., Büchner, T., Dombret, H., Ebert, B.L., Fenaux, P., Larson, R.A., Levine, R.L., Lo-Coco, F., Naoe, T., Niederwieser, D., Ossenkoppele, G.J., Sanz, M., Sierra, J., Tallman, M.S., Tien, H.-F., Wei, A.H., Löwenberg, B., Bloomfield, C.D., 2017. Diagnosis and management of AML in adults: 2017 ELN recommendations from an international expert panel. *Blood* 129 (4), 424–447.
- Cheson, B.D., Fisher, R.I., Barrington, S.F., Cavalli, F., Schwartz, L.H., Zucca, E., Lister, T. A., 2014. Recommendations for initial evaluation, staging, and response assessment of Hodgkin and non-Hodgkin lymphoma: the Lugano classification. *J Clin Oncol* 32 (27), 3059–3067.
- Kumar, S., Paiva, B., Anderson, K.C., Durie, B., Landgren, O., Moreau, P., Munshi, N., Lonial, S., Bladé, J., Mateos, M.-V., Dimopoulos, M., Kastritis, E., Boccadoro, M., Orlowski, R., Goldschmidt, H., Spencer, A., Hou, J., Chng, W.J., Usmani, S.Z., Zamagni, E., Shimizu, K., Jagannath, S., Johnsen, H.E., Terpos, E., Reiman, A., Kyle, R.A., Sonneveld, P., Richardson, P.G., McCarthy, P., Ludwig, H., Chen, W., Cavo, M., Harousseau, J.-L., Lentzsch, S., Hillengass, J., Palumbo, A., Orfao, A., Rajkumar, S.V., Miguel, J.S., Avet-Loiseau, H., 2016. International Myeloma Working Group consensus criteria for response and minimal residual disease assessment in multiple myeloma. *Lancet Oncol* 17 (8), e328–e346.
- Oken, M.M., Creech, R.H., Tormey, D.C., Horton, J., Davis, T.E., McFadden, E.T., Carbone, P.P., 1982. Toxicity and response criteria of the Eastern Cooperative Oncology Group. *AMERICAN JOURNAL OF CLINICAL ONCOLOGY* 5 (6), 649–656.
- Force, A.D.T., Ranieri, V.M., Rubenfeld, G.D., et al., 2012. Acute respiratory distress syndrome: the Berlin Definition. *JAMA* 307 (23), 2526–2533. <https://doi.org/10.1001/jama.2012.5669>.
- Moreno-Torres, V., de la Fuente, S., Mills, P., Muñoz, A., Muñoz, E., Ramos, A., Fernández-Cruz, A., Arias, A., Pintos, I., Vargas, J.A., Cuervas-Mons, V., de Mendoza, C., 2021. Major determinants of death in patients hospitalized with COVID-19 during the first epidemic wave in Madrid, Spain. *Medicine (Baltimore)* 100 (16), e25634.
- Osuchowski, M.F., Winkler, M.S., Skirecki, T., Cajander, S., Shankar-Hari, M., Lachmann, G., Monneret, G., Venet, F., Bauer, M., Brunkhorst, F.M., Weis, S., Garcia-Salido, A., Kox, M., Cavaillon, J.-M., Uhle, F., Weigand, M.A., Flohé, S.B., Wiersinga, W.J., Almansa, R., de la Fuente, A., Martin-Loeches, I., Meisel, C., Spinetti, T., Scheffold, J.C., Cillóniz, C., Torres, A., Giannarellis-Bourboulis, E.J., Ferrer, R., Girardis, M., Cossarizza, A., Netea, M.G., van der Poll, T., Bermejo-Martín, J.F., Rubio, I., 2021. The COVID-19 puzzle: deciphering pathophysiology and phenotypes of a new disease entity. *Lancet Respir Med* 9 (6), 622–642.
- Rhee, C., Kang, J.i., Kim, J., Yoon, H., Kim, S., Kwon, S., Kim, Y., Kim, K., Moon, H., Park, S., Kim, H., Lee, S., Song, J., 2009. Risk factors for acute respiratory distress syndrome during neutropenia recovery in patients with hematologic malignancies. *Crit Care* 13 (6), R173.
- Malek, A.E., Adachi, J.A., Mulanovich, V.E., Sassine, J., Raad, II., McConn, K., Seiler, G. T., Dhal, U., Khawaja, F., Chemaly, R.F., 2021. Immune reconstitution and severity of COVID-19 among hematopoietic cell transplant recipients. *Transpl Infect Dis* 23 (4). <https://doi.org/10.1111/tid.v23.410.1111.tid.13606>.
- Mignard, X., Biard, L., Lemiale, V., Mokart, D., Pène, F., Kouatchet, A., Mayaux, J., Vincent, F., Nyunga, M., Bruneel, P., Rabbat, A., Lebert, C., Perez, P., Meert, A.-P., Benoit, D., Hamidfar, R., Darmon, M., Azoulay, E., Zafrani, L., 2019. Granulocyte colony-stimulating factor and respiratory status of critically ill neutropenic patients with hematologic malignancies. *Leuk Lymphoma* 60 (5), 1156–1163.
- Bonaventura, A., Vecchié, A., Dagna, L., Martinod, K., Dixon, D.L., Van Tassell, B.W., Dentali, F., Montecucco, F., Massberg, S., Levi, M., Abbate, A., 2021. Endothelial dysfunction and immunothrombosis as key pathogenic mechanisms in COVID-19. *Nat Rev Immunol* 21 (5), 319–329.
- O'Donnell, J.S., Peyvandi, F., Martin-Loeches, I., 2021. Pulmonary immuno-thrombosis in COVID-19 ARDS pathogenesis. *Intensive Care Med* 47 (8), 899–902. <https://doi.org/10.1007/s00134-021-06419-w> [published Online First: 2021/05/31].
- Krekre, N., Connors, J.M., 2019. Venous thromboembolism incidence in hematologic malignancies. *Blood Rev* 33, 24–32. <https://doi.org/10.1016/j.blre.2018.06.002> [published Online First: 2018/09/29].

- Taramasso, L., Sepulcri, C., Mikulska, M., Magnasco, L., Lai, A., Bruzzone, B., Dentone, C., Bassetti, M., 2021. Duration of isolation and precautions in immunocompromised patients with COVID-19. *J Hosp Infect* 111, 202–204.
- Pinana, J.L., Martino, R., García-García, I., Parody, R., Morales, M.D., Benzo, G., Gómez-Catalan, I., Coll, R., De La Fuente, I., Luna, A., Merchán, B., Chinea, A., de Miguel, D., Serrano, A., Pérez, C., Diaz, C., Lopez, J.I., Saez, A.J., Bailen, R., Zudaire, T., Martínez, D., Jurado, M., Calbacho, M., Vázquez, L., Garcia-Cadenas, I., Fox, L., Pimentel, A.I., Bautista, G., Nieto, A., Fernandez, P., Vallejo, J.C., Solano, C., Valero, M., Espigado, I., Saldaña, R., Sisinni, L., Ribera, J.M., Jimenez, M.J., Trabazo, M., Gonzalez-Vicent, M., Fernández, N., Talarin, C., Montoya, M.C., Cedillo, A., Sureda, A., 2020. Risk factors and outcome of COVID-19 in patients with hematological malignancies. *Exp Hematol Oncol* 9 (1).
- Saraceni, F., Scortechini, I., Mancini, G., Mariani, M., Federici, I., Gaetani, M., Barbatelli, P., Minnucci, M.L., Bagnarelli, P., Olivieri, A., 2021. Severe COVID-19 in a patient with chronic graft-versus-host disease after hematopoietic stem cell transplant successfully treated with ruxolitinib. *Transpl Infect Dis* 23 (1) [published Online First: 2020/07/07].
- Wijaya, I., Andhika, R., Huang, I., Purwiga, A., Budiman, K.Y., Bashari, M.H., Reniarti, L., Roesli, R.M.A., 2021. The use of Janus Kinase inhibitors in hospitalized patients with COVID-19: Systematic review and meta-analysis. *Clin Epidemiol Glob Health* 11, 100755 [published Online First: 2021/05/11].
- Stack, M., Sacco, K., Castagnoli, R., et al., 2021. BTK inhibitors for Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2): A Systematic Review. *Res Sq.* <https://doi.org/10.21203/rs.3.rs-319342/v1> [published Online First: 2021/04/02].
- Giesen, N., Sprute, R., Rüthrich, M., Khodamoradi, Y., Mellinghoff, S.C., Beutel, G., Lueck, C., Koldehoff, M., Henrich, M., Sandherr, M., von Bergwelt-Baildon, M., Wolf, H.-H., Hirsch, H.H., Wörmann, B., Cornely, O.A., Köhler, P., Schalk, E., von Lilienfeld-Toal, M., 2021. 2021 update of the AGIHO guideline on evidence-based management of COVID-19 in patients with cancer regarding diagnostics, viral shedding, vaccination and therapy. *Eur J Cancer* 147, 154–160 [published Online First: 2021/03/07].