

1 Anti-membrane antibodies persist at least one year and discriminate between past COVID-19  
2 infection and vaccination

3

4 **Running head:** Anti-membrane antibodies in COVID-19

5

6 Maya F. Amjadi<sup>1</sup>, Ryan R. Adyniec<sup>1</sup>, Srishti Gupta<sup>1</sup>, S. Janna Bashar<sup>1</sup>, Aisha M. Mergaert<sup>1,2</sup>,  
7 Katarina M. Braun<sup>3</sup>, Gage K. Moreno<sup>2</sup>, David H. O'Connor<sup>2</sup>, Thomas C. Friedrich<sup>3</sup>, Nasia  
8 Safdar<sup>1</sup>, Sara S. McCoy<sup>1</sup>, Miriam A. Shelef<sup>1,4\*</sup>

9

10 <sup>1</sup> Department of Medicine, University of Wisconsin (UW)-Madison, Madison, 53705 USA

11 <sup>2</sup> Department of Pathology and Laboratory Medicine, UW-Madison, Madison, 53705 USA

12 <sup>3</sup> Department of Pathobiological Sciences, UW-Madison, Madison, 53705 USA

13 <sup>4</sup> William S. Middleton Memorial Veterans Hospital, Madison, 53705 USA

14

15 **\*Corresponding author:**

16 Miriam A. Shelef

17 4130 MFCB

18 1685 Highland Avenue

19 Madison, WI 53705, USA

20 1 (608) 263-5241

21 [mshelef@medicine.wisc.edu](mailto:mshelef@medicine.wisc.edu)

22

23 **Alternate corresponding author:**

24 Maya F. Amjadi

25 Research Services

26 VA Medical Center Rm C3127

27 2500 Overlook Terrace

28 Madison WI 53705-2254

29 608-256-1901x17811

30 mamjadi@wisc.edu

31

32 **(1) Declaration of Conflict of Interest**

33

34 MFA, DHO, and MAS are listed as inventors on a patent filed related to this study  
35 (PCT/US2021/051143; IDENTIFICATION OF SARS-COV-2 EPITOPES DISCRIMINATING  
36 COVID-19 INFECTION FROM CONTROL AND METHODS OF USE). Promega provided  
37 Lumit<sup>TM</sup> SARS-CoV-2 Immunoassay kits.

38

39

40

41

1 **(2) Financial Support**

2  
3 This work was supported by a Wisconsin Partnership Program COVID-19 Response grant  
4 (4647) and a UW Department of Medicine COVID-19 Pilot Award to MAS, by the CDC  
5 (75D30120C09870, 75D30121C11060) to DHO and TCF, and by NIAID (R24OD017850) to  
6 DHO. Additional support includes NIA (T32 AG000213) and MSTP (T32 GM140935) to MFA,  
7 NHLBI (T32 HL007899) to AMM, NCATS (1KL2TR002374) to SSM, and NIAID  
8 (1DP2AI144244-01) to NS. DHO and TCF are members of the Upper Midwest Regional  
9 Accelerator for Genomic Surveillance funded by the Rockefeller Foundation.

10  
11 **(3) Meetings at which all or part of the information has been presented**

12 None

13 **(4) To whom correspondence and requests for reprints should be addressed:**

14 Miriam A. Shelef  
15 4130 MFCB  
16 1685 Highland Avenue  
17 Madison, WI 53705, USA  
18 1 (608) 263-5241  
19 [mshelef@medicine.wisc.edu](mailto:mshelef@medicine.wisc.edu)

20  
21 **(5) Current affiliations for authors whose affiliations have changed since study completion**

22  
23 Aisha M. Mergaert, Labcorp Drug Development, 3301 Kinsman Blvd, Madison, WI 53704  
24 Gage K. Moreno, Broad Institute, 415 Main Street, Cambridge, Massachusetts 02142  
25

1 **Abstract**

2 The consequences of past COVID-19 infection for personal and population health are emerging,  
3 but accurately identifying distant infection is a challenge. Anti-spike antibodies rise after both  
4 vaccination and infection and anti-nucleocapsid antibodies rapidly decline. We evaluated anti-  
5 membrane antibodies in COVID-19 naïve, vaccinated, and convalescent subjects to determine if  
6 they persist and accurately detect distant infection. We found that anti-membrane antibodies  
7 persist for at least a year and are a sensitive and specific marker of past COVID-19 infection.  
8 Thus, anti-membrane and anti-spike antibodies together can differentiate between COVID-19  
9 convalescent, vaccinated, and naïve states to advance public health and research.

10 **Keywords:** antibody, SARS-CoV-2, COVID-19

11

ACCEPTED MANUSCRIPT

## 1 **Background**

2 The cardinal features and challenges of the COVID-19 pandemic have changed. Initially, the  
3 pandemic was defined by SARS-CoV-2 infections in an immunologically naïve population.  
4 Now, immunity from vaccination, infection, or both is common, reducing the severity of future  
5 infection waves. However, identifying infection for research and public health efforts is limited  
6 by the need to perform viral testing during acute infection (which may not occur during  
7 asymptomatic disease or pandemic surges that deplete resources) and by shortcomings in  
8 serologic testing [1]. Anti-nucleocapsid antibodies often decline to seronegativity just months  
9 after infection [2-4]. Anti-spike antibodies persist at least a year post-infection [5, 6] but  
10 typically cannot differentiate between infection and vaccination [7-9]. Anti-membrane antibodies  
11 develop soon after SARS-CoV-2 infection [10-12], but they are rarely assessed and their  
12 longevity is unknown. Here, we evaluated antibodies against the receptor binding domain (RBD)  
13 of spike, nucleocapsid, and membrane antigens in naïve, COVID-19 vaccinated, and COVID-19  
14 convalescent subjects up to a year post symptom resolution to evaluate the persistence of anti-  
15 membrane antibodies and to identify antigens that discriminate between distant infection,  
16 vaccination, and naïve states.

17

## 18 **Methods**

### 19 *Human subjects*

20 Human studies were approved by the UW Institutional Review Board and human subjects  
21 provided written informed consent. Sera and data collected before 2019 from 60 COVID-19  
22 naïve adults without inflammatory disease (exception: one subject with psoriatic arthritis using

1 adalimumab to match two COVID-19 convalescent subjects using adalimumab) were obtained  
2 from the University of Wisconsin (UW) Rheumatology Biorepository [13].  
3  
4 COVID-19 convalescent sera and data were obtained from the UW COVID-19 Convalescent  
5 Biorepository [14]. Briefly, in the spring of 2020, adults with a positive SARS-CoV-2 PCR test  
6 at UW Health were invited to participate until 121 subjects were recruited. Demographic and  
7 clinical information were collected by questionnaire and electronic medical record (EMR)  
8 abstraction. COVID-19 severity was diverse: mild (n=12), moderate (n=86), severe (n=15) and  
9 critical (n=8) as previously defined [14]. Subjects provided blood and clinical information 5  
10 weeks (n=121), 3 months (n=115), 6 months (n=98) and 12 months (n=100) +/- 3 weeks after  
11 symptom resolution. One sample collected >3 weeks from the 3 month timepoint and subjects  
12 for whom the 5 week time point was collected >3 weeks from the intended timepoint (n=1) or  
13 missed >1 blood draw (n=16) were excluded from comparative analyses, generating sample sizes  
14 of 104 (5 weeks), 101 (3 months), 97 (6 months), and 98 (12 months). Based on anti-RBD Ig  
15 elevation timing post-vaccination (Supplementary Figure 1), COVID-19 convalescent subjects at  
16 12 months were considered vaccinated if they received 1 vaccine dose  $\geq 5$  days before sample  
17 collection (n=77) and unvaccinated (n=21) if they received no vaccine (n=17) or their first or  
18 only vaccine dose <5 days before sample collection (n=4).  
19  
20 Vaccinated individuals without past COVID-19 (n=21) were recruited by flyers at UW Health in  
21 summer of 2021. Complete vaccination (>3 weeks after two mRNA vaccine doses or one  
22 Ad26.COV2.S dose) and lack of known COVID-19 was confirmed by questionnaire and EMR  
23 review.

1  
2 Limited clinical data, sera, and SARS-CoV-2 lineages were provided by UW Health Infection  
3 Control for 20 completely vaccinated healthcare workers with breakthrough COVID-19 (positive  
4 PCR and symptoms) in the spring of 2021. Blood collection occurred ~1 day after SARS-CoV-2  
5 PCR (range 0-4 days) and ~3 days after symptom onset (range 0-5 days). Four breakthrough  
6 cases also had PCR positive COVID-19 prior to vaccination completion and 3-6 months prior to  
7 breakthrough infection. Healthcare workers with breakthrough infections were invited to  
8 participate in the longitudinal study, and 3 provided blood ~8 weeks (range 37-70 days) after the  
9 initial collection.

10

#### 11 *Anti-RBD Ig Immunoassay*

12 Anti-RBD Ig was detected by Lumit<sup>TM</sup> SARS-CoV-2 Immunoassay (Promega, Madison, WI)  
13 according to kit instructions using a TEMPEST<sup>®</sup> Liquid Handler (Formulatrix, Bedford, MA)  
14 and a PHERAstar FS plate reader (BMG Labtech, Ortenberg, Germany). Sera were diluted 1:10  
15 to use the kit's recommended sample/calibrator cutoff of 1 for seropositivity. At 1:10, the highest  
16 anti-RBD Ig values were above the linear range, but results were overall similar to a 1:200  
17 dilution (Supplementary Figure 2), at which higher values were within the linear range.

18

#### 19 *Anti-membrane and anti-nucleocapsid IgG ELISA*

20 ELISAs were performed as previously to detect IgG against SARS-CoV-2 membrane (aa 8-23,  
21 ITVEELKKLLEQWNLV-K-biotin) and nucleocapsid (aa 390-405, QTVTLLPAADLDDFSK-  
22 K-biotin) peptides [10] with the following modifications: blocking for >2.5 instead of 1 hour and  
23 serum dilution of 1:50 (nucleocapsid) or 1:500 (membrane), instead of 1:200 to maximally

1 utilize the linear range. Relative absorbance values (IgG binding to uncoated wells subtracted  
2 from coated wells for each subject and values normalized across plates using a serum standard)  
3 of 0 were plotted as 0.0001 to allow a log scale for graphs.

#### 4 5 *Statistics*

6 Statistical analyses were performed using Prism (GraphPad, San Diego, CA) and JMP (SAS  
7 Institute, Cary, NC) software. Antibody levels were compared between naïve or vaccinated  
8 subjects versus all other groups by Kruskal-Wallis One-Way ANOVA with Dunn's multiple  
9 comparisons test. Antibody levels in unvaccinated versus vaccinated 12 month convalescent  
10 samples were compared by Mann-Whitney U test. Matched antibody levels across multiple  
11 timepoints in convalescent subjects were compared by Friedman test with Dunn's multiple  
12 comparisons test. Antibody levels in breakthrough infection subjects were compared at ~3 days  
13 post symptom onset versus ~8 weeks later by Wilcoxon signed rank test. Antibody positivity was  
14 compared between two groups with Fisher's exact test and among multiple groups with a chi-  
15 square test. P values <0.05 were considered significant.

#### 16 17 **Results**

18 We quantified antibodies against RBD, nucleocapsid, and membrane antigens in sera from the  
19 following subjects: naïve, vaccinated with no known COVID-19 infection, COVID-19  
20 convalescent with sera collected 5 weeks, 3 months, 6 months, and 12 months post symptom  
21 resolution (all initially unvaccinated), and vaccinated with subsequent SARS-CoV-2  
22 breakthrough infection. Clinical and demographic information is in Supplementary Table 1.

23

1 For anti-RBD Ig, the chemiluminescent assay had an area under the receiver operator curve  
2 (AUC) of 0.973, 90% sensitivity, and 97% specificity (Supplementary Figure 3A), comparable to  
3 other assays [14]. Convalescent and vaccinated individuals had significantly higher anti-RBD Ig  
4 than naïve subjects, with no significant difference in antibody levels or percent seropositivity  
5 between vaccinated subjects and unvaccinated convalescent subjects (Figure 1A and 1B). In a  
6 matched analysis of the 88 subjects who provided serum at all 4 timepoints (Figure 1C), anti-  
7 RBD Ig levels were statistically different between timepoints, but the extremely small difference  
8 in medians seems unlikely to be biologically meaningful. Further, the percent of seropositive  
9 subjects at 5 weeks (91%) versus 6 months (88%) was not significantly different (Figure 1B).  
10 Because only 21 convalescent subjects remained unvaccinated at 12 months, this timepoint was  
11 not compared to the 5 week timepoint. Not surprisingly, anti-RBD Ig levels in 12 month  
12 convalescent subjects were significantly higher for those who received at least one dose of a  
13 vaccine compared with no vaccine (Figure 1A and Supplementary Figure 1B). Finally, all  
14 vaccinated subjects, with or without past or breakthrough infections, were seropositive for anti-  
15 RBD Ig, with no significant differences in antibody levels between vaccinated subjects with or  
16 without breakthrough infections (Figure 1A and 1B). Overall, these data suggest that anti-RBD  
17 antibodies are detectable in the vast majority of vaccinated and convalescent subjects at least a  
18 year after infection, but cannot differentiate between past infection, vaccination, and vaccination  
19 with breakthrough infection.

20  
21 Next, we evaluated anti-nucleocapsid IgG. The ELISA had an AUC of 0.919, 91% sensitivity,  
22 and 88% specificity (Supplementary Figure 3B). The low specificity of this test is consistent  
23 with other anti-nucleocapsid tests, likely due to cross-reactivity with common cold coronavirus



1 nucleocapsid [14, 15]. Nonetheless, as expected, there was no difference in anti-nucleocapsid  
2 IgG levels between naïve versus vaccinated subjects or between 12 month convalescent  
3 vaccinated versus unvaccinated subjects (Figure 1D). Also as expected, compared to either  
4 vaccinated or naïve subjects, anti-nucleocapsid IgG levels were higher in subjects with past  
5 infection (Figure 1D). However, in a matched analysis of convalescent subjects, anti-  
6 nucleocapsid IgG levels fell significantly over time (Figure 1F) with 34% of subjects  
7 seronegative by 6 months and 48% by 12 months, a significant increase in seronegativity  
8 compared to 8% at 5 weeks (Figure 1E). Interestingly, none of the four breakthrough cases who  
9 also had COVID-19 before vaccination were seropositive for anti-nucleocapsid IgG at the time  
10 of breakthrough infection and only one of three subjects was seropositive eight weeks after  
11 breakthrough infection (Figure 1E). Together, these data highlight the rapid decline of anti-  
12 nucleocapsid antibodies.

13  
14 Last, we evaluated anti-membrane IgG. The ELISA had an AUC of 0.956, 88% sensitivity, and  
15 95% specificity (Supplementary Figure 3C). As expected, anti-membrane IgG levels and percent  
16 seropositivity did not differ between naïve and vaccinated subjects (or between 12 month  
17 convalescent vaccinated and unvaccinated subjects), but were significantly higher in  
18 convalescent subjects (Figure 1G and 1H). In a matched analysis over time (Figure 1I), anti-  
19 membrane IgG levels remained stable at 6 months with an extremely small decline at 12 months.  
20 However, at 12 months, 94% of convalescent samples were seropositive for anti-membrane IgG,  
21 as compared to 88% at 5 weeks (Figure 1H). Interestingly, all four vaccine breakthrough  
22 infection subjects with prior COVID-19 and no breakthrough subjects without prior COVID-19  
23 were seropositive for anti-membrane IgG during acute infection (Figure 1H). Together, these

1 data demonstrate that anti-membrane IgG persists at least a year and can be a sensitive and  
2 specific marker of past COVID-19 infection.

3  
4 Finally, we compared antibody levels at 5 weeks and 12 months post COVID-19 across disease  
5 severity groups. As expected [10, 14], levels of all three antibodies were generally higher in  
6 subjects with more severe COVID-19 at both time points, except anti-nucleocapsid IgG at 12  
7 months (Figure 2).

8  
9 **Discussion**

10 Here, in addition to confirming that anti-RBD antibodies last at least a year and anti-  
11 nucleocapsid IgG declines over months [2-6], we demonstrate that anti-membrane IgG is present  
12 in the vast majority of COVID-19 convalescent patients and persists at least a year. Our findings  
13 are consistent with findings for IgG against a recombinant membrane antigen (polypeptide of aa  
14 1-19 and 101-222) in the early convalescent period [11]. In contrast, Jörrißen and colleagues  
15 found that only ~20% of non-hospitalized COVID-19 convalescent subjects had IgG against a  
16 membrane peptide in the early convalescent period [12]. Our non-hospitalized subjects alone  
17 were 88% positive for anti-membrane IgG at 5 weeks (n=83) and 94% at 12 months (n=77) post  
18 COVID-19. This discrepancy may be due to their smaller sample size (n=30) or use of a different  
19 peptide (aa 1-20).

20  
21 Given the absence of anti-RBD and anti-membrane antibodies in naïve subjects, the presence of  
22 only anti-RBD antibodies in vaccinated subjects, and the presence of both in COVID-19  
23 convalescent subjects up to 12 months after infection, our study suggests that a combination of

1 anti-RBD and anti-membrane antibody testing could be used to detect past COVID-19 infection  
2 and vaccination at a population and individual level. An analogous testing strategy for hepatitis  
3 B uses anti-surface antibodies to detect past infection or vaccination and anti-core antibodies to  
4 detect past infection. While SARS-CoV-2 does not appear to cause persistent infection like  
5 hepatitis B, the long-term consequences of COVID-19 are still emerging and revealing a  
6 previously undetected infection may prove important. At minimum, detecting unknown past  
7 infections may relieve personal anxiety about future infections in some individuals. Moreover,  
8 accurate assessment of past infection in a population could enhance the prediction of and  
9 interpretation of COVID-19 surge outcomes and inform public health policy.

10

11 Limitations of this study include that samples were collected only up to 12 months post COVID-  
12 19 and that we quantified IgG, not IgM or IgA, that binds membrane and nucleocapsid peptides  
13 versus total Ig that binds RBD. Also, subjects were infected by ancestral SARS-CoV-2 lineages  
14 (early 2020) or alpha and delta variants (breakthrough infections, Supplementary Figure 4),  
15 whereas the omicron variant has a single amino acid difference in the membrane peptide  
16 (ITVEELKKLLEEWNLV). Finally, sample sizes for breakthrough infections were small with  
17 samples collected ~3 days after symptom onset, possibly allowing an early antibody response.  
18 Future studies are needed to evaluate later time points, multiple antibody isotypes, larger cohorts,  
19 and antibodies after omicron infections.

20

21 Nonetheless, we demonstrate that anti-membrane antibodies persist at least a year and, together  
22 with anti-RBD antibodies, can accurately identify past-COVID-19 infection and vaccination.

23

1 **Funding**

2 This work was supported by a Wisconsin Partnership Program COVID-19 Response grant  
3 [4647] and a UW Department of Medicine COVID-19 Pilot Award to MAS, by the CDC  
4 [75D30120C09870, 75D30121C11060] to DHO and TCF, and by NIAID [R24OD017850] to  
5 DHO. Additional support includes NIA [T32 AG000213] and MSTP [T32 GM140935] to MFA,  
6 NHLBI [T32 HL007899] to AMM, NCATS [1KL2TR002374] to SSM, and NIAID  
7 [1DP2AI144244-01] to NS. DHO and TCF are members of the Upper Midwest Regional  
8 Accelerator for Genomic Surveillance funded by the Rockefeller Foundation.

9

10 **Acknowledgements**

11 The authors thank Gene Ananiev and Song Guo at UW's Small Molecules Screening Facility  
12 and Laura Perez-Brenner, Samantha Lewis and Melanie Dart from Promega for technical advice,  
13 UW Health Infection Control and Employee Health Services, the UW Clinical Research Office,  
14 and the participating human subjects.

15

16 **Conflicts of Interest**

17 MFA, DHO, and MAS are listed as inventors on a patent filed related to this study  
18 (PCT/US2021/051143; IDENTIFICATION OF SARS-COV-2 EPITOPES DISCRIMINATING  
19 COVID-19 INFECTION FROM CONTROL AND METHODS OF USE). Promega provided  
20 Lumit<sup>TM</sup> SARS-CoV-2 Immunoassay kits.

21

1 **References**

- 2 1. Gong F, Wei HX, Li Q, Liu L, Li B. Evaluation and Comparison of Serological Methods for  
3 COVID-19 Diagnosis. *Front Mol Biosci* **2021**; 8:682405.
- 4 2. Ripperger TJ, Uhrlaub JL, Watanabe M, et al. Orthogonal SARS-CoV-2 Serological Assays  
5 Enable Surveillance of Low-Prevalence Communities and Reveal Durable Humoral  
6 Immunity. *Immunity* **2020**; 53:925-33.e4.
- 7 3. Liu A, Li Y, Peng J, Huang Y, Xu D. Antibody responses against SARS-CoV-2 in COVID-19  
8 patients. *J Med Virol* **2020**.
- 9 4. Bolotin S, Tran V, Osman S, et al. SARS-CoV-2 Seroprevalence Survey Estimates Are  
10 Affected by Anti-Nucleocapsid Antibody Decline. *J Infect Dis* **2021**; 223:1334-8.
- 11 5. Gallais F, Gantner P, Bruel T, et al. Evolution of antibody responses up to 13 months after  
12 SARS-CoV-2 infection and risk of reinfection. *EBioMedicine* **2021**; 71:103561.
- 13 6. Wang Z, Muecksch F, Schaefer-Babajew D, et al. Naturally enhanced neutralizing breadth  
14 against SARS-CoV-2 one year after infection. *Nature* **2021**; 595:426-31.
- 15 7. Wang Z, Schmidt F, Weisblum Y, et al. mRNA vaccine-elicited antibodies to SARS-CoV-2  
16 and circulating variants. *Nature* **2021**; 592:616-22.
- 17 8. Doria-Rose N, Suthar MS, Makowski M, et al. Antibody Persistence through 6 Months after  
18 the Second Dose of mRNA-1273 Vaccine for Covid-19. *N Engl J Med* **2021**; 384:2259-61.
- 19 9. Sadoff J, Gray G, Vandebosch A, et al. Safety and Efficacy of Single-Dose Ad26.COV2.S  
20 Vaccine against Covid-19. *N Engl J Med* **2021**; 384:2187-201.
- 21 10. Heffron AS, McIlwain SJ, Amjadi MF, et al. The landscape of antibody binding in SARS-  
22 CoV-2 infection. *PLoS Biol* **2021**; 19:e3001265.

- 1 11. Lopandić Z, Protić-Rosić I, Todorović A, et al. IgM and IgG Immunoreactivity of SARS-  
2 CoV-2 Recombinant M Protein. *Int J Mol Sci* **2021**; 22.
- 3 12. Jörrißen P, Schütz P, Weiland M, et al. Antibody Response to SARS-CoV-2 Membrane  
4 Protein in Patients of the Acute and Convalescent Phase of COVID-19. *Front Immunol* **2021**;  
5 12:679841.
- 6 13. Holmes CL, Peyton CG, Bier AM, et al. Reduced IgG titers against pertussis in rheumatoid  
7 arthritis: Evidence for a citrulline-biased immune response and medication effects. *PLoS One*  
8 **2019**; 14:e0217221.
- 9 14. Amjadi MF, O'Connell SE, Armbrust T, et al. Specific COVID-19 Symptoms Correlate with  
10 High Antibody Levels against SARS-CoV-2. *Immunohorizons* **2021**; 5:466-76.
- 11 15. Noda K, Matsuda K, Yagishita S, et al. A novel highly quantitative and reproducible assay  
12 for the detection of anti-SARS-CoV-2 IgG and IgM antibodies. *Sci Rep* **2021**; 11:5198.
- 13
- 14

1 **Figure Legend**

2 **Figure 1. Anti-RBD, anti-nucleocapsid, and anti-membrane antibodies after COVID-19**  
3 **vaccination and infection.** Anti-RBD Ig was detected by immunoassay (reported as  
4 sample/calibrator, S/C) and anti-nucleocapsid and anti-membrane IgG were quantified by ELISA  
5 (reported as relative absorbance, rel. abs.) in sera from the following subjects: naive (n=60),  
6 vaccinated with no known COVID-19 infection (Vax, n=21), COVID-19 convalescent 5 weeks  
7 (5w, n=104), 3 months (3m, n=101), 6 months (6m, n=97), and 12 months (12m, n=98) post-  
8 symptom resolution either vaccinated (12m Vax, n=77) or not (12m Unvax, n=21), vaccinated  
9 with breakthrough COVID-19 ~3 days (Vax BT ~3d, n=20) and ~8 weeks (Vax BT ~8w, n=3)  
10 after symptom onset including 4 subjects with previous COVID-19 infection (PI). Anti-RBD Ig  
11 (A), anti-nucleocapsid IgG (D), and anti-membrane IgG (G) levels for all groups were graphed  
12 and compared to naive (blue) or Vax (gray) by Kruskal-Wallis and Dunn's multiple comparisons  
13 tests and 12m Unvax was compared to 12m Vax by Mann-Whitney test (brackets). Vax BT with  
14 both ~3d and ~8w timepoints are represented with triangles or squares and Vax BT subjects with  
15 PI in red symbols. Percent positive (black) and negative (gray) for anti-RBD Ig (B), anti-  
16 nucleocapsid IgG (E), and anti-membrane IgG (H) were graphed and compared between selected  
17 groups by Fisher's exact (line) or chi-square (bracket encompassing compared groups) tests.  
18 Matched anti-RBD Ig (C), anti-nucleocapsid IgG (F) and anti-membrane IgG (I) levels were  
19 compared across timepoints for COVID-19 convalescent subjects (n=88) by Friedman test with  
20 Dunn's multiple comparisons test. For top and bottom panels: bars indicate medians, and dashed  
21 lines indicate cutoffs. For all panels: \*p<0.05, \*\*p<0.01, \*\*\*p<0.001, \*\*\*\*p<0.0001, or not  
22 significant (ns).

23

1 **Figure 2. Anti-SARS-CoV-2 antibody levels are higher after more severe COVID-19.** Anti-  
2 RBD Ig (A), anti-nucleocapsid IgG (B), and anti-membrane IgG (C) were compared across  
3 disease severity groups by Kruskal-Wallis with Dunn's Multiple Comparisons test at indicated  
4 time points (mild: n=11 5w, n=10 6m, n=9 12m; moderate (mod.): n=72 5w, n=68 6m and 12m;  
5 severe: n=15 5w, n=13 6m, n=15 12m; critical: n=6 5w, 6m, and 12m). For all panels: bars  
6 represent medians, dashed lines indicate cutoffs, \*p<0.05, \*\*p<0.01, \*\*\*p<0.001,  
7 \*\*\*\*p<0.0001.

8

ACCEPTED MANUSCRIPT






Figure 1  
25x25 mm (.33 x DPI)

1  
2  
3  
4




Figure 2  
25x25 mm (.33 x DPI)

1  
2  
3