

Addressing inequalities in oral health in India: need for skill mix in the dental workforce

Manu Raj Mathur¹, Ankur Singh², Richard Watt³

¹Public Health Foundation of India, Gurgaon, Haryana, India, ²Australian Research Centre for Population Oral Health, School of Dentistry, The University of Adelaide, Adelaide, Australia, ³Department of Epidemiology and Public Health, University College London, London, United Kingdom

ABSTRACT

Dentistry has always been an under-resourced profession. There are three main issues that dentistry is facing in the modern era. Firstly, how to rectify the widely acknowledged geographical imbalance in the demand and supply of dental personnel, secondly, how to provide access to primary dental care to maximum number of people, and thirdly, how to achieve both of these aims within the financial restraints imposed by the central and state governments. The trends of oral diseases have changed significantly in the last 20 years. The two of the most common oral diseases that affect a majority of the population worldwide, namely dental caries and periodontitis, have been proved to be entirely preventable. Even for life-threatening oral diseases like oral cancer, the best possible available treatment is prevention. There is a growing consensus that appropriate skill mix can prove very beneficial in providing these preventive dental care services to the public and aid in achieving the goal of universal oral health coverage. Professions complementary to dentistry (PCD) have been found to be effective in reducing inequalities in oral health, improving access and spreading the messages of health promotion across entire spectrum of socio-economic hierarchy in various studies conducted globally. This commentary provides a review of the effectiveness of skill mix in dentistry and a reflection on how this can be beneficial in achieving universal oral health care in India.

Keywords: Dentist, health systems, human resource, inequalities, oral health, skill mix

Healthcare systems are responsible for addressing the needs of respective populations without any discrimination. Estimates from Global Burden of Disease demonstrate that oral diseases affect 3.9 billion people and untreated dental caries (tooth decay) is the most prevalent morbid condition among all diseases.^[1] Furthermore, oral diseases significantly affect quality of life and its associated healthcare has a catastrophic effect on the public health budgets.^[1] Linkages between many oral diseases and chronic non communicable diseases are well documented^[2-7] and tooth loss has been reported of being associated with pre-mature mortality.^[1,8,9] Oral diseases may also impact social and psychological wellbeing, consequently leading to social isolation.^[10]

A multi-center study in India reported caries prevalence among 35-44 year olds to be in a range of 48% in Orissa and as high as 86% in Delhi and Maharashtra.^[11] This increase in prevalence

of dental caries is observed parallel to the rapid nutrition transition in the recent decades^[11-14] and may also be one of its consequences. A prevalence as high as 100% was reported for periodontal disease among 35-44 and 65-74 years old in Orissa and Rajasthan^[11] confirming that the public health burden of oral diseases is not only limited to developed countries but a significant problem for a rapidly developing country like India.

The lack of regular national surveillance of oral diseases in India limits estimation of the current prevalence and the study of the trends of oral diseases. However, considering the strength of evidence on sugar intake and dental caries,^[15] and the increasing sugar intake as a consequence of rapid urbanization and subsequent westernization of diet,^[14] epidemic proportions of dental caries can be predicted in near future. Furthermore, India is called as the “oral cancer capital” of the world attributed to its high intake of both smoked and smokeless tobacco products, strongly associated with oral neoplasms.^[16] Most of these highly prevalent oral diseases are largely preventable as they share common risk factors (tobacco, alcohol, unhealthy diet) with other life-threatening chronic diseases^[17] which can

Access this article online

Quick Response Code:



Website:
www.jfmpc.com

DOI:
10.4103/2249-4863.154632

Address for correspondence: Dr. Manu Raj Mathur,
Public Health Foundation of India, Plot No. 47, Sector 44,
Institutional Area, Gurgaon, Haryana - 122 002, India.
E-mail: manu.mathur@phfi.org

be reduced through various health promotion and preventive measures.^[18,19]

The expected rise in burden of oral diseases will pose a massive challenge to the health system in light of increasing unfair and unjust social inequalities in oral health, which have already raised global concerns. The demand for equitable distribution of oral health hence indicates the need to reflect on effective preventive strategies adopted in other countries. A comprehensive dental workforce with appropriate skill mix is one such strategy adopted to address these issues.^[20-24] “Skill mix” is a term used for description of the mix of posts, grades or occupations in any organization. The dental workforce consists of dental surgeons, public health dentists and dental auxiliaries or professionals complementary to dentistry (PCDs). Barnes in his critical review of reorientation of oral health services stated that “*Dental services are now becoming polarized with need towards self-care and minimal intervention on one hand, and high technology care on the other, thus increasing the need to redefine the dentist to auxiliary ratio.*”^[25]

The capacity to which PCDs have been utilized in other health systems is well documented in the literature.^[20,21,23,24] A systematic review^[26] identified five main themes of oral health and examined their competence across these areas. The five identified themes were diagnosis, technical competence, oral health promotion, acceptability and productivity. In regard to the diagnosis of oral diseases, the majority of the studies reported that PCDs who have received appropriate training could diagnose and screen the most prevalent oral diseases similar to a dentist. The review highlighted that 40 of 41 selected studies reporting quality of performing most technical procedures was similar between PCDs and dentists. The review further reported that patients found working by PCDs acceptable and employing of PCDs increased the productivity.^[26] This shows that PCDs are competent to address the basic oral health needs at a population level.

Historically, the first dental college in India was established in 1920 and the recent statistics show that currently 301 colleges offer 25,270 student positions annually.^[27] The underlying reason for such a massive expansion could be the objective of serving oral health care needs of phenomenally increasing Indian population. However, the alarming urban–rural inequalities in dentist to population ratio is quite evident and well documented in the literature.^[27-32] This non-proportional distribution of dental workforce raises concerns as the healthcare needs of rural and severely deprived residents might be ignored while those of urban residents are satisfied. Consequently, social inequalities in oral health by area of residence may increase phenomenally.

PCDs provide a great opportunity to bridge this gap by both providing basic treatment needs and also by carrying out oral health promotion activities in rural areas. This becomes very vital as most of the oral conditions are largely preventable and can be well managed through effective health promotion and acting before they occur. However, this opportunity is untapped as not even one third of all dental colleges train PCDs reflecting the lack of an appropriate

skill mix in India.^[33,34] As dentistry is already an under-resourced profession in India, the ever increasing demands for oral health care is creating an unprecedented pressure on state and central governments to provide equitable health care. Preventive dental services like “atraumatic restorations,” developing oral hygiene skills, tobacco cessation and the basic clinical procedures can be well delivered by PCDs, while the complex clinical procedures that require the presence of a dental surgeon can be taken care through a referral system.^[20-24,26,34] These preventive services would not only reduce the burden of oral diseases, but would also reduce the increasing pressure on the dental profession in India.^[35]

The inadequacy in primary care services for oral health is also highlighted in Universal Health Coverage report of the Planning Commission of India, which may affect the Indian ambition to have universal health coverage.^[35] It is quite evident that a dental equivalent of “inverse care law” operates in India which means that those who require the dental health care most are the ones least likely to receive it.^[36] In conclusion, an appropriate skill mix to carry out more preventative and therapeutic work and an increase in the number of dental auxiliaries is urgently required in India to provide more appropriate and cost-effective care to all population sub-groups across India.

References

1. Marcenes W, Kassebaum NJ, Flaxman E, Naghavi A, Lopez M, Murray A, et al. Global burden of oral conditions in 1990-2010: A systematic analysis. *J Dent Res* 2013;92:592-7.
2. Desvarieux M, Demmer RT, Rundek T, Boden-Albala B, Jacobs DR, Papapanou PN et al. Relationship between periodontal disease, tooth loss, and carotid artery plaque: The Oral Infections and Vascular Disease Epidemiology Study (INVEST). *Stroke* 2003;34:2120-5.
3. Holmlund A, Holm G, Lind L. Severity of periodontal disease and number of remaining teeth are related to the prevalence of myocardial infarction and hypertension in a study based on 4,254 subjects. *J Periodontol* 2006;77:1173-8.
4. Joshipura K. The relationship between oral conditions and ischemic stroke and peripheral vascular disease. *J Am Dent Assoc* 2002;133 Suppl: 23S-30S.
5. Okoro CA, Balluz LS, Eke PI, Ajani UA, Strine TW, Town M, et al. Tooth loss and heart disease: Findings from the Behavioral Risk Factor Surveillance System. *Am J Prev Med* 2005;29 (5 Suppl 1):50-6.
6. Peres MA, Tsakos G, Barbato PR, Silva DA, Peres KG. Tooth loss is associated with increased blood pressure in adults - A multidisciplinary population-based study. *J Clin Periodontol* 2012;39:824-33.
7. Volzke H, Schwahn C, Dorr M, Aumann N, Felix SB, John U, et al. Inverse association between number of teeth and left ventricular mass in women. *J Hypertens* 2007;25:2035-43.
8. Abnet CC, Dawsey SM, Taylor PR, Qiao YL, Dong ZW, Mark SD. Tooth loss is associated with increased risk of total death and death from upper gastrointestinal cancer, heart disease, and stroke in a Chinese population-based cohort. *Int J Epidemiol* 2005;34:467-74.
9. Watt RG, Tsakos G, de Oliveira C, Hamer M. Tooth loss and cardiovascular disease mortality risk—results from the Scottish Health Survey. *PloS one* 2012;7:e30797.

10. Watt RG. Strategies and approaches in oral disease prevention and health promotion. *Bull World Health Organ* 2005;83:711-8.
11. Shah N, Pandey R, Duggal R, Mathur U, Kumar R. Oral health survey in India: A report of multicentric study, WHO – Oral health survey 2004. Geneva, Switzerland: World Health Organization; 2005.
12. Misra A, Singhal N, Sivakumar B, Bhagat N, Jaiswal A, Khurana L. Nutrition transition in India: Secular trends in dietary intake and their relationship to diet-related non-communicable diseases. *J Diabetes* 2011;3:278-92.
13. Shetty PS. Nutrition transition in India. *Public Health Nutr* 2002;5:175-82.
14. Popkin BM, Horton S, Kim S, Mahal A, Shuigao J. Trends in diet, nutritional status, and diet-related noncommunicable diseases in China and India: The economic costs of the nutrition transition. *Nutr Rev* 2001;59:379-90.
15. Moynihan PJ. The role of diet and nutrition in the etiology and prevention of oral diseases. *Bull World Health Organ* 2005;83:694-9.
16. GATS. Global Adult Tobacco Survey (GATS) India Report 2009-2010. Ministry of Health and Family Welfare, India, 2010.
17. Watt RG, Sheiham A. Integrating the common risk factor approach into a social determinants framework. *Community Dent Oral Epidemiol* 2012;40:289-96.
18. Monse B, Benzian H, Naliponguit E, Belizario V, Schratz A, van Palenstein Helderma W. The Fit for School Health Outcome Study-a longitudinal survey to assess health impacts of an integrated school health programme in the Philippines. *BMC Public Health* 2013;13:256.
19. Nakre PD, Harikiran AG. Effectiveness of oral health education programs: A systematic review. *J Int Soc Prev Community Dent* 2013;3:103-15.
20. Gallagher JE, Kleinman ER, Harper PR. Modelling workforce skill-mix: How can dental professionals meet the needs and demands of older people in England? *Br Dent J* 2010;208:E6; discussion 116-7.
21. Gallagher JE, Lim Z, Harper PR. Workforce skill mix: Modelling the potential for dental therapists in state-funded primary dental care. *Int Dent J* 2013;63:57-64.
22. Gallagher JE, Wilson NH. The future dental workforce? *Br Dent J* 2009;206:195-9.
23. Wanyonyi KL, Radford DR, Gallagher JE. Dental skill mix: A cross-sectional analysis of delegation practices between dental and dental hygiene-therapy students involved in team training in the South of England. *Hum Resour Health* 2014;12:65.
24. Wilson NH, Shamsir ZA, Moris S, Slater M, Kok EC, Dunne SM, *et al.* Dental workforce development as part of the oral health agenda for Brunei Darussalam. *Int Dent J* 2013;63:49-55.
25. Barnes D, Tala H. Health manpower out of balance: Conflicts and prospects for oral health. Geneva: CIOMS; 1987.
26. Galloway J, Gorham J, Lambert M, Richards D, Russell D, Russell I, *et al.* The professionals complementary to dentistry: Systematic review and synthesis. London: University College London, Eastman Dental Hospital, Dental Team Studies Unit, 2002.
27. Jaiswal AK, Srinivas P, Suresh S. Dental manpower in India: Changing trends since 1920. *Int Dent J* 2014;64:213-8.
28. Elangovan S, Allareddy V, Singh F, Taneja P, Karimbux N. Indian dental education in the new millennium: Challenges and opportunities. *J Dent Educ* 2010;74:1011-6.
29. Halappa M, Naveen BH, Kumar S, Sreenivasa H. SWOT Analysis of Dental Health Workforce in India: A Dental alarm. *J Clin Diagn Res* 2014;8:Ze03-5.
30. Rao KD, Ryan M, Shroff Z, Vujcic M, Ramani S, Berman P. Rural clinician scarcity and job preferences of doctors and nurses in India: A discrete choice experiment. *PloS one* 2013;8:e82984.
31. Singh A, Purohit BM. Addressing oral health disparities, inequity in access and workforce issues in a developing country. *Int Dent J* 2013;63:225-9.
32. Vundavalli S. Dental manpower planning in India: Current scenario and future projections for the year 2020. *Int Dent* 2014;64:62-7.
33. Mahal AS, Shah N. Implications of the growth of dental education in India. *J Dent Educ* 2006;70:884-91.
34. Mohanty VR, Rajesh GR, Aruna DS. Role of dental institutions in tobacco cessation in India: Current status and future prospects. *Asian Pac J Cancer Prev* 2013;14:2673-80.
35. (HLEG) HLEG. High Level Expert Group Report on Universal Health Coverage for India. Delhi: Planning Commission of India, 2011.
36. Hart JT. The inverse care law. *Lancet* 1971;1:405-12.

How to cite this article: Mathur MR, Singh A, Watt R. Addressing inequalities in oral health in India: Need for skill mix in the dental workforce. *J Fam Med Primary Care* 2015;4:200-2.

Source of Support: Nil. **Conflict of Interest:** None declared.