

Improvement of Preimplantation Development of *In Vitro*-Fertilized Bovine Zygotes by Glucose Supplementation to a Chemically Defined Medium

Nobutada SAKAGAMI^{1)*}, Osamu NISHINO²⁾, Satoshi ADACHI³⁾, Hidenobu UMEKI³⁾, Hiroko UCHIYAMA⁴⁾, Kyoko ICHIKAWA⁵⁾, Kazuhisa TAKESHITA⁶⁾, Etsushi KANEKO¹⁾, Kiyoshi AKIYAMA¹⁾, Shuji KOBAYASHI⁷⁾ and Hiromichi TAMADA⁸⁾

¹⁾Kanagawa Prefectural Livestock Industry Technology Center, Ebina, Kanagawa 243-0417, Japan

²⁾Nara Prefectural Livestock Technology Center, Nara 633-1302, Japan

³⁾Ooita Prefectural Agriculture, Forestry and Fisheries Research Center Livestock Research Institute, Ooita 878-0201, Japan

⁴⁾Miyazaki Prefectural Livestock Research Institute, Miyazaki 889-4411, Japan

⁵⁾Kochi Prefectural Livestock Research Institute, Kochi 789-1233, Japan

⁶⁾Yamaguchi Prefectural Livestock Experiment Station, Yamaguchi 759-2221, Japan

⁷⁾National Livestock Breeding Center, Fukushima 961-8511, Japan

⁸⁾Department of Advanced Pathobiology, Graduate School of Life and Environmental Sciences, Osaka Prefecture University, Sakai, Osaka 598-8531, Japan

(Received 30 October 2013/Accepted 11 June 2014/Published online in J-STAGE 30 June 2014)

ABSTRACT. The influences of glucose supplementation on early development of bovine embryos in BSA-free synthetic oviduct fluid were examined. Among the groups supplemented with 1.5, 2.0, 4.0 or 5.6 mM glucose either at 0, 72 or 144 hr after fertilization, blastocysts yield significantly increased in the group supplemented with 4.0 mM glucose 144 hr after fertilization compared to the controls without glucose supplementation. The results suggest that appropriate amounts of glucose supplemented to the medium at the specific stage of embryo culture may be useful for the production of bovine blastocysts.

KEY WORDS: bovine, chemically defined medium, embryo, embryo culture, glucose

doi: 10.1292/jvms.13-0543; *J. Vet. Med. Sci.* 76(10): 1403-1405, 2014

In general, fetal bovine serum (FBS) is widely employed as a supplement in the media used for culturing *in vitro* fertilized bovine oocytes. Nevertheless, the use of FBS has disadvantages as it contains unknown factors, and the subsequent rate of development of the fertilized oocytes to the blastocyst stage is known to vary among serum lots. Thus, there is a need for a reliable *in vitro* embryo culture protocol that uses a chemically defined medium and yields a high rate of good quality embryos. Many reports showed that the rate of development to the blastocyst stage is lower in chemically defined, serum-free media than in standard culture media containing serum [2, 9, 14] or BSA [1, 15]. Although we have reported significantly improved embryo development to the blastocyst stage by the supplementation of chemically-defined medium with epidermal growth factor (EGF) and insulin-like growth factor I (IGF-I) [14], blastocyst development in this modified chemically-defined medium was still lower than that obtained in medium supplemented with serum.

On the other hand, most of embryo culture systems include glucose as the main energy source. The importance of

glucose as an energy substrate for bovine zygotes has been demonstrated in several studies [4, 7], and its inhibitory effect on bovine embryo development at the early stages has also been described [1, 6, 8, 12]. Although the optimum timing and doses of the glucose addition to the culture media have been addressed in several reports [4, 5, 7, 11], there is limited knowledge on the effect of glucose supplementation on *in vitro* development of bovine embryos under chemically defined conditions [5]. Since the development of embryos to the blastocyst stage in protein-free media was slower than that in media supplemented with serum [2], it is necessary to determine the optimum timing and concentration of glucose supplementation when a chemically defined medium is used. Therefore, the objective of the present study was to determine the optimum timing and concentration of glucose supplementation to a protein free, chemically defined synthetic oviduct fluid (SOF) in order to maximize *in vitro* development of bovine embryos to the blastocyst stage.

The animal experiments in this study were approved by the Institutional Animal Experiment Committee of Kanagawa Prefectural Livestock Industry Technology Center. The experiments were performed in six separate laboratories, using the same experimental reagents, such as the culture media and frozen semen lot, and the results were aggregated. Preliminary experiments were conducted previously at each laboratory site using a protein-free SOF supplemented with 1 mg/ml polyvinyl alcohol (PVA) for embryo culture, and there was no significant difference in the rates of blastocyst development among these laboratories.

*CORRESPONDENCE TO: SAKAGAMI, N., Kanagawa Prefectural Livestock Industry Technology Center, Ebina, Kanagawa 243-0417, Japan. e-mail: sakagami.semi@pref.kanagawa.jp

©2014 The Japanese Society of Veterinary Science

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives (by-nc-nd) License <<http://creativecommons.org/licenses/by-nc-nd/3.0/>>.

Table 1. Effects of timing and concentration of glucose supplementation in SOFaa-E+I medium on development to the blastocyst stage of bovine embryos observed 192 hr after fertilization

Timing of glucose supplementation ^{b)} (hr)	No. of oocytes (replicates)	% of blastocyst at each concentration of glucose (mM) ^{a)}				
		0	1.5	2.0	4.0	5.6
0	1205 (9)	19.2 ± 2.5 ^{c)}	24.0 ± 3.1 ^{c)}	20.2 ± 3.5 ^{c)}	12.6 ± 3.6 ^{cd)A)}	3.1 ± 1.7 ^{d)A)}
72	549 (6)	19.6 ± 4.1	19.7 ± 3.7	20.1 ± 3.7	24.7 ± 5.1 ^{AB)}	16.9 ± 3.4 ^{B)}
144	600 (6)	16.9 ± 2.0 ^{c)}	22.9 ± 3.2 ^{cd)}	27.8 ± 3.7 ^{cd)}	31.3 ± 3.7 ^{d)B)}	29.0 ± 2.3 ^{cd)C)}

a) Data are expressed as the mean ± S.E.M., b) Time of glucose supplementation after *in vitro* fertilization, c–d) Values in the same row with different superscripts are significantly different ($P < 0.05$), A–C) Values in the same column with different superscripts are significantly different ($P < 0.05$).

Bovine ovaries were collected at a slaughterhouse and transported to the laboratory in saline. Cumulus-oocyte complexes (COCs) were collected from follicles with a diameter of approximately 3–5 mm using a 10-ml syringe with an 18-gauge needle. COCs with more than one layer of cumulus cells closely adhered to the zona pellucida, and an oocyte with homogeneous cytoplasm was selected. The selected COCs were washed three times with a maturation medium consisting of TCM-199 (12340-030, Gibco Laboratories, Grand Island, NY, U.S.A.) supplemented with 0.02 AU (Armour Units) porcine follicle stimulating hormone (Antrin R10, Kyoritsu Seiyaku, Tokyo, Japan), 5% FBS (SH30070, Hyclone Laboratories, Logan, UT, U.S.A.), 1 µg/ml estradiol-17β (E1132, Sigma-Aldrich, St. Louis, MO, U.S.A.) and 0.2 mM sodium pyruvate (P2256, Sigma-Aldrich). Fifteen to twenty oocytes were placed in a 100-µl drop of the maturation medium, which was then covered with liquid paraffin (26137-85, Nacalai Tesque, Kyoto, Japan) and incubated in a CO₂ incubator (38.5°C, 5% CO₂ in humidified air) for 20–22 hr. *In vitro* fertilization (IVF) was performed according to our previous report [14]. In this study, semen samples collected from a single Japanese Black bull and frozen on the same day were used. They were thawed in water at 37°C and washed twice by centrifugation (460 × g, 5 min) with 6 ml IVF medium (IVF 100; Research Institute for the Functional Peptides, Yamagata, Japan). Then, the sperm pellet was resuspended to a concentration of 1 × 10⁷ spermatozoa/ml, and the suspension was used to make fertilization drops (100-µl) with a final concentration of 5.0 × 10⁶ spermatozoa/ml. The COCs were washed three times with 350-µl droplets of IVF 100, transferred into the fertilization drops and incubated for 6 hr at 38.5°C under a humidified atmosphere of 5% CO₂ in air. Fifteen oocytes were placed in each fertilization drop. After IVF, the cumulus cells were removed from the oocytes by pipetting through a narrow-bore glass pipette. Cumulus-free presumptive zygotes were placed in a culture medium which was glucose-free modified synthetic oviduct fluid [15] supplemented with 2% (v/v) Basal Medium Eagle essential amino acids (B6766, Sigma-Aldrich), 1% (v/v) Minimum Essential Medium amino acids (11140-050, Gibco) (SO-Faa), 1 mg/ml PVA (P8136, Sigma-Aldrich), 100 ng/ml EGF (E4157, Sigma-Aldrich) and 50 ng/ml IGF-I (I3769, Sigma-Aldrich) (SOFaa-E+I). Twenty presumptive zygotes were incubated in each 100-µl drop covered with liquid paraffin at 38.5°C in 5% CO₂, 5% O₂ and 90% N₂ until 192 hr post-fertilization. Glucose was added to the SOFaa-E+I medium

at different concentrations (0, 1.5, 2.0, 4.0 and 5.6 mM) at different time points (0, 72 and 144 hr) after fertilization, and the developmental rates of embryos were investigated. During glucose addition, the half amount of the culture media was aspirated from each drop with a micropipette under a stereo microscope, and then, preincubated fresh culture media containing the double concentration of glucose were introduced. The percentages of embryos developing to blastocysts were subjected to arcsine transformation before statistical analysis. A two-way analysis of variance (2-way ANOVA) was performed using SPSS (SPSS 16.0 J, User's Guide, SPSS, Tokyo, Japan) followed by Tukey's HSD test. Data are presented as mean ± S.E.M., and a *P* value of less than 0.05 was considered statistically significant.

The results of embryo development are shown in Table 1. There was a significant interaction between the concentration and the timing of glucose supplementation in the developmental rate of blastocysts ($P < 0.05$). The percentage of blastocysts cultured in SOFaa-E+I medium supplemented with 4.0 mM glucose 144 hr after fertilization was significantly increased when compared with controls without glucose supplementation. Moreover, the blastocysts yield in the group supplemented with 4.0 mM glucose 144 hr after fertilization was significantly increased when compared with that in the group supplemented with 4.0 mM glucose 0 hr after fertilization. These results indicated that 4.0 mM glucose supplemented at 144 hr after fertilization is useful for embryo culture in this chemically defined medium. On the other hand, in previous studies using media containing serum, the optimum dose (1.5 or 5.56 mM) and timing (72, 96 or 120 hr after fertilization) of glucose supplementation varied among reports [4, 5, 7, 8]. The type of medium and serum could influence the effect of glucose supplementation on the embryo growth.

When compared among the groups supplemented with various amounts of glucose 0 hr after fertilization, the developmental rate of blastocysts decreased with higher concentration of glucose (4.0 or 5.6 mM), and percentages of blastocysts yield in the group supplemented with 5.6 mM glucose decreased significantly when compared with those with 0, 1.5 or 2.0 mM glucose.

In bovine embryos, Javed and Wright [3] described that glucose does not affect cell growth before the morula stage, and Matsumoto *et al.* [7] reported that glucose supplementation immediately after fertilization at the concentration of 3 mM or less to modified SOF medium containing 5%

superovulated cow serum did not affect blastocyst formation. However, there are some reports showing that addition of high concentration of glucose immediately after *in vitro* fertilization was harmful for early embryonic development [1, 6, 9, 13]. These findings are consistent with our results. Although the reason why high concentrations of glucose obstruct embryonic development is not clear, Rieger [10] showed that glucose increases the intracellular level of oxygen radicals harmful for mammalian embryo development.

Although glucose obstructs the development of the embryos when supplemented in the culture in high concentrations during the early stages, it has important roles on energy metabolism after the compacted morula stage and during blastulation [4, 8, 16]. Prior to the present study, we observed the timing of formation of blastocoels using time-lapse photography as a preliminary test, and blastocoels of embryos cultured in SOFaa-E+I medium could be identified from 132.8 ± 2.7 hr (Mean \pm S.E.M.) after fertilization (unpublished data). In the present study, the rate of embryos developing to the stage of blastocyst was highest in the group supplemented with 4.0 mM glucose 144 hr after fertilization, nearly at the timing of formation of the blastocoel. These results suggest that appropriate amounts of glucose at the later stage of the culture are effective to enhance embryo development. On the other hand, Holm *et al.* [2] reported that bovine embryos developed to early blastocysts at 129.0 ± 2.9 hr after fertilization in serum-free SOFaa medium and at 107.5 ± 3.1 hr after fertilization in SOFaa medium supplemented with serum and suggested that serum in the culture medium accelerated the growth of embryos. It has been reported that in a culture medium containing serum, development of embryos was improved significantly when glucose was supplemented 72 hr [11] after *in vitro* fertilization, which is earlier than the effective timing of glucose supplementation for blastocyst development (144 hr after fertilization) in this study. These findings suggest that the presence of serum in the culture medium might affect the timing of effective glucose supplementation for the embryo development due to its accelerating effect on the embryo growth.

In conclusion, glucose may be useful for blastocyst production, if it is used in appropriate timing and concentration in chemically defined culture media.

ACKNOWLEDGMENTS. The authors would like to thank Dr. Y. Hashiyada at the National Livestock Breeding Center for critical suggestions on the manuscripts and Dr. T. Somfai at the National Institute of Livestock and Grassland Science for proofreading.

REFERENCES

- Furness, C., de Matos, D., Martínez, A. and Matkovic, M. 1997. Effect of glucose on embryo quality and post-thaw viability of *in-vitro*-produced bovine embryos. *Theriogenology* **47**: 481–490. [Medline] [CrossRef]
- Holm, P., Booth, P. J. and Callesen, H. 2002. Kinetics of early *in vitro* development of bovine *in vivo*- and *in vitro*-derived zygotes produced and/or cultured in chemically defined or serum-containing media. *Reproduction* **123**: 553–565. [Medline] [CrossRef]
- Javed, M. H. and Wright, R. W. Jr. 1991. Determination of pentose phosphate and emben-meyerhof pathway activities in bovine embryos. *Theriogenology* **35**: 1029–1037. [Medline] [CrossRef]
- Kim, J. H., Funahashi, H., Niwa, K. and Okuda, K. 1993. Glucose requirement at different developmental stage of *in vitro* fertilized bovine embryos cultured in semi-defined medium. *Theriogenology* **39**: 875–886. [Medline] [CrossRef]
- Kim, J. H., Niwa, K., Lim, J. M. and Okuda, K. 1993. Effects of phosphate, energy substrates, and amino acids on development of *in vitro*-matured, *in vitro*-fertilized bovine oocytes in a chemically defined, protein-free culture medium. *Biol. Reprod.* **48**: 1320–1325. [Medline] [CrossRef]
- Konishi, M., Itakura, H. and Aoyagi, Y. 1994. Effect of glucose in CR1aa on the development into blastocyst of *in vitro* fertilized bovine oocytes co-cultured with cumulus cells. *J. Reprod. Dev.* **40**: j59–j64 (in Japanese). [CrossRef]
- Matsumoto, M., Otoi, T. and Suzuki, T. 1999. Effect of glucose and lactate on development of *in vitro* produced bovine embryos in a modified synthetic oviduct fluid medium. *J. Mamm. Ova Res.* **16**: 73–76 (in Japanese). [CrossRef]
- Matsuyama, K., Miyakoshi, H. and Fukui, Y. 1993. Effect of glucose levels during the *in vitro* culture in synthetic oviduct fluid medium on *in vitro* development of bovine oocytes matured and fertilized *in vitro*. *Theriogenology* **40**: 595–605. [Medline] [CrossRef]
- Pinyopummintr, T. and Bavister, B. D. 1991. *In vitro*-matured/*in vitro*-fertilized bovine oocytes can develop into morulae/blastocysts in chemically defined, protein-free culture media. *Biol. Reprod.* **45**: 736–742. [Medline] [CrossRef]
- Rieger, D. 1992. Relationships between energy metabolism and development of early mammalian embryos. *Theriogenology* **37**: 75–93. [CrossRef]
- Robl, J. M., Dobrinsky, J. R. and DUBY, R. T. 1991. The effect of protein supplements, phosphate and glucose on the *in vitro* development of IVM-IVF bovine oocytes. *Theriogenology* **35**: 263. [CrossRef]
- Rosenkrans, C. F. Jr., Zeng, G. Q., MCNamara, G. T., Schoff, P. K. and First, N. L. 1993. Development of bovine embryos *in vitro* as affected by energy substrates. *Biol. Reprod.* **49**: 459–462. [Medline] [CrossRef]
- Rubessa, M., Boccia, L., Campanile, G., Longobardi, V., Albarella, S., Tateo, A., Zicarelli, L. and Gasparrini, B. 2011. Effect of energy source during culture on *in vitro* embryo development, resistance to cryopreservation and sex ratio. *Theriogenology* **76**: 1347–1355. [Medline] [CrossRef]
- Sakagami, N., Umeki, H., Nishino, O., Uchiyama, H., Ichikawa, K., Takeshita, K., Kaneko, E., Akiyama, K., Kobayashi, S. and Tamada, H. 2012. Normal calves produced after transfer of embryos cultured in a chemically defined medium supplemented with epidermal growth factor and insulin-like growth factor I following ovum pick up and *in vitro* fertilization in Japanese Black cows. *J. Reprod. Dev.* **58**: 140–146. [Medline] [CrossRef]
- Takahashi, Y. and First, N. L. 1992. *In vitro* development of bovine one-cell embryos: Influence of glucose, lactate, pyruvate, amino acids and vitamins. *Theriogenology* **37**: 963–978. [Medline] [CrossRef]
- Thompson, J. G., Partridge, R. J., Houghton, F. D., Cox, C. I. and Leese, H. J. 1996. Oxygen uptake and carbohydrate metabolism by *in vitro* derived bovine embryos. *J. Reprod. Fertil.* **106**: 299–306. [Medline] [CrossRef]