



Rapid Changes in Serum Testosterone in Men With Newly Diagnosed Type 2 Diabetes With Intensive Insulin and Metformin

Yun Hu,¹ Bo Ding,¹ Yun Shen,¹
Reng-Na Yan,¹ Feng-Fei Li,¹ Rui Sun,¹
Ting Jing,¹ Kok-Onn Lee,² and Jian-Hua Ma¹

Diabetes Care 2021;44:1059–1061 | <https://doi.org/10.2337/dc20-1558>

OBJECTIVE

To investigate the effect of metformin on testosterone levels in men with type 2 diabetes mellitus (T2DM).

RESEARCH DESIGN AND METHODS

Seventy men with newly diagnosed drug-naïve T2DM and HbA_{1c} >9.0% (75 mmol/mol) were treated with intensive insulin pump therapy for 5 days to achieve glucose normalization. They were randomized to control (continued on intensive insulin only) and metformin (plus metformin) groups (1:1) for 1 month. Testosterone was measured at baseline, randomization, and after 1-month treatment.

RESULTS

Total, free, and bioavailable testosterone increased significantly within 5 days (all $P < 0.001$). After 1 month, compared with the control group, the metformin group had lower total (12.7 vs. 15.3 nmol/L), free (0.20 vs. 0.24 nmol/L), and bioavailable (4.56 vs. 5.31 nmol/L) testosterone (all $P < 0.05$).

CONCLUSIONS

In men with T2DM, 1-month oral metformin may decrease serum testosterone levels independent of blood glucose control. The effects of long-term metformin on testosterone in men need further study.

Metformin is the most commonly prescribed oral medication in type 2 diabetes mellitus (T2DM). It is inexpensive and effective and may have additional benefits in cancer (1), cardiovascular disease, and other conditions (2).

Low testosterone levels are common in men with T2DM (3). They have been attributed to greater insulin resistance (4) and are associated with higher mortality rates (5). The mechanism remains unclear (6). We previously reported that testosterone levels were significantly lower in men with T2DM using metformin than in men not using metformin in a cross-sectional study (7).

In the current study, we performed a randomized controlled trial in men with newly diagnosed T2DM. While following recommended guidelines in China, treatment was initiated with intensive insulin pump therapy (8). We then compared the effect of the addition of metformin on serum testosterone during 1 month of further intensive insulin therapy. This short duration allowed the assessment of the effect of metformin before significant changes in weight and other factors that might affect blood testosterone.

¹Department of Endocrinology, Nanjing First Hospital, Nanjing Medical University, Jiangsu, China

²Department of Medicine, National University of Singapore, Singapore

Corresponding author: Jian-Hua Ma, majianhua196503@126.com

Received 24 June 2020 and accepted 11 January 2021

Clinical trial reg. no. NCT03982238, clinicaltrials.gov.

This article contains supplementary material online at <https://doi.org/10.2337/figshare.13574474>.

Y.H. and B.D. contributed equally to this article.

© 2021 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <https://www.diabetesjournals.org/content/license>.

RESEARCH DESIGN AND METHODS

We recruited men with newly diagnosed drug-naïve T2DM aged 18–60 years with an $HbA_{1c} > 9.0\%$ (75 mmol/mol) (8). We excluded patients on treatment with lipid-lowering or antihypertensive medications or with any condition affecting testosterone levels.

All patients received insulin pump treatment as inpatients until the blood glucose normalized (premeal 3.9–7.0 mmol/L, postprandial 3.9–10.0 mmol/L, $>80\%$ of values within the range), usually within 3–5 days. The patients were then randomized into control (insulin only) and metformin (insulin + metformin) groups and continued on intensive insulin therapy for 1 month (Supplementary Fig. 1). The original protocol was designed with three groups: control, metformin, and dapagliflozin. Because of funding reasons, the dapagliflozin arm was stopped very soon after recruitment began. Metformin (Merck Serono) was started at 1,500 mg daily. Patients were discharged and seen as outpatients every 3 days to adjust the insulin dose according to the preceding day's 6-point self-monitoring of blood glucose values to maintain near-normal blood glucose. If gastrointestinal symptoms were present, metformin dosage was reduced to 1,000 mg/day and further reduced if symptoms persisted. Patients with a final dose of metformin <500 mg daily or who did not meet the criteria for three consecutive normal range self-monitoring of blood glucose records were withdrawn.

Weight, HbA_{1c} , blood lipids, fasting blood glucose, C-peptide, total testosterone (TT), and sex hormone binding globulin were measured at baseline, randomization, and 30 ± 2 days after randomization (study end). The International Index of Erectile Function (IIEF-5) questionnaire was assessed at baseline and study end. Free testosterone (FT) and bioavailable testosterone (Bio-T) (9) were calculated.

Differences between the groups were examined using the Student unpaired *t* test or the Mann-Whitney *U* test. The data before and after treatment were analyzed by Student paired *t* test or Wilcoxon test. Categorical data were examined with χ^2 test.

RESULTS

Seventy patients completed the study, with 35 in the control group and 35 in the

metformin group. Baseline characteristics were similar between the groups (Supplementary Table 1). Serum testosterone increased significantly after 5 days of intensive insulin therapy in all the patients (baseline vs. randomization: TT 12.07 ± 4.68 vs. 15.35 ± 5.97 nmol/L, FT 0.19 ± 0.08 vs. 0.24 ± 0.10 nmol/L, Bio-T 4.51 ± 1.34 vs. 5.39 ± 1.68 nmol/L, all $P < 0.001$) and each group (Fig. 1 and Supplementary Table 2).

TT, FT, and Bio-T levels significantly decreased after 1 month only in the metformin group compared with the control group ($P < 0.05$) (Fig. 1A–C), and metformin caused superior reductions in TT, FT, and Bio-T versus the control condition (estimated treatment differences: -2.29 [95% CI $-4.10, -0.48$], -0.04 [$-0.08, -0.01$], and -0.73 [$-1.31, -0.15$] nmol/L, $P = 0.014, 0.012,$ and 0.014 , respectively) (Supplementary Table 3). TT values calculated for individual patients as a percentage change from baseline also decreased in the metformin group compared with the control group (-27.4% [95% CI $-47.2, -7.6$], $P = 0.007$) (Fig. 1D).

After adjustment for age, blood lipids, body weight, HOMA2 of insulin resistance, daily metformin, and insulin dose at the end point, only the daily metformin dose had a statistically significant correlation with Δ TT in the stepwise linear regression ($\beta = 0.300, P = 0.029$). The numbers of patients with

testosterone “insufficiency” (TT < 12 nmol/L [346 ng/dL]) (10) decreased after blood glucose normalization compared with baseline in both groups (both $P < 0.05$) (Supplementary Table 2) and were higher in the metformin group than in the control group (57.14 vs. 28.57%, $P = 0.029$) at the end point. The change of IIEF-5 scores did not differ between the two groups ($Z = -0.366, P = 0.715$), despite the changes in testosterone.

CONCLUSIONS

We have demonstrated that 5 days of intensive insulin therapy and normalization of blood glucose increases testosterone significantly in men with newly diagnosed T2DM. We have also shown for the first time in a randomized controlled study that a further 1 month of metformin treatment decreases testosterone significantly in men with T2DM.

This effect of metformin should not be totally unexpected because metformin treatment is associated with lower testosterone concentrations in women with polycystic ovary syndrome (11), despite their lower baseline levels. Studies in men were surprisingly uncommon. We found only one previous study that showed higher semen volume, higher sperm counts, and higher (but not statistically significant) serum testosterone in men with type 1 diabetes on insulin only compared with men with T2DM treated with metformin (12). Other




Figure 1—A–C: TT, FT, and Bio-T levels at baseline (preinsulin therapy), randomization when blood glucose normalized after 5 days of intensive insulin, and the end point after 1-month therapy with intensive insulin with or without metformin. D: Baseline testosterone concentrations calculated individually and expressed as 100%, with the percentage change of TT at randomization and after 1 month of treatment with or without metformin. Data are mean \pm SE. ****** $P < 0.001$, randomization vs. end point; **#** $P < 0.05$, metformin group vs. control group.

previous studies that we are aware of on the effect of metformin on testosterone in men were confounded by changes in blood glucose and body weight (13,14). These studies concluded that metformin increased serum testosterone and attributed this to the improvement in blood glucose control and weight loss.

Poor diabetes control can lead to low testosterone levels and male sexual dysfunction (10). Our results demonstrate for the first time that the blood glucose normalization with intensive insulin treatment was able to increase blood testosterone even within a short period of 5 days. Our study demonstrates that the effect of metformin on testosterone may be independent of blood glucose and weight change because both groups did not differ in blood glucose control and body weight before and after therapy (Supplementary Table 1).

Our 1-month study has some limitations. The longer-term effect of metformin on testosterone and sexual function remains uncertain. In most parts of the world, intensive insulin is not recommended or used as initial treatment for newly diagnosed diabetes, and thus the rapid initial improvement and its subsequent decrease in testosterone with metformin may not be easily discernable or distinguishable from other effects of metformin on weight and insulin resistance. We did not measure luteinizing hormone/follicle-stimulating hormone to determine whether any subjects had secondary hypogonadism, and we lacked a placebo control; these may influence subjective IIEF-5 scores, although none of the investigators were aware of the testosterone values.

Our preliminary results suggest that the long-term effect of metformin should be investigated, with a placebo control, to determine whether measurement of testosterone concentrations may be

useful in men on metformin. We are reminded of the reports on lower vitamin B12 levels in patients on long-term metformin, which are now accepted and therefore monitored (15).

In conclusion, intensive insulin therapy can increase testosterone levels rapidly. With 1-month near normalization of blood glucose, metformin may reduce circulating testosterone levels significantly. The long-term effect of metformin treatment on testosterone needs further study.

Acknowledgments. The authors thank the participants for their involvement and the members of the endocrinology department of Nanjing First Hospital for their support.

Funding. This study was supported by the National Key R&D Program of China (No. 2018YFC1314103), the National Natural Science Foundation of China (No. 81870563) and the Nanjing Municipal Science and Technology Project (No. 201803011).

Duality of Interest. No potential conflicts of interest relevant to this article were reported.

Author Contributions. Y.H. carried out the statistical analysis and wrote the first draft of the manuscript. B.D., Y.S., R.-N.Y., R.S., and T.J. collected and collated data. F.-F.L. and K.-O.L. corrected and approved the revisions and final version of the manuscript. J.-H.M. is responsible for the conception, funding, and design of the study. J.-H.M. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

References

- Margel D, Urbach DR, Lipscombe LL, et al. Metformin use and all-cause and prostate cancer-specific mortality among men with diabetes. *J Clin Oncol* 2013;31:3069–3075
- Holman RR, Paul SK, Bethel MA, Matthews DR, Neil HA. 10-year follow-up of intensive glucose control in type 2 diabetes. *N Engl J Med* 2008;359:1577–1589
- Grossmann M. Low testosterone in men with type 2 diabetes: significance and treatment. *J Clin Endocrinol Metab* 2011;96:2341–2353
- Dhindsa S, Ghanim H, Batra M, et al. Insulin resistance and inflammation in hypogonadotropic

hypogonadism and their reduction after testosterone replacement in men with type 2 diabetes. *Diabetes Care* 2016;39:82–91

5. Muraleedharan V, Marsh H, Kapoor D, Channer KS, Jones TH. Testosterone deficiency is associated with increased risk of mortality and testosterone replacement improves survival in men with type 2 diabetes. *Eur J Endocrinol* 2013;169:725–733

6. Dhindsa S, Ghanim H, Batra M, Dandona P. Hypogonadotropic hypogonadism in men with diabetes. *Diabetes Care* 2018;41:1516–1525

7. Ding B, Sun R, Zhai XF, et al. Association of high circulating testosterone with increased glycaemic variability in type 2 diabetes: a cross-sectional study in China. *Diabetes Metab Res Rev* 2019;35:e3126

8. Weng J, Ji L, Jia W, et al.; Chinese Diabetes Society. Standards of care for type 2 diabetes in China. *Diabetes Metab Res Rev* 2016;32:442–458

9. de Ronde W, van der Schouw YT, Pols HA, et al. Calculation of bioavailable and free testosterone in men: a comparison of 5 published algorithms. *Clin Chem* 2006;52:1777–1784

10. Defeudis G, Mazzilli R, Gianfrilli D, Lenzi A, Isidori AM. The CATCH checklist to investigate adult-onset hypogonadism. *Andrology* 2018;6:665–679

11. Chou KH, von Eye Corleta H, Capp E, Spritzer PM. Clinical, metabolic and endocrine parameters in response to metformin in obese women with polycystic ovary syndrome: a randomized, double-blind and placebo-controlled trial. *Horm Metab Res* 2003;35:86–91

12. Zaidi AA, Khan MA, Sharif A, et al. Comparative study of sperm motility in metformin-using and insulin-dependent diabetics. *Biomed Res Ther* 2017;4:1388–1399

13. Morgante G, Tosti C, Orvieto R, Musacchio MC, Piomboni P, De Leo V. Metformin improves semen characteristics of oligo-terato-asthenozoospermic men with metabolic syndrome. *Fertil Steril* 2011;95:2150–2152

14. Nasrolahi O, Khaneshi F, Rahmani F, Razi M. Honey and metformin ameliorated diabetes-induced damages in testes of rat; correlation with hormonal changes. *Iran J Reprod Med* 2013;11:1013–1020

15. Aroda VR, Edelstein SL, Goldberg RB, et al.; Diabetes Prevention Program Research Group. Long-term metformin use and vitamin B12 deficiency in the Diabetes Prevention Program Outcomes Study. *J Clin Endocrinol Metab* 2016;101:1754–1761