

Disinfection by Ozone Microbubbles Can Cause Morphological Change of *Fusarium oxysporum* f. sp. *melonis* Spores

Masahiko Tamaki¹*, Fumiyuki Kobayashi², Hiromi Ikeura³, and Michio Sato¹

¹School of Agriculture, Meiji University, Kawasaki, Kanagawa 214-8571, Japan

²Faculty of Applied Life Science, Nippon Veterinary and Life Science University, Musashino, Tokyo 180-8602, Japan

³Faculty of Life and Environmental Science, Shimane University, Matsue, Shimane 690-8504, Japan

(Received on November 12, 2017; Revised on March 21, 2018; Accepted on April 23, 2018)

To investigate the difference in the disinfectant efficiency of ozone microbubbles (O₃MB) and ozone millibubbles (O₃MMB), the morphological change of the treated *Fusarium oxysporum* f. sp. *melonis* spores was observed with scanning and transmission electron microscopies (SEM and TEM). The disinfectant efficiency of O₃MB on *F. oxysporum* f. sp. *melonis* spores was greater than that of O₃MMB. On observation with SEM, it was revealed that morphological change of *F. oxysporum* f. sp. *melonis* spores was caused by O₃MB and O₃MMB, and damage to the spore surfaces by O₃MB occurred sooner than that by O₃MMB. On observation with TEM, it was furthermore confirmed that *F. oxysporum* f. sp. *melonis* spores treated with O₃MB induced wavy deformation of cell membrane and the intracellular change different from that with O₃MMB. Therefore, the greater disinfection efficiency of O₃MB was suggested to be caused due to the function of the MB in addition to the oxidative power of O₃.

Keywords : disinfection, *Fusarium oxysporum* f. sp. *melonis* spores, ozone microbubbles

Handling Associate Editor : Jeon, Junhyun

*Corresponding author.

Phone) +81-44-980-5276, FAX) +81-44-980-5276

E-mail) mtamaki@meiji.ac.jp

ORCID

<http://orcid.org/0000-0001-8316-725X>

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Articles can be freely viewed online at www.ppjonline.org.

Hydroponic culture is a plant cultivation technique in which plants are grown in a nutrient solution without soil. It has the potential for high crop productivity in a small area. However, if plant pathogens enter the solution, they can spread rapidly throughout the hydroponic culture facility and cause catastrophic damage. For this reason, disinfection of the solution is essential, although direct administration of pesticides into the solution has been prohibited at law (Ministry of Agriculture, Forestry and Fisheries, 2010). Therefore, it is desired to establish safe and effective alternative disinfection methods, and the various disinfection treatments have been investigated, including UV light, heat, the titanium dioxide photocatalytic reaction, and ozone (O₃) (Bando et al., 2008; Dannehl et al., 2016; Ehret et al., 2001; Igura et al., 2004; Koohakan et al., 2003; Ohtani et al., 2000; Runia, 1995). However, these treatments still are not put into any practical use in terms of the efficiency, treatment time, and running cost. Particularly, O₃ gas is effective disinfectant due to its strong oxidation power, although it is difficult to use in hydroponic cultures because of its extremely low solubility in water (0.105 g 100 ml⁻¹ (0°C)).

Recently, tiny bubbles less than 50 μm in diameter, called microbubble (MB), have been studied and used in many fields. They rise more slowly in water than millibubbles (MMB), which have diameters in the mm to cm range (Takahashi et al., 2003; Takahashi, 2005), and possess additional properties such as the interface charge, long stagnation, slow buoyancy, the shrinkage and the generation of free radicals by their collapsing other than dissolving power (Li and Tsuge, 2006; Li et al., 2009; Zheng et al., 2015). Previously, we focused on long retention time in water and the dissolving power of MB in addition to the strong oxidative power of O₃, and found that O₃MB were more effective than O₃MMB for the disinfection of *Fusarium oxysporum* f. sp. *melonis* spore and *Pectobacterium*

carotovorum subsp. *carotovorum* in nutrient solution for hydroponic culture (Kobayashi et al., 2011a, 2011b, 2011c, 2012). Furthermore, effective disinfection by O₃MB has been reported by other researchers (Chuaqedton et al., 2015; Inatsu et al., 2011), although it is still not clear about the exact mechanism of disinfection by O₃MB. In this study, the morphological change of *F. oxysporum* f. sp. *melonis* spores by O₃MB and O₃MMB was therefore observed by using scanning and transmission electron microscopies (SEM and TEM), and the difference was discussed.

F. oxysporum f. sp. *melonis* NBRC6385 suspension (approximately 1.0×10^7 spores ml⁻¹) was prepared in a manner similar to a previous report (Kobayashi et al., 2011a). For each experiment, 15 l of tap water were collected in a plastic cylindrical container (28 cm dia. × 48 cm height) and kept for 24 h at room temperature to remove chlorine. Water quality test paper (Nissan aquacheck 3; Nissan Chemical Industries, Ltd., Tokyo, Japan) was used to confirm that no residual chlorine remained. O₃ was generated by using an O₃ generator (ED-OG-A10, Ecode-sign Co. Ltd., Saitama, Japan) at a flow rate of 2.5 l min⁻¹. O₃MB and O₃MMB were generated by using a decompression type MB generator (20NEDO4S, Shigen-Kaihatsu Co., Ltd., Kanagawa, Japan) and a commercial air pump, respectively. The concentration of dissolved O₃ in both O₃MB and O₃MMB waters was set to 1.5 ppm at 15°C. The pH of O₃MB and O₃MMB waters was the same at 6.8 and was not changed before and after O₃MB and O₃MMB generation. *F. oxysporum* f. sp. *melonis* spores were added to 100 ml of the O₃MB and O₃MMB waters with final concentrations of 1.0×10^3 - 1.0×10^4 cfu ml⁻¹. Aliquots of the treated waters were collected after 0, 15, 30, 45, 60 and 120 s. Aliquots of 0.1 ml of the collected waters were plated on potato dextrose agar (Difco, Becton Dickinson, Franklin Lakes, NJ, USA) plates, and the plates were incubated at 30°C for 48 h. After incubation, the numbers of surviving spores were measured by counting the colonies formed on the plates. The detection limit was 10 cfu ml⁻¹. Each experiment was performed in duplicate.

F. oxysporum f. sp. *melonis* spores were collected from 5 ml of the suspension by filtration with a cartridge filter (Anotop 10, GE Healthcare UK Ltd., Buckinghamshire, UK). The sample on the cartridge filter was pre-fixed with 2.5% glutaraldehyde solution diluted with a phosphate buffer solution (PBS, pH 7.0). A filter ejected by decomposing the cartridge filter was washed with PBS (pH 7.0), post-fixed with 2% OsO₄ solution for 1 h, and then serially dehydrated for 20 min each in 50%, 70%, 80%, 90%, 95%, 99.5% and dehydrated ethanol. SEM observations were performed as follows: The dehydrated sample was

immersed in a mixture of t-butyl alcohol and dehydrated ethanol (1:1) for about 10 min, transferred to 100% t-butyl alcohol, freeze-dried with a freeze drier (ES-2030, Hitachi High Technologies Co., Tokyo, Japan), and OsO₄-coated with a OsO₄ coater (HPC-1SW, Vacuum device Inc., Mito, Japan) (the thickness of the coating was adjusted to 3 nm). Then the sample was observed with an SEM (JSM-6700F, JEOL Ltd., Akishima, Japan) operated at 3.0 kV. Nine *F. oxysporum* f. sp. *melonis* spores in the SEM photographs were selected at random and the widths was measured with a scale. Significant differences were evaluated by the ANOVA and Fisher's LSD using the Ekuseru-Toukei 2012 for Window statistical software (Social Survey Research Information Co., Ltd., Tokyo, Japan) ($P < 0.05$). TEM observations were performed as follows: The dehydrated samples were serially immersed for 2 h each in 1:1, 2:1, and 3:1 mixtures of Quetol-651 (Cosmo Bio Co., Ltd., Tokyo, Japan) and ethylene glycol diglycidyl ether, and then embedded in 100% Quetol-651 at 60°C. Ultra-thin sections (thickness 70-100 nm) were made from the embedded samples with an ultramicrotome (ULTRA CUT UCT, Leica Microsystems, Wetzlar, Germany). The ultra-thin sections were doubly electron-strained using 4% uranyl acetate for 12 min and lead nitrate for 5 min, and then observed with a TEM (JEM-2010, JEOL Ltd.) operated at 140 kV.

The survival rates of *F. oxysporum* f. sp. *melonis* spores in water treated with O₃MB and O₃MMB are shown in Fig. 1. The disinfectant efficiency of O₃MB on *F. oxysporum* f. sp. *melonis* spores was greater than that of O₃MMB, because the numbers of surviving spores after treatment with O₃MB and O₃MMB reached the detection limit at 45 s and 60 s, respectively. The result agreed with our previous study (Kobayashi et al., 2011a). Amount of hydroxyl




Fig. 1. Disinfection of *F. oxysporum* f. sp. *melonis* spores by O₃MB and O₃MMB. The data presented was the mean of duplicate.




Fig. 2. SEM images of *F. oxysporum* f. sp. *melonis* spores. (A) After treatment with O₃MMB for 30 s, (B) After treatment with O₃MMB for 60 s, (C) After treatment with O₃MB for 30 s, (D) After treatment with O₃MB for 60 s, (E) Non-treated.

radicals generated from O₃ is not enough to have a disinfectant effect (Cho et al., 2003). However, O₃MB may generate more hydroxyl radicals than O₃MMB, because the oxidation-reduction potential and iodine liberation is higher with O₃MB than with O₃MMB (Chuajedton et al., 2015). Furthermore, the use of MB enhances the formation of hydroxyl radicals due to more rapid O₃ decomposition (Tsuge et al., 2009), and the hydroxyl radicals generated from O₃MB accelerate the oxidative power (Chu et al., 2008). The high oxidative power of O₃MB contributes to the oxidative power of O₃, the reactivity of the hydroxyl radicals, the substantivity, ζ surface potential, mass-transfer coefficient, and use efficiency (Zheng et al., 2015). Therefore,

the greater disinfectant efficiency of O₃MB than O₃MMB is likely be due to these synergistic effects.

The SEM images of *F. oxysporum* f. sp. *melonis* spores treated with O₃MB and O₃MMB are shown in Fig. 2. The spores treated with O₃MMB for 30 s showed no obvious surface injury, although spores treated with O₃MMB for 180 s were deformed. On the other hand, spores treated with O₃MB showed obvious surface injury after 30 s and the spores were completely destroyed after 180 s. The widths of *F. oxysporum* f. sp. *melonis* spores treated with O₃MMB for 30 s were lower than those of non-treated spores, and then the spores swelled after 180 s (Fig. 3). However, the widths of spores treated with O₃MB for 30




Fig. 3. The widths of *F. oxysporum* f. sp. *melonis* spores in SEM images. NT: Non-treated. The results indicate the means with standard deviation of 9 spores.

s were the same as those of non-treated spores and diminished in size by 180 s. Furthermore, the TEM images indicated the appearance of liquid foam in the mid-regions of the *F. oxysporum* f. sp. *melonis* spores treated with O₃MMB for 180 s (Fig. 4). Then, the spores had swelled by O₃MMB for 180 s, as water entered the cells through the damaged cell wall. Cho et al. (2010) reported that disinfection of bacterial cells by O₃ was due to injury of the cell wall. Zhang et al. (2011) showed that disinfection of *Pseudomonas aeruginosa* by O₃ was due to increase in cell membrane permeability and coagulation of the intracellular substrate. Thanomsut et al. (2002) concluded that disinfection by O₃ was caused due to the destruction of the cell wall and leakage of the intracellular substrate, followed by




Fig. 4. TEM images of *F. oxysporum* f. sp. *melonis* spores. Top: Non-treated, Middle: After treatment with O₃MMB for 180 s, Bottom: After treatment with O₃MB for 180 s.

cell lysis. Therefore, it is possible that the *F. oxysporum* f. sp. *melonis* spores treated with O₃MMB for 30 s initially shrank due to leakage of the intracellular substrate through the damaged cell wall. On the other hand, the TEM images of the *F. oxysporum* f. sp. *melonis* spores treated with O₃MB for 180 s showed the wavelike deformation of cell membrane and appeared to have a space between the cell membrane and/or wall and the cytoplasm. Diao et al. (2004) confirmed that hydroxyl radicals generated by the Fenton reaction induced the injury of *E. coli* cell membranes greater than O₃ by the observation with SEM. Therefore, it appears that hydroxyl radicals generated from the O₃MB induces the wavy injury of cell membrane of *F. oxysporum* f. sp. *melonis* spores. Furthermore, it was considered that coagulation of the substrate within the spores and leakage of the substrate through the damaged cell membrane were induced by the higher amounts of hydroxyl radicals generated from the tiny O₃MB penetrated into the spores. The result may lead to cell death due to the leakage and/or coagulation of the intracellular substrate, followed by the lysis of the spore.

These results show that disinfection efficiency of O₃MB on *F. oxysporum* f. sp. *melonis* spores is higher than that of O₃MMB and may be due to the action of MB in combination with the high oxidative power of the O₃. Therefore, it is considered that disinfection of *F. oxysporum* f. sp. *melonis* spores by O₃MB causes the leakage and/or coagulation of intracellular components associated with damage to the cell membrane and/or cell wall, and subsequently leads to lysis of the spore.

References

- Bando, K., Kawano, M., Kuroda, Y., Kusakari, S., Yamasaki, M., Maeda, T. and Kourai, H. 2008. Effect of silver-supported photocatalytic sterilization, yield, quality, and nutrient element concentrations in the tomato grown under rockwool culture. *Hortic. Res. (Japan)* 7:309-315 (in Japanese).
- Cho, M., Chung, H. and Yoon, J. 2003. Disinfection of water containing natural organic matter by using ozone-initiated radical reactions. *Appl. Environ. Microbiol.* 69:2284-2291.
- Cho, M., Kim, J., Kim, J. Y., Yoon, J. and Kim, J. H. 2010. Mechanisms of *Escherichia coli* inactivation by several disinfectants. *Water Res.* 44:3410-3418.
- Chu, L. B., Xing, X. H., Yu, A. F., Sun, X. L. and Jurcik, B. 2008. Enhanced treatment of practical textile wastewater by microbubble ozonation. *Process Saf. Environ. Prot.* 86:389-393.
- Chujedton, A., Nuanaon, N., Uthaibutra, J. and Whangchai, K. 2015. Ozone microbubbles disinfection technique to inactivate *penicillium digitatum* in suspension. *Acta Hortic.* 1088:355-358.
- Dannehl, D., Schuch, I., Gao, Y., Cordiner, S. and Schmidt, U. 2016. Effects of hypochlorite as a disinfectant for hydroponic systems on accumulations of chlorate and phytochemical compounds in tomatoes. *Eur. Food Res. Technol.* 242:345-353.
- Diao, H. F., Li, X. Y., Gu, J. D., Shi, H. C. and Xie, Z. M. 2004. Electron microscopic investigation of the bactericidal action of electrochemical disinfection in comparison with chlorination, ozonation and Fenton reaction. *Process Biochem.* 39:1421-1426.
- Ehret, D. L., Alsanius, B., Wohanka, W., Menzies, J. G. and Ut-khede, R. 2001. Disinfection of recirculating nutrient solutions in greenhouse horticulture. *Agronomie* 21:323-339.
- Igura, N., Fujii, M., Shimoda, M. and Hayakawa, I. 2004. Inactivation efficiency of ozonated water for *Fusarium oxysporum* conidia under hydroponic greenhouse conditions. *Ozone Sci. Eng.* 26:517-521.
- Inatsu, Y., Kitagawa, T., Nakamura, N., Kawasaki, S., Nei, D., Bari, L. and Kawamoto, S. 2011. Effectiveness of stable ozone microbubble water on reducing bacteria on the surface of selected leafy vegetables. *Food Sci. Technol. Res.* 17:479-485.
- Kobayashi, F., Ikeura, H., Ohsato, H., Goto, T. and Tamaki, M. 2011a. Disinfection using ozone microbubbles to inactivate *Fusarium oxysporum* f. sp. *melonis* and *Pectobacterium carotovorum* subsp. *carotovorum*. *Crop Prot.* 30:1514-1518.
- Kobayashi, F., Ikeura, H., Ohsato, S. and Tamaki, M. 2011b. Microbicidal effect of microbubbles with ozone, oxygen, and carbon dioxide against *Fusarium oxysporum* f. sp. *melonis* and *Pectobacterium carotovorum* subsp. *carotovorum*. *Journal of the Japanese Society of Agricultural Technology Management* 18:123-128.
- Kobayashi, F., Ikeura, H., Ohsato, S. and Tamaki, M. 2011c. Microbicidal effect of ozone microbubbles generated by different methods on *Fusarium oxysporum* f. sp. *melonis* and *Pectobacterium carotovorum* subsp. *carotovorum*. *Journal of the Japanese Society of Agricultural Technology Management* 18:129-134.
- Kobayashi, F., Ikeura, H., Ohsato, S., Goto, T. and Tamaki, M. 2012. Ozone microbubbles as a disinfection in nutrient solution, and their effects on the composition of fertilizer and the growth of cultivated plants. *Biol. Eng. Trans.* 5:137-146.
- Koohakan, P., Ikeda, H., Kusakari, S., Masuda, T., Mano, K. and Masuda, R. 2003. Effects of TiO₂ photocatalytic sterilizing system on the suppression of tomato root rot disease in the nutrient solution. *Hortic. Res. (Japan)* 2:215-219 (in Japanese).
- Li, P. and Tsuge, H. 2006. Ozone transfer in a new gas-induced contactor with microbubbles. *J. Chem. Eng. Jpn* 39:1213-1220.
- Li, P., Takahashi, M. and Chiba, K. 2009. Degradation of phenol by the collapse of microbubbles. *Chemosphere* 75:1371-1375.
- Ministry of Agriculture, Forestry and Fisheries. 2010. Agricultural

- tural Chemicals Regulation Act. URL http://www.maff.go.jp/j/nouyaku/n_kaisei/zenbun.html/.
- Ohtani, T., Kaneko, A., Fukuda, N., Hagiwara, S. and Sase, S. 2000. Development of a membrane disinfection system for closed hydroponics in a greenhouse. *J. Agric. Eng. Res.* 77:227-232.
- Runia, W. T. 1995. A review of possibilities for disinfection of recirculation water from soilless cultures. *Acta Hortic.* 382:221-229.
- Takahashi, M. 2005. ζ potential of microbubbles in aqueous solutions: electrical properties of the gas-water interface. *J. Phys. Chem. B* 109:21858-21864.
- Takahashi, M., Kawamura, T., Yamamoto, Y., Ohnari, H., Himuro, S. and Shakutsui, H. 2003. Effect of shrinking microbubble on gas hydrate formation. *J. Phys. Chem. B* 107:2171-2173.
- Thanomsub, B., Anupunpisit, V., Chanphetch, S., Watcharachaipong, T., Poonkhum, R. and Srisukonth, C. 2002. Effects of ozone treatment on cell growth and ultrastructural changes in bacteria. *J. Gen. Appl. Microbiol.* 48:193-199.
- Tsuge, H., Li, P., Shimatani, N., Shimamura, Y., Nakata, H. and Ohira, M. 2009. Fundamental study on disinfection effect of microbubbles. *Kagaku Kogaku Ronbunshu* 35:548-552 (in Japanese).
- Zhang, Y. Q., Wu, Q. P., Zhang, J. M. and Yang, X. H. 2011. Effect of ozone on membrane permeability and ultrastructure in *Pseudomonas aeruginosa*. *J. Appl. Microbiol.* 111:1006-1015.
- Zheng, T., Wang, Q., Zhang, T., Shi, Z., Tian, Y., Shi, S., Smale, N. and Wang, J. 2015. Microbubble enhanced ozonation process for advanced treatment of wastewater produced in acrylic fiber manufacturing industry. *J. Hazard. Mater.* 287:412-420.