



POSTER PRESENTATION

Open Access

Factors affecting height in children and adolescents with transfusion-dependent Thalassaemia – results from a Thalassaemia center in Malaysia

Tzer Hwu Ting^{1*}, Bina Sharine Menon², Hishamshah Mohd Ibrahim²

From 7th APPEs Biennial Scientific Meeting
Nusa Dua, Bali. 14-17 November 2012

Thalassaemia, a common inherited haemoglobinopathy in Malaysia¹, causes significant morbidity due to chronic anaemia and complications of iron excess from regular blood transfusion. Short stature is one of the commonest complications in transfusion-dependent thalassaemia with prevalence of 31% to 75%^{2,3}. Causes of growth impairment include iron overload, nutritional deficiency, GH-IGF1 axis dysregulation, abnormal puberty and desferrioxamine toxicity^{4,5}.

We aim to determine the prevalence of short stature, defined as height z-score ≤ -2 , and the factors affecting height in thalassaemia patients attending the Haematology clinic at a tertiary referral center. This observational cross-sectional study involved patients aged 3-19 years, needing blood transfusion at least every 8 weeks, and without co-morbidities or endocrinopathy which affect growth.

Weight & height were measured. Puberty status was clinically assessed. Body mass index (BMI) was calculated as a marker of nutritional status. Mean of ferritin levels measured over the preceding one year was determined as a marker of iron overload.

There were 52 boys (mean age $11.2y \pm 4.8y$, mean height z-score -1.67 ± 1.11 , mean BMI z-score -0.87 ± 1.44 , mean ferritin $3316 \pm 2092 \mu\text{g/L}$) and 47 girls (mean age $11.8y \pm 4.2y$, mean height z-score -1.59 ± 0.97 , mean BMI z-score -0.83 ± 0.95 , mean ferritin $3005 \pm 2356 \mu\text{g/L}$). Overall 35/99 (35%) were short while 64/99 (64%) had normal height. 19/52 (37%) boys and 16/47 (34%) girls were short. 16/19 (84%) short

boys were aged ≥ 9 years, 15/16 (94%) short girls were aged ≥ 8 years. Only one boy (aged 15y) and one girl (aged 14y) had delayed puberty. They were both short with height z-score -3.41 and -2.26 respectively.

Overall there was a moderate positive correlation between height z-score and BMI z-score (Pearson $r = 0.388$, $p=0.00$). There was no significant correlation between height z score and mean ferritin level (Pearson $r = -0.044$, $p= 0.666$).

Significant proportion (35%) of our transfusion-dependent thalassaemia patients have short stature, majority of them in the pubertal age group. Height is significantly associated with nutritional status but not ferritin level. Improvement in nutrition may improve height in these patients.

Authors' details

¹Department of Paediatrics, University Putra Malaysia, Malaysia.

²Haematology & Oncology Unit, Paediatrics Institute, Kuala Lumpur Hospital, Malaysia.

Published: 3 October 2013

doi:10.1186/1687-9856-2013-S1-P61

Cite this article as: Ting et al.: Factors affecting height in children and adolescents with transfusion-dependent Thalassaemia – results from a Thalassaemia center in Malaysia. *International Journal of Pediatric Endocrinology* 2013 **2013**(Suppl 1):P61.

¹Department of Paediatrics, University Putra Malaysia, Malaysia
Full list of author information is available at the end of the article