



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

REVIEW ARTICLE

Messenger ribonucleic acid vaccines for severe acute respiratory syndrome coronavirus-2 – a review



CHRISTINE M. AKAMINE, and HANA M. EL SAHLY

HOUSTON, TEXAS

The mRNA therapeutics have been studied since the 1970s and the currently available mRNA vaccines against COVID-19 are the culmination of decades of scientific research. The mRNA vaccines BNT162b2 and mRNA-1273 have played a key role in our global response to the COVID-19 pandemic as they have demonstrated significant advantages over conventional vaccines and have proven to be highly effective against COVID-19 associated hospitalization and severe illness in large clinical trials and studies using real-world data. (Translational Research 2022; 242:1 – 19)

Abbreviations: COVID-19 = Coronavirus disease of 2019; SARS-CoV-2 = Severe acute respiratory syndrome coronavirus 2; mRNA = Messenger ribonucleic acid; LNP = Liposomal nanoparticle; SAM = Self-amplifying mRNA; dsRNA = Double-stranded RNA; Ad5 = Adenovirus type 5; APCs = Antigen presenting cells; RBD = Receptor-binding domain; MERS-CoV = Middle east respiratory syndrome coronavirus; GMT = Geometric mean titer; BMI = Body mass index; EUA = Emergency use authorization; CDC = Centers for Disease Control; B.1.1.7 = Alpha variant; B.1.351 = Beta variant; COVID-NET = COVID-19-Associated Hospitalization Surveillance Network; VOC = Variants of concern; UTR = Untranslated regions; PAMPs = Pathogen-associated molecular patterns; MHC = Major histocompatibility complex; DCs = Dendritic cells; ACE-2 = Angiotensin converting enzyme receptor; CVnCOV = CureVac; GMC = Geometric mean concentration; FDA = Food and Drug Administration; VAERS = Vaccine adverse event reporting system; BAU = binding antibody units; DNA = deoxyribonucleic acid; tRNA = transfer ribonucleic acid; PRNT₅₀ = plaque reduction neutralization test; Nab = neutralizing antibodies; BLA = Biologics License Application; VE = Vaccine efficacy; VSD = Vaccine Safety Datalink; VA = Department of Veterans Affairs

BACKGROUND

Conventional vaccines, including inactivated, live attenuated, subunit, and conjugated and unconjugated polysaccharides, have long demonstrated great efficacy at preventing many infectious diseases and have made

substantial global impacts to reduce morbidity and mortality due to preventable diseases. With the advent of the Coronavirus disease of 2019 (COVID-19) pandemic caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), there was an urgent need for rapid, large-scale development and deployment of vaccines against this virus to curb the pandemic. However, production of conventional vaccines can be a time-consuming process that is often hampered by dependence on the production of cell lines and constructing protein subunits with high purity and reproducibility, which can often take more than 1 year.¹ Messenger ribonucleic acid (mRNA) vaccines can be manufactured rapidly at low cost due to high yields of in vitro transcription reactions and have additional benefits over traditional available vaccines.^{2,3} For example: mRNA vaccines do not rely

From the Baylor College of Medicine, One Baylor Plaza, Houston, Texas.

Submitted for Publication September 9, 2021; revision submitted December 16, 2021; Accepted for Publication December 17, 2021.

Reprint requests: Christine M. Akamine, Baylor College of Medicine, One Baylor Plaza, Houston, Texas, 77030. e-mail: akamine@bcm.edu.

1931-5244/\$ - see front matter

© 2021 Elsevier Inc. All rights reserved.

<https://doi.org/10.1016/j.trsl.2021.12.007>

on animal products or cell cultures and because mRNA is a noninfectious platform that does not enter the nucleus or incorporate into the genome, thereby avoiding genomic insertion, it provides attractive safety characteristics.^{1,4} For these reasons, mRNA vaccines were considered as one of the frontrunners for vaccine development against SARS-CoV-2 infection, especially in the setting of a pandemic.

The concept of utilizing exogenous mRNA to cause cells to produce proteins of interest in organisms was first described in the early 1970s.^{5,6} In vitro studies involving injection of encephalomyocarditis virus RNA and mRNA encoding rabbit hemoglobin into oocytes from *Xenopus laevis* demonstrated that introduction of RNA molecules into cells could result in protein production.^{5,6} The development of mRNA therapeutics made major strides in 1989 when researchers at the Salk Institute at University of California San Diego first described successful transfection of mRNA in a liposomal nanoparticle (LNP) into the cytoplasm of mouse cells.⁷ A few months later, Wolff et al demonstrated that in vivo transfection of naked mRNA into mouse skeletal muscle resulted in detectable production of the encoded protein.⁸ By 1993, Martinon et al showed that virus-specific cytotoxic T lymphocytes could be elicited in vivo by immunizing mice with liposome-entrapped mRNA encoding the nucleoprotein of the influenza virus.⁹ By 2017, the first proof-of-concept phase 1 human trial of an mRNA vaccine against an infectious disease, the rabies virus, demonstrated the ability of this platform to induce an immune response to the encoded antigen of interest.¹⁰ These early successes supported mRNA technology as a highly versatile and viable contender in ongoing efforts toward innovative vaccine development.

EXPRESSION SYSTEM/MECHANISM

mRNA is a single stranded RNA, and an intermediate transporter of the genetic code, that plays a key role between translation of protein-encoding deoxyribonucleic acid (DNA) to protein production by ribosomes in the cytoplasm.^{2,11} Functional synthetic mRNA is the result of in vitro transcription of a linearized plasmid DNA template by a DNA-dependent RNA polymerase, such as the T7, T3, or Sp6 bacteriophage RNA polymerase, in the presence of nucleoside triphosphates.¹¹⁻¹⁴ The product is a functional synthetic mRNA that resembles mature mRNA found in eukaryotic cells, and the template DNA is subsequently degraded by DNases.^{2,4,11} There are 2 types of mRNA utilized in the current vaccines under development: non-replicating mRNA and self-amplifying mRNA (SAM). Conventional non-replicating mRNA contains a 5' cap, an open reading frame containing the target protein sequence, 5' and 3'

untranslated regions (UTRs), and a poly(A) tail.^{2,4} SAM contains these essential elements, but it also encodes viral replication machinery which enables intracellular RNA amplification thereby increasing protein expression compared with non-replicating RNA.^{2,15}

The process of in vitro transcription results in many byproducts, including contaminating double-stranded RNA (dsRNA), that can provoke an undesirable innate immune response and hasten the degradation of the newly made RNA.¹¹ By acting as pathogen-associated molecular patterns (PAMPs), these byproducts are detected by the immune system and can incite production of type I interferon which can lead to degradation of both the cellular and ribosomal RNA.^{2,16} Additionally, the intrinsically immunogenic nature of RNA itself can provoke such an immune response and thus the process of purification of the mRNA becomes an important step.³ Purification also plays a role in optimizing the expression of mRNA once it enters the cell.³ Several methods of removing the contaminating dsRNA have been described, including fast protein liquid chromatography, high-performance liquid chromatography, and nucleoside modification.¹⁷ Utilization of these techniques facilitates maximal production of the desired protein and avoidance of adverse activation of innate immune responses.

The post-transcription process provides the essential modifications that impart structural stability and enhanced expression of the mRNA code. The 5' cap and poly(A) tail are essential for efficient protein translation and stabilization of the mRNA molecule in the cytoplasm.^{11,18,19} The 5' and 3' UTRs have important structural roles as they provide the mRNA molecule with stability and regulate translation.^{2,18} The 5' and 3' UTRs also prolong the half-life and enhance expression of the mRNA.²⁰ The post-transcription cytoplasmic mRNA then encounters the cell's translation machinery and the protein of interest is produced.¹¹ This protein then undergoes additional post-translational modifications, including proper folding, to make a fully functional protein.¹¹

Other methods of increasing protein translation and expression have been described.² Sequence optimization and modification of codon usage by substituting rare codons for synonymous codons can increase transfer ribonucleic acid (tRNA) in the cytoplasm.^{2,4,21} Another method described involves increasing the guanine and cytosine sequence content of the genetic code, which leads to increased steady-state mRNA levels resulting in increased gene expression.^{2,15,22}

DELIVERY

Once delivered at the site of injection, the mRNA in the vaccine must avoid degradation by nucleases and




Fig 1. Mechanism of nonreplicating mRNA vaccines against COVID-19. For interpretation of the references to color in this figure legend, the reader is referred to the Web version of this article.

cross the cell's plasma membrane to reach the translation machinery in the cytoplasm in order to effect an antigen-specific immune response.^{15,23} Fig 1 summarizes the mechanism of nonreplicating mRNA vaccines against COVID-19. Multiple studies have demonstrated the ability of naked mRNA to be taken up by cells, however carrier systems to protect the mRNA from degradation enhance delivery from the site of injection to the cytoplasm.¹¹ There is also evidence that suggests both humoral and cell-mediated immune responses are improved with intramuscular delivery of LNP-formulated mRNA compared with naked mRNA.³ Several carrier systems have been designed to optimize this process, among which LNP formulations are currently the most promising.^{2,24} Given the intrinsically negative charge of the cell membrane, due to its phospholipid bilayer composed of polar heads and hydrophobic tails, the negatively charged mRNA benefits from the shelter provided by the positively charged LNP delivery systems to efficiently enter the cell.¹⁵ LNPs are generally composed of an ionizable cationic lipid, which facilitates endosomal escape of the mRNA once in the cytoplasm, polyethylene glycol, and zwitterionic lipids, which provide structural support and increase the half-life of the formula, and cholesterol, a stabilizing agent.^{2,24,25} Initially this delivery system was studied for small interfering RNA molecules, however it has subsequently been shown to be a successful and well tolerated

method for both non-replicating mRNA and SAM delivery.^{26,27}

Several studies have evaluated different routes of delivery of mRNA vaccines including intravenous, intradermal and intramuscular.³ When LNP-mRNA is delivered intravenously it binds with high affinity to hepatocytes in the liver, whereas intradermal and intramuscular delivery resulted in longer expression of the antigen at the site of inoculation, with intradermal delivery resulting in the longest lasting expression.^{3,28,29} After intramuscular injection of LNP-mRNA in mice and nonhuman primates, the mRNA is taken up and strongly expressed by local cells including professional antigen presenting cells (APCs), monocytes, granulocytes and dendritic cells (DCs), which may facilitate transport of the antigen to neighboring draining lymph nodes.²⁹ The professional APCs and DCs play a key role in processing the mRNA encoded antigen, presenting it on major histocompatibility complex (MHC) class I and II molecules to CD4⁺ and CD8⁺ T-cells, and facilitating recognition by B cells for the development of a humoral immune response.³

PRECLINICAL DATA

In preclinical studies, mRNA vaccines demonstrated a good safety profile, remarkable efficacy, long-lived

immune responses to infectious pathogens and adaptability for emerging infectious diseases.² Multiple studies in animal models demonstrated that mRNA vaccines were able to elicit strong immunity against a variety of infectious diseases, including Zika virus, influenza, *Toxoplasma gondii*, Ebola, Powassan virus, and rabies.^{2,30-32} mRNA vaccines demonstrated their ability to provoke a robust immune response in the form of both neutralizing antibodies as well as CD8⁺ and CD4⁺ T cell responses. A study of an mRNA vaccine encoding the full-length hemagglutinin of influenza A (PR8HA) demonstrated not only the production of protective humoral immunity but also B and T-cell mediated immune responses in both young and old mice.³³ Few direct comparisons of immunogenicity and durability of protection have been made between mRNA vaccines and other vaccine constructs to date. There is some evidence that mRNA vaccines are more immunogenic with comparable durability of response as other vaccine constructs in preclinical studies. A single dose of the nucleoside-modified pre-membrane and envelope (prM-E) LNP-formulated mRNA vaccine construct against Zika virus (ZIKV prM-E mRNA-LNP) provoked higher neutralizing antibody responses than those elicited by plasmid DNA vaccines in mice and rhesus macaques.^{34,35} Compared with a single 1 mg dose of DNA-vaccine, a single 50 μ g dose of ZIKV prM-E mRNA-LNP induced reporter virus particle neutralizing antibodies (Nab) that were 50 times higher, and over twice as high as 2 doses of DNA vaccine in macaques.³⁶ It also demonstrated that plaque reduction neutralization test (PRNT₅₀) NAb continued to increase over several months and were 50–100 times higher than those provoked by a single dose of purified inactivated virus or plasmid DNA vaccine.³⁴ Neutralizing antibody titers were stable through 12 weeks postimmunization.³⁴ Comparative evaluations of vaccine-induced NAb for ZIKV prM-E mRNA-LNP, an adenovirus vectored vaccine (RhAd52 ZIKV) and PIV in macaques was difficult due to use of different assays.^{34,35} Another study demonstrated a stronger CD4⁺ T cell response after RABV-G mRNA vaccine compared with the inactivated rabies vaccine (Rabipur) in mice.³¹ In newborn piglets, the RABV-G mRNA vaccine lead to prompt seroconversion and comparable Nab titers as the Rabipur vaccine.³¹ Non-human primates who received an LNP-complexed mRNA vaccine against influenza were demonstrated to have equivalent antibodies following 1 dose as compared with an adjuvanted subunit influenza vaccine, higher antibody responses after the second LNP-mRNA dose.²⁹ In rhesus macaques, vaccination with LNP-mRNA compared to adjuvanted HIV-1 Env

recombinant protein vaccination showed similar or higher levels of HIV-1 Env-specific antibodies.³⁷

Since the half-life and durability of nonreplicating mRNA is transient, SAM vaccines aim to increase the expression of the encoded protein via their autoreplicative capabilities.^{38(p7)} Comparisons between SAM and nonreplicating mRNA vaccine efficacy (VE) are few. Vogel et al showed that equivalent levels of protection against influenza A and antigen expression were achieved after just 1.25 μ g of SAM compared with 80 μ g of nonreplicating mRNA (a 64-fold difference in dose), thereby supporting the potency of SAM vaccine constructs.³⁹

Prior studies of coronaviruses demonstrated that the neutralizing antibodies to the spike protein (S protein), which facilitates viral attachment and entry into the host cells, as well as passive transfer of S protein antibodies, provided protection against infection with this family of viruses in challenge models.^{40,41} In SARS-CoV-2, the trimeric spike protein utilizes part of the N-terminal furin cleavage fragment (S1) called the receptor-binding domain (RBD) to bind to the angiotensin converting enzyme 2 (ACE-2) receptor on target cells.^{42,43} Using the fusion machinery of the C-terminal furin cleavage fragment (S2), the S protein undergoes significant conformational changes in order to fuse the virus with the host cell membrane and allow entry of the viral genome, resulting in pre-fusion and postfusion states of the protein.^{1,42,43} Prior studies with a stabilized respiratory syncytial virus (RSV) subunit vaccine candidate demonstrated that utilizing the pre-fusion conformation of the RSV glycoprotein resulted in a significant increase in neutralizing antibody production to key epitopes, and demonstrated that methods to stabilize the S-protein in the pre-fusion state increases S-protein expression.^{44,45} In studies on middle east respiratory syndrome coronavirus (MERS-CoV), SARS-CoV, and human coronavirus HKU1 (CoV-HKU1), it was discovered that 2 proline substitutions (K986P and V987P) in the apex of the central helix of the spike protein prevented postfusion structural changes and effectively stabilized the S proteins in the pre-fusion conformation, termed the 2-SP (or 2P) design.⁴⁵ In a study involving MERS-CoV in a mouse model, immunization with MERS-CoV-S(2P) mRNA-LNP was protective against lethal MERS-CoV challenge in a dose-dependent manner.⁴⁶ To date, 3 mRNA vaccines against SARS-CoV-2 utilize the S-2P design, the mRNA-1273 (Moderna/NIH), BNT162b2 (Pfizer-BioNTech), and CVnCOV (CureVac).

On January 10, 2020 the genomic sequence of SARS-CoV-2 was released and a race to produce a vaccine based on the pathogen sequence began in order to gain a foothold on the rapidly rising global cases of




Fig 2. Timeline of Events in COVID-19 mRNA Vaccine Development.

COVID-19.¹ Fig 2 shows a timeline summarizing key events in COVID-19 mRNA vaccine development. Within 24 hours of the release of the SARS-CoV-2 genomic sequence, the S-2P design was utilized to produce the stabilized pre-fusion SARS-CoV-2 S(2P) protein *in silico* for analysis of the viral structure and development of viral assays.^{1,47-49} Of all the available platforms of vaccine, the manufacturing advantages of mRNA vaccines allowed them to enter promptly into preclinical studies and 2 vaccine candidates, the BNT162b2 and mRNA-1273, were the frontrunner mRNA based COVID-19 vaccine candidates. Other front runner vaccine candidates included Ad26.COV2-S (Janssen Pharmaceuticals/Johnson & Johnson), ChAdOx1 nCoV-19 (Oxford/AstraZeneca), and CoronaVac (Sinovac Biotech). A summary of the preclinical data from the Pfizer-BioNTech and Moderna COVID-19 vaccine products can be found on Table 1 and is discussed below.

Pfizer-BioNTech. The vaccine development program by Pfizer-BioNTech included 4 investigational products, which differed in the nucleotide sequences of the encoded antigens, as part of their mRNA-based vaccine program, named Project Light Speed, in response to the COVID-19 pandemic.⁵⁰ The initial objective was to select a single vaccine candidate and dosing regimen to progress into further studies on safety and efficacy.⁵⁰

The preclinical studies of BNT162b2 in mice and rhesus macaques demonstrated development of potent immunogenicity after vaccination. The BNT162b2 is an LNP-complexed N¹-methyl-pseudouridine (m¹Ψ) nucleoside-modified mRNA encoding the full-length

pre-fusion stabilized SARS-CoV-2 S-2P.⁵¹ These modifications, along with non-coding sequence optimization, were shown to increase RNA translation *in vivo* with the added benefit of decreasing immune sensing.^{28,51,52} Studies in BALB/c mice showed that strong neutralizing antibody production to S1 and the receptor binding domain (RBD) developed in a dose-dependent fashion after administration of 0.2 μg, 1 μg, or 5 μg of BNT162b2, as well as systemic T cell responses.⁵¹ In mice, the prime-only regimen also demonstrated a strong neutralizing antibody and T cell responses.⁵¹ Additionally, studies in rhesus macaques showed that a prime/boost vaccination of 100 μg of BNT162b2 resulted in SARS-CoV-2 neutralizing geometric mean titers (GMT) 18.0 times that seen in convalescent human serum when measured 7 days post dose 2.⁵¹ Prime/boost vaccination of rhesus macaques with 100 μg of BNT162b2 successfully prevented lung infection after rechallenge of SARS-CoV-2 at a high intranasal and intratracheal inoculum (1.05 × 10⁶ PFU of SARS-CoV-2 USA-WA1/2020 isolate).⁵¹ Prime/boost with 30 μg or 100 μg also demonstrated robust neutralizing antibody production as well as CD4⁺ and CD8⁺ T cell responses.⁵¹

Studies on another vaccine investigational project as part of Project Light Speed, BNT162b1, also showed promising results. It was composed of the same LNP-formulated m¹Ψ nucleoside-modified RNA platform as BNT162b2 but expressed a T4 fibrin-derived trimerized version of the RBD to increase immunogenicity.⁵³ Similarly to BNT162b2, BNT162b1 elicited potent humoral and T-cell mediated immune responses

Table 1. Preclinical data summary for mRNA COVID-19 vaccines^a

Study product	Animal model	Dose (50% virus-neutralization GMT (μg))	Cell-mediated immunity	Evidence of protection
BNT162b2 Vogel et al ⁵¹	Rhesus macaques	30 μg (962 in sera drawn 7 or 14 d post dose 2, 285 at 28 d post dose 2) 100 μg (1689 in sera drawn 7 or 14 d post dose 2, 283 at 28 d post dose 2)	CD4 ⁺ T cells (TH1 pre-dominant response) CD8 ⁺ T cells were elicited	No SARS-CoV-2 RNA on BAL samples 3 d after challenge (with intranasal and intratracheal 1.05×10^6 PFU of SARS-CoV-2 USA-WA1/2020 isolate)
BNT162b2 Vogel et al ⁵¹	BALB/c mice	5 μg (296 28 d after 1 dose)	CD4 ⁺ and CD8 ⁺ T cell responses	-
BNT162b1 Vogel et al ⁵¹	Rhesus macaques	30 μg (768 in sera drawn 7 or 14 d post dose 2) 100 μg (1714 in sera drawn 7 or 14 d post dose 2)	CD4 ⁺ T cells (TH1 pre-dominant response) CD8 ⁺ T cells were elicited	SARS-CoV-2 RNA detected in BAL samples of 2/6 immunized macaques 3 d after challenge (with intranasal and intratracheal 1.05×10^6 PFU of SARS-CoV-2 USA-WA1/2020 isolate)
BNT162b1 Vogel et al ⁵¹	BALB/c mice	5 μg (1,056 28 d after 1 dose)	CD4 ⁺ and CD8 ⁺ T cell responses	-
mRNA-1273 ¹	BALB/c mice	1 μg (819 after 2 doses) 10 μg (315 between wk 2 and 4 after 1 dose) 0.0025-20 μg doses trialed with a strong positive dose-dependent production of binding and pseudovirus-neutralizing antibodies	CD4 ⁺ and CD8 ⁺ T cell responses	Protected against viral replication in the lungs, protected and nares in 6 out of 7 mice

^aT helper type 1 TH1; bronchoalveolar lavage BAL; Plaque forming units PFU.

in both mice and macaques in preclinical studies, and protected macaques against rechallenge with SARS-CoV-2.⁵¹ Both the BNT162b1 and BNT162b2 moved forward into phase I clinical trials.

Moderna. The mRNA-1273, a mRNA based vaccine encoding the full-length pre-fusion stabilized SARS-CoV-2 S-2P encapsulated in a LNP (containing 4 lipids including a novel proprietary lipid SM-102), demonstrated protective immune responses and good efficacy in preclinical trials.¹ The ability of S-2P antigen to be delivered by mRNA was confirmed in vitro in mouse models.¹ Assessments of immunogenicity in BALB/c mice after a prime/boost regimen of mRNA-1273, 0.01 μg or 0.1 μg or 1 μg dose, separated by 3 weeks, showed that S-binding and neutralizing antibodies were induced by the vaccine in a dose-dependent manner.¹ A wide dose range utilizing 0.0025–20 μg doses of mRNA-1273 in BALB/c mice was also evaluated to assess immunogenicity and demonstrated a positive dose-dependent production in pseudovirus-neutralizing antibodies.¹ Additionally, a prime-only regimen with 1 μg or 10 μg doses was evaluated and showed binding antibody titers without neutralizing antibody titers in

both regimens.¹ Mice that received prime/boost regimen of 1 μg of mRNA-1273 were completely protected from mouse-adapted SARS-CoV-2 viral replication in the lungs when challenged 5 or 13 weeks after vaccination, and 6 of 7 mice did not have viral replication in the nasal turbinates.¹ Mice that received 0.1 μg prime/boost doses of mRNA-1273 had a reduction of 100-fold in viral load in the lungs, and those that received 0.01 μg prime/boost doses had a 3-fold lung viral load reduction.¹ When re-challenged 7 weeks after a prime-only regimen of 1 μg or 10 μg dose of mRNA-1273, there was no evidence of viral replication in the lungs of immunized mice.¹ The results of the preclinical trials of mRNA-1273 demonstrated that a prime/boost regimen of 1 μg of mRNA-1273 induced pseudovirus-neutralizing antibodies and CD8⁺ T cell responses, as well as more than 3 months protection from viral replication in the respiratory tract.¹

The preclinical studies of mRNA-1273 also evaluated for the theoretical possibility of vaccine associated enhanced disease, which has previously been described with SARS-CoV and MERS-CoV and in some cases of

RSV, measles, dengue and Zika.^{1,54} Subprotective doses of 0.1 μg or 0.01 μg of mRNA-1273 did not demonstrate any evidence of increased mucous production or enhanced lung pathology in mice.¹

CLINICAL DATA OF MRNA THERAPEUTICS FOR SARS COV-2

Given the encouraging results from the preclinical studies of the mRNA-1273, BNT162b1 and BNT162b2 vaccine candidates, these were moved forward into clinical trials to further evaluate dosing regimen, safety, immunogenicity and efficacy.

Pfizer-BioNTech. In a Phase I/II observer-blind, placebo-controlled study of the BNT162b1 vaccine candidate, 45 healthy adults aged 18–55 years old received 2 vaccines at 10 μg or 30 μg separated by 21 days, 100 μg prime only, or placebo.⁵³ RBD-binding IgG titers and SARS-CoV-2 neutralizing antibodies were measured at baseline and days 7, 21, 28, and 35.⁵³ The RBD-binding IgG were seen to persist at high levels after the last day of follow up (day 35) in the 10 μg and 30 μg prime/boost regimen subjects, with levels being 8 and 46.3-times the geometric mean concentration (GMC) seen in convalescent sera, and neutralizing GMT at 1.8 and 2.8-times convalescent sera.⁵³ Subjects who received a single dose of 100 μg did not have increased levels of RBD-binding IgG past day 21, but did demonstrate 3-times the GMC and 0.35-times the GMT of convalescent sera.⁵³ Local and systemic reactions were mainly mild to moderate in severity, transient and dose-dependent after 10 μg and 30 μg prime-boost regimens.⁵³ Fever was more common after the second dose of 30 μg compared with the 10 μg regimen (75% vs 10%), and no severe adverse events were reported.⁵³ There was no increased immunogenicity after prime-only of 100 μg but there was an increase in systemic and local reactions.⁵³

In the phase I/II open-label non-randomized trial in Germany to evaluate immunogenicity and safety of the prime-boost regimen of BNT162b2, healthy participants age 19–55 were given prime-boost doses in 1 μg , 10 μg , 20 μg , or 30 μg dose levels.⁵⁵ Immunogenicity assessments included S1-binding and RBD-binding IgG assays and SARS-CoV-2 neutralizing assays, which were measured at baseline and on days 7, 21, 29, 43, 50, and 85 for all dose levels except 1 μg .⁵⁵ At all doses, a robust immune response, both humoral and T cell mediated, was seen with no serious adverse events reported.⁵⁵

In order to select the final vaccine candidate, an observer-blind, placebo-controlled dose escalation phase I study comparing prime/boost regimens of the 2 front-runner Pfizer-BioNTech vaccine candidates, BNT162b2 and BNT162b1, at 10 μg , 20 μg , and 30 μg dose levels was conducted to evaluate safety and immunogenicity.⁵⁶

One hundred and ninety-five adults 18–55 and 65–85 years old were randomly assigned to receive one of the 2 vaccine candidates or placebo.⁵⁶ Each participant received 2 doses of vaccine 21 days apart or placebo according to their allocation.⁵⁶ Assessments of immunogenicity were conducted at baseline and on days 7, 21, and 35 and included a SARS-CoV-2 serum neutralization assay and RBD-binding or S1-binding IgG assays.⁵⁶ Local reactogenicity was reported frequently after BNT162b1 vaccination, primarily injection site pain within 7 days of injection, in a dose dependent manner and the large majority were mild-moderate in severity.⁵⁶ Similar results were seen after BNT162b2 vaccination, with no grade 4 local reactions reported in either group.⁵⁶ Systemic side effects were milder in the BNT162b2 group, with 17% of the 18–55 age group reporting fever after the second dose and 8% in the 65–85 age group, while in the BNT162b1 group 75% of the 18–55 age group and 33% of the 65–85 group reported fever after the second dose.⁵⁶ Overall, BNT162b2 demonstrated a milder systemic reactogenicity profile compared with BNT162b1, especially in the older adults.⁵⁶ Although both demonstrated equivalent immunogenicity, BNT162b2 at 30 μg was selected over BNT162b1 to continue into phase II/III trials due to better tolerability.⁵⁰ The reasons for the difference seen in reactogenicity between the 2 constructs is unclear, however the BNT162b1 molecule is about 5 times larger in size than BNT162b2 and perhaps this was a contributing factor given the direct immunostimulatory properties of RNA.⁵⁶

The safety and efficacy of BNT162b2 was evaluated in an ongoing phase II/III observer-blinded, placebo-controlled, multinational clinical trial of 43,548 participants 16 years or older who were randomized in a 1:1 allocation to receive 30 μg doses of prime/boost vaccine separated by 21 days.⁵⁷ The primary end point is the VE against laboratory confirmed COVID-19 and the study continues to monitor for safety.⁵⁷ At the first interim analysis, after 2 months of median follow up, 8 cases of PCR confirmed COVID-19 were seen in the BNT162b2 recipients (7 days after second dose), whereas 162 cases of PCR confirmed COVID-19 were seen in the placebo group.⁵⁷ This resulted in a VE of 95% (95% CI, 90.3–97.6).⁵⁷ These results were consistent across age, race/ethnicity, sex, body mass index (BMI) and pre-existing medical comorbidities.⁵⁷ Most local and systemic reactions were mild to moderate in severity, less common in those above age 55, and included transient pain at the injection site, fatigue, and headache.⁵⁷ Serious adverse events occurred in 0.6% of the BNT162b2 recipients and 0.5% of placebo recipients and included a ventricular arrhythmia, shoulder injury due to receipt of the vaccine, axillary

lymphadenopathy, and leg paresthesia.⁵⁷ There were no deaths attributed to the vaccine.⁵⁷ Based on this data, the US Food and Drug Administration (FDA) granted the BNT162b2 vaccine emergency use authorization (EUA) on December 11, 2020 and the study was transitioned to an open-label period. Post-EUA with the subsequent vaccination of millions of individuals, a few rare vaccine-related adverse effects have been described. Severe acute allergic reactions (anaphylaxis) were reported by the Centers for Disease Control (CDC) to have occurred in approximately 11.1 cases per million doses after 21 cases were reported to the Vaccine Adverse Event Reporting System (VAERS) from December 14 to 23, 2020, which occurred at a median interval of 13 minutes after receipt of vaccine (range 2–150 minutes).⁵⁸ No deaths due to anaphylaxis were reported and the CDC issued guidance of vaccine-related anaphylaxis encouraging prompt identification and management.⁵⁸ The secondary efficacy analysis was conducted with up to 6 months of post-vaccination data and continued to demonstrate high efficacy against COVID-19 (91% efficacy, 95% CI 89.0–92.3).⁵⁹ It demonstrated a VE of 96.7% against severe disease (95% CI 80.3–99.9) and in data from South Africa, where the Beta variant was dominant, it demonstrated 100% VE.⁵⁹ No new safety signals were identified and the study participants will continue to be followed for 2 years.⁵⁹ On August 23, 2021 the FDA accepted Pfizer-BioNTech's Biologics License Application (BLA) and granted full approval of the BNT162b2 for people 16 years and older for the prevention of COVID-19 disease.

On September 22, 2021, the FDA amended the EUA to allow for a single 30 μ g booster dose at least 6 months following the completed primary vaccine series for people \geq 65, people 18–64 years at high risk for COVID-19 or with increased institutional or occupational exposure to SARS-CoV-2. In the results from the phase I/II/III trial, VE up to 6 months post-dose 2 waned slightly over time (initial VE 96.2% from 7 days to <2 months postdose 2 to 90.1% from >2 months to <4 months postdose 2, and was 83.7% at >4 months postdose 2).⁶⁰ In light of these data, Pfizer-BioNTech administered a 30 μ g booster dose to a small group of participants during phase 1 of the ongoing clinical trial (11 participants 18–55 years of age and 12 participated 65–85 years of age).⁶¹ A VE of 95.6% against COVID-19 was seen from results of the Phase 3 randomized controlled trial assessing the safety and efficacy of a 30 μ g booster dose with a median of 11 months between the booster and completion of the primary vaccine series.⁶¹

More data are emerging regarding the potential of the mRNA vaccines to decrease asymptomatic

COVID-19 and transmission of COVID-19. A real-world observational study of recipients of BNT162b2 vaccine demonstrated a 2.8–4.5-fold reduction in viral load in infections 12 to 37 days after the first dose of vaccine.⁶² Whether or not this correlates with severity of disease or transmissibility is yet to be determined.

As of May 10, 2021, the US FDA granted EUA authorization of BNT162b2 for adolescents ages 12–15 based on interim results from the ongoing randomized, placebo-controlled, observer blinded US clinical trial of 2,260 participants.⁶³ Local and systemic side effects were common and dose dependent, largely mild to moderate in severity and transient including fatigue, myalgias, arthralgias, fever, and chills.⁶³ Immunogenicity in this age group was evaluated and compared with that demonstrated by the adult cohort ages 16–25 years, and was found to be noninferior (geometric mean ratio 1.76 of SARS-CoV-2 50% neutralizing titers 1 month postdose 2 in 12–15 year olds (GMT 1283.0) relative to 16–25 year olds (GMT 730.8)).⁶³ Among 1,005 vaccine recipients there were no PCR confirmed cases of COVID-19, whereas there were 16 cases of confirmed COVID-19 in 978 placebo recipients, with a VE of 100%.⁶³

Data on children 5–11 years of age has demonstrated a VE 90.7% in preventing COVID-19 in this age group without any serious side effects noted in about 3,100 children in an ongoing study. On October 29, 2021 the US FDA granted EUA for use in children 5–11 years of age. Pfizer-BioNTech is currently conducting a phase 1/2/3 clinical trial evaluating a 2 dose-schedule of BNT162b2 in 3 dosing regimens in children 6 months to 11 years old.

Moderna. In a phase 1 open-label trial involving 45 healthy adults ages 18–55, the mRNA-1273 vaccine candidate was evaluated in a dose-escalation study of a prime/boost regimen of 25 μ g, 100 μ g, or 250 μ g doses separated by 28 days to evaluate safety, immunogenicity, and reactogenicity. Systemic adverse events were largely mild to moderate with greater reactogenicity, in both severity and frequency, demonstrated in a dose dependent fashion where 21% reported 1 or more severe adverse events after the 250 μ g doses.⁶⁴ It demonstrated potent immunogenicity with neutralizing antibodies to S-2P and the RBD in a dose dependent manner after just 1 dose in all participants that was comparable to levels seen in convalescent serum.⁶⁴ Of the 3 doses evaluated, the 100 μ g prime/boost regimen was associated with robust immunogenicity and an improved reactogenicity profile compared with the 250 μ g dose regimen (S-2P ELISA GMT at day 57 was 782,719 in the 100 μ g-group and 1,192,154 in the 250 μ g-group; for comparison, convalescent serum GMT is 142,140).⁶⁴

The phase 2a placebo-controlled, observer-blinded, dose-confirmation trial in 600 healthy adults ages 18–55 and a sentinel group of 55 and older, to evaluate reactogenicity, safety, and immunogenicity of prime/boost regimen of 50 μg , 100 μg , or placebo 28 days apart.⁶⁵ The data on this is still preliminary, however both binding and neutralizing antibodies were seen after both dosing regimens in a dose dependent manner after the first dose but more equivalent after the second dose.⁶⁵

The phase 3 placebo-controlled, observer-blinded study evaluated the prime/boost regimen of 100 μg versus placebo, separated by 28 days, in 30,420 SARS-CoV-2-naïve participants to evaluate for prevention of COVID-19 illness with onset 14 days after second vaccination.⁶⁶ At the interim analysis, with a median follow up of 64 days, there were 11 cases of symptomatic COVID-19 in the mRNA-1273 vaccine recipient group and 189 cases in the placebo group, with a reported VE of 94.1% (95% CI, 89.3%–96.8%; $P < 0.001$).⁶⁶ Reactogenicity was moderate, transient, and more common in the vaccine group.⁶⁶ There were 30 cases of severe COVID-19 and all were in the placebo group.⁶⁶ Severe adverse events were rare with a similar incidence in both groups.⁶⁶ The US FDA granted the mRNA-1273 vaccine EUA on December 18, 2020 and the study design was transitioned to open-label. The secondary efficacy analysis occurred at the completion of the blinded phase of the study, with a median follow up of 148 days, and continued to demonstrate high efficacy at preventing COVID-19 illness (93.2% efficacy, 95% CI 91.0%–94.8%). It also continued to demonstrate high efficacy at preventing severe COVID-19 with only 2 cases of severe COVID-19 in the vaccine group and 106 in the placebo group, a VE of 98.2% (95% CI 92.8%–99.6%).⁶⁷ Asymptomatic SARS-CoV-2 infections were also reduced (VE 63%).⁶⁷ The mRNA-1273 vaccine continues to show an acceptable safety profile and study participants will be followed for 2 years. The safety and efficacy reported in these clinical trials to date is summarized in [Table 2](#) and [Table 3](#) respectively.

In a preprint of an exploratory analysis of the Moderna study, the incidence of COVID-19 infection during delta variant predominate period (July–August 2021) were compared in those who were vaccinated early (14,746 participants vaccinated from July to December 2020, 13 months median follow up, mRNA-1273e) vs later (11,431 participants vaccinated from December 2020 to April 2021, 7.9 months of median follow up, mRNA-1273p).^{68(p19)} During this period of time, 162 COVID-19 cases were seen in the mRNA-1273e group and 88 COVID-19 cases in the mRNA-1273p group (incidence rate 49.0 per 1000 person years in the later vaccinated group compared with 77.1 per

1000 person years in the early vaccinated group).^{68(p19)} Additionally, the severe COVID-19 cases were fewer in the later vaccinated mRNA-1273p compared with the mRNA-1273e group. Most of the sequenced cases were attributed to the delta variant (97% in mRNA-1274e and 99% in mRNA-1273p groups).^{68(p19)}

On October 20, 2021 the FDA amended the EUA to allow for a 50 μg dose of mRNA-1273 as a booster dose at least 6 months after completion of the primary vaccine series for people ≥ 65 , people 18–64 years at high risk for COVID-19 or with increased institutional or occupational exposure to SARS-CoV-2. This was based on review of data from the Phase 2 clinical trial, during which a 50 μg dose of mRNA-1273 as a booster dose was given to 344 interested participants 6–8 months following the second dose of a priming series. Per a Moderna press release on October 20, 2021, waning of neutralizing antibody titers was seen at 6 months, particularly against the variants of concern (VOC), but a booster dose of 50 μg dose of mRNA-1273 significantly elevated neutralizing titers. These effects were seen across all age groups and the reported safety profile was similar to the post-dose 2 profile.⁶⁹

Studies in adolescents have also reported promising results and are currently underway. The phase 2/3 TeenCove study includes 3,732 adolescents from 12 to less than 18 years of age who were randomized 2:1 to receive prime/boost of 100 μg of mRNA-1273 or placebo.⁷⁰ Based on the interim analysis, VE was reported at 100% with 4 cases of COVID-19 in the placebo group and none in the vaccinated group, and no severe adverse events reported.⁷⁰ The analysis of the immunogenicity subset (which included 289 adolescents and 289 young adults (18– ≤ 25)) demonstrated the immune response in adolescents was non-inferior to young adults (geometric mean ratio 1.08 of SARS-CoV-2 50% neutralizing titers 1 month post-dose 2 in 12– < 18 year olds (GMT 807) relative to 18– ≤ 25 year olds (GMT 740)).⁷⁰ Side effects in this adolescent group were comparable to the adult group with the large majority reported as mild to moderate in severity.⁷⁰

Evaluations in children have been initiated. Moderna is currently conducting a phase 2/3 two-part dose escalation, open label, observer blinded, placebo controlled clinical trial of a 2-dose regimen of mRNA-1273 in children 6 months to 11 years old.⁷¹

Special populations. Immunocompromised. Immunocompromised patients, including those on immunosuppressive medications, and those with autoimmune conditions were specifically excluded from the clinical trials. No safety concerns have been identified in this group but current data shows that the mRNA vaccines are less effective at stimulating an immune response in this population.⁷² A large study of 658 transplant

Table 2. Safety analyses data of mRNA COVID-19 vaccines from clinical trials

Adverse Event	BNT162b2 Phase II/III clinical trial after 2 mo of follow-up ⁵⁷	BNT162b2 Phase II/III clinical trial after 6 mo of follow-up ⁵⁹	mRNA-1273 Phase III clinical trial after 2 mo of follow-up ⁶⁶	mRNA-1273 Phase III clinical trial after 5.3 mo of follow-up ⁶⁷
Any Local	-	-	84.2 (N = 15,168)	84.2 (N = 15,166)
Dose 1 (%)			88.6 (N = 14,677)	88.7 (N = 14,691)
Dose 2 (%)				
Any Systemic	-	-	54.9 (N = 15,168)	54.8 (N = 15,166)
Dose 1 (%)			79.4 (N = 14,677)	79.5 (N = 14,691)
Dose 2 (%)				
Any Serious (%)	0.6 (N = 21,621)	0.6 (N = 21,926)	0.6 (N = 15,185)	0.6 (N = 15,184)
Related to study vaccination (%)	26.7 (N = 21,621)	23.9 (N = 21,926)	4.5% (N = 15,185)	13.6% (N = 15,184)
0 (N = 21,621)		0 (N = 21,926)	<0.1 (N = 15,185)	<0.1 (N = 15,184)
Serious (%)	0 (N = 21,621)	0 (N = 21,926)	0 (N = 15,185)	0 (N = 15,184)
Fatal (%)	-	-	0.9 (N = 15,185)	1.4 (N = 15,184)
Medically attended (%)	-	-	0.5 (N = 15,185)	0.5 (N = 15,184)
Severe (%)				

recipients found that only 15% were able to produce measurable antibodies after dose 1 and 2 and 46% did not have an antibody response.⁷² A double-blind, placebo-controlled, randomized study of 120 solid organ transplant recipients demonstrated that a third dose of mRNA-1273 provided increased median percent virus neutralization (71% mRNA-1273 group vs 13% placebo) and improved neutralizing antibodies (60% mRNA-1273 group vs 25% placebo achieved above the 30% threshold).⁷³ In a French study evaluating 101 solid-organ transplant recipients who previously received 2 doses of BNT162b2 vaccine, 44% seroconverted after a third dose of BNT162b2 (26 of 59 patients).⁷⁴ Data continue to emerge emphasizing that solid organ transplant recipients remain a particularly vulnerable population despite receipt of a 2-dose primary vaccination series.^{75,76} Reduced antibody

responses after a 2-dose primary vaccination series has also been observed in those with chronic lymphocytic leukemia (CLL) after vaccination with BNT162b2, with a antibody response rate of 39.5% reported in one study.⁷⁷ The study noted that those in clinical remission were most likely to respond to vaccination (79.2%) whereas those who hadn't yet received treatment for their malignancy responded at a lower rate (55.2%).⁷⁷ The responses in those actively receiving treatment were poor (16%) and none of those who received an anti-CD20 agent within 12 months of vaccination responded.⁷⁷ Not much is known yet about the magnitude of the role of cell-mediated immune responses to COVID-19 in this population, however recent data suggests that CD4⁺ T-cell responses to the S-protein may positively correlate with anti-SARS-CoV-2 IgG and IgA titers.⁷⁸ Data from the VISION Network compared

Table 3. Efficacy of the mRNA COVID-19 vaccines from clinical trials

Trial	SARS-CoV-2 infection	COVID-19*	Severe COVID-19*	Death due to COVID-19
BNT162b2 Phase II/III clinical trial after 2 months of follow-up ⁵⁷	-	94.6 (95% CI 89.9–97.3)	75% (95% CI –152.6, 99.5)	100.0
mRNA-1273 Phase III clinical trial after 2 mo of follow-up ⁶⁶	-	94.1 (95% CI 89.3–96.8)	100.0% (95% CI could not be estimated)	100.00
BNT162b2 Phase II/III clinical trial after 6 mo of follow-up ⁵⁹	58.4% (95% CI 40.8–71.2)	91.3 (95% CI 89.0–92.3)	96.7% (95% CI 80.3–99.9) *as defined by the FDA	100.0
mRNA-1273 Phase III clinical trial after 5.3 mo of follow-up ⁶⁷	63.0% (95% CI 56.6–68.5)	93.2% (95% CI 91.0–94.8)	98.2% (95% CI 92.8–99.6)	100.00

*See reference for definition of COVID-19 and severe COVID-19.

VE against COVID-19 associated hospitalization after receipt of a 2-dose vaccination series in 20,101 immunocompromised adults and 69,116 immunocompetent adults.⁷⁹ VE was found to be lower in the immunocompromised patients (77%) compared with the immunocompetent patients (90%) irrespective of age, delta variant predominance, and mRNA vaccine product.⁷⁹ VE point estimates demonstrated a VE of 79% in those with solid malignancy, 74% with hematologic malignancy, 81% with a rheumatologic or inflammatory disorder, 73% with an intrinsic immune or immunodeficiency, 59% in solid organ or stem cell transplant recipients. Generally, the VE point estimates were higher for those who received 2 doses of Moderna compared with Pfizer-BioNTech except in those with a rheumatologic or inflammatory disorder (they were the same).⁷⁹ On August 12, 2021, the FDA amended the EUA for the Pfizer-BioNTech and Moderna COVID-19 vaccines, which allowed moderately to severely immunocompromised individuals to receive an additional dose of COVID-19 vaccination for a 3 total dose primary vaccine series. One day later the CDC issued a statement supporting the FDA's amendment. Several studies have demonstrated that 33%–50% of immunocompromised people developed detectible antibodies after a 3rd dose when they had not after a 2-dose primary vaccination series.^{74,80-82} The CDC has since allowed for this population to receive a 4th dose of mRNA COVID-19 vaccination. Additional data regarding the VE after a 3-dose primary vaccination series are needed.

Pregnancy. Preliminary findings evaluating safety of mRNA-1273 and BNT162b2 in 35,691 pregnant women from ages 16 to 54 via the V-safe surveillance data system did not show any concerning safety signals in pregnancy or neonatal outcomes.⁸³ A descriptive, exploratory study in a cohort of 103 women including 30 pregnant and 16 lactating women, evaluated the immunogenicity after mRNA COVID-19 vaccination.⁸⁴ There was demonstrated immunogenicity in this population of women, including humoral and T cell mediated immunity, and vaccine associated antibodies were conferred via cord blood and breast milk to the infant.⁸⁴ Both binding and neutralizing antibodies were also seen against 2 variants, B.1.1.7 (Alpha) and B.1.351 (Beta).⁸⁴ Data from Norway found no evidence to suggest early pregnancy loss after COVID-19 vaccination, with a reported adjusted odds ratio of 0.8 (95% CI 0.67–0.96) after Pfizer-BioNTech and 0.84 (95% CI 0.56–1.25) after Moderna within the prior 5 weeks of vaccination.⁸⁵ Another study including 105,466 pregnancies showed that 8.0% of women with ongoing pregnancies had received a mRNA COVID-19 vaccine in the prior 28 days relative to an index date vs

8.6% of spontaneous abortions and the spontaneous abortions were not associated with an increased odds of COVID-19 vaccine exposure compared with ongoing pregnancies (adjusted odds ratio 1.02; 95% CI 0.96–1.08).⁸⁶ Emerging data continues to show that COVID-19 vaccination (including BNT162b2 and mRNA-1273) in pregnant women is safe, effective, and immunogenic, allowing transfer of antibodies to the neonates.⁸⁷⁻⁸⁹ In August 2021 the CDC recommended COVID-19 vaccination for pregnant women.

Adolescent and Pediatric populations are discussed in their respective sections above.

Side effects. Although the phase 3 studies of both mRNA-1273 and BNT162b2 included rigorous evaluations for safety and side effects, the post-authorization administration of mRNA vaccines to millions of people has identified rare events that were not seen in the clinical trials as they were not powered to detect these. In the most rigorous vaccine safety monitoring effort in US history, several vaccine safety-monitoring systems have been implemented to further identify and track adverse events during the mRNA vaccine rollout. The Vaccine Safety Datalink (VSD) includes 9 healthcare sites in collaboration with the CDC and links patient vaccination and EMR data. VAERS collects reports of AEs postvaccination nationwide. V-Safe lets users use their smartphone to report post-vaccination symptoms and health concerns, it also has a pregnancy registry to monitor those who received a vaccine just before or during pregnancy. The Clinical Immunization Safety Assessment Project (CISA) is a panel of vaccine safety experts from the CDC who collaborate with 7 research centers across the US to monitor vaccine safety.⁹⁰ VAERS and V-Safe are self-reporting modalities that are available to the public and allow for rapid identification of potential safety signals. The limitation is that this information is subject to reporting bias and potentially incomplete data.

Rare cases of anaphylaxis after COVID-19 mRNA vaccination were reported via VAERS with a reporting rate of 4.7 cases/million doses administered after Pfizer-BioNTech vaccine and 2.5 cases/million doses administered after Moderna vaccine.⁹¹ A larger study of 64,900 hospital employees reported severe allergic reactions consistent with anaphylaxis at a rate of 2.47 per 10,000 mRNA COVID-19 vaccination.⁹² Although the mechanism of anaphylaxis is still unclear, 31% of those with anaphylactic reactions had a history of anaphylaxis.⁹² Rare cases of myocarditis and pericarditis have been seen post-mRNA COVID-19 vaccination.^{93,94} Although the incidence is difficult to estimate with data rapidly accumulating, VAERS data as of October 6, 2021 showed 3,336 cases of myocarditis and pericarditis (2,459 myopericarditis, 877

pericarditis alone), primarily in younger males within 7 days after the second dose of mRNA vaccine (reporting rates of myocarditis per 1 million doses administered within 7 days after the second dose of an mRNA vaccine ranged from 2.8 to 37.4 in those 12–39 years of age).⁹⁵ From a retrospective review of data from Israel, the incidence of myocarditis remained rare but was increased after vaccination, particularly in males 16–19 years of age (rate ratio 30 days after second vaccination 8.96 in comparison to unvaccinated persons).⁹⁶ Another large database study from Israel estimated the incidence of myocarditis after at least 1 dose of mRNA vaccine was 2.13 cases per 100,000 people, with the highest in males 16–29 years of age (10.69 cases per 100,000 persons).⁹⁷ Data from the VSD demonstrated an adjusted rate ratio of 1.72 for myocarditis/pericarditis in the 21 day interval after any dose of an mRNA vaccine.⁹⁸ A subgroup analysis further demonstrated in data through October 9, 2021 that those age 12–39 are at increased risk, especially through 7 days post vaccination, with significantly higher rates after Moderna than after Pfizer-BioNTech (8.0 excess cases in the risk period per 1 million doses of Moderna versus Pfizer-BioNTech within 2 days after either vaccine, 13.3 within 7 days of the second dose).⁹⁸ Other studies which show a greater risk of myocarditis/pericarditis after Moderna than Pfizer-BioNTech include studies from Canada⁹⁹ and Scandinavia.¹⁰⁰ However, other US vaccine safety monitoring systems including VAERS, FDA Biologics Effectiveness and Safety Systems, and the Department of Veterans Affairs (VA), did not show a difference in rate between the 2 mRNA vaccines. The incidence of myocarditis after dose 3 of mRNA vaccines is not yet available as data is only available in a small number of volunteers from the clinical trials, but is forthcoming.

Effectiveness of mRNA vaccines. The clinical trials continue to demonstrate the efficacy and safety of the COVID-19 vaccines and emerging real-world data corroborates these findings. Table 4 summarizes the real-world effectiveness data from selected key studies. Data from Israel after launching its COVID-19 vaccination program on December 20, 2020, showed that VE of the BNT162b2 vaccine was 92% for any SARS-CoV-2 infection, 94% for symptomatic COVID-19, 87% for COVID-19 related hospitalization, and 92% for severe COVID-19.¹⁰¹ Real-world data of vaccine effectiveness in 7,280 older adults ≥ 65 years from the COVID-19-Associated Hospitalization Surveillance Network (COVID-NET) re-demonstrated high effectiveness in this population.¹⁰² Of fully vaccinated adults aged 65–74 years, VE at preventing COVID-19-associated hospitalization was 96% for both BNT162b2 and mRNA-1273 recipients.¹⁰² In those

who were fully vaccinated but ≥ 75 years, the efficacy was 91% for BNT162b2 and 96% for mRNA-1273 recipients.¹⁰²

Ongoing assessments of vaccine effectiveness remains important as SARS-CoV-2 VOC continue to emerge in the setting of concern for waning vaccine-induced immunity.¹⁰³ The data on the durability of vaccine-induced immunity and VE against the current VOC, including the Alpha, Beta, and Delta variants, is variable. This is likely due to differences in study population and design including variation in methods of data analysis, end points of interest, diagnostic testing utilized, and prevalence of SARS-CoV-2 variants.^{103,104} There has been suggestion of reduced neutralization in vitro against the Beta variant in individuals vaccinated with BNT162b2.¹⁰⁵ A case-controlled observational study in Qatar, which assessed the vaccine effectiveness of BNT162b2 against any infection with the Beta or Alpha variants, estimated vaccine effectiveness against the Beta variant was 75% and 89.5% against the Alpha variant.¹⁰⁶ However, demonstrated effectiveness against severe, critical, or fatal disease with COVID-19 remained high at 97.4%.¹⁰⁶ Several additional studies have continued to demonstrate that despite the apparent in vitro reduction of neutralization against the Beta variant, vaccine effectiveness does not appear to diminish.¹⁰⁷ In an observational study evaluating the effectiveness of the mRNA vaccines against infection with SARS-CoV-2 in residents of nursing homes, vaccine effectiveness demonstrated a notable decline, from 74.7% (74.2% for Pfizer-BioNTech, 74.7 for Moderna) to 53.1% (52.4% for Pfizer-BioNTech, 50.6% for Moderna), in the pre-Delta period compared with the Delta-predominant period in the US.¹⁰³ Data from the UK found vaccine effectiveness against symptomatic COVID-19 after BNT162b2 to be 88.0% against the Delta variant (compared with 93.7% against Alpha).¹⁰⁸ Retrospective data from the US demonstrated a VE of 73% (95% CI 72–74) against SARS-CoV-2 infection after receipt of BNT162b2 vaccine series, but did show a decline from 1 month to 5 months after vaccination (88% and 47% VE respectively).¹⁰⁹ Another study conducted across 5 US VA hospitals showed a VE against COVID-19 related hospitalizations of 87% even during a period of widespread Delta variant prevalence. (Pfizer-BioNTech 83.4%; 95% CI 74–89.4; Moderna 91.6%; 95% CI 83.5–95.7)¹¹⁰ Data from the HEROS-RECOVER Cohort showed a VE of 80% (95% CI 69–88) against SARS-CoV-2 infection during the study period of December 14, 2020 to August 14, 2021.¹¹¹ Additional studies using realworld data continue to support these findings (Table 4).^{103,112–115} Both the BNT162b2 and mRNA-1273 continue to demonstrate evidence of high and sustained protection against severe

Table 4. Real world effectiveness of mRNA COVID-19 vaccines from selected studies

Study	Vaccine	Study period	Study population	Vaccine efficacy (%)
Tartof et al ¹⁰⁹	BNT162b2	12/14/20-8/8/21	≥ 12 y old Kaiser Permanente (California) patients	Symptomatic COVID-19 73 (95% CI 72–74) COVID-19 associated hospitalization 90 (95% CI 89–92)
Bajema et al ¹¹⁰ SUPERNOVA Network	BNT162b2, mRNA-1273	2/1/21-8/6/21	US veterans ≥ 18 y	COVID-19 associated hospitalization 87 (95% CI 90.4–91.1) BNT162b2 83.4 (95% CI 74–89.4) mRNA-1273 91.6 (95% CI 83.5–95.7)
Fowlkes et al ¹¹¹ HEROS-RECOVER Cohort	BNT162b2, mRNA-1273, Ad26.COV2.S	12/14/20-8/14/21	Healthcare workers	Any SARS-CoV-2 Infection 80 (95% CI 69–88)
Rosenberg et al ¹¹²	BNT162b2, mRNA-1273, Ad26.COV2.S	5/3/21-7/19/21	Adults ≥ 18 y (New York)	Any SARS-CoV-2 Infection 91.8–75.0 COVID-19 associated hospitalization >89%
Thompson et al ¹¹³	BNT162b2, mRNA-1273	1/1/21-6/22/21	Adults ≥ 50 y/o	COVID-19 associated hospitalization 89 (95% CI 87–91) 90 (95% CI 86–93) for ICU admission 91 (95% CI 89-93) for ER/urgent care
Grannis et al ¹¹⁴ VISION Network	BNT162b2, mRNA-1273, Ad26.COV2.S	June-August 2021	Adults ≥ 18	COVID-19 associated hospitalization 86 (95% CI 82–89) BNT162b2 80 Moderna 95 82 for COVID-19 ED/UC encounters
Study	Vaccine	Study Period	Study Population	Vaccine Efficacy (%)
Nanduri et al ¹⁰³ NHSN	BNT162b2, mRNA-1273	March - August 2021	Nursing home residents	Any SARS-CoV-2 Infection 74.7 (95% CI 70.0–78.8) during pre-Delta (BNT162b2 74.2, mRNA-1273 74.7) 53.1 (95% CI 49.1–56.7) during Delta (BNT162b2 52.4, mRNA-1273 50.6)
Goldberg et al ¹¹⁵ Israeli national database	BNT162b2	7/11/21-7/31/21	≥ 16 y	Any SARS-CoV-2 Infection 52-83 Severe COVID-19 85–98

SUPERNOVA Network: Surveillance Platform for Enteric and Respiratory Infectious Organisms at the VA; HEROS-RECOVER Cohort: Arizona Healthcare, Emergency Response and Other Essential Workers Surveillance Study (HEROES); Research on the Epidemiology of SARS-CoV-2 in Essential Response Personnel (RECOVER); VISION Network: VISION network includes Columbia University Irving Medical Center (New York), HealthPartners (Minnesota and Wisconsin), Intermountain Healthcare (Utah), Kaiser Permanente Northern California (California), Kaiser Permanente Northwest (Oregon and Washington), Regenstrief Institute (Indiana), and University of Colorado (Colorado); NHSN: National Healthcare Safety Network, Emergency Department (ED); Urgent Care (UC).

COVID-19 disease, even in high risk populations, through 24 weeks after vaccination.¹¹⁶ Additionally, studies continue to show a reduction in severe COVID-19-associated illness in highly vaccinated populations.¹¹⁷ One study reported that cumulative

hospitalizations were 17 times higher in the unvaccinated from January 24, 2021 to July 24, 2021 compared with vaccinated people.¹¹⁸ Moderna has developed 2 additional vaccine candidates, mRNA-1273.351 targeting the Beta variant and mRNA-1273.211 with a 50/50

mix of mRNA-1273 and mRNA-1273.351, and has entered into clinical trials with these constructs.¹¹⁹ Pre-clinical studies from mice show that mRNA-1273.351 successfully increased neutralizing antibodies to the Beta variant, while the mRNA-1273.211 induced broad neutralization against multiple VOC (D614G, B.1.351, P.1, B.1.427/B.1.429).¹¹⁹ Whether future third dose booster vaccinations will target VOC or the initial reference strain is an area of ongoing evaluation. In a multi-state study of 3,089 hospitalized adults, VE against COVID-19-associated hospitalization was found to be sustained with 86% at 2–12 weeks post vaccination and 84% at 13–24 weeks post vaccination for patients fully vaccinated against SARS-CoV-2 with a COVID-19 mRNA vaccine (mRNA-1273 or BNT162b2).¹¹⁶ These data were consistent even across subgroups at high risk for severe disease, although adults with an immunocompromising condition had an overall lower VE.¹¹⁶ The secondary analysis data of Pfizer-BioNTech's BNT162b2 vaccine demonstrated a peak VE against lab-confirmed COVID-19 of 96.2% from 7 days to less than 2 months after the second vaccine dose, however went on to demonstrate that VE declined an average of 6% every 2 months thereafter.⁵⁹ Some emerging data have suggested that the VE for Moderna is higher than for Pfizer-BioNTech (93% vs 88%) based on realworld data among US adults.¹²⁰

Realworld VE data of Pfizer-BioNTech against COVID-19 associated hospitalization in adolescents has so far demonstrated a VE of 93% among 464 hospitalized persons aged 12–18.¹²¹

Heterologous boost. Given the available data on homologous booster vaccinations has been shown to be safe and immunogenic, evaluations of heterologous booster vaccinations are underway. In a preprint of a US phase 1/2 nonrandomized, open-label clinical trial, previously vaccinated adults received 1 of 3 vaccines (Moderna 100 μ g, Janssen Ad26.COVS.2.S 5×10^{10} virus particles, or Pfizer-BioNTech BNT 162b2 30 μ g).¹²² The initial data demonstrated that heterologous booster vaccines resulted in similar or higher neutralizing antibody titers than homologous booster vaccines (6.2–76 fold vs 4.6–56 fold), with mRNA vaccines demonstrating the higher antibody titers within 28 days of the boost.¹²²

CureVac. The mRNA vaccine candidate from CureVac (CVnCoV) encoding the stabilized full-length spike protein encapsulated in a LNP was initially a promising contender of very similar design to the mRNA-1273 and BNT162b2.¹²³ In the phase 1 assessment, all investigated doses provoked an immune response with 56%–77% demonstrating virus neutralization test seroconversion 2 weeks after lower doses (2–8 μ g dose level) and 100% after the 12 μ g dose

level.¹²⁴ In a phase 2b/3 international randomized, double-blinded, placebo-controlled study of 40,000 adults 18 years of age or older, the interim VE was 47% against COVID-19 disease of any severity.¹²³ Given these disappointing efficacy results, the vaccine did not move forward. The reasons for the failure of CVnCoV despite being very similar to the mRNA vaccine constructs from Pfizer-BioNTech and Moderna are unclear but may be related to the use of modified mRNA. The CVnCoV vaccine contained uridine instead of the modified nucleoside pseudouridine found in the Pfizer-BioNTech and Moderna products.^{107,123} This may have resulted in differences in mRNA translation or innate immunity responses.¹⁰⁷ Several other mRNA vaccine constructs are currently in phase I to III clinical trials but at this time the majority lack clinical data and therefore will not be discussed in this review.

DISCUSSION

New and emerging pathogens continue to threaten human health globally and preparedness and pandemic response are key for early containment of outbreaks. Coronaviruses are zoonotic diseases that have previously been predicted to be highly capable of causing pandemics.¹ In the last 20 years SARS-CoV-2 is the third novel betacoronavirus to cause significant disease after SARS-CoV, first reported in China in 2002, and MERS-CoV, first reported in Saudi Arabia in 2012.¹

Since the first studies exploring mRNA therapeutics in the 1970s, the collective scientific effort to evaluate and develop mRNA as a viable preventative therapeutic for infectious diseases has made tremendous progress and demonstrated promising results. Despite the common misconception that mRNA vaccines are a new development, they are actually the culmination of decades of scientific research and have been under research for years for various infectious diseases including Ebola, CMV and Zika viruses. The mRNA vaccines BNT162b2 and mRNA-1273 have played a key role in our global response to the COVID-19 pandemic as they have demonstrated significant advantages over conventional vaccines. Compared with the other available vaccines against SARS-CoV-2, the mRNA vaccines have demonstrated higher antibody responses.¹⁰⁷ While many of the studies to date utilize convalescent serum as a comparator, the correlate of protection is still unknown and this is an area of ongoing investigation although there is mounting evidence that neutralizing antibodies to the spike protein play an important role.^{125,126} Data from the UK evaluating humoral immune responses to with the ChAdOx1 nCoV-19 vaccine suggests that an 80% VE against

symptomatic SARS-CoV-2 (in the Alpha variant dominated setting) was demonstrated with 264 anti-spike binding antibody units (BAU/mL), 506 BAU/mL for anti-RBD antibodies, 26 IU/mL pseudovirus neutralization, and 247 normalized live-virus neutralization titers.¹²⁷ It is possible that these responses are dependent on different vaccine platforms, therefore conclusions should be put into the appropriate context until more data is available. Data from the mRNA vaccines is still in the preprint form, however appears to show a similar trend.¹²⁸ While the role of T cell responses in protective immunity against SARS-CoV-2 is still an ongoing area of research, some evidence suggests that CD4⁺ T cell responses to the S-protein were vigorous after vaccination and correlated with anti-SARS-CoV-2 IgG and IgA levels.⁷⁸ T cells have been demonstrated to play a role in prevention of progression of disease and duration of RNA shedding, however their role as a correlate of protection is still unknown.¹²⁹ Goel et al. showed that SARS-CoV-2 specific memory B cells increased in the 3-6 months after mRNA vaccination, despite a decline in antibody levels.¹³⁰ Several studies support the finding that both antigen specific CD4⁺ and CD8⁺ T cells are induced by mRNA vaccination against COVID-19 and appear to be durable through 6 months post-vaccination.¹³⁰⁻¹³² Given the observation that protection against severe COVID-19 and death persists despite evidence of waning antibodies, it is reasonable to hypothesize that cellular immunity including memory responses likely significantly contribute, however further studies are required. In line with this hypothesis, the afforded protection against severe Covid-19 has been challenged with 2 viral variants so far, Delta and Omicron, which were associated with a ~3 and 25–50- fold reduction in neutralization by antibodies raised against the vaccine strain.^{133,134} A large study from Kaiser Permanente has shown that the emergence of the delta variant resulted in diminution of protection against infection with the Delta variant but not against Covid-19 hospitalizations, which remained at 94.5% 6-8 months after vaccination.¹³⁵ Early epidemiologic data from the US indicate that infections with the Omicron variant occur after previous infection, 2 doses of mRNA Covid-19 vaccines, and boosting albeit with characteristically mild outcomes.¹³⁶

This is a rapidly evolving area of study and data continue to emerge. The COVID-19 mRNA vaccines from Pfizer-BioNTech and Moderna have proven to be highly efficacious against COVID-19-associated hospitalization and severe disease. Their efficacy has been demonstrated across different age groups, racial and ethnic groups, and coexisting medical comorbidities. Due to the success of the mRNA vaccines during the

COVID-19 pandemic, Pfizer-BioNTech and Moderna are exploring their application to other infectious disease pathogens and COVID-19 may be just the beginning. The application of mRNA vaccines to a wide variety of other infectious disease pathogens is an area of rapid growth, including HIV, influenza, RSV, CMV, hepatitis C, Nipah virus, Zika virus, human metapneumovirus, and parainfluenza type 3. Some of these recent mRNA vaccine constructs are expected to enter phase 1 clinical trials as early as this year.

ACKNOWLEDGMENTS

Conflicts of Interest: All authors have read the journal's policy on disclosure of potential conflicts of interest. CMA has no conflicts of interest. HMES has no conflicts of interest.

There was no funding for this work.

All authors have read the journal's authorship agreement and the manuscript has been reviewed and approved by all authors.

REFERENCES

1. Corbett KS, Edwards DK, Leist SR. SARS-CoV-2 mRNA vaccine design enabled by prototype pathogen preparedness. *Nature* 2020;586(7830):567–71. <https://doi.org/10.1038/s41586-020-2622-0>.
2. Pardi N, Hogan MJ, Porter FW, Weissman D. mRNA vaccines — a new era in vaccinology. *Nat Rev Drug Discov* 2018;17:261–79. <https://doi.org/10.1038/nrd.2017.243>.
3. Rauch S, Jasny E, Schmidt KE, Petsch B. New vaccine technologies to combat outbreak situations. *Front Immunol* 2018;9. <https://doi.org/10.3389/fimmu.2018.01963>.
4. Sahin U, Karikó K, Türeci Ö. mRNA-based therapeutics—developing a new class of drugs. *Nat Rev Drug Discov* 2014;13:759–80. <https://doi.org/10.1038/nrd4278>.
5. Laskey RA, Gurdon JB, Crawford LV. Translation of encephalomyocarditis viral RNA in oocytes of *Xenopus laevis*. *Proc Natl Acad Sci U S A*. 1972;69:3665–9. <https://doi.org/10.1073/pnas.69.12.3665>.
6. Gurdon JB, Lane CD, Woodland HR, Marbaix G. Use of frog eggs and oocytes for the study of messenger RNA and its translation in living cells. *Nature* 1971;233:177–82. <https://doi.org/10.1038/233177a0>.
7. Malone RW, Felgner PL, Verma IM. Cationic liposome-mediated RNA transfection. *Proc Natl Acad Sci U S A*. 1989;86:6077–81. <https://doi.org/10.1073/pnas.86.16.6077>.
8. Wolff JA, Malone RW, Williams P. Direct gene transfer into mouse muscle in vivo. *Science* 1990;247:1465–8. <https://doi.org/10.1126/science.1690918> 4949 Pt 1.
9. Martinon F, Krishnan S, Lenzen G. Induction of virus-specific cytotoxic T lymphocytes in vivo by liposome-entrapped mRNA. *Eur J Immunol* 1993;23:1719–22. <https://doi.org/10.1002/eji.1830230749>.
10. Alberer M, Gnad-Vogt U, Hong HS. Safety and immunogenicity of a mRNA rabies vaccine in healthy adults: an open-label, non-randomised, prospective, first-in-human phase 1 clinical trial. *Lancet Lond Engl* 2017;390:1511–20. [https://doi.org/10.1016/S0140-6736\(17\)31665-3](https://doi.org/10.1016/S0140-6736(17)31665-3).

11. Schlake T, Thess A, Fotin-Mleczek M, Kallen KJ. Developing mRNA-vaccine technologies. *RNA Biol* 2012;9:1319–30. <https://doi.org/10.4161/rna.22269>.
12. Pardi N, Muramatsu H, Weissman D, Karikó K. In vitro transcription of long RNA containing modified nucleosides. *Methods Mol Biol Clifton NJ* 2013;969:29–42. https://doi.org/10.1007/978-1-62703-260-5_2.
13. Krieg PA, Melton DA. Functional messenger RNAs are produced by SP6 in vitro transcription of cloned cDNAs. *Nucleic Acids Res* 1984;12:7057–70. <https://doi.org/10.1093/nar/12.18.7057>.
14. Pascolo S. Vaccination with messenger RNA (mRNA). *Handb Exp Pharmacol* 2008;221:35. https://doi.org/10.1007/978-3-540-72167-3_11.
15. Jackson NAC, Kester KE, Casimiro D, Gurnathan S, DeRosa F. The promise of mRNA vaccines: a biotech and industrial perspective. *Npj Vaccines* 2020;5:1–6. <https://doi.org/10.1038/s41541-020-0159-8>.
16. Liang SL, Quirk D, Zhou A. RNase L: its biological roles and regulation. *IUBMB Life* 2006;58:508–14. <https://doi.org/10.1080/15216540600838232>.
17. Karikó K, Muramatsu H, Ludwig J, Weissman D. Generating the optimal mRNA for therapy: HPLC purification eliminates immune activation and improves translation of nucleoside-modified, protein-encoding mRNA. *Nucleic Acids Res* 2011;39. <https://doi.org/10.1093/nar/gkr695> e142.
18. Gallie DR. The cap and poly(A) tail function synergistically to regulate mRNA translational efficiency. *Genes Dev* 1991;5:2108–16. <https://doi.org/10.1101/gad.5.11.2108>.
19. Banerjee AK. 5'-terminal cap structure in eucaryotic messenger ribonucleic acids. *Microbiol Rev* 1980;44:175–205.
20. Holtkamp S, Kreiter S, Selmi A. Modification of antigen-encoding RNA increases stability, translational efficacy, and T-cell stimulatory capacity of dendritic cells. *Blood* 2006;108:4009–17. <https://doi.org/10.1182/blood-2006-04-015024>.
21. Gustafsson C, Govindarajan S, Minshull J. Codon bias and heterologous protein expression. *Trends Biotechnol* 2004;22:346–53. <https://doi.org/10.1016/j.tibtech.2004.04.006>.
22. Kudla G, Lipinski L, Cuffin F, Helwak A, Zyllicz M. High guanine and cytosine content increases mRNA levels in mammalian cells. *PLoS Biol* 2006;4:e180. <https://doi.org/10.1371/journal.pbio.0040180>.
23. Houseley J, Tollervey D. The many pathways of RNA degradation. *Cell* 2009;136:763–76. <https://doi.org/10.1016/j.cell.2009.01.019>.
24. Kowalski PS, Rudra A, Miao L, Anderson DG. Delivering the messenger: advances in technologies for therapeutic mRNA delivery. *Mol Ther J Am Soc Gene Ther* 2019;27:710–28. <https://doi.org/10.1016/j.ymthe.2019.02.012>.
25. Yanez Arteta M, Kjellman T, Bartesaghi S. Successful reprogramming of cellular protein production through mRNA delivered by functionalized lipid nanoparticles. *Proc Natl Acad Sci U S A* 2018;115. <https://doi.org/10.1073/pnas.1720542115> E3351–E3360. doi:.
26. Geall AJ, Verma A, Otten GR. Nonviral delivery of self-amplifying RNA vaccines. *Proc Natl Acad Sci U S A* 2012;109:14604–9. <https://doi.org/10.1073/pnas.1209367109>.
27. Guan S, Rosenecker J. Nanotechnologies in delivery of mRNA therapeutics using nonviral vector-based delivery systems. *Gene Ther* 2017;24:133–43. <https://doi.org/10.1038/gt.2017.5>.
28. Pardi N, Tuyishime S, Muramatsu H. Expression kinetics of nucleoside-modified mRNA delivered in lipid nanoparticles to mice by various routes. *J Control Release Off J Control Release Soc* 2015;217:345–51. <https://doi.org/10.1016/j.jconrel.2015.08.007>.
29. Lutz J, Lazzaro S, Habbeddine M. Unmodified mRNA in LNPs constitutes a competitive technology for prophylactic vaccines. *NPJ Vaccines* 2017;2:29. <https://doi.org/10.1038/s41541-017-0032-6>.
30. Chahal JS, Khan OF, Cooper CL. Dendrimer-RNA nanoparticles generate protective immunity against lethal Ebola, H1N1 influenza, and *Toxoplasma gondii* challenges with a single dose. *Proc Natl Acad Sci U S A* 2016;113:E4133–42. <https://doi.org/10.1073/pnas.1600299113>.
31. Schnee M, Vogel AB, Voss D. An mRNA Vaccine Encoding Rabies Virus Glycoprotein Induces Protection against Lethal Infection in Mice and Correlates of Protection in Adult and Newborn Pigs. *PLoS Negl Trop Dis* 2016;10. <https://doi.org/10.1371/journal.pntd.0004746> e0004746.
32. VanBlargan LA, Himansu S, Foreman BM, Ebel GD, Pierson TC, Diamond MS. An mRNA vaccine protects mice against multiple tick-transmitted flavivirus infections. *Cell Rep* 2018;25:3382–92. <https://doi.org/10.1016/j.celrep.2018.11.082> e3.
33. Petsch B, Schnee M, Vogel AB. Protective efficacy of in vitro synthesized, specific mRNA vaccines against influenza A virus infection. *Nat Biotechnol* 2012;30:1210–6. <https://doi.org/10.1038/nbt.2436>.
34. Pardi N, Hogan MJ, Pelc RS. Zika virus protection by a single low-dose nucleoside-modified mRNA vaccination. *Nature* 2017;543:248–51. <https://doi.org/10.1038/nature21428>.
35. Abbink P, Larocca RA, De La, Barrera RA. Protective efficacy of multiple vaccine platforms against Zika virus challenge in rhesus monkeys. *Science* 2016;353:1129–32. <https://doi.org/10.1126/science.aah6157>.
36. Dowd, et al. Rapid development of a DNA vaccine for Zika virus. *Science* 2016;354(6309):237–40. Accessed August 27, 2021.
37. Saunders KO, Pardi N, Parks R, et al. Lipid nanoparticle encapsulated nucleoside-modified mRNA vaccines elicit polyfunctional HIV-1 antibodies comparable to proteins in nonhuman primates. *npj Vaccines* 2021;6:50. <https://doi.org/10.1038/s41541-021-00307-6>. Accessed August 30, 2021.
38. Brito LA, Kommareddy S, Maione D. Chapter Seven - Self-Amplifying mRNA Vaccines. In: Huang L, Liu D, Wagner E, eds. *Advances in Genetics*. Vol 89. *Nonviral Vectors for Gene Therapy*. Cambridge, Massachusetts, USA: Academic Press, 2015:179–233.
39. Vogel AB, Lambert L, Kinnear E. Self-amplifying rna vaccines give equivalent protection against influenza to mRNA vaccines but at much lower doses. *Mol Ther J Am Soc Gene Ther* 2018;26:446–55. <https://doi.org/10.1016/j.ymthe.2017.11.017>.
40. Wang L, Shi W, Joyce MG. Evaluation of candidate vaccine approaches for MERS-CoV. *Nat Commun* 2015;6:7712. <https://doi.org/10.1038/ncomms8712>.
41. Wang L, Shi W, Chappell JD. Importance of neutralizing monoclonal antibodies targeting multiple antigenic sites on the middle east respiratory syndrome coronavirus spike glycoprotein to avoid neutralization escape. *J Virol* 2018;92. <https://doi.org/10.1128/JVI.02002-17>.
42. Yan R, Zhang Y, Li Y, Xia L, Guo Y, Zhou Q. Structural basis for the recognition of SARS-CoV-2 by full-length human ACE2. *Science* 2020;367:1444–8. <https://doi.org/10.1126/science.abb2762>.
43. Yi C, Sun X, Ye J. Key residues of the receptor binding motif in the spike protein of SARS-CoV-2 that interact with ACE2 and neutralizing antibodies. *Cell Mol Immunol* 2020;17:621–30. <https://doi.org/10.1038/s41423-020-0458-z>.
44. Crank MC, Ruckwardt TJ, Chen M. A proof of concept for structure-based vaccine design targeting RSV in humans. *Science* 2019;365:505–9. <https://doi.org/10.1126/science.aav9033>.

45. Dai L, Gao GF. Viral targets for vaccines against COVID-19. *Nat Rev Immunol* 2021;21:73–82. <https://doi.org/10.1038/s41577-020-00480-0>.
46. Cockrell AS, Yount BL, Scobey T. A mouse model for MERS coronavirus-induced acute respiratory distress syndrome. *Nat Microbiol* 2016;2:16226. <https://doi.org/10.1038/nmicrobiol.2016.226>.
47. Wrapp D, Wang N, Corbett KS. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* 2020;367:1260–3. <https://doi.org/10.1126/science.abb2507>.
48. Freeman B, Lester S, Mills L. Validation of a SARS-CoV-2 spike protein ELISA for use in contact investigations and serosurveillance. *bioRxiv*. Published online April 25, 2020. <https://doi.org/10.1101/2020.04.24.057323>.
49. Klumpp-Thomas C, Kalish H, Drew M. Standardization of enzyme-linked immunosorbent assays for serosurveys of the SARS-CoV-2 pandemic using clinical and at-home blood sampling. *MedRxiv Prepr Serv Health Sci*. Published online May 25, 2020. <https://doi.org/10.1101/2020.05.21.20109280>.
50. Walsh EE, Frenck R, Falsey AR. RNA-based COVID-19 vaccine BNT162b2 selected for a pivotal efficacy study. *medRxiv*. Published online August 20, 2020: 2020.08.17.20176651. <https://doi.org/10.1101/2020.08.17.20176651>.
51. Vogel AB, Kanevsky I, Che Y. BNT162b vaccines protect rhesus macaques from SARS-CoV-2. *Nature* 2021;592:283–9. <https://doi.org/10.1038/s41586-021-03275-y>.
52. Karikó K, Muramatsu H, Welsh FA. Incorporation of pseudouridine into mRNA yields superior nonimmunogenic vector with increased translational capacity and biological stability. *Mol Ther J Am Soc Gene Ther* 2008;16:1833–40. <https://doi.org/10.1038/mt.2008.200>.
53. Mulligan MJ, Lyke KE, Kitchin N. Phase I/II study of COVID-19 RNA vaccine BNT162b1 in adults. *Nature* 2020;586:589–93. <https://doi.org/10.1038/s41586-020-2639-4>.
54. Halstead SB, O'Rourke EJ. Dengue viruses and mononuclear phagocytes. I. Infection enhancement by non-neutralizing antibody. *J Exp Med* 1977;146:201–17. <https://doi.org/10.1084/jem.146.1.201>.
55. Sahin U, Muik A, Vogler I. BNT162b2 induces SARS-CoV-2-neutralising antibodies and T cells in humans. *medRxiv*. Published online December 11, 2020: 2020.12.09.20245175. <https://doi.org/10.1101/2020.12.09.20245175>.
56. Walsh EE, Frenck RW, Falsey AR. Safety and immunogenicity of two RNA-Based Covid-19 vaccine candidates. *N Engl J Med* 2020. <https://doi.org/10.1056/NEJMoa2027906> Published online October 14.
57. Polack FP, Thomas SJ, Kitchin N. Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine. *N Engl J Med* 2020;383:2603–15. <https://doi.org/10.1056/NEJMoa2034577>.
58. Shimabukuro T, Nair N. Allergic reactions including anaphylaxis after receipt of the first dose of Pfizer-BioNTech COVID-19 vaccine. *JAMA* 2021;325:780. <https://doi.org/10.1001/jama.2021.0600>.
59. Thomas SJ, Moreira ED, Kitchin N. Six month safety and efficacy of the BNT162b2 MRNA COVID-19 vaccine, 2021:2021.07.28.21261159. <https://doi.org/10.1101/2021.07.28.21261159>.
60. Thomas SJ, Moreira ED, Kitchin N. Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine through 6 months. *N Engl J Med* 2021;385:1761–73. <https://doi.org/10.1056/NEJMoa2110345>.
61. Falsey AR, Frenck RW, Walsh EE. SARS-CoV-2 Neutralization with BNT162b2 vaccine dose 3. *N Engl J Med* 2021;385:1627–9. <https://doi.org/10.1056/NEJMc2113468>.
62. Levine-Tiefenbrun M, Yelin I, Katz R. Initial report of decreased SARS-CoV-2 viral load after inoculation with the BNT162b2 vaccine. *Nat Med* 2021;27:790–2. <https://doi.org/10.1038/s41591-021-01316-7>.
63. Frenck RW, Klein NP, Kitchin N. Safety, immunogenicity, and efficacy of the BNT162b2 Covid-19 vaccine in adolescents. *N Engl J Med* 2021;385(3):239–50. <https://doi.org/10.1056/NEJMoa2107456>.
64. Jackson LA, Anderson EJ, Roupael NG. An mRNA vaccine against SARS-CoV-2 — preliminary report. *N Engl J Med* 2020;383:1920–31. <https://doi.org/10.1056/NEJMoa2022483>.
65. Chu L, McPhee R, Huang W. A preliminary report of a randomized controlled phase 2 trial of the safety and immunogenicity of mRNA-1273 SARS-CoV-2 vaccine. *Vaccine* 2021;39:2791–9. <https://doi.org/10.1016/j.vaccine.2021.02.007>.
66. Baden LR, El Sahly HM, Essink B. Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. *N Engl J Med* 2020;0. <https://doi.org/10.1056/NEJMoa2035389> null.
67. El Sahly HM, Baden LR, Essink B. Efficacy of the mRNA-1273 SARS-CoV-2 vaccine at completion of blinded phase. *N Engl J Med* 2021;0. <https://doi.org/10.1056/NEJMoa2113017>.
68. Baden LR, El Sahly HM, Essink B. Covid-19 in the phase 3 trial of mRNA-1273 during the delta-variant surge. *MedRxiv Prepr Serv Health Sci* 2021. <https://doi.org/10.1101/2021.09.17.21263624> Published online September 22 2021.09.17.21263624.
69. Moderna Announces FDA Authorization of a Booster Dose of Moderna's COVID-19 Vaccine in the U.S. Moderna, Inc. <https://investors.modernatx.com/news-releases/news-release-details/moderna-announces-fda-authorization-booster-dose-modernas-covid/>. Accessed November 3, 2021.
70. Ali K, Berman G, Zhou H. Evaluation of mRNA-1273 SARS-CoV-2 vaccine in adolescents. *N Engl J Med* 2021;0. <https://doi.org/10.1056/NEJMoa2109522>.
71. Moderna Reports Second Quarter Fiscal Year 2021 Financial Results and Provides Business Updates Moderna, Inc. <https://investors.modernatx.com/news-releases/news-release-details/moderna-reports-second-quarter-fiscal-year-2021-financial/>. Accessed September 8, 2021.
72. Boyarsky BJ, Werbel WA, Avery RK. Antibody response to 2-Dose SARS-CoV-2 mRNA vaccine series in solid organ transplant recipients. *JAMA* 2021;325:2204–6. <https://doi.org/10.1001/jama.2021.7489>.
73. Hall VG, Ferreira VH, Ku T. Randomized trial of a third dose of mRNA-1273 vaccine in transplant recipients. *N Engl J Med* 2021. <https://doi.org/10.1056/NEJMc2111462> Published online August 11.
74. Kamar N, Abravanel F, Marion O, Couat C, Izopet J, Del Bello A. Three doses of an mRNA Covid-19 vaccine in solid-organ transplant recipients. *N Engl J Med* 2021;385:661–2. <https://doi.org/10.1056/NEJMc2108861>.
75. Grupper A, Rabinowich L, Schwartz D. Reduced humoral response to mRNA SARS-CoV-2 BNT162b2 vaccine in kidney transplant recipients without prior exposure to the virus. *Am J Transplant Off J Am Soc Transplant Am Soc Transpl Surg* 2021;21:2719–26. <https://doi.org/10.1111/ajt.16615>.
76. Rabinowich L, Grupper A, Baruch R. Low immunogenicity to SARS-CoV-2 vaccination among liver transplant recipients. *J Hepatol* 2021;75:435–8. <https://doi.org/10.1016/j.jhep.2021.04.020>.
77. Herishanu Y, Avivi I, Aharon A. Efficacy of the BNT162b2 mRNA COVID-19 vaccine in patients with chronic lymphocytic leukemia. *Blood* 2021;137:3165–73. <https://doi.org/10.1182/blood.2021011568>.

78. Grifoni A, Weiskopf D, Ramirez SI. Targets of T Cell responses to SARS-CoV-2 coronavirus in humans with COVID-19 disease and unexposed individuals. *Cell*. 2020;181:1489–501. <https://doi.org/10.1016/j.cell.2020.05.015> e15.
79. Embi PJ. Effectiveness of 2-dose vaccination with mRNA COVID-19 vaccines against COVID-19-associated hospitalizations among immunocompromised adults — Nine States January–September 2021 MMWR Morb Mortal Wkly Rep 2021;70. <https://doi.org/10.15585/mmwr.mm7044e3>.
80. Longlune N, Nogier MB, Miedougé M. High immunogenicity of a messenger RNA-based vaccine against SARS-CoV-2 in chronic dialysis patients. *Nephrol Dial Transplant Off Publ Eur Dial Transpl Assoc - Eur Ren Assoc* 2021;36:1704–9. <https://doi.org/10.1093/ndt/gfab193>.
81. Espi M, Charmetant X, Barba T. Justification, safety, and efficacy of a third dose of MRNA vaccine in maintenance hemodialysis patients: a prospective observational study, 2021: 2021.07.02.21259913. <https://doi.org/10.1101/2021.07.02.21259913>.
82. Werbel WA, Boyarsky BJ, Ou MT. Safety and immunogenicity of a third dose of SARS-CoV-2 vaccine in solid organ transplant recipients: a case series. *Ann Intern Med* 2021;174:1330–2. <https://doi.org/10.7326/L21-0282>.
83. Shimabukuro TT, Kim SY, Myers TR. Preliminary findings of mRNA Covid-19 vaccine safety in pregnant persons. *N Engl J Med* 2021;0. <https://doi.org/10.1056/NEJMoa2104983> null.
84. Collier ARY, McMahan K, Yu J. Immunogenicity of COVID-19 mRNA vaccines in pregnant and lactating women. *JAMA* 2021;13. <https://doi.org/10.1001/jama.2021.7563> Published online May.
85. Magnus MC, Gjessing HK, Eide HN, Wilcox AJ, Fell DB, Häberg SE. Covid-19 vaccination during pregnancy and first-trimester miscarriage. *N Engl J Med* 2021;0. <https://doi.org/10.1056/NEJMc2114466> null.
86. Kharbanda EO, Haapala J, DeSilva M. Spontaneous abortion following COVID-19 vaccination during pregnancy. *JAMA* 2021;326:1629–31. <https://doi.org/10.1001/jama.2021.15494>.
87. Perl SH, Uzan-Yulzari A, Klainer H. SARS-CoV-2-specific antibodies in breast milk after COVID-19 vaccination of breastfeeding women. *JAMA* 2021;325:2013–4. <https://doi.org/10.1001/jama.2021.5782>.
88. Dagan N, Barda N, Biron-Shental T. Effectiveness of the BNT162b2 mRNA COVID-19 vaccine in pregnancy. *Nat Med* 2021;27:1693–5. <https://doi.org/10.1038/s41591-021-01490-8>.
89. Gray KJ, Bordt EA, Atyeo C. Coronavirus disease 2019 vaccine response in pregnant and lactating women: a cohort study. *Am J Obstet Gynecol* 2021;225(3). <https://doi.org/10.1016/j.ajog.2021.03.023> 303.e1–303.e17.
90. Vaccine Safety Monitoring Vaccine Safety CDC. Published August 28, 2020. <https://www.cdc.gov/vaccinesafety/ensuring-safety/monitoring/index.html>. Accessed August 28, 2020.
91. Shimabukuro TT, Cole M, Su JR. Reports of Anaphylaxis after receipt of mRNA COVID-19 vaccines in the US–December 14, 2020–January 18, 2021. *JAMA* 2021;325:1101–2. <https://doi.org/10.1001/jama.2021.1967>.
92. Blumenthal KG, Robinson LB, Camargo CA Jr. Acute allergic reactions to mRNA COVID-19 vaccines. *JAMA* 2021;325(15):1562–5. <https://doi.org/10.1001/jama.2021.3976>.
93. Shay DK, Shimabukuro TT, DeStefano F. Myocarditis occurring after immunization with mRNA-Based COVID-19 vaccines. *JAMA Cardiol* 2021. <https://doi.org/10.1001/jamacardio.2021.2821> Published online June 29.
94. Hause AM. COVID-19 vaccine safety in adolescents aged 12–17 years — United States, December 14, 2020–July 16, 2021. *MMWR Morb Mortal Wkly Rep*. 2021;70. <https://doi.org/10.15585/mmwr.mm7031e1>.
95. VAERS - Data. Accessed November 5, 2021. <https://vaers.hhs.gov/data.html>.
96. Mevorach D, Anis E, Cedar N. Myocarditis after BNT162b2 mRNA vaccine against Covid-19 in Israel. *N Engl J Med* 2021;0. <https://doi.org/10.1056/NEJMoa2109730> null.
97. Witberg G, Barda N, Hoss S. Myocarditis after Covid-19 vaccination in a large health care organization. *N Engl J Med* 2021;0. <https://doi.org/10.1056/NEJMoa2110737> null.
98. Vaccine Safety Datalink (VSD) VSD Monitoring Ensuring Safety Vaccine Safety CDC. Published September 30, 2021. <https://www.cdc.gov/vaccinesafety/ensuringsafety/monitoring/vsd/index.html>. Accessed November 5, 2021.
99. Canada PHA of. Summary: NACI recommendation on the use of mRNA COVID-19 vaccines in adolescents. Published August 27, 2021. <https://www.canada.ca/en/public-health/services/immunization/national-advisory-committee-on-immunization-naci/recommendations-use-covid-19-vaccines/mrna-adolescents/summary.html>. Accessed November 5, 2021.
100. Paterlini M. Covid-19: Sweden, Norway, and Finland suspend use of Moderna vaccine in young people “as a precaution”. *BMJ* 2021;375:n2477. <https://doi.org/10.1136/bmj.n2477>.
101. Dagan N, et al. BNT162b2 mRNA Covid-19 vaccine in a nationwide mass vaccination setting. *N. Engl. J. Med.* 2021;384:1412–23. Accessed August 31, 2021.
102. Moline HL. Effectiveness of COVID-19 vaccines in preventing hospitalization among adults aged ≥65 years — COVID-NET, 13 States February–April 2021 MMWR Morb Mortal Wkly Rep 2021;70. <https://doi.org/10.15585/mmwr.mm7032e3>.
103. Nanduri S. Effectiveness of Pfizer-BioNTech and Moderna Vaccines in Preventing SARS-CoV-2 Infection Among Nursing Home Residents Before and During Widespread Circulation of the SARS-CoV-2 B.1.617.2 (Delta) Variant — National Healthcare Safety Network, March 1–August 1, 2021. *MMWR Morb Mortal Wkly Rep*. 2021;70. <https://doi.org/10.15585/mmwr.mm7034e3>.
104. Haas EJ, Angulo FJ, McLaughlin JM. Impact and effectiveness of mRNA BNT162b2 vaccine against SARS-CoV-2 infections and COVID-19 cases, hospitalisations, and deaths following a nationwide vaccination campaign in Israel: an observational study using national surveillance data. *Lancet Lond Engl* 2021;397:1819–29. [https://doi.org/10.1016/S0140-6736\(21\)00947-8](https://doi.org/10.1016/S0140-6736(21)00947-8).
105. Planas D, Bruel T, Grzelak L. Sensitivity of infectious SARS-CoV-2 B.1.1.7 and B.1.351 variants to neutralizing antibodies. *Nat Med* 2021;27:917–24. <https://doi.org/10.1038/s41591-021-01318-5>.
106. Abu-Raddad LJ, Chemaitelly H, Butt AA. Effectiveness of the BNT162b2 Covid-19 Vaccine against the B.1.1.7 and B.1.351 Variants. *N Engl J Med* 2021;0. <https://doi.org/10.1056/NEJMc2104974> null.
107. Tregoning JS, Flight KE, Higham SL, Wang Z, Pierce BF. Progress of the COVID-19 vaccine effort: viruses, vaccines and variants versus efficacy, effectiveness and escape. *Nat Rev Immunol*. Published online August 2021;9:1–11. <https://doi.org/10.1038/s41577-021-00592-1>.
108. Lopez Bernal J, Andrews N, Gower C. Effectiveness of Covid-19 vaccines against the B.1.617.2 (Delta) Variant. *N Engl J Med* 2021;385:585–94. <https://doi.org/10.1056/NEJMoa2108891>.

109. Tartof SY, Slezak JM, Fischer H. Effectiveness of mRNA BNT162b2 COVID-19 vaccine up to 6 months in a large integrated health system in the USA: a retrospective cohort study. *The Lancet* 2021;398:1407–16. [https://doi.org/10.1016/S0140-6736\(21\)02183-8](https://doi.org/10.1016/S0140-6736(21)02183-8).
110. Bajema KL, Dahl RM, Prill MM. Effectiveness of COVID-19 mRNA vaccines against COVID-19-associated hospitalization - five veterans affairs medical centers United States, February 1-August 6, 2021 *MMWR Morb Mortal Wkly Rep* 2021;70:1294–9. <https://doi.org/10.15585/mmwr.mm7037e3>.
111. Fowlkes A. Effectiveness of COVID-19 Vaccines in Preventing SARS-CoV-2 Infection Among Frontline Workers Before and During B.1.617.2 (Delta) Variant Predominance — Eight U.S. Locations December 2020-August 2021 *MMWR Morb Mortal Wkly Rep* 2021;70. <https://doi.org/10.15585/mmwr.mm7034e4>.
112. Rosenberg ES. New COVID-19 cases and hospitalizations among adults, by vaccination status — New York, May 3-July 25, 2021. *MMWR Morb Mortal Wkly Rep* 2021;70. <https://doi.org/10.15585/mmwr.mm7037a7>.
113. Thompson MG, Stenehjem E, Grannis S. Effectiveness of Covid-19 vaccines in ambulatory and inpatient care settings. *N Engl J Med* 2021;385:1355–71. <https://doi.org/10.1056/NEJMoa2110362>.
114. Grannis SJ. Interim Estimates of COVID-19 Vaccine Effectiveness Against COVID-19-Associated Emergency Department or Urgent Care Clinic Encounters and Hospitalizations Among Adults During SARS-CoV-2 B.1.617.2 (Delta) Variant Predominance — Nine States, June-August 2021. *MMWR Morb Mortal Wkly Rep* 2021;70. <https://doi.org/10.15585/mmwr.mm7037e2>.
115. Goldberg Y, Mandel M, Bar-On YM. Waning Immunity after the BNT162b2 Vaccine in Israel. *N Engl J Med* 2021;0. <https://doi.org/10.1056/NEJMoa2114228> null.
116. Tenforde MW. Sustained Effectiveness of Pfizer-BioNTech and Moderna Vaccines Against COVID-19 Associated Hospitalizations Among Adults — United States, March-July 2021. *MMWR Morb Mortal Wkly Rep* 2021;70. <https://doi.org/10.15585/mmwr.mm7034e2>.
117. Christie A. Decreases in COVID-19 Cases, Emergency Department Visits, Hospital Admissions, and Deaths Among Older Adults Following the Introduction of COVID-19 Vaccine — United States, September 6, 2020-May 1, 2021. *MMWR Morb Mortal Wkly Rep* 2021;70. <https://doi.org/10.15585/mmwr.mm7023e2>.
118. Havers FP, Pham H, Taylor CA. COVID-19-Associated Hospitalizations among Vaccinated and Unvaccinated Adults ≥18 Years - COVID-NET, 13 States, January 1 - July 24, 2021.; 2021:2021.08.27.21262356. <https://doi.org/10.1101/2021.08.27.21262356>.
119. Wu K, Choi A, Koch M. Variant SARS-CoV-2 MRNA Vaccines Confer Broad Neutralization as Primary or. Booster Series in Mice 2021. <https://doi.org/10.1101/2021.04.13.439482>.
120. Self WH, Tenforde MW, Rhoads JP, et al. Comparative Effectiveness of Moderna, Pfizer-BioNTech, and Janssen (Johnson & Johnson) Vaccines in Preventing COVID-19 Hospitalizations Among Adults Without Immunocompromising Conditions — United States, March–August 2021. *MMWR Morb Mortal Wkly Rep* 2021;70:1337–43. <http://dx.doi.org/10.15585/mmwr.mm7038e1>.
121. Olson SM. Effectiveness of Pfizer-BioNTech mRNA Vaccination Against COVID-19 Hospitalization Among Persons Aged 12-18 Years — United States, June-September 2021. *MMWR Morb Mortal Wkly Rep* 2021;70. <https://doi.org/10.15585/mmwr.mm7042e1>.
122. Atmar RL, Lyke KE, Deming ME. Heterologous SARS-CoV-2 Booster Vaccinations - Preliminary Report, 2021:2021.10.10.21264827. doi:10.1101/2021.10.10.21264827
123. CureVac COVID vaccine let-down spotlights mRNA design challenges. <https://www.nature.com/articles/d41586-021-01661-0>. Accessed August 19, 2021.
124. Kremsner P, Mann P, Bosch J. Phase 1 Assessment of the Safety and Immunogenicity of an MRNA- Lipid Nanoparticle Vaccine Candidate Against SARS-CoV-2 in Human Volunteers, 2020:2020.11.09.20228551. <https://doi.org/10.1101/2020.11.09.20228551>.
125. Earle KA, Ambrosino DM, Fiore-Gartland A. Evidence for antibody as a protective correlate for COVID-19 vaccines. *Vaccine* 2021;39:4423–8. <https://doi.org/10.1016/j.vaccine.2021.05.063>.
126. Khoury DS, Cromer D, Reynaldi A. Neutralizing antibody levels are highly predictive of immune protection from symptomatic SARS-CoV-2 infection. *Nat Med* 2021;27:1205–11. <https://doi.org/10.1038/s41591-021-01377-8>.
127. Feng S, Phillips DJ, White T. Correlates of protection against symptomatic and asymptomatic SARS-CoV-2 infection. *Nat Med* 2021;29:1–9. <https://doi.org/10.1038/s41591-021-01540-1> Published online September.
128. Gilbert PB, Montefiori DC, McDermott A. Immune Correlates Analysis of the mRNA-1273 COVID-19 Vaccine Efficacy Trial. *MedRxiv Prepr Serv Health Sci* 2021;15. <https://doi.org/10.1101/2021.08.09.21261290> Published online August 2021.08.09.21261290.
129. Shrotri M, Schalkwyk MCI, van Post N. T cell response to SARS-CoV-2 infection in humans: a systematic review. *PLOS ONE* 2021;16:e0245532. <https://doi.org/10.1371/journal.pone.0245532>.
130. Goel RR, Painter MM, Apostolidis SA. mRNA vaccines induce durable immune memory to SARS-CoV-2 and variants of concern, 2020. *Science*. 0(0):eabm0829. <https://doi.org/10.1126/science.abm0829>.
131. Oberhardt V, Luxenburger H, Kemming J. Rapid and stable mobilization of CD8+ T cells by SARS-CoV-2 mRNA vaccine. *Nature* 2021;597:268–73. <https://doi.org/10.1038/s41586-021-03841-4>.
132. Tarke A, Sidney J, Methot N. Impact of SARS-CoV-2 variants on the total CD4+ and CD8+ T cell reactivity in infected or vaccinated individuals. *Cell Rep Med* 2021;2(7):100355. <https://doi.org/10.1016/j.xcrm.2021.100355>.
133. Planas D, Veyer D, Baidaliuk A. Reduced sensitivity of SARS-CoV-2 variant Delta to antibody neutralization. *Nature* 2021;596:276–80.
134. Doria-Rose N, Shen X, Schmidt SD. Booster of mRNA-1273 reduces SARS-CoV-2 Omicron escape from neutralizing antibodies. *MedRxiv Prepr Serv Health Sci*. Published online December 15, 2021. 12.15.21267805 <https://doi.org/10.1101/2021.12.15.21267805>.
135. Florea A, Sy LS, Luo Y. Durability of mRNA-1273 against COVID-19 in the time of Delta: Interim results from an observational study. *MedRxiv Prepr Serv Health Sci* 2021. <https://doi.org/10.1101/2021.12.13.21267620> Published online December 15 2021.12.13.21267620.
136. CDC COVID-19 Response Team. SARS-CoV-2 B.1.1.529 (Omicron) variant-United States, December 1-8, 2021. *Morb Mortal Weekl Rep*. 2021; 70:1731-1734