



Cytogenetic analysis of three species of *Pseudacteon* (Diptera, Phoridae) parasitoids of the fire ants using standard and molecular techniques

Mónica G. Chirino¹, Patricia J. Folgarait¹, Lawrence E. Gilbert², Silvia Lanzavecchia³ and Alba G. Papeschi⁴

¹Centro de Estudios e Investigaciones, Universidad Nacional de Quilmes, Bernal, Buenos Aires, Argentina.

²Brackenridge Field Laboratory, University of Texas, Austin, USA.

³Instituto de Genética, INTA, Castelar, Buenos Aires, Argentina.

⁴Laboratorio de Citogenética y Evolución, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Buenos Aires, Argentina.

Abstract

Pseudacteon flies, parasitoids of worker ants, are being intensively studied as potentially effective agents in the biological control of the invasive pest fire ant genus *Solenopsis* (Hymenoptera: Formicidae). This is the first attempt to describe the karyotype of *P. curvatus* Borgmeier, *P. nocens* Borgmeier and *P. tricuspis* Borgmeier. The three species possess $2n = 6$; chromosomes I and II were metacentric in the three species, but chromosome pair III was subtelocentric in *P. curvatus* and *P. tricuspis*, and telocentric in *P. nocens*. All three species possess a C positive band in chromosome II, lack C positive heterochromatin on chromosome I, and are mostly differentiated with respect to chromosome III. *P. curvatus* and *P. tricuspis* possess a C positive band, but at different locations, whereas this band is absent in *P. nocens*. Heterochromatic bands are neither AT nor GC rich as revealed by fluorescent banding. *In situ* hybridization with an 18S rDNA probe revealed a signal on chromosome II in a similar location to the C positive band in the three species. The apparent lack of morphologically distinct sex chromosomes is consistent with proposals of environmental sex determination in the genus. Small differences detected in chromosome length and morphology suggests that chromosomes have been highly conserved during the evolutionary radiation of *Pseudacteon*. Possible mechanisms of karyotype evolution in the three species are suggested.

Key words: *Pseudacteon*, karyotype evolution, C-banding, fluorescent banding, FISH.

Received: December 19, 2008; Accepted: May 4, 2009.

Introduction

The Phoridae, one of the most diverse families within the order Diptera, includes more than 26,000 min species, varying in life style from scavengers and predators to parasitoids (Disney, 1994; Gilbert and Jervis, 1998). The phorid genus *Pseudacteon* Coquillett are parasitoids, most of the species parasitizing *Solenopsis* Westwood fire-ants (Hymenoptera: Formicidae) (Disney, 1994). *Pseudacteon* flies are solitary parasitoids (Disney, 1994). The females inject a single egg into each worker ant, the third larval instar of which migrating into and later pupating within the host's cephalic capsule (Porter *et al.*, 1995; Porter, 1998).

Solenopsis has a cosmopolitan distribution (Disney, 1994; Patrock *et al.*, 2009). In South America, 22 *Pseudacteon* species have been reported as parasitizing fire-ants of the *Solenopsis saevissima* complex (Porter and Pesquero, 2001; Brown *et al.*, 2003; Folgarait *et al.*, 2005b;

Calcaterra, 2007; Kronforst *et al.*, 2007). Currently, these flies are used as biological control agents against *Solenopsis invicta* Buren and *S. richteri* Forel fire-ants, native to Argentina (Mescher *et al.*, 2003; Ross *et al.*, 2008) and exotic pests in the United States (Porter *et al.*, 2004; Thead *et al.*, 2005; Morrison and Porter, 2006).

Despite numerous studies on the life-history traits of the genus *Pseudacteon*, including apparent environmental sex-determination in some species, cytogenetic studies are lacking. Dipteran species generally possess low diploid chromosome numbers, these ranging from $2n = 4$ to 20, with modal numbers at 6 and 8 (due to several families of "lower" Diptera *i.e.* SO Nematocera) and 12 (with many species of the SO Brachycera and belonging to calyptrate and acalyptrate families) (White, 1973). Almost all species of Sarcophagidae and Calliphoridae (SO Brachycera, superfamily Oestroidea) have $2n = 12$ with a pair of sex chromosomes XY (Parise-Maltempi and Avancini, 2000; 2001). In the family Drosophilidae as a whole (SO Brachycera, superfamily Acalyptrata) the primitive chromosome number appears to be $2n = 12$ with five pairs of

long acrocentric chromosomes (including the sex chromosome pair XY) and a small dot-like pair. Reduction in chromosome number has occurred many times independently by fusion of these elements (White, 1973). Many derived sex chromosome systems have been described and the evolutionary mechanisms of sex chromosome differentiation has been analyzed in *Drosophila* species (Steinemann and Steinemann, 2005). In Muscidae (SO Brachycera, superfamily Muscoidea) most species also possess $2n = 12$ with XY sex chromosomes in the male. However, muscid species have been described with $2n = 10$ chromosomes and lacking differentiated sex chromosomes (Parise-Maltempo and Avancini, 2001). In Phoridae (SO Brachycera, superfamily Phoroidea), *Megaselia scalaris* Loew and *M. spiracularis* Schmitz, the only 2 species that have been studied cytogenetically, possess the diploid number $2n = 6$, with 2 metacentric and 1 telocentric chromosome pairs. They do not possess heteromorphic sex chromosomes and sex is determined by a single *Maleness* factor that can be located in any of the three linkage groups (Mainx, 1964; Traut and Willhoeft, 1990).

In order to evaluate whether the well-marked differences among *P. curvatus* Borgmeier, *P. nocens* Borgmeier and *P. tricuspis* Borgmeier in size, morphology, behavior and phylogenetic relations are also reflected in their karyotypes, our aim was to describe the karyotype of these species by using standard and molecular cytogenetic techniques. Descriptive values of the karyotype, heterochromatin content and distribution, and the location of nucleolus organizer regions (NORs) were analyzed. A comparative analysis of cytogenetic results also provided information on the possible karyotype evolution within the genus. Furthermore, we analyzed the presence of differentiated sex chromosomes, since it has been suggested, in the case of other species in the genus *Pseudacteon*, that sex determination is related to host size, and thus could be environmentally determined (Morrison *et al.*, 1999).

Material and Methods

Fly rearing

Experiments were carried out between August, 2006 and November, 2007. *Pseudacteon* species were reared in colonies of *S. invicta* ants collected near Mercedes, Corrientes province (29° 47' S, 58° 03' W). Adult males and females of *P. curvatus* were captured in the Reserva Ecológica Costanera Sur (RECS), Buenos Aires province (34° 37' S, 58° 22' W), those of *P. tricuspis* near San Javier, Santa Fe province (30° 58' S, 59° 94' W), and those of *P. nocens* from Mercedes, Corrientes province (29° 11' S, 58° 5' W), all in Argentina.

Slide preparation

Different oviposition assays were performed in the lab following the same methodology used by Chirino *et al.*

(2009). The “attacked” ants were placed in a rearing room at 28 ± 1 °C with a 12:12 (light: darkness) photoperiod and $80 \pm 10\%$ RH.

In Diptera, cytogenetic analyses are mainly carried out with third instar larvae or early pupae (Traut and Willhoeft, 1990; Cevallos and Nation, 2004). Since development in *Pseudacteon* takes place within its host (Porter *et al.*, 1995), it is difficult to obtain larvae at the appropriate developmental stages for cytogenetic studies. Thus we dissected pupae of 3-5 days, which lacked differentiated tissues, but showed high mitotic indices.

Since the females of *P. tricuspis* and *P. nocens* emerge from larger ant-heads than males, parasitized worker heads were measured and developed pupae were classified as either potential females or males. In contrast, both sexes of *P. curvatus* develop in similar sized hosts, so they could not be classified by sex (Morrison *et al.*, 1997; Folgarait *et al.*, 2006).

Pseudacteon pupae were dissected in a saline solution, swollen in a hypotonic solution (0.075 M KCl) for 10-20 min, and fixed for 15-30 min in freshly prepared Carnoy fixative (ethanol: chloroform: acetic acid, 6:3:1). Slides were prepared according to Traut (1976).

Chromosome morphology

Suitable cells at the mitotic prometaphase were selected and the mean descriptive values of the karyotype were calculated using information obtained from at least 3 cells per slide for each individual of each species analyzed (20.78 ± 9.76 cells). Each slide represented different individual pupae. Mitotic prometaphases were obtained from *P. tricuspis* pupae (17 males and 24 females); from *P. nocens* pupae (27 males and 21 females), and 35 specimens of *P. curvatus*. Slides were stained with the fluorescent dye DAPI (Rebagliati *et al.*, 2003) for morphological studies. The nomenclature of Levan *et al.* (1964) was used to describe chromosome morphology.

Banding techniques

Heterochromatin content and distribution was analyzed by means of the C-banding technique and sequential DAPI and CMA₃ banding. C-bands were performed according to Sumner (1972) with slight modifications. Slides were hydrolyzed in HCl 0.2N at room temperature for 15 min, briefly washed in distilled water and then incubated at room temperature for 35 min in a 5% Ba(OH)₂ solution. Slides were thoroughly washed in tap water for 1 min and incubated for 1 h in 2XSSC at 60 °C. After washing in distilled water, the slides were stained with DAPI (Rebagliati *et al.*, 2003), since band resolution improves when pre-treated slides are stained with DAPI instead of Giemsa. Fluorescent staining with GC specific chromomycin A₃ (CMA₃) and AT specific 4'-diamidino-2-phenylindole (DAPI) was carried out according to Rebagliati *et al.* (2003).

For location of NORs, Ag-NOR staining was applied according to Howell and Black (1980). Fluorescent *in situ* hybridization (FISH) with 18S rDNA was also undertaken.

Fluorescence *in situ* hybridization

Unlabelled 18S rDNA probes were generated by PCR, using primers described by Fuková *et al.* (2005) and sequenced using an Automated Sequencer ABI 3100 (DNA Sequencer, PE, Applied Biosystems), thereby confirming their being a section of 18S rDNA of *Ceratitis capitata* (Weidemann) (Diptera: Tephritidae). PCR was done in a Mastercycler[®] Gradient Eppendorf thermal cycler (Eppendorf AG, Hamburg, Germany). Reactions were carried out with template genomic DNA extracted from *C. capitata* by standard procedure according to Baruffi *et al.* (1995). The PCR product showed a single band of about 900 bp on a 1% agarose gel. The band was recovered from the gel and purified by using Wizard SV gel and PCR – clean up (Promega). The rDNA probe was labeled by nick translation with biotin 14-dUTP (BioNick Labeling System, Invitrogen Life Technologies Inc., Buenos Aires, Argentina). FISH with a biotinylated probe was carried out as described by Sahara *et al.* (1999) with several modifications as described by Fuková *et al.* (2005). Hybridization signals were detected with streptavidin-Cy3 conjugate (Sigma, Saint Louis, USA). The preparations were counterstained with 0.5 µg/mL DAPI in PBS, 1% Triton X-100, and mounted in anti-fade (Vectashield Mounting Medium, Vector Laboratories, Inc., Burlingame, USA).

Photographs

Preparations were observed in a Leica DMLB epifluorescence microscope. Black-and-white images of chromosomes were recorded with a CCD camera (Leica DFC350 FX, Leica IM50 Version 4.0, Leica Microsystems Imaging Solutions Ltd. Cambridge, UK) separately for each fluorescent dye. Images were pseudocolored (light blue for DAPI and red for Cy3), and, when necessary, superimposed with the aid of AdobePhotoshop version 6.0.

Statistical analysis

The following chromosome measurements were performed with MicroMeasure for Windows, version 3.3: centromeric index CI (length of short arm as a percentage of the whole chromosome), arm ratio *r* (relationship between the long and the short arms of each chromosome), relative length % (length of a chromosome as a percentage of the total chromosome length), and total chromosome length – TCL (the sum of the lengths of all the chromosomes of the complement). Comparisons of data within and between species were undertaken by analyses of variance (ANOVA), adjusted by the Bonferroni method. Statistical comparisons were done using the STATISTIX Program (Analytic Software 1998, Tallahassee, FL).

Results

Analysis of mitotic chromosomes of the pupae of *Pseudacteon tricuspis*, *P. curvatus* and *P. nocens* revealed a $2n = 6$ diploid number (Figure 1 a-b). Chromosome pairs I and II were metacentric in the three species, but chromosome pair III was subtelocentric in *P. tricuspis* and *P. curvatus*, and telocentric in *P. nocens* (Table 1). Neither *P. tricuspis* nor *P. nocens* possessed a distinguishable heteromorphic pair in potential males or females ($t = 0.43$, $df = 39$, $p = 0.67$ for *P. tricuspis* and $t = 0.07$, $df = 46$, $p = 0.95$ for *P. nocens*, 2-tailed *t*-Test). The three species did differ in total chromosome length (TCL) ($F_{2,99} = 2.38$, $p = 0.0453$) which was higher in *P. curvatus*, lower in *P. nocens* and intermediate in *P. tricuspis* (Table 2). Chromosome I in *P. nocens* was relatively larger than in *P. tricuspis*




Figure 1 - Cells in the mitotic metaphase (a) and anaphase (b) of *P. curvatus* stained with DAPI. Fluorescent banding with DAPI (c,e) and CMA₃ (d,f) in *P. curvatus* (c-d) and *P. tricuspis* (e-f). Bar = 10 µm.

Table 1 - Comparison of the somatic complement of the *Pseudacteon* species analyzed. %: relative length, r: arm ratio, CI: centromeric index, N: number of slides analyzed.

<i>Pseudacteon curvatus</i>						
Chromosomes	Average length (μm) ¹	% ¹	r	CI	Classification	N
I	47.11 \pm 17.53a	19.96a	1.14 \pm 0.10	46.87	m	
II	42.19 \pm 14.89a	17.75b	1.15 \pm 0.09	46.63	m	35
III	30.01 \pm 12.14b	12.29c	3.99 \pm 0.89	21.18	st	
<i>Pseudacteon tricuspis</i>						
Chromosomes	Average length (μm)	%	r	CI	Classification	N
I	44.69 \pm 11.14a	19.88a	1.20 \pm 0.17	45.78	m	
II	40.43 \pm 9.26a	18.07b	1.18 \pm 0.16	46.20	m	41
III	27.20 \pm 7.77b	12.05c	5.20 \pm 0.74	14.53	st	
<i>Pseudacteon nocens</i>						
Chromosomes	Average length (μm)	%	r	CI	Classification	N
I	41.21 \pm 10.12a	20.76a	1.14 \pm 0.08	46.94	m	
II	36.64 \pm 8.88b	18.48b	1.13 \pm 0.07	46.98	m	48
III	21.53 \pm 5.56c	10.75c	4.91 \pm 0.60	12.20	t	

¹The comparisons of chromosomal lengths and the relative lengths (%) were made by means of one-way ANOVA. Different letters indicate significant differences ($p < 0.05$) adjusted by Bonferroni.

and *P. curvatus* ($F_{2,123} = 8.68$, $p = 0.0004$). Chromosome II in *P. nocens* was also larger than in *P. curvatus* ($F_{2,123} = 5.86$, $p = 0.0039$), whereas in *P. tricuspis* it was of intermediate length. Finally, chromosome III was smaller in *P. nocens* than in *P. tricuspis* and *P. curvatus* ($F_{2,123} = 29.14$, $p < 0.0001$) (Table 2).

Different banding techniques were applied in order to reveal heterochromatin and analyze its location and composition. DAPI and CMA₃ sequential fluorescent banding did not reveal any regions either in AT rich (DAPI bright) or GC rich (CMA₃ bright) for any chromosome pair in any of the three species (Figure 1 c-f). After C-banding treatment, slides were stained with DAPI, allowing a better analysis of the C-heterochromatin pattern (Figure 2). C positive heterochromatin was scarce in all species, with a small C positive (C+) band in the long arm (q) of chromosome II and a completely C negative (C-) chromosome I (Figure 2a-f). "Somatic pairing" was evident from prophase up to meta-

phase (Figure 2a, d-i), but was lost during anaphase (Figure 2b-c). The three species varied as to the C-banding pattern in chromosome III. Thus, in *P. curvatus* a C+ band present in the short arm (p) extended over 81% of its length (Figure 2a-c). In *P. tricuspis*, a C+ band observed in the long (q) arm near the centromere corresponded to approximately 4% of its length (Figure 2d-e). Finally, the telocentric pair in *P. nocens* was completely C- (Figure 2f).

Silver impregnation revealed only one nucleolus in interphase nuclei and two signals in early prophase cells in the three species. However, no positive results were obtained in metaphase chromosomes (Figure 3). *In situ* hybridization with the 18S rDNA probe on metaphase cells showed a single hybridization signal in chromosome II in the three species at a location similar to that of the C+ band (Figure 2g-i; Figure 4). This suggests that in the three *Pseudacteon* species a single cluster of rRNA genes is located in the long arm (q) of chromosome II near the centromere.

Table 2 - Comparisons of total chromosomal lengths (TCL) and relative lengths (%) among the *Pseudacteon* species analyzed.

Species	TCL (μm) ¹	%		
		I	II	III
<i>P. curvatus</i>	225.31 \pm 70.74 A	19.96 B	17.75 B	12.29 A
<i>P. tricuspis</i>	220.11 \pm 54.69 AB	19.88 B	18.07 AB	12.05 A
<i>P. nocens</i>	198.45 \pm 46.01 B	20.76 A	18.48 A	10.75 B

¹The comparisons of the total chromosomal lengths (TCL) and the relative lengths (%) were analyzed by means of one-way ANOVA. Different letters indicate significant differences ($p < 0.05$) adjusted by Bonferroni.

Discussion

The *Pseudacteon* species analyzed presented $2n = 6$, with chromosomes I and II being metacentric and chromosome III subtelocentric in *P. curvatus* and *P. tricuspis* but telocentric in *P. nocens*. Total chromosome length (TCL) varied among *Pseudacteon* species, with *P. curvatus* \geq *P. tricuspis* \geq *P. nocens*, probably representing variations in DNA content that do not affect chromosome morphology. A similar situation has been reported in other dipteran fami-




Figure 2 - Somatic mitosis of *Pseudacteon curvatus* (a-c, g), *P. tricuspis* (d-e, h) and *P. nocens* (f,i) after C banding (a-f) and *in situ* hybridization with a 18S rDNA probe (g-i). At the top right-hand corner the names of species are indicated as PC = *P. curvatus*, PT = *P. tricuspis*, and PN = *P. nocens*. Arrows (a-f) indicate blocks of heterochromatin and head arrows (g-i), rDNA sites. Bar = 10 µm.




Figure 3 - Interphase nuclei of *P. curvatus* (a) and *P. tricuspis* (b), and the early prophase (c) in which one nucleolus was observed after silver nitrate NOR banding. d) Metaphase of *P. tricuspis*. No Ag-NOR band was detected on the condensed chromosomes. Bar = 10 µm.

lies (Rafael and Tadei, 1998; Parise-Maltempi and Avancini, 2001; Selivon *et al.*, 2005).

Heterochromatin content was very scarce in these *Pseudacteon* species. Chromosome I was completely C-, with a single C+ band situated at the location of the 18S rDNA hybridization signal on chromosome II. Major differences were observed in chromosome III, with a C+ band in the p-arm of the subtelocentric pair in *P. curvatus* and the q-arm in *P. tricuspis*, although this was absent in the telocentric pair of *P. nocens* (Figure 4). The divergence between the former two species could be explained by pericentric inversion in chromosome III during their evolutionary history (Figure 5). Its absence in *P. nocens* suggests that it could have diverged before the differentiation of *P. tricuspis* and *P. curvatus*. Furthermore, the smaller size of chromosome III in *P. nocens*, as compared to the other two species, could be explained by the absence of a heterochromatin block. Although it would be necessary to analyze other *Pseudacteon* species, we propose here a possible hypothesis on karyotype evolution within the genus. We propose that after the divergence of *P. nocens*, and through the acquisition of heterochromatin, there was an increase in DNA across the genome of the ancestor of the other two species, but principally in chromosome III. In spite of the fragmented and unresolved phylogenies of some *Pseudacteon* species (Kronforst *et al.*, 2007; Calcaterra *et al.*, 2008), several conclusions support our hypothesis. The recent common ancestor of *P. nocens* is different from the most recent common ancestor of *P. curvatus* and *P. tricuspis* (Calcaterra *et al.*, 2008), and phylogenies based on the mitochondrial genes *Cytochrome Oxidase I* and *II* and the nuclear gene *Wingless* (Kronforst *et al.*, 2007) suggest that *P. tricuspis* could have derived from the ancestor from which *P. curvatus* diverged.

In reference to the studied *Pseudacteon* species, Ag-NOR banding revealed the presence of one nucleolus but not on metaphase chromosomes. Similar observations have been reported in other dipterans in which results from silver staining were unsatisfactory, and only centromeric cores could be differentiated (Motara *et al.*, 1985; Wallace and Newton, 1987; Bedo and Webb, 1989; Marchi and Pili, 1994).

Cytogenetic characterization of *Pseudacteon* species failed to reveal morphologically differentiated sex chromosomes. The only other two species in the Phoridae in which cytogenetical data has come under analysis were *Megaselia scalaris* and *M. spiracularis*. Accordingly, both possess the same diploid number $2n = 6$ and a similar chromosome morphology, with two metacentric and one telocentric chromosomes (Mainx, 1964). In this genus sex determination depends on the presence of a single *Maleness* factor M that triggers off male development (Mainx, 1964; Willhoeft and Traut, 1990; Traut and Wollert, 1998; Traut *et al.*, 1999). Through molecular cytogenetic techniques and molecular markers it has been established that the chromo-




Figure 4 - Idiograms of *Pseudacteon curvatus* (a), *P. tricuspis* (b) and *P. nocens* (c). C-bands are shown as black segments, and arrows point out the location of the hybridization signal corresponding to 18S rDNA.

some *X* and *Y* in *Megaselia* are in a very early state of molecular differentiation (Traut, 1994).

In most of the studied *Pseudacteon* species, the sex of the offspring seems to be facultatively determined by the size of the host, with females emerging from the larger ones (Morrison *et al.*, 1999; Folgarait *et al.*, 2005a, 2006). From an evolutionary point of view, this makes sense since 1) fire-ant workers are highly variable in size (Tschinkel *et al.*, 2003), 2) female flies have a higher fitness if they emerge from larger hosts, exhibiting a higher efficiency in development and larger sizes (Chirino, Gilbert, Folgarait, unpublished), and 3) females emerge from a narrower range of ant-sizes than males (Folgarait *et al.*, 2005a, 2006; Chirino, Gilbert, Folgarait, unpublished). However, in two small species, *P. cultellatus* Borgmeier (Folgarait *et al.*, 2002) and *P. curvatus* (Chirino *et al.*, 2009), females and males develop in similar-sized hosts, and emerged adults are not sexually dimorphic in size, so that the above mentioned pattern is difficult to detect. The number of different species of *Pseudacteon* that develop on *S. invicta* exceeds the diversity observed in other ants (Disney, 1994). This diversity could be influenced by *Solenopsis* size polymorphism since there is a correlation between the sizes of both parasitoid and host (Morrison *et al.*, 1997). Different *Pseudacteon*

species oviposit at different times during the day, at different locations (close to mounds or the foraging trails of ants) and during different seasons. *P. nocens* in particular is crepuscular and abundant in shady conditions, but *P. curvatus* and *P. tricuspis* are more abundant at midday and in sunny conditions (Folgarait *et al.*, 2007a; 2007b). Furthermore, *P. curvatus* mainly attacks small workers (Chirino *et al.*, 2009), *P. tricuspis* larger ants (Morrison *et al.*, 1997), and *P. nocens*, medium-sized ones. Finally, *P. tricuspis* females exhibit competitive, aggressive and territorial behaviors, which are not observed in either *P. curvatus* or *P. nocens*. In this work we have shown that the well-known differences among *P. curvatus*, *P. nocens* and *P. tricuspis* as regards size, morphology, behavior and phylogenetic relations are also reflected in their karyotypes.

The exact mechanism of sex determination in *Pseudacteon* flies is perplexing (Porter, 1998). Environmental sex determination (ESD) has been suggested as a possible mechanism for the observed pattern (Morrison *et al.*, 1999). Typically, taxa having non-differentiated sex chromosomes usually show labile sex-determination, and sex can be determined by environmental stimuli (Solari, 1994). Our demonstration of non-differentiated sex chromosomes in *Pseudacteon* phorids is thus consistent with the earlier suggestion of environmental sex determination in these parasitoid flies, and should encourage further investigation of how sex is determined in this genus.




Figure 5 - Hypothesized karyotype evolution in *Pseudacteon curvatus* and *P. tricuspis*. Differences in chromosome III can be explained by pericentric inversion, although it is not possible to determine the direction of evolutionary change.

Acknowledgments

We thank Lic. Gloria Albioni Montenegro for her help in several aspects of the experiments. This research was supported by grants from the Programa de Investigación en Interacciones Biológicas (PPUNQ-0340/03), Universidad Nacional de Quilmes; the R.J. Kleberg & H.C. Kleberg Foundation; Universidad de Buenos Aires (X164) and PICT 2007-00635 of ANPCyT. M.G.Ch., P.J.F. and

A.G.P. thank the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET).

References

- Baruffi L, Damiani G, Guglielmino C, Bandi C, Malacrida A and Gasperi G (1995) Polymorphism within and between population of *Ceratitis capitata*. Comparison between RAPD and Multilocus enzyme electrophoresis data. *Heredity* 74:524-537.
- Bedo DG and Webb GC (1989) Conservation of nucleolar structure in polytene tissues of *Ceratitis capitata* (Diptera, Tephritidae). *Chromosoma* 98:443-449.
- Brown BV, Folgarait PJ and Gilbert L (2003) A new species of *Pseudacteon* attacking *solenopsis* fire ants (Hymenoptera, Formicidae) in Argentina. *Sociobiology* 41:685-688.
- Calcaterra LA (2007) *Pseudacteon calderensis*, a new fly species (Diptera, Phoridae) attacking the fire ants *Solenopsis interrupta* (Hymenoptera, Formicidae) in northwestern Argentina. *Ann Entomol Soc Am* 100:470-473.
- Calcaterra LA, Delgado A and Tsutsui ND (2008) Activity patterns and parasitism rates of fire ant-decapitating flies (Diptera, Phoridae, *Pseudacteon* spp.) in their native Argentina. *Ann Entomol Soc Am* 101:539-550.
- Cevallos VE and Nation JL (2004) Chromosomes of the Caribbean fruit fly (Diptera, Tephritidae). *Fla Entomol* 87:361-364.
- Chirino MG, Gilbert LE and Folgarait PJ (2009) Behavior and development of *Pseudacteon curvatus* (Diptera, Phoridae) varies according to the social form of its host *Solenopsis invicta* (Hymenoptera, Formicidae) in its native range. *Environ Entomol* 38:198-206.
- Disney RHL (1994) *Scuttle Flies: The Phoridae*. Chapman & Hall, London, 480 pp.
- Folgarait PJ, Bruzzone OA and Gilbert LE (2002) Development of *Pseudacteon cultellatus* (Diptera, Phoridae) on *Solenopsis invicta* and *Solenopsis richteri* fire ants (Hymenoptera, Formicidae). *Environ Entomol* 31:403-410.
- Folgarait PJ, Chirino MG, Wilson Patrock RJ and Gilbert LE (2005a) Development of *Pseudacteon obtusus* (Diptera, Phoridae) on *Solenopsis invicta* and *Solenopsis richteri* fire ants (Hymenoptera, Formicidae). *Environ Entomol* 34:308-316.
- Folgarait PJ, Bruzzone OA, Porter SD, Pesquero MA and Gilbert LE (2005b) Biogeography and macroecology of phorid flies that attack fire ants in southeastern Brazil and Argentina. *J Biogeogr* 32:353-367.
- Folgarait PJ, Wilson Patrock RJ and Gilbert LE (2006) Development of *Pseudacteon nocens* (Diptera, Phoridae) on *Solenopsis invicta* and *Solenopsis richteri* fire ants (Hymenoptera, Formicidae). *J Econ Entomol* 99:295-307.
- Folgarait PJ, Patrock RJW and Gilbert LE (2007a) Associations of fire ant phorids and microhabitats. *Environ Entomol* 36:731-742.
- Folgarait PJ, Patrock RJW and Gilbert LE (2007b) The influence of ambient conditions and space on the phenological patterns of a *Solenopsis* phorid guild in an arid environment. *Biol Control* 42:262-273.
- Fuková I, Nguyen P and Marec F (2005) Codling moth cytogenetics: Karyotype, chromosomal location of rDNA, and molecular differentiation of sex chromosomes. *Genome* 48:1083-1092.
- Gilbert F and Jervis M (1998) Functional, evolutionary and ecological aspects of feeding-related mouthpart specializations in parasitoid flies. *Biol J Linn Soc* 63:495-535.
- Howell WM and Black AD (1980) Controlled silver-staining of nucleolus organizer regions with a protective colloidal developer: A 1-step method. *Experientia* 36:1014-1015.
- Kronforst MR, Folgarait PJ, Patrock RJW and Gilbert LE (2007) Genetic differentiation between body size biotypes of the parasitoid fly *Pseudacteon obtusus* (Diptera, Phoridae). *Mol Phylogenet Evol* 43:1178-1184.
- Levan A, Fredga K and Sandberg A (1964) Nomenclature for centromeric position on chromosomes. *Hereditas* 52:201-220.
- Mainx F (1964) The genetics of *Megaselia scalaris* Loew (Phoridae): A new type of sex determination in Diptera. *Am Nat* 98:415-430.
- Marchi A and Pili E (1994) Ribosomal RNA genes in mosquitoes: Localization by fluorescence *in situ* hybridization (FISH). *Heredity* 72:599-605.
- Mescher MC, Ross KG, Shoemaker DD, Keller L and Krieger MJB (2003) Distribution of the two social forms of the fire ant *Solenopsis invicta* (Hymenoptera, Formicidae) in the native South American range. *Ann Entomol Soc Am* 96:810-817.
- Morrison LW and Porter SD (2006) Post-release host-specificity testing of *Pseudacteon tricuspis*, a phorid parasitoid of *Solenopsis invicta* fire ants. *Biocontrol* 51:195-205.
- Morrison LW, Dall'aglio-Holvorcem CG and Gilbert L (1997) Oviposition behavior and development of *Pseudacteon* flies (Diptera, Phoridae), parasitoids of *Solenopsis* fire ants (Hymenoptera, Formicidae). *Environ Entomol* 26:14-22.
- Morrison LW, Porter SD and Gilbert LE (1999) Sex ratio variations as a function of host size in *Pseudacteon* flies (Diptera, Phoridae), parasitoids of *Solenopsis* fire ants (Hymenoptera, Formicidae). *Biol J Linn Soc* 66:257-267.
- Motara MA, Pathak S, Satya-Prakash KL and Hsu TC (1985) Argentophilic structures of spermatogenesis in the yellow fever mosquito. *J Hered* 76:295-300.
- Parise-Maltempi PP and Avancini RMP (2000) Cytogenetics of the neotropical flesh fly *Pattonella intermutans* (Diptera, Sarcophagidae). *Genet Mol Biol* 23:563-567.
- Parise-Maltempi PP and Avancini RMP (2001) C-banding and FISH in chromosomes of the blow flies *Chrysomia megacephala* and *Chrysomia putoria* (Diptera, Calliphoridae). *Mem Inst Oswaldo Cruz* 96:371-377.
- Patrock RJW, Porter SD, Gilbert LE and Folgarait PJ (2009) Distributional patterns of *Pseudacteon* associated with the *Solenopsis saevissima* complex in South America. *J Insect Sci* (in press).
- Porter SD (1998) Biology and behavior of *Pseudacteon* decapitating flies (Diptera, Phoridae) that parasitize *Solenopsis* fire ants (Hymenoptera, Formicidae). *Fla Entomol* 81:292-308.
- Porter SD and Pesquero MA (2001) Illustrated key to *Pseudacteon* decapitating flies (Diptera, Phoridae) that attack *Solenopsis saevissima* complex fire ants in South America. *Fla Entomol* 90:135-138.
- Porter SD, Nogueira de Sá LA and Morrison LW (2004) Establishment and dispersal of the fire ant decapitating fly

- Pseudacteon tricuspis* in North Florida. *Biol Control* 29:179-188.
- Porter SD, Pesquero MA, Campiolo S and Fowler HG (1995) Growth and development of *Pseudacteon* phorid fly maggots (Diptera, Phoridae) in the head of *Solenopsis* fire ant workers (Hymenoptera, Formicidae). *Environ Entomol* 24:475-479.
- Rafael MS and Tadei WP (1998) Metaphase karyotypes of *Anopheles (Nyssorhynchus) darlingi* Root and *A. (N.) nuneztovari* Gabaldón (Diptera, Culicidae). *Genet Mol Biol* 21:351-354.
- Rebagliati P, Papeschi AG and Mola LM (2003) Meiosis and fluorescent banding in *Edessa mediatubunda* and *E. rufomarginata* (Heteroptera, Pentatomidae, Edessinae). *Eur J Entomol* 100:11-18.
- Ross KG, Krieger MJB, Keller L and Shoemaker DD (2008) Genetic variation and structure in native populations of the fire ant *Solenopsis invicta*: Evolutionary and demographic implications. *Biol J Linn Soc* 92:541-560.
- Sahara K, Marec F and Traut W (1999) TTAGG telomeric repeats in chromosomes of some insects and other arthropods. *Chromosome Res* 7:449-460.
- Selivon D, Perondini ALP and Rocha D (2005) Karyotype characterization of *Anastrepha* fruit flies (Diptera, Tephritidae). *Neotrop Entomol* 34:273-279.
- Solari AJ (1994) Sex chromosomes and sex determination in vertebrates. CRC Press, Boca Raton, 336 pp.
- Steinemann S and Steinemann M (2005) Retroelements: Tools for sex chromosome evolution. *Cytogenet Genome Res* 110:134-143.
- Sumner AT (1972) A simple technique for demonstrating centromeric heterochromatin. *Exp Cell Res* 75:304-306.
- Thead LG, Vogt JT and Streett DA (2005) Dispersal of the fire ant decapitating fly, *Pseudacteon curvatus* (Diptera, Phoridae) in northeast Mississippi. *Fla Entomol* 88:214-216.
- Traut W (1976) Pachytene mapping in the female silkworm *Bombyx mori* L. (Lepidoptera). *Chromosoma* 58:275-284.
- Traut W (1994) Sex determination in the fly *Megaselia scalaris*, a model system for primary steps of sex chromosome evolution. *Genetics* 136:1097-1104.
- Traut W and Willhoeft U (1990) A jumping sex determining factor in the fly *Megaselia scalaris*. *Chromosoma* 99:407-412.
- Traut W and Wollert B (1998) An X/Y DNA segment from an early stage of sex chromosome differentiation in the fly *Megaselia scalaris*. *Genome* 41:289-294.
- Traut W, Sahara K, Otto TD and Marec F (1999) Molecular differentiation of sex chromosomes probed by comparative genomic hybridization. *Chromosoma* 108:173-180.
- Tschinkel WR, Mikheyev AS and Storz SR (2003) Allometry of workers of the fire ant, *Solenopsis invicta*. *J Insect Sci* 3:e2.
- Wallace AJ and Newton ME (1987) Heterochromatin diversity and cyclic responses to selective silver staining in *Aedes aegypti* (L.). *Chromosoma* 95:89-93.
- White MJD (1973) *Animal Cytology and Evolution*. 3rd edition. Cambridge University Press, London, 961 pp.
- Willhoeft U and Traut W (1990) Molecular differentiation of the homomorphic sex chromosomes in *Megaselia scalaris* (Diptera) detected by random DNA probes. *Chromosoma* 99:237-242.

Internet Resources

MicroMeasure Web site, <http://www.colostate.edu/Depts/Biology/MicroMeasure>, for MicroMeasure for Windows, version 3.3.

Associate Editor: Yatiyo Yonenaga-Yassuda

License information: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.