

Vitamin D supplementation to prevent acute respiratory infections: systematic review and meta-analysis of aggregate data from randomised controlled trials

David A Jolliffe^{1,2†}

Carlos A Camargo Jr³

John D Sluyter⁴

Mary Aglipay⁵

John F Aloia⁶

Davaasambuu Ganmaa⁷

Peter Bergman⁸

Arturo Borzutzky⁹

Camilla T Damsgaard¹⁰

Gal Dubnov-Raz¹¹

Susanna Esposito¹²

Clare Gilham¹³

Adit A Ginde¹⁴

Inbal Golan-Tripto¹⁵

Emma C Goodall¹⁶

Cameron C Grant¹⁷

Christopher J Griffiths^{1,2}

Anna Maria Hibbs^{18,19}

Wim Janssens²⁰

Anuradha Vaman Khadilkar²¹

Ilkka Laaksi²²

Margaret T Lee²³

Mark Loeb²⁴

Jonathon L Maguire⁵

Paweł Majak²⁵

David T Mauger²⁶

Semira Manaseki-Holland²⁷

David R Murdoch²⁸

Akio Nakashima²⁹

Rachel E Neale³⁰

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.
Hai Pham³⁰

Christine Rake¹³

Judy R Rees³¹

Jenni Rosendahl³²

Robert Scragg⁴

Dheeraj Shah³³

Yoshiki Shimizu³⁴

Steve Simpson-Yap^{35,36}

Geeta Trilok Kumar³⁷

Mitsuyoshi Urashima²⁹

Adrian R Martineau^{1,2†}

¹Institute for Population Health Sciences, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, UK

²Asthma UK Centre for Applied Research, Queen Mary University of London, London, UK

³Department of Emergency Medicine, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

⁴School of Population Health, University of Auckland, Auckland, New Zealand

⁵Department of Pediatrics, St Michael's Hospital, Toronto, Ontario, Canada

⁶Bone Mineral Research Center, Winthrop University Hospital, Mineola, NY, USA

⁷Department of Nutrition, Harvard School of Public Health, Boston, MA, USA

⁸Department of Laboratory Medicine, Karolinska Institutet, Stockholm, Sweden

⁹Department of Pediatric Infectious Diseases and Immunology, School of Medicine, Pontificia Universidad Católica de Chile, Santiago, Chile

¹⁰Department of Nutrition, Exercise and Sports, University of Copenhagen, Frederiksberg, Denmark

¹¹Exercise, Lifestyle and Nutrition Clinic, Edmond and Lily Safra Children's Hospital, Tel Hashomer, Israel

¹²Pediatric Clinic, Pietro Barilla Children's Hospital, Department of Medicine and Surgery, University of Parma, Parma, Italy

¹³Department of Non-Communicable Disease Epidemiology, London School of Hygiene and Tropical Medicine, London, UK

¹⁴Department of Emergency Medicine, University of Colorado School of Medicine, Aurora, CO, USA

¹⁵Saban Pediatric Medical Center, Soroka University Medical Center, Faculty of Health Sciences, Ben-Gurion University, Beer Sheva, Israel

¹⁶Department of Health Research Methods, Evidence and Impact, McMaster University, Hamilton, ON, Canada

¹⁷Department of Paediatrics: Child & Youth Health, Faculty of Medical and Health Sciences, University of Auckland, Auckland, New Zealand

¹⁸Department of Pediatrics, Case Western Reserve University School of Medicine, Cleveland, OH, USA

¹⁹University Hospitals Rainbow Babies and Children's Hospital, Cleveland, OH, USA

²⁰Universitair ziekenhuis Leuven, Leuven, Belgium

²¹Hirabai Cowasji Jehangir Medical Research Institute, Maharashtra, India

²²Faculty of Medicine and Health Technology, University of Tampere, Tampere, Finland

²³Division of Pediatric Hematology/Oncology/Stem Cell Transplantation, Columbia University Medical Center, New York, NY USA

²⁴Department of Pathology and Molecular Medicine, McMaster University, Hamilton, ON, Canada

²⁵Department of Pediatric Pulmonology, Medical University of Lodz, Lodz, Poland

²⁶Department of Statistics, The Pennsylvania State University, Hershey, PA, USA

²⁷Department of Public Health, Epidemiology and Biostatistics, Institute of Applied Health Sciences, College of Medical and Dental Sciences, University of Birmingham, Birmingham, UK

²⁸Department of Pathology, University of Otago, Christchurch, New Zealand

²⁹Jikei University School of Medicine, Tokyo, Japan

³⁰Population Health Department, QIMR Berghofer Medical Research Institute, Queensland, Australia

³¹Department of Epidemiology, Geisel School of Medicine at Dartmouth, Lebanon, NH, USA

³²Children's Hospital, Pediatric Research Centre, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

³³Department of Paediatrics, University College of Medical Sciences, Delhi, India

³⁴FANCL Research Institute, FANCL Corporation, Yokohama, Japan

³⁵Neuroepidemiology Unit, Melbourne School of Population & Global Health, The University of Melbourne, Melbourne, VIC, Australia

³⁶Menzies Institute for Medical Research, University of Tasmania, Hobart, TAS,
Australia

³⁷Institute of Home Economics, University of Delhi, New Delhi, India

† To whom correspondence should be addressed at the Institute for Population Health
Sciences, Barts and The London School of Medicine and Dentistry, Queen Mary
University of London, 58 Turner St, London E1 2AB, UK

Tel: +44 207 882 2551 | Fax: +44 207 882 2552 | Email: d.a.jolliffe@qmul.ac.uk or
a.martineau@qmul.ac.uk

Abstract

Background: A 2017 meta-analysis of data from 25 randomised controlled trials of vitamin D supplementation for the prevention of acute respiratory infections revealed a protective effect of the intervention. Since then, 20 new RCTs have been completed.

Methods: Systematic review and meta-analysis of data from randomised controlled trials (RCTs) of vitamin D for ARI prevention using a random effects model. Pre-specified sub-group analyses were done to determine whether effects of vitamin D on risk of ARI varied according to baseline 25-hydroxyvitamin D (25[OH]D) concentration or dosing regimen. We searched MEDLINE, EMBASE, the Cochrane Central Register of Controlled Trials (CENTRAL), Web of Science and the ClinicalTrials.gov registry from inception to 1st May 2020. Double-blind RCTs of supplementation with vitamin D or calcidiol, of any duration, were eligible if they were approved by a Research Ethics Committee and if ARI incidence was collected prospectively and pre-specified as an efficacy outcome. Aggregate data, stratified by baseline 25(OH)D concentration, were obtained from study authors. The study was registered with PROSPERO (no. CRD42020190633).

Findings: We identified 45 eligible RCTs (total 73,384 participants). Data were obtained for 46,331 (98.0%) of 47,262 participants in 42 studies, aged 0 to 95 years. For the primary comparison of vitamin D supplementation vs. placebo, the intervention reduced risk of ARI overall (Odds Ratio [OR] 0.91, 95% CI 0.84 to 0.99; P for heterogeneity 0.01). No statistically significant effect of vitamin D was seen for any of the sub-groups defined by baseline 25(OH)D concentration. However, protective effects were seen for trials in which vitamin D was given using a daily dosing regimen (OR 0.75, 95% CI 0.61 to 0.93); at daily dose equivalents of 400-1000 IU (OR 0.70, 95% CI 0.55 to 0.89); and for a duration of ≤ 12 months (OR 0.82, 95% CI 0.72 to 0.93). No significant interaction was seen between allocation to vitamin D vs. placebo and dose frequency, dose size, or study duration. Vitamin D did not influence the proportion of participants experiencing at least one serious adverse event (OR 0.97, 95% CI 0.86 to 1.09). Risk of bias within individual studies was assessed as being low for all but three trials. A funnel plot showed left-sided asymmetry ($P=0.008$, Egger's test).

Interpretation: Vitamin D supplementation was safe and reduced risk of ARI, despite evidence of significant heterogeneity across trials. Protection was associated with administration of daily doses of 400-1000 IU vitamin D for up to 12 months. The relevance of these findings to COVID-19 is not known and requires investigation.

Funding: None

Research in context

Evidence before this study

The active vitamin D metabolite, 1,25-dihydroxyvitamin D, induces innate immune responses to respiratory viruses and bacteria. A previous meta-analysis of individual participant data from 10,933 participants in 25 randomised controlled trials of vitamin D supplementation for the prevention of acute respiratory infection demonstrated an overall protective effect (adjusted Odds Ratio [aOR] 0.88, 95% confidence interval 0.81 to 0.96). Sub-group analysis revealed most benefit in those with the lowest vitamin D status at baseline who received daily or weekly supplementation (aOR 0.30, 0.17 to 0.53).

Added value of this study

Our meta-analysis of aggregate data from 46,331 participants in 42 randomised controlled trials, stratified by baseline 25(OH)D concentration, provides an updated estimate of the protective effects of vitamin D against acute respiratory infection overall, and in sub-groups defined by baseline vitamin D status and dosing frequency, amount and duration.

Implications of all the available evidence

Overall, vitamin D reduced the risk of having one or more acute respiratory infections (OR 0.91, 0.84 to 0.99), but there was evidence of significant heterogeneity across trials (P for heterogeneity 0.01). A funnel plot showed left-sided asymmetry, which may reflect publication bias and/or heterogeneity of effect across trials. No statistically significant effect of vitamin D was seen for any of the sub-groups defined by baseline 25(OH)D concentration. However, protective effects were seen in trials where vitamin D was given using a daily dosing regimen (OR 0.75, 0.61 to 0.93); at daily dose equivalents of 400-1000 IU (OR 0.70, 0.55 to 0.89); and for a duration of ≤ 12 months (OR 0.82, 0.72 to 0.93). The relevance of these findings to COVID-19 is not known and requires investigation.

Introduction

Interest in the potential for vitamin D supplementation to reduce risk of acute respiratory infections (ARI) has increased since the emergence of the COVID-19 pandemic.¹ This stems from findings of laboratory studies, showing that vitamin D metabolites support innate immune responses to respiratory viruses,² together with observational studies reporting independent associations between low circulating levels of 25-hydroxyvitamin D (25[OH]D, the widely accepted biomarker of vitamin D status) and increased risk of ARI caused by other pathogens.^{3,4} Randomised controlled trials (RCTs) of vitamin D for the prevention of ARI have produced heterogeneous results, with some showing protection, and others reporting null findings. We previously meta-analysed individual participant data from 10,933 participants in 25 RCTs⁵⁻²⁹ and showed a protective overall effect that was stronger in those with lower baseline 25(OH)D levels, and in trials where vitamin D was administered daily or weekly rather than in more widely spaced bolus doses.³⁰ Since the date of the final literature search performed for that study (December 2015), 20 RCTs with 62,063 participants fulfilling the same eligibility criteria have been completed.³¹⁻⁵⁰ We therefore sought data from these more recent studies for inclusion in an updated meta-analysis of stratified aggregate (trial-level) data to determine whether vitamin D reduced ARI risk overall, and to evaluate whether effects of vitamin D on ARI risk varied according to baseline 25(OH)D concentration and/or dosing regimen (frequency, dose size, and trial duration).

Methods

Protocol, Registration and Ethical Approvals

Methods were pre-specified in a protocol that was registered with the PROSPERO International Prospective Register of Systematic Reviews (https://www.crd.york.ac.uk/PROSPERO/display_record.php?RecordID=190633). Research Ethics Committee approval to conduct this meta-analysis was not required in the UK; local ethical permission to contribute data from primary trials was required and obtained for studies by Camargo *et al*¹³ (The Ethics Review Committee of the Mongolian Ministry of Health), Murdoch *et al*¹⁴ (Southern Health and Disability Ethics Committee, ref. URB/09/10/050/AM02), Rees *et al*¹⁷ (Committee for the Protection of Human Subjects, Dartmouth College, USA; Protocol # 24381), Tachimoto *et al*⁸ (Ethics committee of the Jikei University School of Medicine, ref 26-333: 7839), Tran *et al*¹⁸ (QIMR Berghofer Medical Research Institute Human Research Ethics Committee, P1570) and Urashima *et al*^{6,20} (Ethics committee of the Jikei University School of Medicine, ref 26-333: 7839).

Eligibility Criteria

Randomised, double-blind, trials of supplementation with vitamin D₃, vitamin D₂ or 25(OH)D of any duration, with a placebo or low-dose vitamin D control, were eligible for inclusion if they had been approved by a Research Ethics Committee and if data on incidence of ARI were collected prospectively and pre-specified as an efficacy outcome. The latter requirement was imposed in order to minimise misclassification bias (prospectively designed instruments to capture ARI events were deemed more likely to be sensitive and specific for this outcome). Studies reporting results of long-term follow-up of primary RCTs were excluded.

Study Identification and Selection

Two investigators (ARM and DAJ) searched MEDLINE, EMBASE, the Cochrane Central Register of Controlled Trials (CENTRAL), Web of Science and the ClinicalTrials.gov registry using the electronic search strategies described in the Methods Section of Supplementary Material. Searches were regularly updated up to and including 1st May

2020. No language restrictions were imposed. These searches were supplemented by searching review articles and reference lists of trial publications. Collaborators were asked if they knew of any additional trials. Three investigators (DAJ, CAC and ARM) determined which trials met the eligibility criteria.

Data Collection Processes

Summary data from trials which contributed to our previous meta-analysis of individual participant data³⁰ were extracted from our central database, with permission from the Principal Investigators. Summary data relating to the primary outcome (overall and by sub-group) and secondary outcomes (overall only) from newly identified trials were requested from Principal Investigators. On receipt, they were assessed for consistency with associated publications. Study authors were contacted to provide missing data and to resolve any queries arising from these consistency checks. Once queries had been resolved, clean summary data were uploaded to the study database, which was held in STATA IC v14.2 (StataCorp, College Station, TX).

Data relating to study characteristics were extracted for the following variables: study setting, eligibility criteria, 25(OH)D assay and levels, details of intervention and control regimens, trial duration, case definitions for ARI and number entering primary analysis (after randomisation). Follow-up summary data were requested for the proportions of participants experiencing one or more ARI during the trial, both overall and stratified by baseline serum 25(OH)D concentration, where this was available. We also requested summary data on the proportions of participants who experienced one or more of the following events during the trial: upper respiratory infection (URI); lower respiratory infection (LRI); Emergency Department attendance and/or hospital admission for ARI; death due to ARI or respiratory failure; use of antibiotics to treat an ARI; absence from work or school due to ARI; a serious adverse event; death due to any cause; and potential adverse reactions to vitamin D (hypercalcaemia and renal stones).

Risk of Bias Assessment for Individual Studies

We used the Cochrane Collaboration Risk of Bias tool⁵¹ to assess the following variables: sequence generation, allocation concealment, blinding of participants,

personnel and outcome assessors, completeness of outcome data, evidence of selective outcome reporting and other potential threats to validity. Study quality was assessed independently by two investigators (ARM and DAJ), except for the five trials for which DAJ and/or ARM were investigators, which were assessed by CAC. Discrepancies were resolved by consensus.

Definition of outcomes

The primary outcome of the meta-analysis was the proportion of participants experiencing one or more ARIs, with the definition of ARI encompassing events classified as URI, LRI and ARI of unclassified location (i.e. infection of the upper and/or lower respiratory tract). Secondary outcomes were incidence of URI and LRI, analysed separately; incidence of Emergency Department attendance and/or hospital admission for ARI; death due to ARI or respiratory failure; use of antibiotics to treat an ARI; absence from work or school due to ARI; incidence of serious adverse events; death due to any cause; and incidence of potential adverse reactions to vitamin D (hypercalcaemia and renal stones).

Synthesis Methods

Data were analysed by DAJ; results were checked and verified by JDS. Our meta-analysis approach followed published guidelines.⁵² The primary comparison was of participants randomised to vitamin D vs. placebo: this was performed for all of the outcomes listed above. For trials that included higher-dose, lower-dose and placebo arms, data from higher-dose and lower-dose arms were pooled for analysis of the primary comparison. A secondary comparison of participants randomised to higher vs. lower doses of vitamin D was performed for the primary outcome only. A log odds ratio and its standard error was calculated for each outcome within each trial from the proportion of participants experiencing one or more events in the intervention vs. control arm. These were meta-analysed in a random effects model using the Metan package⁵³ within STATA IC v14.2 to obtain a pooled odds ratio with a 95% confidence interval and a measure of heterogeneity summarized by the I^2 statistic and its corresponding P value.

Exploration of variation in effects

To explore reasons for heterogeneity of effect of the intervention between trials we performed a stratified analysis according to baseline vitamin D status (serum 25[OH]D <25 vs. 25-49.9 vs. 50-74.9 vs. ≥75 nmol/L) and according to age at baseline (<1 vs. 1-15.99 vs. 16-64.99 vs. ≥65 yrs). We additionally conducted sub-group analyses according to vitamin D dosing regimen (administration of daily vs. weekly vs. monthly or less frequent doses), dose size (daily equivalent <400 IU vs. 400-1000 IU vs. 1001-2000 IU vs. >2,000 IU), trial duration (≤12 months vs. >12 months) and presence of airway disease (trial restricted to participants with asthma vs. those restricted to participants with COPD vs. those in which participants without airway disease were eligible). The thresholds for baseline 25(OH)D concentration used in sub-group analyses were selected *a priori* on the basis that they represent cut-offs that are commonly used to distinguish profound vitamin D deficiency (<25 nmol/L), moderate vitamin D deficiency (25-49.9 nmol/L) and sub-optimal vitamin D status (50-74.9 nmol/L).⁵⁴ An exploratory analysis restricted to studies with optimal dosing frequency, dose size and duration was also performed.

To investigate factors associated with heterogeneity of effect between subgroups of trials, we performed multivariable meta-regression analysis on trial-level characteristics, namely, dose frequency, dose size and trial duration, to produce an adjusted odds ratio, a 95% confidence interval and a P value for interaction for each factor. Independent variables were dichotomised to create a more parsimonious model (serum 25(OH)D of <25 vs. ≥25 nmol/L; administration of daily vs. non-daily doses; daily equivalent of ≤1000 IU vs. >1000 IU; and trial duration of ≤12 vs. >12 months). The meta-regression analysis excluded data from two trials that included higher-dose, lower-dose and placebo arms,^{18,47} since the higher-dose and lower-dose arms spanned the 1,000 IU/day cut-off, rendering them unclassifiable for the purposes of this analysis.

Quality Assessment Across Studies

For the primary analysis, the likelihood of publication bias was investigated through the construction of a contour-enhanced funnel plot.⁵⁵ We used the five GRADE considerations (study limitations, consistency of effect, imprecision, indirectness and

publication bias)⁵⁶ to assess the quality of the body of evidence contributing to analyses of the primary efficacy outcome and major secondary outcomes of our meta-analysis.

Sensitivity analyses

We conducted three exploratory sensitivity analyses for the primary comparison of the primary outcome: one excluded RCTs where risk of bias was assessed as being unclear; one excluded RCTs in which incidence of ARI was not the primary or co-primary outcome; and one substituted diary-defined ARI events (available for 2598 participants) for survey-defined ARI events (available for n=16,000 participants) from the trial by Pham et al.⁴⁴

Role of the funding source

This study was conducted without external funding.

Results

Study selection and data obtained

The study selection process is illustrated in Figure 1. Our search identified a total of 1,528 unique studies that were assessed for eligibility, of which 45 studies with a total of 73,384 randomised participants fulfilled eligibility criteria. Studies for which full text was reviewed prior to exclusion due to ineligibility are listed in Table S1. Of the 45 eligible studies identified, 34 compared effects of a single vitamin D regimen vs. placebo only,^{5-17,19,20,22,23,25-28,31,33,36,38,39,41-45,48-50} 5 compared effects of higher-dose, lower-dose and placebo arms,^{18,21,24,40,47} and 6 compared effects of higher- vs. lower-dose regimens of vitamin D only.^{29,32,34,35,37,46} Stratified aggregate data were sought and obtained for all but 3 eligible studies.⁴⁸⁻⁵⁰ Data for the primary outcome (proportion of participants with one or more ARI) were obtained for 46,331 (98.0%) of 47,262 participants in 42 studies.^{5-29,31-47}

Study and participant characteristics

Characteristics of the 42 studies contributing data to this meta-analysis and their participants are presented in Table 1. Trials were conducted in 18 different countries on 5 continents, and enrolled participants of both sexes from birth to 95 years of age. Baseline serum 25(OH)D concentrations were determined in 34 of 42 trials: mean baseline 25(OH)D concentration ranged from 18.9 to 90.9 nmol/L (to convert to ng/ml, divide by 2.496). Forty-one studies administered oral vitamin D₃ to participants in the intervention arm, while 1 study administered oral 25(OH)D. Vitamin D was given as monthly to 3-monthly bolus doses in 13 studies; as weekly doses in 6 studies; as daily doses in 21 studies; and as a combination of bolus and daily doses in 2 studies. Trial duration ranged from 8 weeks to 5 years. Incidence of ARI was primary or co-primary outcome for 22 studies, and a secondary outcome for 20 studies.

Risk of Bias Within Studies

Details of the risk of bias assessment are provided in supplementary Table S2. Four trials were assessed as being at unclear risk of bias due to high loss to follow-up. In the trial by Laaksi and colleagues,⁸ 37% of randomised participants were lost to follow-up. In the trial by Dubnov-Raz and colleagues,²⁶ 52% of participants did not complete all symptom questionnaires. In the unpublished trial by Reyes and colleagues,⁴⁷ loss to follow-up ranged from 33% to 37% across the three study arms, and in the unpublished trial by Golan-Tripto and colleagues,⁴⁶ 50% of participants were lost to follow-up. All other trials were assessed as being at low risk of bias for all seven aspects assessed.

Overall Results, Primary Outcome

For the primary comparison of vitamin D vs. placebo control, supplementation resulted in a statistically significant reduction in the proportion of participants experiencing at least one ARI (Odds Ratio [OR] 0.91, 95% Confidence Interval [CI] 0.84 to 0.99; 44,009 participants in 36 studies; Figure 2, Table 2; Cates Plot, Figure S1). Heterogeneity of effect was moderate (I^2 37.2%, P for heterogeneity 0.01).

For the secondary comparison of higher- vs. lower-dose vitamin D, we observed no statistically significant difference in the proportion of participants with at least one ARI (OR 0.87, 95% CI 0.73 to 1.04; 3,047 participants in 11 studies; I^2 0.0%, P for heterogeneity 0.50; Figure S2).

Sub-group analyses, Primary Outcome

To investigate reasons for the observed heterogeneity of effect for the primary comparison of vitamin D vs. placebo control, we stratified this analysis by two participant-level factors (baseline vitamin D status and age) and by four trial-level factors (dose frequency, dose size, trial duration, and airway disease comorbidity). Results are presented in Table 2 and Figures S3-S8. No statistically significant effect of vitamin D was seen for participants with baseline 25(OH)D <25 nmol/L (OR 0.78, 95% CI 0.53 to 1.16; 3,617 participants in 19 studies), 25-49.9 nmol/L (OR 1.03, 95% CI 0.92 to 1.15; 9,167 participants in 28 studies), 50-74.9 nmol (OR 0.90, 95% CI 0.76 to 1.06; 5,417 participants in 29 studies), or ≥ 75 nmol/L (OR 0.97, 95% CI 0.81 to 1.16; 3,014 participants in 25 studies; Figure S3). A statistically significant protective effect of vitamin

D was seen for participants aged 1-15.9 years (OR 0.71, 95% CI 0.57 to 0.90; 11,871 participants in 15 studies), but not in participants aged <1 year (OR 0.95, 95% CI 0.82 to 1.10; 5,697 participants in 5 studies), 16-64.99 years (OR 0.97, 95% CI 0.93 to 1.09; 9,603 participants in 21 studies), or ≥ 65 years (OR 0.96, 95% CI 0.90 to 1.02; 16,983 participants in 16 studies; Figure S7).

With regard to dosing frequency, a statistically significant protective effect was seen for trials where vitamin D was given daily (OR 0.75, 95% CI 0.61 to 0.93; 4,005 participants in 18 studies), but not for trials in which it was given weekly (OR 0.97, 95% CI 0.88 to 1.06; 12,756 participants in 6 studies), or monthly to 3-monthly (OR 0.98, 95% CI 0.93 to 1.03; 27,248 participants in 12 studies; Figure S4). Statistically significant protective effects of the intervention were also seen in trials where vitamin D was administered at daily equivalent doses of 400-1000 IU (OR 0.70, 95% CI 0.55 to 0.89; 2,305 participants in 10 studies), but not where the daily dose equivalent was <400 IU (OR 0.65, 95% CI 0.31 to 1.37; 2,308 participants in 2 studies), 1001-2000 IU (OR 0.97, 95% CI 0.92 to 1.03; 31,702 participants in 15 studies), or >2000 IU (OR 1.05, 95% CI 0.84 to 1.31; 6,906 participants in 7 studies; Figure S5). Statistically significant protective effects were also seen for trials with a duration of ≤ 12 months (OR 0.82, 95% CI 0.72 to 0.93; 9,255 participants in 29 studies) but not in those lasting >12 months (OR 0.99, 95% CI 0.95 to 1.04; 34,754 participants in 7 studies; Figure S6).

Finally, statistically significant protective effects were also seen for trials that were not restricted to participants with asthma or COPD (OR 0.91, 95% CI 0.84 to 0.99; 42,799 participants in 30 studies), but not in trials that exclusively enrolled participants with asthma (OR 0.73, 95% CI 0.36 to 1.49; 795 participants in 4 studies), or COPD (OR 1.01, 95% CI 0.68 to 1.51; 415 participants in 2 studies; Figure S8).

An exploratory analysis restricted to eight placebo-controlled trials investigating effects of daily dosing at doses of 400-1000 IU/day with duration ≤ 12 months (for which mean baseline 25(OH)D level ranged from 54.8 nmol/L to 88.9 nmol/L) showed a statistically significant reduction in the proportion of participants experiencing at least one ARI (OR 0.58, 95% CI 0.45 to 0.75; 1,232 participants in 8 studies; Figure S9; Cates Plot, Figure S1). Heterogeneity of effect for this exploratory analysis was low (I^2 0.0%, P for heterogeneity 0.67).

Multivariable Meta-Regression Analysis

Multivariable meta-regression analysis of trial-level sub-groups did not identify a statistically significant interaction between allocation to vitamin D vs. placebo and dose frequency, size or trial duration (Table S3).

Secondary outcomes

Meta-analysis of secondary outcomes was performed for results of placebo-controlled trials only; results are presented in Table 3. Overall, without consideration of participant- or trial-level factors, vitamin D supplementation did not have a statistically significant effect on the proportion of participants with one or more URI, LRI, courses of antimicrobials for ARI, work/school absences due to ARI, hospitalisations or emergency department attendances for ARI, serious adverse events of any cause, death due to ARI or respiratory failure, death due to any cause, or episodes of hypercalcaemia or renal stones.

Risk of bias across studies

A funnel plot for the proportion of participants experiencing at least one ARI (Figure S10) showed left-sided asymmetry, confirmed with an Egger's regression test⁵⁷ ($P=0.008$). This might reflect heterogeneity of effect across trials, or publication bias arising from omission of small trials showing non-protective effects of vitamin D from the meta-analysis.⁵⁸ Given the latter possibility, the quality of the body of evidence contributing to analyses of the primary efficacy outcome and major secondary outcomes was downgraded to moderate (Table S4).

Sensitivity Analyses

Results of exploratory sensitivity analyses are presented in Table S5. Meta-analysis of the proportion of participants in placebo-controlled trials experiencing at least one ARI, excluding 4 studies assessed as being at unclear risk of bias,^{8,26,46,47} revealed protective

effects of vitamin D supplementation consistent with the main analysis (OR 0.93, 95% CI 0.86 to 1.00; 43,626 participants in 33 studies). Sensitivity analysis for the same outcome, excluding 18 placebo-controlled trials that investigated ARI as a secondary outcome, did not show a statistically significant protective effect (OR 0.89, 95% CI 0.77 to 1.03; 7,537 participants in 18 studies). A sensitivity analysis for the same outcome, substituting diary-defined ARI events (available for 2598 participants) for survey-defined ARI events (available for n=16,000 participants) in the trial by Pham et al⁴⁴ revealed protective effects of vitamin D supplementation consistent with the main analysis (OR 0.90, 95% CI 0.83 to 0.99; 30,607 participants in 36 studies).

Discussion

This updated meta-analysis of RCTs of vitamin D supplementation for the prevention of ARI includes data from an additional 17 studies completed since December 2015, when we performed the final literature search for our prior individual participant data meta-analysis.³⁰ For expediency during the COVID-19 pandemic, we used a trial-level approach for this update, which includes data from a total of 46,331 participants in 42 trials. Overall, we report a modest statistically significant protective effect of vitamin D supplementation, as compared with placebo (OR 0.91, 95% CI 0.84 to 0.99). As expected, there was significant heterogeneity ($P=0.01$) across trials, which might have led to an under-estimate of the protective effect, and contributed to the asymmetry observed in the funnel plot.⁵⁸ Alternatively, left-sided asymmetry in the funnel plot may reflect publication bias, which might have led to an over-estimate of the protective effect. In contrast to findings of our previous meta-analysis,³⁰ we did not observe enhanced protection in those with the lowest 25(OH)D levels at baseline. However, there was evidence that efficacy of vitamin D supplementation varied according to dosing regimen and trial duration, with protective effects associated with daily administration of doses of 400-1000 IU vitamin D given for ≤ 12 months. An exploratory analysis restricted to data from 8 trials fulfilling these design criteria revealed a larger protective effect (OR 0.58, 95% CI 0.45 to 0.75) without significant heterogeneity across trials (P for heterogeneity 0.67).

The magnitude of the overall protective effect seen in the current analysis (OR 0.91, 95% CI 0.84 to 0.99) is modest, and similar to the value reported in our previous meta-

analysis of individual participant data (adjusted OR 0.88, 95% CI 0.81 to 0.96).³⁰ In keeping with our previous study, the point estimate for this effect was lower among those with baseline 25(OH)D <25 nmol/L than in those with higher baseline vitamin D status. However, in contrast to our previous finding, a statistically significant protective effect of vitamin D was not seen in those with the lowest 25(OH)D concentrations. This difference reflects the inclusion of null data from four new RCTs in which vitamin D was given in relatively high doses at weekly or monthly intervals over 2-5 years.^{41,42,44,45} Null results of these studies contrast with protective effects reported from earlier trials in which smaller daily doses of vitamin D were given over shorter periods.^{8,9,13,16} These differing findings suggest that the frequency, amount and duration of vitamin D supplementation may be key determinants of its protective efficacy. In keeping with this hypothesis, statistically significant protective effects of vitamin D were seen for meta-analysis of trials where vitamin D was given daily; where it was given at doses of 400-1000 IU/day; and where it was given for 12 months or less. When results of trials that investigated daily administration of 400-1000 IU over ≤ 12 months were pooled in an exploratory meta-analysis, a protective effect was seen (OR 0.58, 95% CI 0.45 to 0.75) with low heterogeneity (I^2 0.0%, P for heterogeneity 0.67). Greater protective efficacy of lower vs higher doses of vitamin D might reflect deleterious effects of higher-dose vitamin D on its own metabolism, or on host responses to respiratory pathogens: head-to-head mechanistic studies in individuals randomised to different regimens of vitamin D supplementation are needed to investigate this issue.

The current study has several strengths: it contains the very latest RCT data available in this fast-moving field, including findings from three large phase 3 trials published in 2020^{41,42,44} as well as some as-yet unpublished studies.^{46,47} The inclusion of additional studies allowed us to analyse results of placebo-controlled studies vs. high-dose / low-dose studies separately, and gave us the power to investigate reasons for heterogeneity of effect observed across trials. For example, we could distinguish the effects of daily vs. weekly dosing, which were previously pooled.³⁰

Our work also has limitations. Given the need to generate a rapid update of our previous work in the context of the COVID-19 pandemic, we meta-analysed aggregate (trial-level) data, rather than individual participant data; this allowed us to proceed rapidly, without

the delays introduced by the need to establish multiple data sharing agreements. However, we did contact authors to get unpublished estimates of effect that were stratified by pre-defined baseline 25(OH)D levels, harmonised across studies: thus, we were able to provide accurate data for the major participant-level effect-modifier of interest. Despite the large number of trials overall, relatively few compared effects of lower- vs. higher-dose vitamin D: our power for this secondary comparison was therefore limited. We lacked the individual participant data to investigate race/ethnicity and obesity as potential effect-modifiers. We also could not account for other factors that might influence the efficacy of vitamin D supplements for ARI prevention (e.g., taking the supplement with or without food) or secular trends that would influence trials, such as the increased societal use of vitamin D supplements;⁵⁹ concurrent use of standard dose vitamin D supplements or multivitamins in the “placebo” group would effectively render these as high- vs. low-dose trials and potentially drive results toward the null. A final limitation relates to the funnel plot, which suggests that the overall effect size may have been over-estimated due to publication bias; we have mitigated this by inclusion of data from unpublished studies identified by searching clinicaltrials.gov where this was obtainable.

In summary, this updated meta-analysis of data from RCTs of vitamin D for the prevention of ARI showed a statistically significant overall protective effect of the intervention. The protective effect was heterogenous across trials; it also may have been over-estimated due to publication bias. In contrast to findings of our previous meta-analysis of individual participant data, we did not see a protective effect of vitamin D supplementation among those with the lowest baseline vitamin D status. The vitamin D dosing regimen of most benefit was daily and used standard doses (e.g., 400 to 1000 IU) for up to 12 months. The relevance of these findings to COVID-19 is not known and requires investigation.

Acknowledgements

This study was conducted without external funding. DAJ is supported by a Barts Charity Lectureship (ref MGU045). ARM is supported by the United Kingdom Office for Students. The views expressed are those of the authors and not necessarily those of Barts Charity or the Office for Students. Sources of support for individual trials are detailed in Supplementary Material. We thank all the people who participated in primary randomised controlled trials, and the teams who conducted them.

Author Contributions

DAJ and ARM wrote the study protocol and designed statistical analyses. DAJ, CAC and ARM assessed eligibility of studies for inclusion and performed risk of bias assessments. Statistical analyses were done by DAJ; results were checked and verified by JDS. DAJ and ARM wrote the first draft of the report. All authors revised it critically for important intellectual content, gave final approval of the version to be published, and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work were appropriately investigated and resolved.

Competing Interests

All authors have completed the ICMJE uniform disclosure form. No author has had any financial relationship with any organisations that might have an interest in the submitted work in the previous three years. No author has had any other relationship, or undertaken any activity, that could appear to have influenced the submitted work.

Transparency Declaration

DAJ and ARM are the manuscript's guarantors and they affirm that this is an honest, accurate, and transparent account of the study being reported and that no important aspects of the study have been omitted. All analyses were pre-specified in the study protocol, other than the exploratory analyses whose results are presented in Table 2 (sub-group analyses by age and presence of asthma/COPD, requested by reviewers), Table S5 and Figure S7.

Data Sharing: the study dataset is available from d.a.jolliffe@qmul.ac.uk.

Table 1: Characteristics of the 42 eligible trials and their participants

Study first author, year	Setting	Participants	Mean age, years (s.d.) [range]	Male: Female	25(OH)D assay, EQA scheme	Mean baseline 25(OH)D, nmo/L (s.d.)	Baseline 25(OH)D <25 nmol/L (%)	Mean attained 25(OH)D, intervention arm, nmo/L (s.d.)	Intervention: Control (total)	Oral dose of vitamin D ₃ , intervention arm	Control	Trial duration	ARI definition	ARI primary or secondary outcome?	N contributing data / N randomised (%)
Li-Ng 2009 ⁵	USA	Healthy adults	57.9 (13.6) [21.4 – 80.6]	34:128	RIA (DiaSorin), DEQAS	63.7 (25.5)	3/150 (2.0)	88.5 (23.2)	84:78 (162)	50 µg daily	Placebo	3 mo	URI: ≥2 URI symptoms in absence of allergy symptoms	Primary	157/162 (96.9)
Urashima 2010 ⁶	Japan	Schoolchildren	10.2 (2.3) [6.0 – 15.0]	242:188	Not determined	Not determined	Not determined	Not determined	217:213 (430)	30 µg daily	Placebo	4 mo	URI: influenza A/B diagnosed by RIDT or RIDT-negative ILI	Primary	334/430 (77.7)
Manaseki-Holland 2010 ⁷	Afghanistan	Pre-school children with pneumonia	1.1 (0.8) [0.1 – 3.3]	257:196	Not determined	Not determined	Not determined	Not determined	224:229 (453)	2.5 mg bolus once	Placebo	3 mo	LRI: repeat episode of pneumonia – age-specific tachypnoea without wheeze	Secondary	453/453 (100.0)
Laaksi 2010 ⁸	Finland	Military conscripts	19.1 (0.6) [18.0 – 21.0]	164:0	EIA (IDS OCTEIA)	75.9 (18.7)	0/73 (0.0)	71.6 (22.9)	80:84 (164)	10 µg daily	Placebo	6 mo	ARI: medical record diagnosis	Primary	164/164 (100.0)
Majak 2011 ⁹	Poland	Children with asthma	10.9 (3.3) [6.0 – 17.0]	32:16	RIA (BioSource Europe), RIQAS	88.9 (38.2)	0/48 (0.0)	37.6 (13.1)	24:24 (48)	12.5 µg daily	Placebo	6 mo	ARI: self-report	Secondary	48/48 (100.0)
Trilok-Kumar 2011 ¹⁰	India	Low birthweight infants	0.1 (0.0) [0.0 – 0.3]	970:1109	--	Not determined	Not determined	55.0 (22.5)	1039:1040 (2079)	35 µg weekly	Placebo	6 mo	ARI: medical record diagnosis of events causing hospitalisation	Secondary	2064/2079 (99.3)
Lehouck 2012 ¹¹	Belgium	Adults with COPD	67.9 (8.3) [48.0 – 86.0]	145:37	RIA (Diasorin), DEQAS	49.8 (29.2)	31/182 (17.0)	130.0 (44.7)	91:91 (182)	2.5 mg bolus monthly	Placebo	1 yr	URI: self-report	Secondary	175/182 (96.2)
Manaseki-Holland 2012 ¹²	Afghanistan	Infants	0.5 (0.3) [0.0 – 1.0]	1591:1455	--	Not determined	Not determined	32.7 (17.1)	1524:1522 (3046)	2.5 mg bolus 3-monthly	Placebo	1.5 yr	LRI: pneumonia confirmed by chest radiograph	Primary	3011/3046 (98.9)
Camargo 2012 ¹³	Mongolia	3 rd /4 th grade schoolchildren	10.0 (0.9) [7.0 – 12.7]	129:118	LC-MS/MS, DEQAS	18.9 (9.7)	192/245 (78.4)	49.1 (15.1)	143:104 (247)	7.5 µg daily	Placebo	7 wk	ARI: parent-reported 'chest infections or colds'	Secondary	244/247 (98.8)
Murdoch 2012 ¹⁴	New Zealand	Healthy adults	48.1 (9.7) [18.0 – 67.6]	81:241	LC-MS/MS, DEQAS	72.1 (22.1)	5/322 (1.6)	123.6 (27.5)	161:161 (322)	2 x 5 mg bolus monthly then 2.5 mg bolus monthly	Placebo	1.5 yr	URI: assessed with symptom score	Primary	322/322 (100.0)
Bergman 2012 ¹⁵	Sweden	Adults with increased susceptibility to ARI	53.1 (13.1) [20.0 – 77.0]	38:102	CLA (DiaSorin), DEQAS	49.3 (23.2)	15/131 (11.45)	94.9 (38.1)	70:70 (140)	100 µg daily	Placebo	1 yr	URI: assessed with symptom score	Secondary	124/140 (88.6)
Marchisio 2013 ¹⁶	Italy	Children with recurrent acute otitis media	2.8 (1.0) [1.3 – 4.8]	64:52	CLA (DiaSorin), ISO9001	65.3 (17.3)	2/116 (1.7)	90.3 (21.1)]	58:58 (116)	25 µg daily	Placebo	6 mo	URI: doctor-diagnosed acute otitis media	Primary	116/116 (100.0)
Rees 2013 ¹⁷	USA	Adults with previous colorectal adenoma	61.2 (6.6) [47.1 – 77.9]	438:321 ^(a)	RIA (IDS), DEQAS	62.5 (21.3)	0/759 (0.0)	186.9 (455.1)	399:360 (759)	25 µg daily	Placebo	13 mo (average)	URI: assessed from daily symptom diary	Secondary	759/759 (100.0)
Tran 2014 ¹⁸	Australia	Healthy older adults	71.7 (6.9) [60.3 – 85.2]	343:301	CLA (DiaSorin), DEQAS	41.7 (13.5)	66/643 (10.3)	71.0 (19.6)	430:214 (644)	0.75 mg bolus vs. 1.5 mg bolus monthly	Placebo	1 yr	URI: self-reported cold	Secondary	594/644 (92.2)
Goodall 2014 ¹⁹	Canada	Healthy university students	19.6 (2.2) [17.0 – 33.0]	218:382	Not determined	Not determined	Not determined	Not determined	300:300 (600)	0.25 mg weekly (factorial with gargling)	Placebo	8 wk	URI: self-reported cold	Primary	492/600 (82.0)
Urashima 2014 ²⁰	Japan	High school students	16.5 (1.0) [15.0 – 18.0]	162:85	Not determined	Not determined	Not determined	Not determined	148:99 (247)	50 µg daily	Placebo	2 mo	URI: influenza A diagnosed by RIDT or RIDT-negative ILI	Primary	247/247 (100.0)
Grant 2014 ²¹	New Zealand	Pregnant women and offspring	Offspring unborn at baseline	0:260 (pregnant women) 121:128 (offspring)	LC-MS/MS, DEQAS	54.8 (25.8)	30/200 (15.0)	92.9 (41.6)	173:87 (pregnant women, 260) 164:85 (offspring, 249)	Pregnant women: 25 µg vs. 50 µg daily. Offspring: 10 µg vs. 20 µg daily	Placebo	9 mo (3 mo in pregnancy + 6 mo in infancy)	ARI: doctor-diagnosed ARI precipitating primary care consult	Secondary	236/260 (90.8)

Study first author, year	Setting	Participants	Mean age, years (s.d.) [range]	Male: Female	25(OH)D assay, EQA scheme	Mean baseline 25(OH)D, nmo/L (s.d.)	Baseline 25(OH)D <25 nmo/L (%)	Mean attained 25(OH)D, intervention arm, nmo/L (s.d.)	Intervention: Control (total)	Oral dose of vitamin D ₃ , intervention arm	Control	Trial duration	ARI definition	ARI primary or secondary outcome?	N contributing data / N randomised (%)
Martineau 2015a ²² [ViDiCO]	UK	Adults with COPD	64.7 (8.5) [40.0 – 85.0]	144:96	LC-MS/MS, DEQAS	46.1 (25.7)	50/240 (20.8)	67.3 (27.5)	122:118 (240)	3 mg bolus 2-monthly	Placebo	1 yr	URI: assessed from daily symptom diary	Co-primary	240/240 (100.0)
Martineau 2015b ²³ [ViDIAs]	UK	Adults with asthma	47.9 (14.4) [16.0 – 78.0]	109:141	LC-MS/MS, DEQAS	49.6 (24.7)	36/250 (14.4)	69.4 (21.0)	125:125 (250)	3 mg bolus 2-monthly	Placebo	1 yr	URI: assessed from daily symptom diary	Co-primary	250/250 (100.0)
Martineau 2015c ²⁴ [ViDiFlu]	UK	Older adults and their carers	67.1 (13.0) [21.4 – 94.0]	82:158	LC-MS/MS, DEQAS	42.9 (23.0)	60/240 (25.0)	84.8 (24.1)	137:103 (240)	Older adults: 2.4 mg bolus 2-monthly + 10 µg daily Carers: 3 mg 2-monthly	Older adults: placebo + 10 µg daily Carers: placebo	1 yr	URI & LRI, both assessed from daily symptom diary	Co-primary	240/240 (100.0)
Simpson 2015 ²⁵	Australia	Healthy adults	32.2 (12.2) [18.0 – 52.0]	14:20	LC-MS/MS, DEQAS	67.9 (23.0)	0/33 (0.0)	Not determined	18:16 (34)	0.5 mg weekly	Placebo	17 wk	ARI assessed with symptom score	Primary	34/34 (100.0)
Dubnov-Raz 2015 ²⁶	Israel	Adolescent swimmers with vitamin D insufficiency	15.2 (1.6) [12.9 – 18.6]	34:20	RIA (DiaSorin), DEQAS	60.4 (11.9)	0/54 (0.0)	73.7 (16.6)	27:27 (54)	50 µg daily	Placebo	12 wk	URI assessed with symptom score	Primary	25/54 (46.3)
Denlinger 2016 ²⁷	USA	Adults with asthma	39.2 (12.9) [18.0 – 85.0]	130:278	CLA (DiaSorin), VDSP	47.0 (16.9)	55/408 (13.5)	104.3 (32.4)	201:207 (408)	2.5 mg bolus then 100 µg daily	Placebo	28 wk	URI assessed with symptom score	Secondary	408/408 (100.0)
Tachimoto 2016 ²⁸	Japan	Children with asthma	9.9 (2.3) [6.0 – 15.0]	50:39	RIA (DiaSorin), CAP	74.9 (24.6)	1/89 (1.1)	85.7 (24.5)	54:35 (89)	20 µg daily, first 2 mo.	Placebo	6 mo	URI: assessed with symptom score	Secondary	89/89 (100.0)
Ginde, 2016 ²⁹	USA	Institutionalised older adults	80.7 (9.9) [60.0 – 95.0]	45:62	LC-MS/MS, VDSP	57.3 (22.7)	12/107 (11.2)	Not determined	55:52 (107)	2.5 mg bolus monthly + ≤25 µg per day equivalent	Placebo + 10-25 µg per day equivalent	1 yr	ARI: medical record diagnosis	Primary	107/107 (100.0)
Gupta 2016 ³¹	India	Children with pneumonia	1.4 (1.1) [0.5 – 5.0]	226:98	RIA (Immunotech SAS/ DiaSorin)	43.9 (33.4)	104/312 (33.3)	64.1 (43.9)	162:162 (324)	2.5 mg bolus, single dose	Placebo	6 mo	Physician confirmed recurrent pneumonia	Co-primary	314/324 (96.9)
Aglipay 2017 ³²	Canada	Healthy children	2.7 (1.5) [1.0 – 5.0]	404:296	CLA (Roche ELECSYS)	90.9 (20.9)	1/703 (0.1)	High dose: 121.6 (2.2); Low dose: 91.9 (1.7)	349:354	50 µg daily	10 µg daily	4-8 mo (mean 6.3 mo)	URI: lab confirmed	Primary	699/703 (99.4)
Arihiro 2018 ³³	Japan	Adults with diagnosis of inflammatory bowel disease	44.7 (1.3) [18.0 – 82.0]	136:87	RIA (Diasorin)	58.6 (22.0)	5/223 (2.2)	80.4 (21.5)	119:118 (237)	12.5 µg daily	Placebo	6 mo	Lab confirmed influenza	Primary	223/237 (94.1)
Hibbs 2018 ³⁴	USA	African American preterm infants	Offspring unborn at baseline	166:133 ^[a]	RIA	55.4 (22.2)	0/300 (0.0)	95.0 (21.2)	153:147 (300)	10 µg daily, regardless of dietary intake	10 µg daily, only if dietary intake was <5 µg daily	1 yr	ARI: self-reported URI/LRI	Secondary	300/300 (100.0)
Lee 2018 ³⁵	USA	Children and young adults with sickle cell disease	9.9 (3.9) [3.0 – 20.0]	30:32	LC-MS/MS, DEQAS	35.7 (16.5)	18/62 (29.0)	92.4 (23.7)	31:31 (62)	2.5 mg bolus monthly	0.3 mg monthly	2 yrs	Self-reported respiratory events, including ARI	Primary	62/62 (100.0)
Loeb 2018 ³⁶	Vietnam	Healthy children and adolescents	8.5 (4.0) [3.0 – 17.0]	621:679	CLA (DiaSorin), DEQAS	65.5 (16.8)	5/1153 (43.4)	91.8 (23.6)	650:650 (1300)	0.35 mg weekly	Placebo	8 mo	RT-PCR confirmed influenza A or B	Primary	1153/1300 (88.7)
Rosendahl 2018 ³⁷	Finland	Healthy infants	Offspring unborn at baseline	495:492	CLA (IDS-iSYS) VDSP	81.5 (25.9)	0/879 (0.0)	117.7 (26.1)	492:495 (987)	30 µg daily	10 µg daily	2 yrs	Parent reported infections, including ARI	Co-primary	897/987 (90.9)
Shimizu 2018 ³⁸	Japan	Healthy adults	52.7 (6.5) [45.0 – 74.0]	66:149	RIA (DiaSorin)	48.9 (13.5)	1/214 (0.5)	114.6 (32.7)	126:126 (252)	10 µg daily (25[OH] D) ^[c]	Placebo	4 mo	URI: self-reported	Primary	215/252 (85.3)
Aloia 2019 ³⁹	USA	Healthy African American women aged over 60 years	69.0 (5.3) [65.4 – 72.5]	0:260	LC-MS/MS, NIST	54.4 (16.7)	9/258 (3.5)	117.0 (28.0)	130:130 (260)	50 µg daily	Placebo	3 mo	ARI: self-reported cold/flu	Secondary	260/260 (100.0)
Hauger 2019 ⁴⁰	Denmark	Healthy children	6.6 (1.5) [4.0 – 8.0]	61:69	LC-MS/MS, DEQAS	56.8 (12.5)	0/118 (0.0)	20 µg arm: 75.8 (11.5) 10 µg arm: 61.8 (10.6)	43:44:43 (130)	20 µg / 10 µg daily	Placebo	5 mo	ARI: self-reported	Secondary	118/130 (90.8)

Study first author, year	Setting	Participants	Mean age, years (s.d.) [range]	Male: Female	25(OH)D assay, EQA scheme	Mean baseline 25(OH)D, nmol/L (s.d.)	Baseline 25(OH)D <25 nmol/L (%)	Mean attained 25(OH)D, intervention arm, nmol/L (s.d.)	Intervention: Control (total)	Oral dose of vitamin D ₃ , intervention arm	Control	Trial duration	ARI definition	ARI primary or secondary outcome?	N contributing data / N randomised (%)
Camargo 2020 ⁴¹	New Zealand	Older adults	66.4 (8.3) [50.0 – 84.0]	2,935:2,121	LC-MS/MS, DEQAS	63.4 (23.6)	89/5056 (1.8)	135.0 (39.9)	2558:2552 (5110)	5 mg bolus loading dose; then 2.5 mg bolus monthly	Placebo	3 yrs	ARI: self-reported cold/flu	Secondary	5056/5110 (98.9)
Ganmaa, 2020 ⁴²	Mongolia	Healthy school children	9.4 (1.6) [6.0 – 13.0]	4485:4366	EIA (Biomerieux), DEQAS	29.7 (10.5)	2813/8851 (31.8)	77.4 (22.7)	4418:4433 (8851)	0.35 mg weekly	Placebo	3 yrs	ARI: self-reported	Secondary	8851/8851 (100.0)
Mandlik 2020 ⁴³	India	Healthy children	8.1 (1.2) [6.0 – 12.0]	158:127	EIA (DLD diagnostics)	58.9 (10.9)	0/237 (0.0)	80 (23.3)	135:150 (285)	25 µg daily + 500 mg calcium	Placebo	6 mo	URI: self-reported	Secondary	244/285 (85.6)
Pham 2020 ⁴⁴	Australia	Older adults	69.3 (5.5) [60.0 – 86.0]	8678:7322	LC-MS/MS, VDSP	Not determined	Not determined	114.8 (30.3) ^[d]	8000:8000 (16000)	1.5 mg bolus monthly	Placebo	5 yrs	ARI: self-reported	Secondary	16,000/16,000 (100.0)
Rake 2020 ⁴⁵	England	Healthy older adults	72.2 (4.9) [65.0 – 84.0]	408:379	CLA (Cobas 6000 Roche)	50.2 (27.1)	127/787 (16.1)	109.2 (33.9)	395:392 (787)	2.5 mg bolus monthly	Placebo	2 yrs	URI/LRI: GP recorded	Secondary	787/787 (100.0)
Golan-Tripto, unpublished ⁴⁶	Israel	Prematurely born infants	0 (0)	21:29	CLA (DiaSorin)	33.6 (29.7)	19/46 (41.3)	20 µg arm: 78.0 (75.0) 10 µg arm: 81.0 (73.0)	25:25 (50)	20 µg daily	10 µg daily	1 yr	ARI: GP recorded	Secondary	25/50 (50.0)
Reyes, unpublished ⁴⁷	Chile	Healthy pre-school children	2.2 (0.5) [1.3 – 3.3]	168:135	LC-MS/MS	62.2 (15.5)	1/194 (0.5)	0.14 mg arm: 82.4 (24.5) 0.28 mg arm: 104.6 (52.9)	99:103:101 (303)	0.14 mg / 0.28 mg weekly	Placebo	6 mo	ARI: self-reported	Primary	194/303 (64.0)

[a] Sex missing for two participants randomised to intervention arm and subsequently excluded from analysis due to lack of outcome data. [b] Sex missing for one participant. [c] equivalent to 30 ug vitamin D₃.⁶⁰ 1 µg vitamin D₃ = 40 international units (IU); 25(OH)D concentrations reported in ng/ml were converted to nmol/L by multiplying by 2.496. [d] from subset of participants randomised to intervention; for comparison, mean 25(OH)D at follow-up in subset of participants randomised to placebo was 77.5 nmol/L (sd 25.2 nmol/L); 25(OH)D, 25-hydroxyvitamin D; RIDT, rapid influenza diagnostic test; COPD, chronic obstructive pulmonary disease; D₃, vitamin D₃ (cholecalciferol); p.o., *per os* (orally); mo, month; yr, year; wk, week. ARI, acute respiratory infection; CAP, College of American Pathologists, CLA, chemiluminescent assay; DEQAS, Vitamin D External Quality Assessment Scheme; EIA, enzyme immunoassay; EQA, external quality assessment; GP, general practitioner; LC-MS/MS, liquid chromatography tandem-mass spectrometry, RIA, radio-immunoassay; URI, upper respiratory infection; LRI, lower respiratory infection; ILI, influenza-like illness; RIQAS, Randox International Quality Assessment Scheme; VDSP, Vitamin D Standardisation Program of the Office of Dietary Supplements, National Institutes of Health, USA

Table 2: Placebo controlled RCTs: Proportion of participants experiencing at least one acute respiratory infection, overall and stratified by potential effect-modifiers

Potential effect-modifier	No of trials	Proportion with ≥ 1 ARI, intervention group (%)	Proportion with ≥ 1 ARI, control group (%)	Odds ratio (95% CI)	I ² %	P for heterogeneity
Overall	36	13685/22288 (61.4)	13565/21721 (62.5)	0.91 (0.84 to 0.99)	37.2	0.01
Baseline 25(OH)D, nmol/L^[a]						
<25	19	1348/1798 (75.0)	1388/1819 (76.3)	0.78 (0.53 to 1.16)	47.2	0.01
25 – 49.9	28	3439/4666 (73.7)	3347/4501 (74.4)	1.03 (0.92 to 1.15)	0.4	0.46
50 – 74.9	29	1679/2839 (59.1)	1565/2578 (60.7)	0.90 (0.76 to 1.06)	11.2	0.29
≥ 75	25	945/1543 (61.2)	908/1471 (61.7)	0.97 (0.81 to 1.16)	0.0	0.78
Dosing frequency						
Daily	18	1056/2134 (49.5)	1020/1871 (54.5)	0.75 (0.61 to 0.93)	52.5	0.005
Weekly	6	4482/6421 (69.8)	4447/6335 (70.2)	0.97 (0.88 to 1.06)	0.0	0.48
Monthly or less frequently	12	8147/13733 (59.3)	8098/13515 (59.9)	0.98 (0.93 to 1.03)	0.0	0.57
Daily dose equivalent, IU^[b]						
<400	2	482/1175 (41.0)	511/1133 (45.1)	0.65 (0.31 to 1.37)	86.3	0.007
400-1000	10	656/1236 (53.1)	627/1069 (58.7)	0.70 (0.55 to 0.89)	31.2	0.16
1001-2000	15	9946/15885 (62.6)	10022/15817 (63.4)	0.97 (0.92 to 1.03)	1.0	0.44
>2000	7	2291/3462 (66.2)	2250/3444 (65.3)	1.05 (0.84 to 1.31)	37.1	0.15
Trial duration, months						
≤ 12	29	1977/4887 (40.5)	1866/4368 (42.7)	0.82 (0.72 to 0.93)	38.1	0.02
>12	7	11708/17401 (67.3)	11699/17353 (67.4)	0.99 (0.95 to 1.04)	0.0	0.91
Age, yrs^[a]						
<1	5	875/2901 (30.2)	839/2796 (30.0)	0.95 (0.82 to 1.10)	18.7	0.30
1-15.9	15	4297/5994 (71.7)	4303/5877 (73.2)	0.71 (0.57 to 0.90)	46.0	0.03
16-64.9	21	3137/4876 (64.3)	3087/4727 (65.3)	0.97 (0.93 to 1.09)	11.5	0.31
≥ 65	16	5376/8589 (62.6)	5352/8394 (63.4)	0.96 (0.90 to 1.02)	0.0	0.67
Airway disease						
Asthma only	4	203/404 (50.2)	202/391 (51.7)	0.73 (0.36 to 1.49)	71.7	0.01
COPD only	2	106/208 (51.0)	104/207 (50.2)	1.01 (0.68 to 1.51)	0.0	0.71
Unrestricted	30	13376/21676 (61.7)	13259/21123 (62.8)	0.91 (0.84 to 0.99)	35.0	0.03

[a] The number of trials in each category for this variable adds up to more than 36, since this is a participant-level variable, i.e. some trials contributed data from participants who fell into more than one category

[b] Data from two trials that included higher-dose, lower-dose and placebo arms^{18,47} are excluded from this sub-group analysis, since the higher-dose and lower-dose arms spanned the 1,000 IU/day cut-off, rendering them unclassifiable

Table 3: Placebo-controlled studies: Secondary outcomes

Variables	No of trials	Proportion with ≥1 event, intervention group (%)	Proportion with ≥1 event, control group (%)	Odds ratio (95% CI)	I ² %	P for heterogeneity
Efficacy outcomes						
Upper respiratory infection*	28	7931/13493 (58.8)	7823/13034 (60.0)	0.96 (0.90 to 1.02)	4.4	0.40
Lower respiratory infection*	14	3583/12167 (29.4)	3601/12027 (29.9)	0.98 (0.93 to 1.04)	0.0	0.56
Emergency department attendance and/or hospital admission due to ARI	18	117/9887 (1.2)	125/9769 (1.3)	0.90 (0.70 to 1.16)	0.0	1.00
Death due to ARI or respiratory failure	33	14/13612 (0.1)	11/13058 (0.1)	1.04 (0.61 to 1.78)	0.0	1.00
Use of antibiotics to treat an ARI*	13	2035/7562 (26.9)	2082/7423 (28.0)	0.91 (0.82 to 1.02)	13.4	0.31
Absence from work or school due to ARI	10	378/1527 (24.7)	364/1044 (34.9)	0.91 (0.69 to 1.20)	35.3	0.13
Safety outcomes						
Serious adverse event of any cause*	35	545/13861 (3.9)	554/13326 (4.2)	0.97 (0.86 to 1.09)	0.0	1.00
Death due to any cause	34	117/13854 (0.8)	97/13293 (0.7)	1.15 (0.90 to 1.49)	0.0	1.00
Hypercalcaemia	21	50/9294 (0.5)	39/8919 (0.4)	1.20 (0.81 to 1.79)	0.0	1.00
Renal stones	20	110/11540 (1.0)	128/11138 (1.1)	0.85 (0.66 to 1.10)	0.0	1.00

* This analysis includes a subset of participants in the trial by Pham et al, who completed symptom diaries.

Figure 1: Flow chart of study selection





Figure 2: Forest plot of placebo-controlled RCTs reporting proportion of participants experiencing 1 or more acute respiratory infection.



NOTE: Weights are from random effects analysis

*This analysis includes data from the subset of ViDiFlu trial participants who were randomised to vitamin D vs. placebo control. **For this trial, participants were asked to report the occurrence of ARI during the one month prior to completing each annual survey (max surveys=5). The numerator is the number of participants who reported an ARI on at least one survey. The ARI outcomes for participants who completed fewer than 5 surveys and who did not report an ARI (N=2239; 14%) were estimated based on the % affected among those who completed all 5 surveys (N=12152; 76%).

References

1. Lanham-New SA, Webb AR, Cashman KD, et al. Vitamin D and SARS-CoV-2 virus/COVID-19 disease. *BMJ Nutrition, Prevention & Health* 2020.
2. Greiller CL, Martineau AR. Modulation of the Immune Response to Respiratory Viruses by Vitamin D. *Nutrients* 2015; **7**(6): 4240-70.
3. Jolliffe DA, Griffiths CJ, Martineau AR. Vitamin D in the prevention of acute respiratory infection: systematic review of clinical studies. *J Steroid Biochem Mol Biol* 2013; **136**: 321-9.
4. Pham H, Rahman A, Majidi A, Waterhouse M, Neale RE. Acute Respiratory Tract Infection and 25-Hydroxyvitamin D Concentration: A Systematic Review and Meta-Analysis. *International Journal of Environmental Research and Public Health* 2019; **16**(17).
5. Li-Ng M, Aloia JF, Pollack S, et al. A randomized controlled trial of vitamin D3 supplementation for the prevention of symptomatic upper respiratory tract infections. *Epidemiol Infect* 2009; **137**(10): 1396-404.
6. Urashima M, Segawa T, Okazaki M, Kurihara M, Wada Y, Ida H. Randomized trial of vitamin D supplementation to prevent seasonal influenza A in schoolchildren. *Am J Clin Nutr* 2010; **91**(5): 1255-60.
7. Manaseki-Holland S, Qader G, Isaq Masher M, et al. Effects of vitamin D supplementation to children diagnosed with pneumonia in Kabul: a randomised controlled trial. *Trop Med Int Health* 2010; **15**(10): 1148-55.
8. Laaksi I, Ruohola JP, Mattila V, Auvinen A, Ylikomi T, Pihlajamaki H. Vitamin D supplementation for the prevention of acute respiratory tract infection: a randomized, double-blinded trial among young Finnish men. *J Infect Dis* 2010; **202**(5): 809-14.
9. Majak P, Olszowiec-Chlebna M, Smejda K, Stelmach I. Vitamin D supplementation in children may prevent asthma exacerbation triggered by acute respiratory infection. *J Allergy Clin Immunol* 2011; **127**(5): 1294-6.
10. Trilok Kumar G, Sachdev HS, Chellani H, et al. Effect of weekly vitamin D supplements on mortality, morbidity, and growth of low birthweight term infants in India up to age 6 months: randomised controlled trial. *BMJ* 2011; **342**: d2975.
11. Lehouck A, Mathieu C, Carremans C, et al. High doses of vitamin D to reduce exacerbations in chronic obstructive pulmonary disease: a randomized trial. *Ann Intern Med* 2012; **156**(2): 105-14.
12. Manaseki-Holland S, Maroof Z, Bruce J, et al. Effect on the incidence of pneumonia of vitamin D supplementation by quarterly bolus dose to infants in Kabul: a randomised controlled superiority trial. *Lancet* 2012; **379**(9824): 1419-27.
13. Camargo CA, Jr., Ganmaa D, Frazier AL, et al. Randomized trial of vitamin D supplementation and risk of acute respiratory infection in Mongolia. *Pediatrics* 2012; **130**(3): e561-7.
14. Murdoch DR, Slow S, Chambers ST, et al. Effect of vitamin D3 supplementation on upper respiratory tract infections in healthy adults: the VIDARIS randomized controlled trial. *JAMA* 2012; **308**(13): 1333-9.
15. Bergman P, Norlin AC, Hansen S, et al. Vitamin D3 supplementation in patients with frequent respiratory tract infections: a randomised and double-blind intervention study. *BMJ Open* 2012; **2**(6): e001663.
16. Marchisio P, Consonni D, Baggi E, et al. Vitamin D supplementation reduces the risk of acute otitis media in otitis-prone children. *Pediatr Infect Dis J* 2013; **32**(10): 1055-60.
17. Rees JR, Hendricks K, Barry EL, et al. Vitamin D3 Supplementation and Upper Respiratory Tract Infections in a Randomized, Controlled Trial. *Clin Infect Dis* 2013.
18. Tran B, Armstrong BK, Ebeling PR, et al. Effect of vitamin D supplementation on antibiotic use: a randomized controlled trial. *Am J Clin Nutr* 2014; **99**(1): 156-61.
19. Goodall EC, Granados AC, Luinstra K, et al. Vitamin D3 and gargling for the prevention of upper respiratory tract infections: a randomized controlled trial. *BMC Infectious Diseases* 2014; **14**: 273.

20. Urashima M, Mezawa H, Noya M, Camargo CA, Jr. Effects of vitamin D supplements on influenza A illness during the 2009 H1N1 pandemic: a randomized controlled trial. *Food & Function* 2014; **5**(9): 2365-70.
21. Grant CC, Kaur S, Waymouth E, et al. Reduced primary care respiratory infection visits following pregnancy and infancy vitamin D supplementation: a randomised controlled trial. *Acta Paediatr* 2014.
22. Martineau AR, James WY, Hooper RL, et al. Vitamin D3 supplementation in patients with chronic obstructive pulmonary disease (ViDiCO): a multicentre, double-blind, randomised controlled trial. *The Lancet Respiratory Medicine* 2015; **3**(2): 120-30.
23. Martineau AR, MacLaughlin BD, Hooper RL, et al. Double-blind randomised placebo-controlled trial of bolus-dose vitamin D3 supplementation in adults with asthma (ViDiAs). *Thorax* 2015; **70**(5): 451-7.
24. Martineau AR, Hanifa Y, Witt KD, et al. Double-blind randomised controlled trial of vitamin D3 supplementation for the prevention of acute respiratory infection in older adults and their carers (ViDiFlu). *Thorax* 2015; **70**(10): 953-60.
25. Simpson SJ, van der Mei I, Stewart N, Blizzard L, Tettey P, Taylor B. Weekly cholecalciferol supplementation results in significant reductions in infection risk among the vitamin D deficient: results from the CIPRIS pilot RCT. *BMC Nutrition* 2015; **1**(7).
26. Dubnov-Raz G, Rinat B, Hemila H, Choleva L, Cohen AH, Constantini NW. Vitamin D supplementation and upper respiratory tract infections in adolescent swimmers: a randomized controlled trial. *Pediatric Exercise Science* 2015; **27**(1): 113-9.
27. Denlinger LC, King TS, Cardet JC, et al. Vitamin D Supplementation and the Risk of Colds in Patients with Asthma. *Am J Respir Crit Care Med* 2016; **193**(6): 634-41.
28. Tachimoto H, Mezawa H, Segawa T, Akiyama N, Ida H, Urashima M. Improved Control of Childhood Asthma with Low-Dose, Short-Term Vitamin D Supplementation: A Randomized, Double-Blind, Placebo-Controlled Trial. *Allergy* 2016.
29. Ginde AA, Blatchford P, Breese K, et al. High-Dose Monthly Vitamin D for Prevention of Acute Respiratory Infection in Older Long-Term Care Residents: A Randomized Clinical Trial. *J Am Geriatr Soc* 2017; **65**(3): 496-503.
30. Martineau AR, Jolliffe DA, Hooper RL, et al. Vitamin D supplementation to prevent acute respiratory tract infections: systematic review and meta-analysis of individual participant data. *BMJ* 2017; **356**: i6583.
31. Gupta P, Dewan P, Shah D, et al. Vitamin D Supplementation for Treatment and Prevention of Pneumonia in Under-five Children: A Randomized Double-blind Placebo Controlled Trial. *Indian Pediatr* 2016; **53**(11): 967-76.
32. Aglipay M, Birken CS, Parkin PC, et al. Effect of High-Dose vs Standard-Dose Wintertime Vitamin D Supplementation on Viral Upper Respiratory Tract Infections in Young Healthy Children. *JAMA* 2017; **318**(3): 245-54.
33. Arihiro S, Nakashima A, Matsuoka M, et al. Randomized Trial of Vitamin D Supplementation to Prevent Seasonal Influenza and Upper Respiratory Infection in Patients With Inflammatory Bowel Disease. *Inflamm Bowel Dis* 2019; **25**(6): 1088-95.
34. Hibbs AM, Ross K, Kerns LA, et al. Effect of Vitamin D Supplementation on Recurrent Wheezing in Black Infants Who Were Born Preterm: The D-Wheeze Randomized Clinical Trial. *JAMA* 2018; **319**(20): 2086-94.
35. Lee MT, Kattan M, Fennoy I, et al. Randomized phase 2 trial of monthly vitamin D to prevent respiratory complications in children with sickle cell disease. *Blood Adv* 2018; **2**(9): 969-78.
36. Loeb M, Dang AD, Thiem VD, et al. Effect of Vitamin D supplementation to reduce respiratory infections in children and adolescents in Vietnam: A randomized controlled trial. *Influenza Other Respir Viruses* 2019; **13**(2): 176-83.
37. Rosendahl J, Valkama S, Holmlund-Suila E, et al. Effect of Higher vs Standard Dosage of Vitamin D3 Supplementation on Bone Strength and Infection in Healthy Infants: A Randomized Clinical Trial. *JAMA Pediatrics* 2018; **172**(7): 646-54.
38. Shimizu Y, Ito Y, Yui K, Egawa K, Orimo H. Intake of 25-Hydroxyvitamin D3 Reduces Duration and Severity of Upper Respiratory Tract Infection: A Randomized, Double-Blind,

Placebo-Controlled, Parallel Group Comparison Study. *J Nutr Health Aging* 2018; **22**(4): 491-500.

39. Aloia JF, Islam S, Mikhail M. Vitamin D and Acute Respiratory Infections-The PODA Trial. *Open Forum Infectious Diseases* 2019; **6**(9): ofz228.

40. Hauger H, Ritz C, Mortensen C, et al. Winter cholecalciferol supplementation at 55 degrees N has little effect on markers of innate immune defense in healthy children aged 4-8 years: a secondary analysis from a randomized controlled trial. *European Journal of Nutrition* 2019; **58**(4): 1453-62.

41. Camargo CA, Sluyter J, Stewart AW, et al. Effect of monthly high-dose vitamin D supplementation on acute respiratory infections in older adults: A randomized controlled trial. *Clin Infect Dis* 2020; **71**: 311-7.

42. Ganmaa D, Uyanga B, Zhou X, et al. Vitamin D Supplements and Prevention of Tuberculosis Infection and Disease. *N Engl J Med* 2020; **383**: 359-68.

43. Mandlik R, Mughal Z, Khadilkar A, et al. Occurrence of infections in schoolchildren subsequent to supplementation with vitamin D-calcium or zinc: a randomized, double-blind, placebo-controlled trial. *Nutr Res Pract* 2020; **14**(2): 117-26.

44. Pham H, Waterhouse M, Baxter C, et al. The effect of vitamin D supplementation on acute respiratory tract infection in older Australian adults: an analysis of data from the D-Health Trial: a randomised controlled trial. *The Lancet Diabetes & Endocrinology* 2020; **in press**.

45. Rake C, Gilham C, Bukasa L, et al. High-dose oral vitamin D supplementation and mortality in people aged 65-84 years: the VIDAL cluster feasibility RCT of open versus double-blind individual randomisation. *Health Technol Assess* 2020; **24**(10): 1-54.

46. Golan-Tripto I. The Effect of Vitamin D Administration to Premature Infants on Vitamin D Status and Respiratory Morbidity (NCT02404623). <https://clinicaltrials.gov/ct2/show/NCT02404623>.

47. Reyes ML, Vizcaya C, Borzutzky A, et al. Study of Vitamin D for the Prevention of Acute Respiratory Infections in Children (NCT02046577). <https://clinicaltrials.gov/ct2/show/NCT02046577>.

48. Ducharme FM, Jensen M, Mailhot G, et al. Impact of two oral doses of 100,000 IU of vitamin D3 in preschoolers with viral-induced asthma: a pilot randomised controlled trial. *Trials* 2019; **20**(1): 138.

49. Moreno A. Effect of Supplementation With Vitamin D on the Acute Bronchitis Prevention During the First Year of Life (NCT01875757). <https://clinicaltrials.gov/ct2/show/NCT01875757>.

50. Camargo CA. Effects of Vitamin D and Omega-3 Fatty Acids on Infectious Diseases and hCAP18 (VITAL Infection; NCT01758081).

51. Higgins JP, Altman DG, Gotzsche PC, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011; **343**: d5928.

52. Higgins JPT TJ, Chandler J, Cumpston M, Li T, Page MJ, Welch VA (editors). Cochrane Handbook for Systematic Reviews of Interventions version 6.0 (updated July 2019). *Cochrane* 2019.

53. Harris R, Bradburn M, Deeks J, et al. METAN: Stata module for fixed and random effects meta-analysis, revised 23 Sep 2010. Statistical Software Components S456798. Boston College Department of Economics; 2006.

54. Pearce SH, Cheetham TD. Diagnosis and management of vitamin D deficiency. *BMJ* 2010; **340**: b5664.

55. Peters JL, Sutton AJ, Jones DR, Abrams KR, Rushton L. Contour-enhanced meta-analysis funnel plots help distinguish publication bias from other causes of asymmetry. *J Clin Epidemiol* 2008; **61**(10): 991-6.

56. Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008; **336**(7650): 924-6.

57. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997; **315**(7109): 629-34.

58. Sterne JA, Sutton AJ, Ioannidis JP, et al. Recommendations for examining and interpreting funnel plot asymmetry in meta-analyses of randomised controlled trials. *BMJ* 2011; **343**: d4002.

59. Kantor ED, Rehm CD, Du M, White E, Giovannucci EL. Trends in Dietary Supplement Use Among US Adults From 1999-2012. *JAMA* 2016; **316**(14): 1464-74.
60. Jetter A, Egli A, Dawson-Hughes B, et al. Pharmacokinetics of oral vitamin D(3) and calcifediol. *Bone* 2014; **59**: 14-9.

Vitamin D supplementation to prevent acute respiratory infections: systematic review and meta-analysis of data from randomised controlled trials

Supplementary Material

Methods

Search Strategies

A. Medline

Cochrane Highly Sensitive Search Strategy for identifying randomised controlled trials

- #1. randomized controlled trial [pt] OR controlled clinical trial [pt] OR randomized [tiab] OR placebo [tiab] OR drug therapy [sh] OR randomly [tiab] OR trial [tiab] OR groups [tiab]
- #2. animals [mh] NOT humans [mh]
- #3. #1 NOT #2

Terms specific to vitamin D

- #4. Vitamin D OR vitamin D2 OR vitamin D3 OR cholecalciferol OR ergocalciferol OR alphacalcidol OR alfacalcidol OR calcitriol OR paricalcitol OR doxerocalciferol

Terms specific to acute respiratory infection

- #5. Acute Respiratory Infection OR Upper Respiratory Infection OR Lower Respiratory Infection OR Respiratory Tract Infection OR Common Cold OR Sinusitis OR Pharyngitis OR Laryngitis OR Laryngotracheobronchitis OR Tonsillitis OR peritonsillar abscess OR Croup OR Epiglottitis OR supraglottitis OR Otitis Media OR Pneumonia OR Bronchopneumonia OR Bronchitis OR Pleurisy OR Pleuritis

Combination of terms to identify randomised controlled trials of vitamin D for the prevention of acute respiratory infection

- #3 AND #4 AND #5

B. EMBASE

Terms for identifying randomised controlled trials

#1 'randomized controlled trial'/exp OR 'single blind procedure'/exp OR 'double blind procedure'/exp OR 'crossover procedure'/exp

#2 random*:ab,ti OR placebo*:ab,ti OR crossover*:ab,ti OR 'cross over':ab,ti OR allocat*:ab,ti OR ((singl* OR doubl*) NEXT/1 blind*):ab,ti OR trial:ti

#3. #1 OR #2

Terms specific to vitamin D

#4. vitamin AND d OR vitamin AND d2 OR vitamin AND d3 OR cholecalciferol OR ergocalciferol OR alphacalcidol OR alfacalcidol OR calcitriol OR paricalcitol OR doxerocalciferol

Terms specific to acute respiratory infection

#5. acute AND respiratory AND infection OR upper AND respiratory AND infection OR lower AND respiratory AND infection OR respiratory AND tract AND infection OR common AND cold OR sinusitis OR pharyngitis OR laryngitis OR laryngotracheobronchitis OR tonsillitis OR peritonsillar AND abscess OR croup OR epiglottitis OR supraglottitis OR otitis AND media OR pneumonia OR bronchopneumonia OR bronchitis OR pleurisy OR pleuritis

Combination of terms to identify randomised controlled trials of vitamin D for the prevention of acute respiratory infection

#3 AND #4 AND #5

C. Cochrane Central

Terms specific to vitamin D

#1. Vitamin D OR vitamin D2 OR vitamin D3 OR cholecalciferol OR ergocalciferol OR alphacalcidol OR alfacalcidol OR calcitriol OR paricalcitol OR doxerocalciferol

Terms specific to acute respiratory infection

#2. Acute Respiratory Infection OR Upper Respiratory Infection OR Lower Respiratory Infection OR Respiratory Tract Infection OR Common Cold OR Sinusitis OR Pharyngitis OR Laryngitis OR Laryngotracheobronchitis OR Tonsillitis OR peritonsillar abscess OR Croup OR Epiglottitis OR supraglottitis OR Otitis Media OR Pneumonia OR Bronchopneumonia OR Bronchitis OR Pleurisy OR Pleuritis

Combination of terms to identify randomised controlled trials of vitamin D for the prevention of acute respiratory infection

#1 AND #2

D. Web of Science

TS =(Vitamin D OR vitamin D2 OR vitamin D3 OR cholecalciferol OR ergocalciferol OR alphacalcidol OR alfacalcidol OR calcitriol OR paricalcitol OR doxerocalciferol) AND TS =(Acute Respiratory Infection OR Upper Respiratory Infection OR Lower Respiratory Infection OR Respiratory Tract Infection OR Common Cold OR Sinusitis OR Pharyngitis OR Laryngitis OR Laryngotracheobronchitis OR Tonsillitis OR peritonsillar abscess OR Croup OR Epiglottitis OR supraglottitis OR Otitis Media OR Pneumonia OR Bronchopneumonia OR Bronchitis OR Pleurisy OR Pleuritis) AND TS =(placebo* or random* or clinical trial* or double blind* or single blind* or rct)

E. ClinicalTrials.gov

Vitamin D AND respiratory AND infection

Sources of support for participating trials

The trial by Aglipay and colleagues was supported by the competitive grants from the Canadian Institutes of Health Research Institutes of Human Development, Child and Youth Health and Nutrition, Metabolism and Diabetes (grant number MOP-114945) and the Thrasher Research Fund (award number 9113).

The trial by Aloia and colleagues was supported by the National Institute of Aging (grant number R01-AG032440-01A2).

The trial by Arihiro and colleagues was supported by the Ministry of Education, Culture, Sports, Science, and Technology in the Japan-Supported Program for the Strategic Research Foundation at Private Universities and funding from the Department of Gastroenterology and Hepatology, Jikei University of Medicine, Tokyo, Japan.

The trial by Bergman et al was supported by grants from the Swedish Research Council, the Strategic Research Foundation (SSF), the Swedish Heart and Lung foundation, Karolinska Institutet, Stockholm County Council and the Swedish Cancer Society as well as by the Magnus Bergwall and Åke Wiberg foundations.

The trials by Camargo and colleagues were supported by a grant for the Blue Sky Study from an anonymous foundation and the Massachusetts General Hospital; and the Health Research Council of New Zealand (grant number 10/400) and the Accident Compensation Corporation of New Zealand.

The trial by Ganmaa and colleagues was supported by the National Institutes of Health (Grant Number 1R01HL122624-01).

The trial by Ginde and colleagues was supported by NIH/NIA grant K23AG040708, NIH/NCATS Colorado CTSA Grant UL1TR001082, and the American Geriatrics Society Jahnigen Career Development Scholars Award.

The trial by Golan-Tripto and colleagues was supported by the Soroka JNF UK Clinical Research Scholar Program.

The trial by Goodall and colleagues was supported in part by the Canadian Institutes of Health Research [OPP 86940] and with in-kind support from Copan Italia, Brescia Italy.

The trial by Grant and colleagues was supported by the Health Research Council of New Zealand, Grant Number 09/215R.

The trial by Gupta and colleagues was supported by the Indian Council of Medical Research, New Delhi.

The trial by Hauger and colleagues was supported by Lundbeckfonden (grant number R180-2014-3481), by Brødrene Hartmann's Fund (A26842), and by the European

Commission (FP7/2007–2013) under Grant Agreement 613977 for the ODIN Integrated Project (Food-based solutions for optimal vitamin D nutrition and health through the life cycle).

The trial by Hibbs and colleagues was supported by the National Heart, Lung, and Blood Institute and Office of Dietary Supplements (grant number R01HL109293).

The trial by Lee and colleagues was supported by the US Food and Drug Administration Orphan Product Development (grant number R01FD003894).

The trial by Loeb and colleagues was supported by the Institute for Infectious Diseases Research at McMaster University.

The trials by Manaseki-Holland and colleagues were supported by New Zealand Aid (ref: GRA/470/2) and The Wellcome Trust (ref: 082476).

The trial by Mandlik and colleagues was supported by a core grant from the Hirabai Cowasji Jehangir Medical Research Institute.

The trials by Martineau and colleagues were supported by the National Institute for Health Research under its Programme Grants for Applied Research Programme (Reference Number RP-PG-0407-10398).

The trial by Murdoch and colleagues was supported by the Health Research Council of New Zealand, grant number 09/302.

The trial by Pham and colleagues was supported by project grants from the National Health and Medical Research Council (grant numbers GNT1046681 and GNT1120682).

The trial by Reyes and colleagues was supported by Fondo Nacional de Investigacion y Desarrollo en Salud (ref. SA13I20173).

The trial by Rosendahl and colleagues was supported by the Foundation for Pediatric Research, the Finnish Medical Foundation, Governmental Subsidy for Clinical Research, the Päivikki and Sakari Sohlberg Foundation, the Academy of Finland, the Sigrid Jusélius Foundation, the Folkhälsan Research Foundation, the Novo Nordisk Foundation, the Orion Research Foundation, and Barncancerfonden.

The trial by Rake and colleagues was supported by the National Institute for Health Research Health Technology Assessment programme (ref. HTA 08/116/48).

The trial by Rees and colleagues was supported by the National Cancer Institute at the National Institutes of Health (grant numbers CA098286 and CA098286-S).

The trial by Shimizu and colleagues was supported by the FANCL Corporation.

The trial by Simpson and colleagues was supported by the Royal Hobart Hospital Research Foundation.

The trial by Tachimoto and colleagues was supported by the Ministry of Education, Culture, Sports, Science, and Technology in the Japan-Supported Program for the Strategic Research Foundation at Private Universities, JSPH KAKENHI Grant Number 23591553, and funding from Jikei University of Medicine, Tokyo, Japan.

The trial by Tran and colleagues was supported by the National Health and Medical Research Council of Australia, grant 613655.

The trial by Trilok Kumar and colleagues was supported by the Department of Biotechnology, Government of India (ref BT/PR-PR7489/PID/20/285/2006), Nutrition Third World and Sight and Life.

Table S1: Reasons for exclusion of potentially relevant studies

First author, year	Reason for exclusion
Somnath, 2017 ¹	Ineligible: open-label trial
Jung, 2018 ²	Ineligible: ARI outcome not pre-specified
Ramos-Martínez, 2018 ³	Ineligible: intervention was administration of 1,25-dihydroxyvitamin D
Zhou, 2018 ⁴	Ineligible: open-label trial
Hueniken, 2019 ⁵	Ineligible: same trial as Aglipay et al ⁶
Singh, 2019 ⁷	Ineligible: open-label trial

Table S2: Risk of Bias Assessment

	Sequence generation	Allocation concealment	Blinding of participants and personnel	Blinding of outcome assessment	Incomplete outcome data	Selective reporting	Other bias
Li-Ng 2009 ⁸	✓	✓	✓	✓	✓	✓	✓
Urashima 2010 ⁹	✓	✓	✓	✓	✓	✓	✓
Manaseki-Holland 2010 ¹⁰	✓	✓	✓	✓	✓	✓	✓
Laaksi 2010 ¹¹	✓	✓	✓	✓	?	✓	✓
Majak 2011 ¹²	✓	✓	✓	✓	✓	✓	✓
Trilok-Kumar 2011 ¹³	✓	✓	✓	✓	✓	✓	✓
Lehouck 2012 ¹⁴	✓	✓	✓	✓	✓	✓	✓
Manaseki-Holland 2012 ¹⁵	✓	✓	✓	✓	✓	✓	✓
Camargo 2012 ¹⁶	✓	✓	✓	✓	✓	✓	✓
Murdoch 2012 ¹⁷	✓	✓	✓	✓	✓	✓	✓
Bergman 2012 ¹⁸	✓	✓	✓	✓	✓	✓	✓
Marchisio 2013 ¹⁹	✓	✓	✓	✓	✓	✓	✓
Rees 2013 ²⁰	✓	✓	✓	✓	✓	✓	✓
Tran 2014 ²¹	✓	✓	✓	✓	✓	✓	✓
Goodall 2014 ²²	✓	✓	✓	✓	✓	✓	✓
Urashima 2014 ²³	✓	✓	✓	✓	✓	✓	✓
Grant 2014 ²⁴	✓	✓	✓	✓	✓	✓	✓
Martineau 2015a ²⁵ [ViDiCO]	✓	✓	✓	✓	✓	✓	✓
Martineau 2015b ²⁶ [ViDiAs]	✓	✓	✓	✓	✓	✓	✓
Martineau 2015c ²⁷ [ViDiFlu]	✓	✓	✓	✓	✓	✓	✓
Simpson 2015 ²⁸	✓	✓	✓	✓	✓	✓	✓
Dubnov-Raz 2015 ²⁹	✓	✓	✓	✓	?	✓	✓
Denlinger 2016 ³⁰	✓	✓	✓	✓	✓	✓	✓
Tachimoto 2016 ³¹	✓	✓	✓	✓	✓	✓	✓
Ginde 2016 ³²	✓	✓	✓	✓	✓	✓	✓
Gupta 2016 ³³	✓	✓	✓	✓	✓	✓	✓
Aglipay 2017 ⁶	✓	✓	✓	✓	✓	✓	✓
Arihiro 2018 ³⁴	✓	✓	✓	✓	✓	✓	✓
Hibbs 2018 ³⁵	✓	✓	✓	✓	✓	✓	✓
Lee 2018 ³⁶	✓	✓	✓	✓	✓	✓	✓
Loeb 2018 ³⁷	✓	✓	✓	✓	✓	✓	✓
Rosendahl 2018 ³⁸	✓	✓	✓	✓	✓	✓	✓
Shimizu 2018 ³⁹	✓	✓	✓	✓	✓	✓	✓
Aloia 2019 ⁴⁰	✓	✓	✓	✓	✓	✓	✓
Hauger 2019 ⁴¹	✓	✓	✓	✓	✓	✓	✓
Camargo 2020 ⁴²	✓	✓	✓	✓	✓	✓	✓
Ganmaa 2020 ⁴³	✓	✓	✓	✓	✓	✓	✓
Mandlik 2020 ⁴⁴	✓	✓	✓	✓	✓	✓	✓
Pham 2020 ⁴⁵	✓	✓	✓	✓	✓	N/A	✓
Rake 2020 ⁴⁶	✓	✓	✓	✓	✓	✓	✓
Golan-Tripto (unpublished) ⁴⁷	✓	✓	✓	✓	?	N/A	✓
Reyes (unpublished) ⁴⁸	✓	✓	✓	✓	?	N/A	✓

✓ = low risk of bias; ? = unclear risk of bias N/A = not applicable (unpublished)

Table S3: Multivariate meta-regression model for proportion of participants experiencing at least one acute respiratory tract infection, by trial-level subgroups.

Variables	No of trials ^[a]	Proportion with ≥1 ARI, intervention group (%)	Proportion with ≥1 ARI, control group (%)	Odds ratio (95% CI) ^[b]	I ² %	P value for heterogeneity	Adjusted odds ratio (95% CI) ^[c]	P value for interaction ^[c]
Dosing frequency								
Daily	18	1056/2134 (49.5)	1020/1871 (54.5)	0.75 (0.61 to 0.93)	52.5	0.005	0.94 (0.74 to 1.18)	0.57
Not daily	16	12319/19624 (62.8)	12390/19592 (63.2)	0.98 (0.93 to 1.02)	0.0	0.56	Referent	
Daily dose equivalent, IU								
≤1000	12	1138/2411 (47.2)	1138/2202 (51.7)	0.70 (0.56 to 0.87)	48.6	0.03	0.82 (0.64 to 1.04)	0.09
>1000	22	12237/19347 (63.2)	12272/19261 (63.7)	0.99 (0.92 to 1.06)	16.0	0.25	Referent	
Trial duration, months								
≤12	27	1667/4357 (38.3)	1711/4110 (41.6)	0.81 (0.70 to 0.93)	41.9	0.01	0.89 (0.72 to 1.10)	0.28
>12	7	11708/17401 (67.3)	11699/17353 (67.4)	0.99 (0.95 to 1.04)	0.0	0.91	Referent	

[a] Data from two trials that included higher-dose, lower-dose and placebo arms^{21,48} are excluded, since the higher-dose and lower-dose arms in these studies spanned the 1,000 IU/day cut-off, rendering them unclassifiable for the purposes of this analysis. [b] Within-sub-group odds ratios from random effects model adjusting for study weights. [c] adjusted odds ratios and P values for interaction from multivariable meta-regression model including dichotomised variables for dose frequency, size and trial duration.

Table S4: Summary of Findings Table

Vitamin D₃ compared to placebo for prevention of acute respiratory infection (ARI)

Population: children and adults of any age, sex or ethnic origin, with or without co-morbidity

Setting: Eighteen countries on four continents (Asia, Australasia, Europe, North America)

Intervention: oral vitamin D₃ (cholecalciferol) supplementation

Comparison: oral placebo

Outcomes	Anticipated absolute effects* (95% CI)		Relative effect (95% CI)	No of participants (studies)	Quality of the evidence (GRADE)
	Risk with placebo	Risk with Vitamin D			
Proportion with at least one ARI, all participants	625 per 1,000	597 per 1,000 (574 to 620)	OR 0.91 (0.84 to 0.99)	44009 (36 RCTs)	⊕⊕⊕ MODERATE
Proportion with at least one ARI, participants in placebo-controlled trials with duration ≤12 months investigating daily dosing with 400-1,000 IU vitamin D ₃ /day	572 per 1,000	436 per 1,000 (375 to 500)	OR 0.58 (0.45 to 0.75)	1232 (8 RCTs)	⊕⊕⊕ MODERATE
Proportion with at least one hospital admission or emergency department attendance due to ARI	13 per 1,000	12 per 1,000 (9 to 15)	OR 0.90 (0.70 to 1.16)	19656 (18 RCTs)	⊕⊕⊕ MODERATE
Proportion with serious adverse event, any cause	42 per 1,000	40 per 1,000 (36 to 45)	OR 0.97 (0.86 to 1.09)	27187 (35 RCTs)	⊕⊕⊕ MODERATE
Proportion of deaths due to ARI or respiratory failure	1 per 1,000	1 per 1,000 (1 to 1)	OR 1.04 (0.61 to 1.78)	26670 (33 RCTs)	⊕⊕⊕ MODERATE

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

CI: Confidence interval; OR: Odds ratio

GRADE Working Group grades of evidence

High quality: We are very confident that the true effect lies close to that of the estimate of the effect

Moderate quality: We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different

Low quality: Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect

Very low quality: We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

Table S5: Results of exploratory sensitivity analyses excluding placebo-controlled trials at unclear risk of bias and studies investigating ARI incidence as a secondary outcome.

Reason for exclusion (number of excluded studies)	No. of studies included	Proportion with ≥ 1 ARI, intervention group (%)	Proportion with ≥ 1 ARI, control group (%)	Odds ratio (95% CI)	I ² %	P value for heterogeneity
Studies at unclear risk of bias (n=3)	33	13511/22061 (61.2)	13442/21565 (62.3)	0.93 (0.86 to 1.00)	36.5	0.02
Studies with ARI incidence as secondary outcome (n=18)	18	1155/3851 (30.0)	1107/3686 (30.0)	0.89 (0.77 to 1.03)	15.6	0.27
Substituting diary-defined ARI events (available for 2598 participants) for survey-defined ARI events (available for n=16,000 participants) in the trial by Pham et al ⁴⁵	36	8842/15606 (56.7)	8659/15001 (57.7)	0.90 (0.83 to 0.99)	37.1	0.02

Figure S1: Cates plot illustrating reduction in risk of one or more acute respiratory infections with vitamin D supplementation vs. placebo (primary comparison), **A)** Overall, and **B)** in trials with duration ≤ 12 months where vitamin D₃ was administered using daily doses of 400-1000 IU/day.


A



B




Figure S2: Forest plot of RCTs comparing effects of higher- vs. lower-dose vitamin D, reporting proportion of participants experiencing at least one acute respiratory infection.




*This analysis includes data from the subset of ViDiFlu trial participants who were randomised to higher- vs. lower-dose vitamin D.

Figure S4: Forest plot of RCTs comparing effects of vitamin D vs. placebo, reporting proportion of participants experiencing at least one acute respiratory infection, by frequency of supplementation (daily vs. weekly vs. monthly to 3-monthly)




*This analysis includes data from the subset of ViDiFlu trial participants who were randomised to vitamin D vs. placebo control. **For this trial, participants were asked to report the occurrence of ARTI during the one month prior to completing each annual survey (max surveys=5). The numerator is the number of people who reported an ARTI on at least one survey. The ARTI outcomes for people who completed fewer than 5 surveys and who did not report an ARTI (N=2239; 14%) were estimated based on the % affected among those who completed all 5 surveys (N=12,152; 76%).

Figure S5: Forest plot of RCTs comparing effects of vitamin D vs. placebo, reporting proportion of participants experiencing at least one acute respiratory infection, by daily dose equivalents (<400 IU/day vs. 400-1000 IU/day vs. 1001-2000 IU/day vs. >2000 IU/day).




*This analysis includes data from the subset of ViDiFlu trial participants who were randomised to vitamin D vs. placebo control. **For this trial, participants were asked to report the occurrence of ARTI during the one month prior to completing each annual survey (max surveys=5). The numerator is the number of people who reported an ARTI on at least one survey. The ARTI outcomes for people who completed fewer than 5 surveys and who did not report an ARTI (N=2239; 14%) were estimated based on the % affected among those who completed all 5 surveys (N=12,152; 76%).

Figure S6: Forest plot of RCTs comparing effects of vitamin D vs. placebo, reporting proportion of participants experiencing at least one acute respiratory infection, by trial duration (≤ 12 months vs. >12 months).




*This analysis includes data from the subset of ViDiFlu trial participants who were randomised to vitamin D vs. placebo control. **For this trial, participants were asked to report the occurrence of ARTI during the one month prior to completing each annual survey (max surveys=5). The numerator is the number of people who reported an ARTI on at least one survey. The ARTI outcomes for people who completed fewer than 5 surveys and who did not report an ARTI (N=2239; 14%) were estimated based on the % affected among those who completed all 5 surveys (N=12,152; 76%).

Figure S7: Forest plot of RCTs comparing effects of vitamin D vs. placebo, reporting proportion of participants experiencing at least one acute respiratory infection, by the following age strata: **A)** <1 year; **B)** 1-15.99 years; **C)** 16-64.99 years, and **D)** ≥65 years



*This analysis includes data from the subset of ViDiFlu trial participants who were randomised to vitamin D vs. placebo control. **For this trial, participants were asked to report the occurrence of ARTI during the one month prior to completing each annual survey (max surveys=5). The numerator is the number of people who reported an ARTI on at least one survey. The ARTI outcomes for people who completed fewer than 5 surveys and who did not report an ARTI (N=2239; 14%) were estimated based on the % affected among those who completed all 5 surveys (N=12,152; 76%).

Figure S8: Forest plot of RCTs comparing effects of vitamin D vs. placebo, reporting proportion of participants experiencing at least one acute respiratory infection, by presence or absence of airway disease comorbidity.



*This analysis includes data from the subset of ViDiFlu trial participants who were randomised to vitamin D vs. placebo control. **For this trial, participants were asked to report the occurrence of ARTI during the one month prior to completing each annual survey (max surveys=5). The numerator is the number of people who reported an ARTI on at least one survey. The ARTI outcomes for people who completed fewer than 5 surveys and who did not report an ARTI (N=2239; 14%) were estimated based on the % affected among those who completed all 5 surveys (N=12,152; 76%).

Figure S9 Forest plot of sub-set of RCTs with duration ≤ 12 months comparing effects of daily vitamin D at a dose of 400-1000 IU/day vs. placebo, reporting proportion of participants experiencing at least one acute respiratory infection





Figure S10: Funnel plot of placebo-controlled RCTs reporting proportion of participants experiencing 1 or more acute respiratory infection.



Egger's test for publication bias: P=0.008

References

1. Somnath SH, Biswal N, Chandrasekaran V, Jagadisan B, Bobby Z. Therapeutic effect of vitamin D in acute lower respiratory infection: A randomized controlled trial. *Clin Nutr ESPEN* 2017; **20**: 24-8.
2. Jung HC, Seo M-W, Lee S, Kim SW, Song JK. Vitamin D₃ Supplementation Reduces the Symptoms of Upper Respiratory Tract Infection during Winter Training in Vitamin D-Insufficient Taekwondo Athletes: A Randomized Controlled Trial. *Int J Environ Res Public Health* 2018; **15**(9): 2003.
3. Ramos-Martinez E, Lopez-Vancell MR, Fernandez de Cordova-Aguirre JC, et al. Reduction of respiratory infections in asthma patients supplemented with vitamin D is related to increased serum IL-10 and IFN γ levels and cathelicidin expression. *Cytokine* 2018; **108**: 239-46.
4. Zhou J, Du J, Huang L, Wang Y, Shi Y, Lin H. Preventive Effects of Vitamin D on Seasonal Influenza A in Infants: A Multicenter, Randomized, Open, Controlled Clinical Trial. *The Pediatric Infectious Disease Journal* 2018; **37**(8): 749-54.
5. Hueniken K, Aglipay M, Birken CS, et al. Effect of High-Dose Vitamin D Supplementation on Upper Respiratory Tract Infection Symptom Severity in Healthy Children. *Pediatr Infect Dis J* 2019; **38**(6): 564-8.
6. Aglipay M, Birken CS, Parkin PC, et al. Effect of High-Dose vs Standard-Dose Wintertime Vitamin D Supplementation on Viral Upper Respiratory Tract Infections in Young Healthy Children. *JAMA* 2017; **318**(3): 245-54.
7. Singh N, Kamble D, Mahantshetti NS. Effect of Vitamin D Supplementation in the Prevention of Recurrent Pneumonia in Under-Five Children. *Indian Journal of Pediatrics* 2019; **86**(12): 1105-11.
8. Li-Ng M, Aloia JF, Pollack S, et al. A randomized controlled trial of vitamin D₃ supplementation for the prevention of symptomatic upper respiratory tract infections. *Epidemiol Infect* 2009; **137**(10): 1396-404.
9. Urashima M, Segawa T, Okazaki M, Kurihara M, Wada Y, Ida H. Randomized trial of vitamin D supplementation to prevent seasonal influenza A in schoolchildren. *Am J Clin Nutr* 2010; **91**(5): 1255-60.
10. Manaseki-Holland S, Qader G, Isaq Masher M, et al. Effects of vitamin D supplementation to children diagnosed with pneumonia in Kabul: a randomised controlled trial. *Trop Med Int Health* 2010; **15**(10): 1148-55.
11. Laaksi I, Ruohola JP, Mattila V, Auvinen A, Ylikomi T, Pihlajamaki H. Vitamin D supplementation for the prevention of acute respiratory tract infection: a randomized, double-blinded trial among young Finnish men. *J Infect Dis* 2010; **202**(5): 809-14.
12. Majak P, Olszowiec-Chlebna M, Smejda K, Stelmach I. Vitamin D supplementation in children may prevent asthma exacerbation triggered by acute respiratory infection. *J Allergy Clin Immunol* 2011; **127**(5): 1294-6.
13. Kumar GT, Sachdev HS, Chellani H, et al. Effect of weekly vitamin D supplements on mortality, morbidity, and growth of low birthweight term infants in India up to age 6 months: randomised controlled trial. *BMJ* 2011; **342**: d2975.
14. Lehouck A, Mathieu C, Carremans C, et al. High doses of vitamin D to reduce exacerbations in chronic obstructive pulmonary disease: a randomized trial. *Ann Intern Med* 2012; **156**(2): 105-14.
15. Manaseki-Holland S, Maroof Z, Bruce J, et al. Effect on the incidence of pneumonia of vitamin D supplementation by quarterly bolus dose to infants in Kabul: a randomised controlled superiority trial. *Lancet* 2012; **379**(9824): 1419-27.

16. Camargo CA, Jr., Ganmaa D, Frazier AL, et al. Randomized trial of vitamin D supplementation and risk of acute respiratory infection in Mongolia. *Pediatrics* 2012; **130**(3): e561-7.
17. Murdoch DR, Slow S, Chambers ST, et al. Effect of vitamin D3 supplementation on upper respiratory tract infections in healthy adults: the VIDARIS randomized controlled trial. *JAMA* 2012; **308**(13): 1333-9.
18. Bergman P, Norlin AC, Hansen S, et al. Vitamin D3 supplementation in patients with frequent respiratory tract infections: a randomised and double-blind intervention study. *BMJ Open* 2012; **2**(6): e001663.
19. Marchisio P, Consonni D, Baggi E, et al. Vitamin D supplementation reduces the risk of acute otitis media in otitis-prone children. *Pediatr Infect Dis J* 2013; **32**(10): 1055-60.
20. Rees JR, Hendricks K, Barry EL, et al. Vitamin D3 Supplementation and Upper Respiratory Tract Infections in a Randomized, Controlled Trial. *Clin Infect Dis* 2013.
21. Tran B, Armstrong BK, Ebeling PR, et al. Effect of vitamin D supplementation on antibiotic use: a randomized controlled trial. *Am J Clin Nutr* 2014; **99**(1): 156-61.
22. Goodall EC, Granados AC, Luinstra K, et al. Vitamin D3 and gargling for the prevention of upper respiratory tract infections: a randomized controlled trial. *BMC Infectious Diseases* 2014; **14**: 273.
23. Urashima M, Mezawa H, Noya M, Camargo CA, Jr. Effects of vitamin D supplements on influenza A illness during the 2009 H1N1 pandemic: a randomized controlled trial. *Food & Function* 2014; **5**(9): 2365-70.
24. Grant CC, Kaur S, Waymouth E, et al. Reduced primary care respiratory infection visits following pregnancy and infancy vitamin D supplementation: a randomised controlled trial. *Acta Paediatr* 2014.
25. Martineau AR, James WY, Hooper RL, et al. Vitamin D3 supplementation in patients with chronic obstructive pulmonary disease (ViDiCO): a multicentre, double-blind, randomised controlled trial. *The Lancet Respiratory Medicine* 2015; **3**(2): 120-30.
26. Martineau AR, MacLaughlin BD, Hooper RL, et al. Double-blind randomised placebo-controlled trial of bolus-dose vitamin D3 supplementation in adults with asthma (ViDiAs). *Thorax* 2015; **70**(5): 451-7.
27. Martineau AR, Hanifa Y, Witt KD, et al. Double-blind randomised controlled trial of vitamin D3 supplementation for the prevention of acute respiratory infection in older adults and their carers (ViDiFlu). *Thorax* 2015.
28. Simpson SJ, van der Mei I, Stewart N, Blizzard L, Tettey P, Taylor B. Weekly cholecalciferol supplementation results in significant reductions in infection risk among the vitamin D deficient: results from the CIPRIS pilot RCT. *BMC Nutrition* 2015; **1**(7).
29. Dubnov-Raz G, Rinat B, Hemila H, Choleva L, Cohen AH, Constantini NW. Vitamin D supplementation and upper respiratory tract infections in adolescent swimmers: a randomized controlled trial. *Pediatric Exercise Science* 2015; **27**(1): 113-9.
30. Denlinger LC, King TS, Cardet JC, et al. Vitamin D Supplementation and the Risk of Colds in Patients with Asthma. *Am J Respir Crit Care Med* 2016; **193**(6): 634-41.
31. Tachimoto H, Mezawa H, Segawa T, Akiyama N, Ida H, Urashima M. Improved Control of Childhood Asthma with Low-Dose, Short-Term Vitamin D Supplementation: A Randomized, Double-Blind, Placebo-Controlled Trial. *Allergy* 2016.
32. Ginde AA, Blatchford P, Breese K, et al. High-Dose Monthly Vitamin D for Prevention of Acute Respiratory Infection in Older Long-Term Care Residents: A Randomized Clinical Trial. *J Am Geriatr Soc* 2017; **65**(3): 496-503.
33. Gupta P, Dewan P, Shah D, et al. Vitamin D Supplementation for Treatment and Prevention of Pneumonia in Under-five Children: A Randomized Double-blind Placebo Controlled Trial. *Indian Pediatr* 2016; **53**(11): 967-76.
34. Arihiro S, Nakashima A, Matsuoka M, et al. Randomized Trial of Vitamin D Supplementation to Prevent Seasonal Influenza and Upper Respiratory Infection in Patients With Inflammatory Bowel Disease. *Inflamm Bowel Dis* 2019; **25**(6): 1088-95.

35. Hibbs AM, Ross K, Kerns LA, et al. Effect of Vitamin D Supplementation on Recurrent Wheezing in Black Infants Who Were Born Preterm: The D-Wheeze Randomized Clinical Trial. *JAMA* 2018; **319**(20): 2086-94.
36. Lee MT, Kattan M, Fennoy I, et al. Randomized phase 2 trial of monthly vitamin D to prevent respiratory complications in children with sickle cell disease. *Blood Adv* 2018; **2**(9): 969-78.
37. Loeb M, Dang AD, Thiem VD, et al. Effect of Vitamin D supplementation to reduce respiratory infections in children and adolescents in Vietnam: A randomized controlled trial. *Influenza Other Respir Viruses* 2019; **13**(2): 176-83.
38. Rosendahl J, Valkama S, Holmlund-Suila E, et al. Effect of Higher vs Standard Dosage of Vitamin D3 Supplementation on Bone Strength and Infection in Healthy Infants: A Randomized Clinical Trial. *JAMA Pediatrics* 2018; **172**(7): 646-54.
39. Shimizu Y, Ito Y, Yui K, Egawa K, Orimo H. Intake of 25-Hydroxyvitamin D3 Reduces Duration and Severity of Upper Respiratory Tract Infection: A Randomized, Double-Blind, Placebo-Controlled, Parallel Group Comparison Study. *J Nutr Health Aging* 2018; **22**(4): 491-500.
40. Aloia JF, Islam S, Mikhail M. Vitamin D and Acute Respiratory Infections-The PODA Trial. *Open Forum Infectious Diseases* 2019; **6**(9): ofz228.
41. Hauger H, Ritz C, Mortensen C, et al. Winter cholecalciferol supplementation at 55°N has little effect on markers of innate immune defense in healthy children aged 4–8 years: a secondary analysis from a randomized controlled trial. *European Journal of Nutrition* 2019; **58**(4): 1453-62.
42. Camargo CA, Sluyter J, Stewart AW, et al. Effect of monthly high-dose vitamin D supplementation on acute respiratory infections in older adults: A randomized controlled trial. *Clin Infect Dis* 2020; **71**: 311-7.
43. Ganmaa D, Uyanga B, Zhou X, et al. Vitamin D Supplements and Prevention of Tuberculosis Infection and Disease. *N Engl J Med* 2020; **383**: 359-68.
44. Mandlik R, Mughal Z, Khadilkar A, et al. Occurrence of infections in schoolchildren subsequent to supplementation with vitamin D-calcium or zinc: a randomized, double-blind, placebo-controlled trial. *Nutr Res Pract* 2020; **14**(2): 117-26.
45. Pham H, Waterhouse M, Baxter C, et al. The effect of vitamin D supplementation on acute respiratory tract infection in older Australian adults: an analysis of data from the D-Health Trial: a randomised controlled trial. *The Lancet Diabetes & Endocrinology* 2020; **in press**.
46. Rake C, Gilham C, Bukasa L, et al. High-dose oral vitamin D supplementation and mortality in people aged 65-84 years: the VIDAL cluster feasibility RCT of open versus double-blind individual randomisation. *Health Technol Assess* 2020; **24**(10): 1-54.
47. Golan-Tripto I. The Effect of Vitamin D Administration to Premature Infants on Vitamin D Status and Respiratory Morbidity (NCT02404623). <https://clinicaltrials.gov/ct2/show/NCT02404623>.
48. Reyes ML, Vizcaya C, Borzutzky A, et al. Study of Vitamin D for the Prevention of Acute Respiratory Infections in Children (NCT02046577). <https://clinicaltrials.gov/ct2/show/NCT02046577>.