



ELSEVIER

Contents lists available at ScienceDirect

## Data in Brief

journal homepage: [www.elsevier.com/locate/dib](http://www.elsevier.com/locate/dib)



### Data Article

# Data from an integrative approach decipher the surface proteome of *Propionibacterium freudenreichii*



Caroline Le Maréchal<sup>a,b</sup>, Vincent Peton<sup>a,b</sup>, Coline Plé<sup>c</sup>,  
Christophe Vroland<sup>a,b</sup>, Julien Jardin<sup>a,b</sup>, Valérie Briard-Bion<sup>a,b</sup>,  
Gaël Durant<sup>a,b</sup>, Victoria Chuat<sup>a,b,d</sup>, Valentin Loux<sup>e</sup>,  
Benoit Foligné<sup>c</sup>, Stéphanie-Marie Deutsch<sup>a,b</sup>,  
Hélène Falentin<sup>a,b,\*</sup>, Gwénaél Jan<sup>a,b,\*</sup>

<sup>a</sup> INRA, UMR1253 STLO, Science et Technologie du Lait et de l'Œuf, F-35042 Rennes, France

<sup>b</sup> GROCAMBUS OUEST, UMR1253 STLO, F-35042 Rennes, France

<sup>c</sup> Lactic acid Bacteria & Mucosal Immunity, Center for Infection and Immunity of Lille, Institut Pasteur de Lille, U 1019, UMR 8204 Université Lille Nord de France, 1 rue du Pr Calmette, BP 245, F-59019 Lille, France

<sup>d</sup> INRA, UMR1253 STLO, CIRM-BIA, F-35042 Rennes, France

<sup>e</sup> INRA, UR MIG, F-78352 Jouy-en-Josas, France

### ARTICLE INFO

#### Article history:

Received 26 August 2014

Accepted 26 August 2014

Available online 22 September 2014

### ABSTRACT

The surface proteins of the probiotic *Propionibacterium freudenreichii* were inventoried by an integrative approach that combines *in silico* protein localization prediction, surface protein extraction, shaving and fluorescent CyDye labeling. Proteins that were extracted and/or shaved and/or labeled were identified by nano-LC-MS/MS following trypsinolysis. This method's combination allowed to confirm detection of true surface proteins involved in host/probiotic interactions. The data, supplied in this article, are related to the research article entitled "Surface proteins of *P. freudenreichii* are involved in its anti-inflammatory properties" (Le Maréchal et al., 2014 [6]).

© 2014 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>).

DOI of original article: <http://dx.doi.org/10.1016/j.jprot.2014.07.018>

\* Corresponding authors.

E-mail address: [gwenael.jan@rennes.inra.fr](mailto:gwenael.jan@rennes.inra.fr) (G. Jan).

<http://dx.doi.org/10.1016/j.dib.2014.08.009>

2352-3409/© 2014 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>).

## Specifications Table

Subject area	<i>Microbiology, Food Technology</i>
More specific subject area	<i>Probiotics, fermented foods</i>
Type of data	<i>Tables and figures</i>
How data was acquired	<i>2-D PAGE: Multiphor II and Ettan DALTtwelve (GE Healthcare) Image analysis: Image-Master 2D Mass spectrometry: QSTAR XL (Applied Biosystem)</i>
Data format	<i>Analyzed</i>
Experimental factors	<i>Dairy propionibacteria were grown in milk aqueous phase and directly subjected to guanidine extraction, trypsin shaving or CyDye labeling (see Fig. 1)</i>
Experimental features	<i>All the samples were analyzed by nano-LC coupled to MS/MS</i>
Data source location	<i>City, Country and/or Latitude &amp; Longitude (&amp; GPS coordinates) for collected samples/data if applicable</i>
Data accessibility	<i>Within this article</i>

## Value of the data

- Probiotic bacteria are analyzed after growth in milk aqueous phase, not in laboratory medium as it is usually done.
- A combination of 3 proteomic methods is used to validate surface localization of the identified proteins.
- The data point out candidate genes for functional genomics.
- The data open new perspectives to identify molecular mechanisms of probiotic/host interactions.

## 1. Data, experimental design, materials and methods [6]

### 1.1. Data

Fig. 1 summarizes the 3 proteomic methods used to acquire data on the surface proteins of the probiotic *Propionibacterium freudenreichii*. Data acquired by guanidine hydrochloride extraction are presented in [Supplementary Table 1](#). Data acquired by enzymatic shaving using trypsin are presented in [Supplementary Table 2](#). Data acquired by in situ fluorescent labeling using CyDye DIGE Fluor minimal Dye are presented in [Supplementary Table 3](#).

### 1.2. Genome sequencing, annotation and bio-informatics


The genome of *P. freudenreichii* ITG P20 was previously sequenced and annotated [4] and the draft assembly deposited in the European Nucleotide Archive (EMBL-EBI accession number: CONTIGS: CCBE010000001–CCBE010000111; SCAFFOLDS:HG975453–HG975511). Prediction of subcellular localization of encoded proteins were done in this work using SurfG+ [2].

### 1.3. Whole-cell protein extracts

Whole cell SDS extracts were prepared by disruption of cells in SDS according to a procedure modified from one previously described [7].

### 1.4. Extraction of surface proteins non-covalently bound to the cell wall using guanidine hydrochloride

Surface layer proteins were extracted according to a procedure modified from one previously described [8]. 100 mL of stationary phase culture (see above) were harvested by centrifugation (6000g, 10 min, 4 °C) and washed in an equal volume of PBS prior to resuspension in 5 M guanidine hydrochloride to a final OD<sub>650</sub> of 20. The suspension was incubated 15 min at 50 °C prior to



**Fig. 1.** Experimental flowchart of the surface protein analysis procedure.

centrifugation (21,000g, 20 min, 30 °C) to eliminate cells. The supernatant was then dialyzed exhaustively against 0.1% SDS in distilled water during 24 h at 4 °C using 10,000 kDa cutoff Slide-A-Lyer<sup>®</sup> Dialysis Cassette (ThermoScientific, Rockford, USA) prior to proteomic investigations.

### 1.5. Enzymatic shaving of surface proteins

100 mL of stationary phase culture (see above) were harvested by centrifugation (6000g, 10 min, 4 °C) and washed in an equal volume of PBS [pH 8.5] containing 5 mM DTT prior to resuspension in 1/10 volume of the same buffer. Sequencing grade modified trypsin (V5111, Promega, Madison, USA) was

dissolved in the same buffer (qsp 0.2 g/L) and added to the bacterial suspension. “Shaving” was performed for 1 h at 37 °C in a 0.5 mL reaction volume containing  $5 \times 10^9$  bacteria and 4 µg of trypsin, with gentle agitation (180 rpm). Bacteria were removed by centrifugation (8000g, 10 min, 20 °C) and the supernatant filtered (0.2 µm, Nalgene) prior to addition of 1 µg of trypsin to complete digestion of released peptides (16 h, 37 °C). Trypsin digestion of the supernatant was stopped by adding trifluoroacetic acid to a final concentration of 0.15% (v/v). The supernatants containing peptides were then concentrated in a Speed-Vac concentrator prior to nano-LC–MS/MS analysis.

### 1.6. Nano-LC–MS/MS analyses

For trypsinolyzed proteins, nano-LC experiments were performed using an on-line liquid chromatography tandem mass spectrometry (MS/MS) setup using a Dionex U3000-RSLC nano-LC system fitted to a QSTAR XL (MDS SCIEX, Ontario, Canada) equipped with a nano-electrospray ion source (ESI) (Proxeon Biosystems A/S, Odense, Denmark). Samples were first concentrated on a PepMap 100 reverse-phase column (C18, 5 µm particle size, 300-µm inner diameter (i.d.) by 5 mm length) (Dionex, Amsterdam, The Netherlands). Peptides were separated on a reverse-phase PepMap 100 column (C18, 3 µm particle size, 75 µm i.d. by 150 mm length) (Dionex) at 35 °C, using solvent A (2% (vol/vol) acetonitrile, 0.08% (vol/vol) formic acid, and 0.01% (vol/vol) TFA in deionized water) and solvent B (95% (vol/vol) acetonitrile, 0.08% (vol/vol) formic acid, and 0.01% (vol/vol) TFA in deionized water). A linear gradient from 10% to 50% of solvent B in 40 min was applied for the elution at a flow rate of 0.3 µL/min. Eluted peptides were directly electrosprayed into the mass spectrometer operated in positive mode. A full continuous MS scan was carried out followed by three data-dependent MS/MS scans. Spectra were collected in the selected mass range 400–2000  $m/z$  for MS and 60–2000  $m/z$  for MS/MS spectra. The three most intense ions from the MS scan were selected individually for collision-induced dissociation (1+ to 4+ charged ions were considered for the MS/MS analysis). The mass spectrometer was operated in data-dependent mode automatically switching between MS and MS/MS acquisition using Analyst QS 1.1 software. The instrument was calibrated by multipoint calibration using fragment ions that resulted from the collision-induced decomposition of a peptide from  $\beta$ -casein,  $\beta$ -CN (193–209). The proteins present in the samples were identified from MS and MS/MS data by using MASCOT v.2.2 software for search into two concatenated databases: (i) a homemade database containing all the predicted proteins of the *P. freudenreichii* strain CIRM-BIA 129 used in this study and (ii) a portion of the UniProtKB database corresponding to *P. freudenreichii*. Search parameters were set as follows. A trypsin enzyme cleavage was used, the peptide mass tolerance was set to 0.2 Da for both MS and MS/MS spectra, and two variable modifications (oxidation of methionine and deamidation of asparagine and glutamine residues) were selected. For each protein identified in NanoLC-ESI-MS/MS, a minimum of two peptides with MASCOT score corresponding to a *P* value below 0.05 were necessary for validation with a high degree of confidence. For automatic validation of the peptides from MASCOT search results, the 1.19.2 version of the IRMa software was used [3].

### 1.7. In situ surface labeling

The surface labeling procedure was adapted from Hagner-McWhirter et al. [5]. Bacteria were grown and harvested as described above and washed in an equal volume of ice-cold PBS containing 33 mM Tris–HCl [pH 8.5] prior to centrifugation. Bacteria were resuspended in 1/10 volume of ice-cold labeling buffer (PBS containing, 1 M urea, 33 mM Tris–HCl [pH 8.5]). Labeling was performed on ice, in the dark, in a 1 mL reaction volume containing  $10^{10}$  live and intact bacteria, and 200 pmol of CyDye DIGE Fluor Cy5 minimal dye (GE Healthcare, Orsay, France). Labeling was stopped by adding 1 µmol of Lysine to quench the dye. Labeled bacteria were centrifuged and washed in PBS (pH 7.4), centrifuged and resuspended in SDS lysis buffer (50 mM Tris–HCl [pH 7.5], 0.3% SDS, 200 mM DTT) prior to whole cell protein extraction as described above.

### 1.8. Two-dimensional imaging and spot picking

Whole-cell protein SDS extracts of labeled bacteria were precipitated using the 2D Clean-Up Kit (GE Healthcare) prior to dissolution in destreak rehydration solution (100 µl per sample) added with 2% (w/v) ampholyte containing buffer (IPG-Buffer 4–7, GE Healthcare). Isoelectric focusing was carried out using pH 4–7, 18 cm, Immobiline Dry Strips on a Multiphor II electrophoresis system (GE Healthcare) for a total of 60 kV h using a standard procedure described previously [1]. The second dimensional separation was performed on the Ettan™ DALTtwelve electrophoresis system (GE Healthcare) using 14% acrylamide separating gels without a stacking gel at a voltage of 50 V for 1 h and 180 V for about 7 h. Fluorescent images of the gels were immediately acquired on a Typhoon PhosphorImager (GE Healthcare) using the appropriate laser excitation for Cy5 fluorescence. Gels were then fixed and Coomassie Blue-stained as described above. Visible images were acquired on an ImageScanner III (GE Healthcare). Images were further analyzed using Image- Master 2D software. Fluorescent profiles of 2-DE-separated proteins were reproducible in at least three individual experiments. Fluorescent and Coomassie-Blue visible images of the 2D electrophoresis gels were matched to detect surface-exposed proteins. Fluorescent spots corresponding to surface-exposed proteins were excised from 2-DE gels as previously described [7] when detectable by Coomassie-Blue staining. Proteins were identified by tandem mass spectrometry (MS/MS) after an in-gel trypsin digestion adapted from Shevchenko [9]. Briefly, gel pieces were washed with acetonitrile and ammonium bicarbonate solution, and then dried under vacuum in a SpeedVac concentrator (SVC100H-200; Savant, Thermo Fisher Scientific, Waltham, MA, USA). In-gel trypsin digestion was performed overnight at 37 °C and stopped with spectrophotometric-grade trifluoroacetic acid (TFA) (Sigma-Aldrich). The supernatants containing peptides were then vacuum dried in a Speed-Vac concentrator and stored at –20 °C until mass spectrometry analysis. Nano-LC–MS/MS analysis was as described above.

## Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.dib.2014.08.009>.

## References

- [1] R. Anastasiou, P. Leverrier, I. Krestas, A. Rouault, G. Kalantzopoulos, P. Boyaval, E. Tsakalidou, G. Jan, Changes in protein synthesis during thermal adaptation of *Propionibacterium freudenreichii* subsp. *shermanii*, *Int. J. Food Microbiol.* 108 (2006) 301–314.
- [2] A. Barinov, V. Loux, A. Hammani, P. Nicolas, P. Langella, D. Ehrlich, E. Maguin, M. van de Guchte, Prediction of surface exposed proteins in *Streptococcus pyogenes*, with a potential application to other Gram-positive bacteria, *Proteomics* 9 (2009) 61–73.
- [3] V. Dupierris, C. Masselon, M. Court, S. Kieffer-Jaquinod, C. Bruley, A toolbox for validation of mass spectrometry peptides identification and generation of database: IRMa, *Bioinformatics* 25 (2009) 1980–1981.
- [4] Falentin, H., S.M. Deutsch, V. Loux, A. Hammani, J. Buratti, S. Parayre, V. Chuat, V. Barbe, J.M. Aury, G. Jan, and Y. Le Loir. 2014. Permanent draft genome of the probiotic strain *Propionibacterium freudenreichii* CIRM-BIA 129 (ITG P20). *Stand. Genomic Sci.*, submitted for publication.
- [5] A. Hagner-McWhirter, M. Winkvist, S. Bourin, R. Marouga, Selective labelling of cell-surface proteins using CyDye DIGE Fluor minimal dyes, *J. Vis. Exp.* (2008).
- [6] M.C. Le, V. Peton, C. Ple, C. Vroland, J. Jardin, V. Briard-Bion, G. Durant, V. Chuat, V. Loux, B. Foligne, S.M. Deutsch, H. Falentin, G. Jan, Surface proteins of *Propionibacterium freudenreichii* are involved in its anti-inflammatory properties, *J. Proteomics* (2014).
- [7] P. Leverrier, J.P. Vissers, A. Rouault, P. Boyaval, G. Jan, Mass spectrometry proteomic analysis of stress adaptation reveals both common and distinct response pathways in *Propionibacterium freudenreichii*, *Arch. Microbiol.* 181 (2004) 215–230.
- [8] S. Lortal, A. Rouault, B. Cesselin, U.B. Sleytr, Paracrystalline surface layers of dairy propionibacteria, *Appl. Environ. Microbiol.* 59 (1993) 2369–2374.
- [9] A. Shevchenko, S. Sunyaev, A. Liska, P. Bork, A. Shevchenko, Nano-electrospray tandem mass spectrometry and sequence similarity searching for identification of proteins from organisms with unknown genomes, *Methods Mol. Biol.* 211 (2003) 221–234.